

# Genome-Wide Association Analysis Reveals Genetic Variations and Candidate Genes Associated with Salt Tolerance Related Traits in *Gossypium Hirsutum*

**Peng Xu**

Jiangsu Academy of Agricultural Sciences

**Qi Guo**

Jiangsu Academy of Agricultural Sciences

**Shan Meng**

Jiangsu Academy of Agricultural Sciences

**Xianggui Zhang**

Jiangsu Academy of Agricultural Sciences

**Zhenzhen Xu**

Jiangsu Academy of Agricultural Sciences

**Wangzhen Guo**

Nanjing Agricultural University

**Xinlian Shen** (✉ [xlshen68@126.com](mailto:xlshen68@126.com))

Chinese Academy of Agricultural Sciences <https://orcid.org/0000-0002-0111-9850>

---

## Research article

**Keywords:** *Gossypium hirsutum*, Genome-wide association mapping, Genotyping By-Sequencing, salt tolerance, Virus-induced gene silencing assay

**Posted Date:** September 8th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-66236/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Genomics on January 6th, 2021. See the published version at <https://doi.org/10.1186/s12864-020-07321-3>.

# Abstract

**Background:** As a pioneer industrial crop in saline-alkali lands, cotton is more resistant to salt and drought stresses. However, abiotic stresses still have significant negative effects on its growth and development. It is therefore important to breed salt tolerance varieties which can help accelerate the improvement of cotton production. The development of molecular markers linked to causal genes has provided an effective and efficient approach for improving salt tolerance.

**Results:** In this study, a genome-wide association study of salt tolerance related traits at seedling stage was performed based on two years of phenotype identification for 217 representative upland cotton cultivars by Genotyping By-Sequencing platform. A total of 51,060 polymorphic SNPs unevenly distributed in 26 chromosomes were screened across the cotton cultivars, and 25 associations with 27 SNPs scattered over 12 chromosomes were detected significantly ( $-\log_{10}P > 4$ ) associated with three salt tolerance related traits in 2016 and 2017. Among these, the associations on chromosome A13 and D08 for RPH, A07 for RSFW, A08 and A13 for RSDW were expressed in both two environments, indicating that they were likely to be stable QTLs. A total of 12 salt-induced candidate genes were identified differentially expressed by the combination of GWAS and transcriptome analysis. Three promising genes were selected to conduct preliminary function verification of salt tolerance. The increase of GH\_A13G0171-silenced plants in salt related traits under salt stress indicated its negative function in regulating the salt stress response.

**Conclusions:** These results provided important genetic variations and candidate genes for accelerate the improvement of salt tolerance in cotton.

## Background

The competition for arable land between food crops and cotton (*Gossypium* spp.) has existed for a long time in China. However, as population continues to grow, the emphasis on food security has inadvertently moved cotton production to more marginal soils in saline-alkali areas and coastal beaches. Cotton as the largest source of textiles fiber in the world is more resistant to abiotic stresses such as high salt and drought stresses than other crop. However, excessive salt in the soil can severely affects its growth and development of cotton plants [1], with injury resulting in a reduction in fiber yield by as much as 60% [2]. Therefore, breeding for cotton varieties with improved salt tolerance could alleviate the conflict between food crops and cotton by reclaiming and utilizing saline-alkali coastal lands for production. Similarly, in the northwestern inland cotton production region of China, the availability of salt tolerant varieties would expand the area of cotton production by promoting the synchronous growth of food crops and cotton.

The genetic architecture of salt tolerance is one of the most important subjects in plant science. Salt tolerance is a complex quantitative trait, which is controlled by multiple genes and involves a variety of physiological and biochemical metabolic pathways in cotton. In addition, the expression of each gene is sensitive to external environment. In the current study, the main conventional breeding approaches for

developing salt tolerant varieties are screening and collecting salt-tolerant germplasm resources, then transferring elite loci by hybridization, composite hybridization and backcrossing methods. The progress of conventional breeding of salt tolerance in upland cotton is impeded due to the lack of high salt-tolerant resources [3], low heritability [4], genetic complexity, and the difficulties in phenotyping [5].

The development of molecular markers linked to causal genes for a trait has provided an effective and efficient approach for improving quantitative traits. Once identified, markers linked to a quantitative trait locus (QTL) such as for salt tolerance can then serve as a selection tool for rapid and efficient marker-assisted selection (MAS). QTL mapping via bi-parental populations is an important method for quantitative trait research, and has been widely employed to map a number of traits including salt tolerance in various crops [6]. In addition to QTL mapping, association mapping based on linkage disequilibrium (LD) is another approach for detecting molecular markers tightly linked with quantitative traits in a natural germplasm population. Association analysis is time- and cost-effective, and can mine for genetic variations existing in the natural population, and more importantly, it takes advantage of the recombination information that has accumulated in the natural population over the long-term evolution process, thus achieving a higher mapping resolution, possibly identifying the causative genes [7, 8]. Because of its effectiveness in QTL identification, association mapping has been widely used in crop species and plays an important role in molecular breeding [9–14].

Genotyping By-Sequencing (GBS) is a reduced representation genotyping platform, and has a broad application in crop genetics and breeding [15–18]. In general, the GBS approach begins with an enzymatic digestion to reduce genome complexity using barcoding restriction enzymes, then performs multiple sequencing of barcoding DNA fragments on the high-throughput Next-Generation Sequencing platform. A bioinformatics analysis of indexed sequence reads is followed to identify genetic variants. Finally, genetic diversity analysis is carried out based on a sample-by-variant matrix [19]. GBS is a powerful and cost-effective tool to assess variations across populations.

Since the first molecular marker-based genetic map of cotton published in 1994 [20], cotton scientists have identified a large number of important agronomic traits QTLs, including fiber quality traits, yield traits, and disease resistance. However, only limited reports have been published on salt tolerance in cotton [21–25]. Herein, we reported the genome-wide association mapping of salt tolerant QTLs during the seedling stage performed over two years of phenotyping on 217 representative upland cotton cultivars. These results provided important genetic variations and candidate genes for accelerating the improvement of salt tolerance in cotton.

## Results

### Phenotypic diversity analysis

In order to evaluate the phenotypic variation of salt tolerance related traits in the association mapping population, we analyzed 3 traits related to salt tolerance in two years. To determine the salt tolerance, the

217 upland cotton cultivars (Additional file 1: Table S1) were treated with 350 mM NaCl and clean water in 2016 and 2017, respectively. Three traits related to salt tolerance were determined, including relative plant height (RPH), relative shoot fresh matter weight (RSFW) and relative shoot dry matter weight (RSDW). The average trait values in each year were used for phenotypic analysis and association mapping in single environment. The mean values, ranges, standard deviations, and coefficients of variation for these salt tolerance related traits in each year are shown in Table 1. Great differences among the variation coefficient (CV) in both years were found for the three traits. Overall, the upland cotton cultivars in this association mapping panel clearly exhibited considerable natural variation in the three traits to salt tolerance and displayed very high genetic diversity.

Table 1  
Statistics of various traits related to salt tolerance

Year	Trait	Range	Mean	SD	CV%
2016	RPH	0.189–0.760	0.440	0.145	0.330
	RSFW	0.136–0.762	0.411	0.150	0.365
	RSDW	0.137–0.756	0.497	0.143	0.288
2017	RPH	0.169–0.780	0.445	0.143	0.321
	RSFW	0.129–0.794	0.420	0.167	0.398
	RSDW	0.172–0.776	0.521	0.158	0.303

RPH: relative plant height; RSFW: relative shoot fresh matter weight; RSDW: relative shoot dry matter weight; SD: Standard Deviation; CV: Coefficient of Variation.

## Genetic Diversity Analysis Based On Snp

GBS produced 53,321 high-quality polymorphic SNPs among the 217 upland cotton cultivars, with 41,733 SNPs located in the intergenic intervals and 6,188 SNPs in the coding regions. Of these, 95.8% of the loci (51,060 SNPs) were mapped onto 26 chromosomes of the cotton genome, and were selected for the GWAS analysis (Additional file 2: Table S2). These SNPs were not evenly distributed (Additional file 3: Fig.S1) with an average of 1964 SNPs per chromosome ranging from 863 to 5209. Chromosome A08 had the most SNPs (5209) with the highest SNP density (199 Kb/SNP). D04 had the least SNPs (863), but A02 had the lowest SNP density with 680 Kb/SNP. The genetic diversity of this population varied from 0.224 (A08) to 0.389 (A13). The polymorphism information content (PIC) varied from 0.192 (A08) to 0.305 (A13) (Table 2). The above results indicated a relatively large span in genetic diversity index, and polymorphism information content in the cotton genome.

Table 2  
Summary of polymorphic SNPs mapped in 26 chromosomes of *Gossypium hirsutum*

Chr.	Chr length (Mb)	No. of SNPs	SNP density (Kb/SNP)	Gene diversity	PIC	LD Decay (Kb) $r^2 = 0.1$	$r^2 = 0.2$
A01	99.9	2111	473	0.294	0.235	1145	990
A02	83.5	1228	680	0.344	0.275	1215	903
A03	100.3	1490	673	0.321	0.26	1385	1035
A04	62.9	1138	553	0.314	0.255	1104	974
A05	92.1	2202	418	0.346	0.277	1354	1060
A06	103.2	3924	263	0.232	0.199	1377	1157
A07	78.3	1897	413	0.319	0.259	1258	959
A08	103.7	5209	199	0.224	0.192	1136	891
A09	75	1639	458	0.312	0.252	1299	1028
A10	100.9	2335	432	0.308	0.254	1191	957
A11	93.3	2027	460	0.257	0.215	1194	970
A12	87.5	1498	584	0.306	0.247	1183	915
A13	80	2996	267	0.389	0.305	1358	1058
D01	61.5	2385	258	0.318	0.261	880	435
D02	67.3	1914	352	0.321	0.26	712	599
D03	46.7	934	500	0.27	0.225	1046	803
D04	51.5	863	597	0.331	0.266	953	844
D05	61.9	1206	513	0.308	0.25	1204	816
D06	64.3	2454	262	0.281	0.235	1275	917
D07	55.3	2293	241	0.324	0.264	1124	817
D08	61.9	2463	251	0.306	0.248	909	523
D09	51	1643	310	0.286	0.239	1002	732
D10	63.4	1517	418	0.294	0.241	733	663
D11	66.1	1080	612	0.317	0.257	846	801
D12	59.1	1407	420	0.276	0.228	913	754

Chr.: Chromosome; PIC: Polymorphism information content.

Chr.	Chr length (Mb)	No. of SNPs	SNP density (Kb/SNP)	Gene diversity	PIC	LD Decay (Kb) $r^2 = 0.1$	$r^2 = 0.2$
D13	60.5	1207	501	0.287	0.236	1325	988
Total	1931.1	51060	427	0.303	0.248	1120	869

Chr.: Chromosome; PIC: Polymorphism information content.

## Population Structure And Linkage Disequilibrium Analysis

It is important for genome-wide association mapping to control the effect of population structure, because population stratification could eliminate spurious associations between genotypes and phenotypes [26, 27]. STRUCTURE software was used to calculate the Bayesian clustering from  $K = 1$  to 10 for five repetitions.  $\ln P(D)$  value continued to increase from  $K = 1$  to  $K = 10$  without a significant inflection point (Fig. 1a). However, there was an obvious spike at the value of  $\Delta K = 3$  (Fig. 1b), suggesting that the population could be divided into 3 subgroups (Fig. 1c). Taking the corresponding Q matrix at  $k = 3$  as the covariate could reasonably eliminate spurious association effects and improve the association mapping accuracy.

The LD distribution among chromosomes of the 217 upland cotton cultivars was shown in Table 2. The LD decay distance was 869 Kb and 1120 Kb when the  $r^2$  dropped to 0.2 and 0.1, respectively. The LD decay distance was not evenly distributed among chromosomes, ranged from 435 Kb (D01) to 1157 Kb (A06) when  $r^2$  was set at 0.2 and ranged from 712 Kb (D02) to 1377 Kb (A06) when  $r^2$  was set at 0.1, the overall LD decay in the At subgenome was significantly higher than that in the Dt subgenome.

## Association Analysis

In order to explore the genetic factors underlying salt tolerance, the mixed linear models (MLMs) was performed by simultaneously accounting for population structure and relative kinship matrix to conduct a GWAS. A total of 25 significant associations ( $-\log_{10} P > 4$ ) with 27 significant SNPs located on chromosomes A05, A07, A08, A09, A10, A11, A12, A13, D02, D03, D06, and D09 were detected for the three salt tolerance related traits in the 2016 and 2017 dataset. Eleven associations with 12 SNPs were detected in 2016 (Fig. 2) and nine associations with 9 SNPs were detected in 2017 (Fig. 3). In addition, five associations with 6 SNPs were detected in both 2016 and 2017 dataset. The phenotypic variance explained (PVE) by individual QTL ranged from 1.29–7.00% (Table 3).

Table 3  
Summary of SNPs significantly associated with salt tolerance related traits

Traits	Chr.	Site	Allele	MAF	-Log10(P)	PVE	Environment
RPH	A10	84786908	C/T	0.15(T)	4.27	5.67	2016
	A10	84851927	G/A	0.20(A)	4.10	4.31	2016
	A13	1869056	C/G	0.33(C)	4.22, 4.36	4.41,4.93	2016, 2017
	A07	90682411	C/T	0.12(T)	4.35	4.58	2017
	A09	62356818	A/G	0.19(A)	4.13	7.00	2017
	A12	74098493	G/A	0.49(A)	4.58	5.51	2017
	D02	38449756	T/A	0.47(T)	4.01	1.50	2017
	D03	18961034	T/G	0.11(G)	4.18	2.54	2017
	D08	49014753	C/T	0.13(T)	4.34, 4.16	4.35,3.83	2016, 2017
	D08	49080865	G/A	0.13(A)	4.34, 4.16	4.35,3.83	2016, 2017
RSFW	A08	5515194	C/T	0.10(T)	4.79	4.23	2016
	A10	84786908	C/T	0.15(T)	4.19	2.95	2016
	D06	19966743	C/G	0.5(G)	4.09	3.56	2016
	A07	90532061	C/G	0.13(G)	4.73	5.23	2017
	A07	90682411	C/T	0.12(T)	4.13, 4.73	2.40, 5.25	2016, 2017
	A08	41804418	T/A	0.09(T)	4.23	2.74	2017
	A08	53916411	G/A	0.088(G)	4.15	2.35	2017
	A11	117938337	G/A	0.32(A)	4.09	3.08	2017
RSDW	A05	94406259	C/T	0.33(T)	4.17	2.92	2016
	A08	5589045	G/A	0.19(A)	4.67	4.85	2016
	A08	6254890	T/G	0.22(G)	4.36,4.66	3.52,4.33	2016,2017
	A13	42992196	A/T	0.48(T)	4.27,4.28	2.79,3.03	2016,2017
	D08	47703450	C/T	0.096(T)	4.35	1.93	2016
	D08	48527413	C/A	0.12(A)	4.23	1.29	2016
	D08	48911757	C/G	0.13(G)	4.44	2.62	2016

Chr.: Chromosome; RPH: relative plant height; RSFW: relative shoot fresh matter weight; RSDW: relative shoot dry matter weight; MAF: Minor Allele Frequency; PVE: phenotypic variance explained.

Traits	Chr.	Site	Allele	MAF	-Log10(P)	PVE	Environment
	D08	49014753	C/T	0.13(T)	5.20	3.88	2016
	D08	49080865	G/A	0.13(A)	5.20	3.88	2016

Chr.: Chromosome; RPH: relative plant height; RSFW: relative shoot fresh matter weight; RSDW: relative shoot dry matter weight; MAF: Minor Allele Frequency; PVE: phenotypic variance explained.

For RPH, nine significant associations with 10 SNPs were identified on chromosomes A07, A09, A10, A12, A13, D02, D03, and D08. The PVE ranged from 1.50–7.00%. Moreover, the association with one SNPs on chromosome A13 and the association with two SNPs on chromosome D08 were detected in both two years.

For RSDW, eight significant associations with 9 SNPs on A05, A08, A13, and D08 were detected. The PVE ranged from 1.29–4.85%. The association with one SNPs on chromosome A08 and the association with one SNP on chromosome A13 were detected in both two years.

For RSFW, eight associations with 8 significant SNPs on A07, A08, A10, A11 and D06 were detected in 2016 and 2017. The PVE ranged from 2.35–5.25%. The association with three SNPs on chromosome A07 was detected in both two years.

Pleiotropy was also found in our association analysis. For example, the significant SNP A07\_90682411 was simultaneously detected associated with RPH in 2017 and RSFW in both two environments. The significant SNP A10\_84786908 was simultaneously detected associated with RPH in 2016 and RSFW in 2016. The significant SNP D08\_49014753 and D08\_49080865 simultaneously detected associated with RPH in 2016, 2017, and RSDW in 2016. In addition, the associations on A08 for RSFW and RSDW in 2016 were only 70 kb apart. Their confidence interval overlapped for each other.

## Identification And Preliminary Function Verification Of Candidate Genes

In general, the LD decay could be used as confidence interval to identify candidate genes. Because the cotton genome has a large LD decay [28, 29], we extracted potential candidate genes within 100 Kb of flanking significant markers on the basis of the published upland cotton genome sequencing database [30]. A total of 156 genes were identified in these intervals (Additional file 4: Table S3). Gene ontology (GO) enrichment analysis indicated that “dioxygenase activity”, “oxidoreductase activity, acting on single donors with incorporation of molecular oxygen, incorporation of two atoms of oxygen” and “oxidoreductase activity, acting on single donors with incorporation of molecular oxygen” were significantly enriched using an FDR adjusted P-value of  $\leq 0.05$  as the cutoff. Of the 156 genes, 12 were differentially expressed between salt tolerance variety Miscott7913-83 and salt sensitive variety Su 12 according to the previous transcriptome sequencing results [31] (Additional file 4: Table S3). Some of

these genes may be associated with salt stress, such as GH\_A08G0488 and GH\_A10G1620 encoding protein kinase. Protein kinase has been proved to play an important role in salt tolerance in cotton [32]. Another gene GH\_A13G0171, which encodes aquaporin, was also likely to regulate the salt stress response [33]. The confidence interval contains GH\_A13G0171 was simultaneously detected in both years. Through a *t* test, we found that the salt tolerance of upland cotton cultivars with the G haplotype was significantly higher than that of cultivars with the C haplotype in both years (Fig. 4).

The three promising genes (GH\_A08G0488, GH\_A10G1620 and GH\_A13G0171) were selected to conduct preliminary function verification of salt tolerance in cotton. Analysis of gene expression patterns could provide important clues for gene function determination. Quantitative RT-PCR was performed to analyze the expression levels of GH\_A08G0488, GH\_A10G1620 and GH\_A13G0171 in roots and leaves under salt stress treatment in salt tolerance variety Miscott7913-83 and salt sensitive variety Su 12. As shown in Fig. 5, the three genes were induced by salt stress and displayed distinct expression patterns in response to salt stress in salt tolerance variety Miscott7913-83. The three genes had a much higher expression level in roots than in leaves. The gene GH\_A13G0171 exhibited a significantly down-regulated expression in both root and leaf tissues after salt stress. The gene GH\_A08G0488 exhibited a significantly up-regulated expression in both root and leaf tissues. The expression level of GH\_A10G1620 showed an increase in leaf and no significant changes in the root tissues. As shown in Fig. 6, the three genes were also displayed distinct expression patterns in response to salt stress in sensitive variety Su 12. The gene GH\_A13G0171 exhibited an identical expression pattern in Miscott7913-83 and Su 12. The expression levels of GH\_A10G1620 and GH\_A08G0488 were not significant different.

To confirm the functional roles of GH\_A08G0488, GH\_A10G1620 and GH\_A13G0171 genes under salt stress, virus-induced gene silencing (VIGS) assay was used to repress expression of these genes in salt tolerance variety Miscott2111 plants. The inoculated seedlings were grown in three light incubators at 23 °C under a 16-h light and 8-h dark cycle as three biological replicates. At the developmental period when two leaves had formed, the pTRV2 :: GH\_A08G0488, pTRV2 :: GH\_A10G1620, pTRV2::GH\_A13G0171 and pTRV2::00 inoculated plants were treated with 350 mM NaCl. After 15 days, the plant height, fresh and dry shoot matter weight were determined and the corresponding relative values were calculated. The transcripts of the three genes in the VIGS leaves were significantly reduced compared to the untreated Miscott2111 plants, indicating that they were effectively silenced in VIGS plants (Additional file 5: Fig.S2). Compared with the control pTRV2::00, no significance effect on their phenotypes was observed in pTRV2 :: GH\_A08G0488 and pTRV2::GH\_A10G1620 inoculated plants, and the plant height, fresh and dry shoot matter weigh of GH\_A13G0171-silenced plants were significantly higher than that of pTRV2::00 inoculated plants (Table 4; Fig. 7), indicating that GH\_A13G0171 could reduce seedling tolerance to salt stress. The corresponding markers SNP\_A13\_1869056 could be applied for improvement of cotton salt tolerance.

Table 4

Statistics of traits related to salt tolerance for the VIGS lines and its control at the seedling stage

	RPH	RSFW	RSDW
pTRV2::00	0.512 ± 0.033	0.497 ± 0.027	0.534 ± 0.032
pTRV2::GH_A13G0171	0.591 ± 0.022*	0.614 ± 0.032*	0.657 ± 0.029*
pTRV2::GH_A08G0488	0.481 ± 0.029	0.474 ± 0.033	0.506 ± 0.036
pTRV2::GH_A10G1620	0.473 ± 0.026	0.494 ± 0.024	0.515 ± 0.030
The data are mean ± SE of three biological replicates. *indicate statistical significance at the 0.05 probability level.			
pTRV2::00: Negative control; pTRV2::GH_A13G0171: GH_A13G0171 knockout seedlings; pTRV2::GH_A08G0488: GH_A08G0488 knockout seedlings; pTRV2::GH_A10G1620: GH_A10G1620 knockout seedlings.			

## Discussion

An ideal association mapping population should encompass sufficient phenotypic and genotypic diversity [34]. Thus, selection of a suitable panel with abundant genetic diversity is especially critical for association mapping. In this study, the population panel consisted of 217 cultivars from different ecological regions (the Yellow River, the Yangtze River, the Northwestern inland region and the Northern China region) of China and from abroad (America and Australia) and had abundant variation in salt tolerance related traits. Phenotypic analysis showed that the salt tolerance related traits in two environments had wide variations and stable changing trends in different environments. Genotyping analysis revealed an average genetic diversity and PIC of 0.303 and 0.248, respectively. As a result, the 217 cotton accessions with broad geographic and genetic diversity coverage in our research provided sufficient detection power for association mapping.

Recent years, association mapping as an important method for the identification of complex quantitative traits has been widely used in cotton for various traits including salt tolerance [35–40]. Jia et al. [21] identified 3 SSR markers significantly associated with the relative survival rate under salt stress through association mapping methods using 323 *G. hirsutum* germplasms. Du et al. [22] performed association analysis of 304 upland cotton cultivars and identified 95 significant associations for 10 salt tolerance related traits at the germination and seedling stages. These studies were conducted based on the SSR markers with lower mapping resolution and smaller genome coverage. Therefore, the genetic factors of salt tolerance related traits discovered with high-density SNP markers could provide more precise insight into salt tolerance in cotton than those previous studies. Sun et al. [24] performed a genome-wide association study to identify marker-trait associations under salt stress using 713 upland cotton accessions through the Illumina Infinium CottonSNP63K array. A total of 23 SNPs were significantly associated with two salt-tolerance-related traits. All of the above reported associations were only detected in a single environment. QTL mapping of important agronomic traits of cotton has been carried out for

nearly two decades, but there are few successful examples of breeding new varieties by marking-assisted breeding [41]. An important reason is the insufficient of available information on multi-environmental detected QTLs. Salt tolerance as a complex quantitative trait controlled by multiple genes needs repeated phenotypic evaluations in multiple environments. In this study, a total of 25 significant associations with 27 SNPs scattered over 12 chromosomes were detected for salt tolerance related traits. Among these, the associations on chromosome A13 and D08 for RPH, A07 for RSFW, A08 and A13 for RSDW were expressed in both two environments, indicating that they were likely to be stable QTLs. These results provided important genetic variations for accelerating the improvement of salt tolerance in cotton.

By comparing our association mapping results with those of previous reports showed that most of the 25 associations were novel reported loci. The physical location of the SSR marker NAU2561 on A05 significantly associated with the relative MDA content reported by Du et al. (2016) [22] was only 1.7 Mb from SNP\_A05\_94406377, which was significantly associated RSDW in 2016. The physical distance between BNL1231, which was reported by Du et al. (2016) [22] to be associated with relative SOD activity, and SNP\_A11\_117938337 which was associated with RSFW in 2017 was only 1 Mb. Considering larger LD decay distances in the cotton genome, it is reasonable to presume that these two loci could be the same loci.

Many studies have reported the positive effects of AQPs in improving plant tolerance to salt stresses. In this study, both gene expression and silencing analyses showed that GH\_A13G0171, which encodes aquaporin (AQP) PIP-type, was likely to regulate the salt stress response. AQPs play an important role in maintaining water homeostasis in the responses of plants to environmental stress by regulating water movement across vacuolar membranes [42]. Overexpression of *MaPIP2-7* in banana improved tolerance to drought, cold, and salt stresses [43]. Overexpression of *SpAQP1* promoted seed germination and root growth under salt stress in transgenic tobaccos [33]. The negative effect of AQP also has been reported. Wang et al. [44] found that overexpression of *GsTIP2;1* in *Arabidopsis thaliana* depressed salt tolerance and dehydration stress. Plants with overexpressed *GsTIP2;1* exhibited higher dehydration rate, suggesting that this gene may mediate stress response by increasing water loss. Our data showed that GH\_A13G0171-silenced plants were significantly increased in salt tolerance related traits, indicating its negative effect in regulating the salt stress response. The exact role of this gene still needs further verification by RNA interference or CRISPR technology. Nonetheless, this study adds further credence to the relationship between AQPs and plant stress response.

## Conclusions

It is of great significance to explore the genetic variants and the underlying candidate genes in upland cotton for salt tolerance improvement. In this study, we reported the genome-wide association mapping of salt tolerant QTLs during the seedling stage by GBS platform. A total of 25 associations with 27 SNPs scattered over 12 chromosomes were detected significantly associated with three salt tolerance related traits in two environments. Among of which, five associations were simultaneously expressed in both two environments. The VIGS assay revealed that GH\_A13G0171, which encodes aquaporin PIP-type, was

negative to regulate the salt stress response. These results will contribute to the genetic improvement of salt tolerance in cotton.

## Materials And Methods

### Plant materials

A collection of 217 upland cotton cultivars (*Gossypium hirsutum* L.) were selected to assemble an association mapping panel in this study. These cultivars were collected from the Institute of Industrial Crops, Jiangsu Academy of Agricultural Sciences, kindly supplied by the Institute of Cotton Research, Chinese Academy of Agricultural Science (CRI-CAAS) and Cotton Research Center of Shandong Academy of Agricultural Sciences. Of these upland cotton cultivars, 162 were collected from China and 55 were introduced from other countries. Chinese cultivars were from four ecological areas: The Yellow River (73), the Yangtze River (62), the Northwestern inland region (23), and the early maturation area in Northern China (4).

In 2015, the 217 upland cotton cultivars were planted at the Lishui Plant Experiment Station, Jiangsu Academy of Agricultural Sciences, China. Genomic DNA extraction referred to the CTAB method described by Paterson et al. (1993) [45]. The central natural open bolls of the 217 upland cotton cultivars were hand-harvested and ginned. The delinted seeds were then used for the subsequent determination of salt tolerance assay.

### Investigation of salt tolerance related traits

The cotton seeds of each cultivar were surface sterilized, transferred into 9 cm sterile Petri dishes on filter paper, and then wetted with 35 mL distilled water for germination. The Petri dishes were incubated in a growth chamber at 32°C with no light. The germinated seeds were then grown in cups of dimension 220 ml with equal amounts of 1:1 mix of the matrix and vermiculite, and placed in 57cm × 41cm plastic pallets in a greenhouse. At the developmental period when two leaves had formed, the seedlings for each cultivar with the same growth vigor were divided into three groups of ten plants. The first ten seedlings with two replications per cultivar were selected before treatment, their plant height, fresh and dry shoot matter weight was determined as the 0-day control; the second ten seedlings with two replications were selected for 350 mM NaCl treatment, and the third ten seedlings with two replications were treated with clean water. After 7 days, the plant height, fresh and dry shoot matter weight of NaCl and clean water treated seedlings was determined respectively. The phenotype increments under 350 mM NaCl in 7 days was calculated using the following formula: (phenotypic effects value under salt stress after 7 days - phenotypic effects value at 0 day). The phenotype increments under clean water in 7 days was calculated using the following formula: (phenotypic effects value under well-watered control after 7 days - phenotypic effects value at 0 day). The ratio of the phenotype increments under 350 mM NaCl and clean water was considered as the relative values. The phenotype of three salt tolerance related traits were calculated using the following formula: (phenotype increments under 350 mM NaCl / (phenotype increments under clean water) × 100%.

## **Library preparation and Illumina sequencing for GBS analysis**

Firstly, a GBS pre-design experiment was performed. The type of enzymes and their fragments size was evaluated based on three criteria. (1) The number of tags must be appropriate. (2) The enzymatic tags must be evenly distributed. (3) Repeated tags must be avoided. Tags about 50 bp was selected to maintain the sequence depth uniformity of different fragments. The GBS library was constructed in accordance with the pre-designed scheme and referred to the method described by Fan et al. (2018) [46].

Then, pair-end sequencing was performed on the selected tags using an Illumina high-throughput sequencing platform. The sequences were sorted according to the barcodes. To make sure that reads were reliable and without artificial bias (low quality paired reads, which mainly resulted from base-calling duplicates and adapter contamination) in the following analyses. C scripts were conducted to processed raw fastq format reads through a series of quality control (QC) procedures as followed: (1) removing reads with unidentified nucleotides (N)  $\geq 10\%$ ; (2) removing reads with  $> 50\%$  bases having phred quality  $< 5$ ; (3) removing reads aligned to the adapter more than 10 nucleotides.

## **SNP detection and annotation**

The clean reads were anchored to the cotton reference genome [30] using Burrows-Wheeler Aligner (BWA) [47]. The SAM tools software [48] was used to convert alignment files to BAM files. Variants calling were performed by using the GATK software. SNP was filtered by the perl script called ANNOVAR [49].

## **Population structure analysis and Linkage disequilibrium**

The population structure of our mapping panel was inferred from the GBS data with STRUCTURE software [50]. Five independent runs were performed; the number of populations (K) was set from 1 to 10; the burn-in time and Markov-chain Monte Carlo replication numbers was set to 10,000. The optimal K value was determined by comparing the LnP (D) and  $\Delta k$  based on the rate of change in LnP (D) [51]. A Q-matrix produced by STRUCTURE listed the estimated membership coefficients in a cluster for the subsequent association analysis. The relative kinship matrix was calculated by SPAGeDi [52]. The software TASSEL 3.0 package was employed to estimate the Linkage disequilibrium (LD) parameters ( $r^2$ ) of adjacent loci within the same LD chromosome for each pair of SNP loci [53].

## **Association mapping**

The TASSEL 3.0 software package [53] was employed to construct association tests of salt tolerance related traits. The Mixed linear models (MLMs) was performed by simultaneously accounting for multiple levels of population structure (Q-matrix) and relative kinship matrix (K-matrix) according to the methods described by Yu et al. (2006) [54].

## **Expression analysis of salt-induced genes**

Total RNA was extracted using an improved CTAB method [55]. For quantitative real-time PCR, PrimeScript RT Master Mix (Perfect Real Time) from TaKaRa (Dalian, China) was used to synthesize first-strand cDNA. The qRT-PCR program was performed on an ABI QuantStudio 5 RealTime PCR System. The qRT-PCR reaction system referred to the method described by Xu et al. (2017) [56]. Quantitative real-time PCR was carried out with three biological replicates. The cotton actin gene was employed as a reference gene. The specific primers for qRT-PCR are listed in Additional file 6: Table S4.

### **Virus-induced gene silencing assay**

About 300-500 bp fragments overlapped the C-terminal and 3' UTR from 3 differentially expressed genes (GH\_A08G0488, GH\_A10G1620 and GH\_A13G0171) were isolated in Moscote2111. The specific primers were listed in Additional file 6: Table S4. These fragments were sequenced and cloned into double enzyme (EcoRI and BamHI) digested pTRV2, generating a new vector named pTRV2::gene. The pTRV1, pTRV2::gene, pTRV2::CLA1 (positive control) and pTRV2::00 (negative control) were introduced into *Agrobacterium tumefaciens* strain GV3101 by heat shock method. The VIGS assay was performed according to the protocol described by Gao et al. (2011) [57]. A ratio of 1:1 mixed pTRV1 and pTRV2::gene were injected into the seedlings with mature cotyledons but without a visible true leaf via a syringe infiltration. The inoculated seedlings were grown in a light incubator at 23°C under a 16-h light and 8-h dark cycle. Two weeks later, the inoculated plants RNA was isolated to detect whether the target gene was still expressed. For each gene, a total of 100 seedlings were knocked out for subsequent salt tolerance identification.

## **Abbreviations**

GWAS: Genome-wide association study; GBS: Genotyping By-Sequencing; BWA: Burrows-Wheeler Aligner; LD: Linkage disequilibrium; VIGS: Virus-induced gene silencing; CV: Coefficients of variation; GO: Gene Ontology; MLM: Mixed linear model; PIC: Polymorphism information content; SNPs: Single nucleotide polymorphisms; RPH: relative plant height; RSFW: relative shoot fresh matter weight; RSDW: relative shoot dry matter weight;

## **Declarations**

### **Acknowledgments**

We would like to thank the Institute of Cotton Research, Chinese Academy of Agricultural Science and Cotton Research Center of Shandong Academy of Agricultural Sciences for providing cotton accessions used in this study. We would like to thank Dr. Peng W. Chee from Molecular Cotton Breeding Laboratory, University of Georgia for helpful comments.

### **Authors' contributions**

Experiments were designed by XLS and WZG. Experiments were performed by PX,

QG, SM, XGZ and ZZX. PX drafted the manuscript, XLS revised the manuscript. All authors read and approved the final manuscript.

## Funding

This work was supported by grants from the Natural Science Foundation of Jiangsu Province (BK20160580), the Major Project of the National Transgene (2016ZX08005-004)

## Ethics approval and consent to participate

Not applicable.

## Consent for publication

Not applicable.

## Competing interests

The authors declared that they had no competing interests.

## Availability of data and materials

All data used in the current study are included in this published article or are available from the corresponding author on reasonable request.

## References

1. Qadir M, Shams M. Some agronomic and physiological aspects of salt tolerance in cotton (*Gossypium hirsutum* L.). *Journal Agronomy Crop Science*. 1997;179:101–6.
2. Greenway H, Munns R. Mechanisms of salt tolerance in nonhalophytes. *Annual Review Plant Physiology*. 1980;31:149–90.
3. Ye W, Liu J. Identification technology on salt tolerance of cotton germplasm and its application. *China Cotton*. 1998;25(9):37–8.
4. Tiwari RS, Picchioni GA, Steiner RL, Jones DC, Hughs SE, Zhang J. Genetic variation in salt tolerance at the seedling stage in an interspecific backcross inbred line population of cultivated tetraploid cotton. *Euphytica*. 2013;194:1–11.
5. Flowers TJ. Improving crop salt tolerance. *J Exp Bot*. 2004;55:307–19.
6. Ashraf M, Foolad MR. Crop breeding for salt tolerance in the era of molecular markers and marker-assisted selection. *Plant Breeding*. 2013;132:10–20.
7. Zhang D, Song HN, Chen H, Hao DR, Wang H, Kan GZ, Jin HX, Yu DY. The acid phosphatase-encoding gene *GmACP1* contributes to soybean tolerance to low-phosphorus stress. *PLOS Genetics*. 2014;10(1):e1004061.


8. Chu S, Wang J, Zhu Y, Liu S, Zhou X, Zhang H, Wang CE, Yang W, Tian Z, Cheng H, Yu D. An R2R3-type MYB transcription factor, *GmMYB29*, regulates isoflavone biosynthesis in soybean. *PLOS Genetics*. 2017;13(5):e1006770.
9. Iquiria E, Humira S, François B. Association mapping of QTLs for sclerotinia stem rot resistance in a collection of soybean plant introductions using a genotyping by sequencing (GBS) approach. *BMC Plant Biol*. 2015;15:5.
10. Nambeesan SU, Mandel JR, Bowers JE, Marek L, Ebert D, Corbi J, Rieseberg L, Knapp SJ, Burke JM. Association mapping in sunflower (*Helianthus annuus* L.) reveals independent control of apical vs. basal branching. *BMC Plant Biol*. 2015;15:84.
11. Zanke CD, Rodemann B, Ling J, Muqaddasi QH, Plieske J, Polley A, Kollers S, Ebmeyer E, Korzun V, Argillier O, Stiewe G, Zschäckel T, Ganai MW, Röder MS. Genome-wide association mapping of resistance to eyespot disease (*Pseudocercospora herpotrichoides*) in European winter wheat (*Triticum aestivum* L.) and fine-mapping of Pch1. *Theor Appl Genet*. 2017;130:505–14.
12. Zhao Y, Wang H, Chen W, Zhao P, Gong H, Sang X, Cui Y. Regional association analysis-based fine mapping of three clustered QTL for *verticillium wilt* resistance in cotton (*G. hirsutum* L.). *BMC Genom*. 2017;18:661.
13. Rahimi Y, Bihamta MR, Taleei A, Alipour H, Ingvarsson PK. Genome-wide association study of agronomic traits in bread wheat reveals novel putative alleles for future breeding programs. *BMC Plant Biol*. 2019;19(1):541.
14. Thapa R, Tabien RE, Thomson MJ, Septiningsih EM. Genome-wide association mapping to identify genetic loci for cold tolerance and cold recovery during germination in rice. *Front Genet*. 2020;11:22.
15. Edae EA, Olivera PD, Jin Y, Poland JA, Rouse MN. Genotype-by-sequencing facilitates genetic mapping of a stem rust resistance locus in *Aegilops umbellulata*, a wild relative of cultivated wheat. *BMC Genom*. 2016;17:1039.
16. Navarro J, Wilcox M, Burgueño J, Romay C, Swarts K, Trachsel S, Preciado E, Terron A, Delgado HV, Vidal V, Ortega A, Banda AE, Montiel N, Ortiz-Monasterio I, Vicente FS, Espinoza AG, Atlin G, Wenzl P, Hearne S, Bucker ES. A study of allelic diversity underlying flowering-time. *Nat Genet*. 2017;49:476–80.
17. Wu X, Guo X, Wang A, Liu P, Wu W, Zhao Q, Zhao M, Zhu Y, Chen Z. Quantitative trait loci mapping of plant architecture-related traits using the high-throughput genotyping by sequencing method. *Euphytica*, 2019, 215 (12).
18. Saxena RK, Kale S, Mir RR, Mallikarjuna N, Yadav P, Das RR, Molla J, Sonnappa M, Ghanta A, Narasimhan Y, Rathore A, Kumar CVS, Varshney RK. Genotyping-by-sequencing and multilocation evaluation of two interspecific backcross populations identify QTLs for yield-related traits in pigeonpea. *Theor Appl Genet*. 2020;133(3):737–49.
19. Peterson GW, Dong YB, Horbach C, Fu YB. Genotyping by-sequencing for plant genetic diversity analysis: a lab guide for SNP genotyping. *Diversity*. 2014;6:665–80.

20. Reinisch AJ, Dong JM, Brubaker CL, et al. A detailed RFLP map of cotton, *Gossypium hirsutum* × *Gossypium barbadense*: Chromosome organization and evolution in a disomic polyploid genome. *Genetics*. 1994;138:829–47.
21. Jia Y, Sun J, Wang X, Zhou Z, Pan Z, He S, Pan B, Wang L, Du X. Molecular diversity and association analysis of drought and salt tolerance in *Gossypium hirsutum* L. Germplasm. *Journal of Integrative Agriculture*. 2014;13(9):1845–53.
22. Du L, Cai C, Wu S, Zhang F, Hou S, Guo W. Evaluation and exploration of favorable QTL alleles for salt stress Related traits in cotton cultivars (*G. hirsutum* L.). *PLOS One*. 2016;11(3):e0151076.
23. Zhao YL, Wang HM, Shao BX, Chen W, Guo ZJ, Gong HY, Sang XH, Wang JJ, Ye WW. SSR-based association mapping of salt tolerance in cotton (*Gossypium hirsutum* L.). *Genetics and molecular research*, 2016, 15(2): gmr.15027370.
24. Sun Z, Li H, Zhang Y, Li Z, Ke H, Wu L, Zhang G, Wang X, Ma Z. Identification of SNPs and candidate genes associated with salt tolerance at the seedling stage in cotton (*Gossypium hirsutum* L.). *Front Plant Sci*. 2018;9:1011.
25. Yasir M, He S, Sun G, Geng X, Pan Z, Gong W, Jia Y, Du X. A genome-wide association study revealed key SNPs/genes associated with salinity stress tolerance in upland cotton. *Genes*. 2019;10:829.
26. Stich B, Maurer HP, Melchinger AE, Frisch M, Heckenberger M, van der Voort JR, Peleman J, Sorensen AP, Reif JC. Comparison of linkage disequilibrium in elite European maize inbred lines using AFLP and SSR markers. *Mol Breeding*. 2006;17:217–26.
27. Yang XH, Yan JB, Zheng YP, Yu JM, Li JS. Reviews of association analysis for quantitative traits in plants. *Acta Agronomica Sinica*. 2007;33:523–30.
28. Sun Z, Wang X, Liu Z, Gu Q, Zhang Y, Li Z, Ke H, Yang J, Wu J, Wu L, Zhang G, Zhang C, Ma Z. Genome-wide association study discovered genetic variation and candidate genes of fibre quality traits in *Gossypium hirsutum* L. *Plant biotechnology journal*. 2017;15(8):982–96.
29. Huang C, Nie X, Shen C, You C, Li W, Zhao W, Zhang X, Lin Z. Population structure and genetic basis of the agronomic traits of upland cotton in China revealed by a genome-wide association study using high-density SNPs. *Plant Biotechnol J*. 2017;15:1374–86.
30. Hu Y, Chen J, Fang L, Zhang Z, Ma W, Niu Y, Ju L, Deng J, Zhao T, Lian J, Baruch K, Fang D, Liu X, Ruan YL, Rahman M, Han J, Wang K, Wang Q, Wu H, Mei G, Zang Y, Han Z, Xu C, Shen W, Yang D, Si Z, Dai F, Zou L, Huang F, Bai Y, Zhang Y, Brodt A, Ben-Hamo H, Zhu X, Zhou B, Guan X, Zhu S, Chen X, Zhang T. *Gossypium barbadense* and *Gossypium hirsutum* genomes provide insights into the origin and evolution of allotetraploid cotton. *Nat Genet*. 2019;51(4):739–48.
31. Xu P, Yang Y, Guo Q, Zhang X, Xu Z, Shen X. Development of EST-SSR and EST-InDel markers associated with salt tolerance in upland cotton. *Cotton Science*. 2016;28(1):65–74.
32. Zhao J, Gao Y, Zhang Z, Chen T, Guo W, Zhang T. A receptor-like kinase gene (*GbRLK*) from *Gossypium barbadense* enhances salinity and drought-stress tolerance in Arabidopsis. *BMC Plant Biol*. 2013;13:110.

33. Chang W, Liu X, Zhu J, Fan W, Zhang Z. An aquaporin gene from halophyte *Sesuvium portulacastrum*, *SpAQP1*, increases salt tolerance in transgenic tobacco. *Plant Cell Rep.* 2016;35:385–95.
34. Flint-Garcia SA, Thuillet AC, Yu JM, Pressoir G, Romero SM, Mitchell SE, Doebley J, Kresovich S, Goodman MM, Buckler ES. Maize association population: a high-resolution platform for quantitative trait locus dissection. *Plant J.* 2005;44:1054–64.
35. Nie X, Huang C, You C, Li W, Zhao W, Shen C, Zhang B, Wang H, Yan Z, Dai B, Wang M, Zhang X, Lin Z. Genome-wide SSR-based association mapping for fiber quality in nation-wide upland cotton inbred cultivars in China. *BMC Genom.* 2016;17:352.
36. Fang L, Wang Q, Hu Y, Jia Y, Chen J, Liu B, Zhang Z, Guan X, Chen S, Zhou B, Mei G, Sun J, Pan Z, He S, Xiao S, Shi W, Gong W, Liu J, Ma J, Cai C, Zhu X, Guo W, Du X, Zhang T. Genomic analyses in cotton identify signatures of selection and loci associated with fiber quality and yield traits. *Nat Genet.* 2017;49:1089–98.
37. Wang M, Tu L, Lin M, Lin Z, Wang P, Yang Q, Ye Z, Shen C, Li J, Zhang L, Zhou X, Nie X, Li Z, Guo K, Ma Y, Huang C, Jin S, Zhu L, Yang X, Min L, Yuang D, Zhang Q, Lindsey K, Zhang X. Asymmetric subgenome selection and cis-regulatory divergence during cotton domestication. *Nat Genet.* 2017;49:579–87.
38. Dong C, Wang J, Chen Q, Yu Y, Li B. Detection of favorable alleles for yield and yield components by association mapping in upland cotton. *Genes Genomics.* 2018;40:725–34.
39. Ma J, Liu J, Pei W, Ma Q, Wang N, Zhang X, Cui Y, Li D, Liu G, Wu M, Zhang X, Song J, Zhang J, Yu S, Yu J. Genome-wide association study of the oil content in upland cotton (*Gossypium hirsutum* L.) and identification of *GhPRXR1*, a candidate gene for a stable QTL *qOC-Dt5-1*. *Plant Science*, 2019, 286:89–97.
40. Zhu G, Gao W, Song X, Sun F, Hou S, Liu N, Huang Y, Zhang D, Ni Z, Chen Q, Guo W. Genome-wide association reveals genetic variation of lint yield components under salty field conditions in cotton (*Gossypium hirsutum* L.). *BMC Plant Biol.* 2020;20(1):23.
41. Guo WZ, Zhang TZ, Zhu XF, Pan JJ. Modified backcross pyramiding breeding with molecular marker-assisted selection and its applications in cotton. *Acta Agronomica Sinica.* 2005;31(8):963–70.
42. Kapilan R, Vaziri M, Zwiazek JJ. Regulation of aquaporins in plants under stress. *Biol Res.* 2018;51:4.
43. Xu Y, Hu W, Liu J, Song S, Hou X, Jia C, Li J, Miao H, Wang Z, Tie W, Xu B, Jin Z. An aquaporin gene *MaPIP2-7* is involved in tolerance to drought, cold and salt stresses in transgenic banana (*Musa acuminata* L.). *Plant Physiol Biochem.* 2020;147:66–76.
44. Wang X, Li Y, Ji W, Bai X, Cai H, Zhu D, Sun X, Chen L, Zhu Y. A novel *Glycine soja* tonoplast intrinsic protein gene responds to abiotic stress and depresses salt and dehydration tolerance in transgenic *Arabidopsis thaliana*. *J Plant Physiol.* 2011;168:1241–8.
45. Paterson AH, Brubaker CL, Wendel JF. A rapid method for extraction of cotton (*Gossypium spp.*) genomic DNA suitable for RFLP or PCR analysis. *Plant Molecular Biology Reporter.* 1993;11(2):122–7.


46. Fan L, Wang L, Wang X, Zhang H, Zhu Y, Guo J, Geng H, Chen Q, Qu Y. A high-density genetic map of extra-long staple cotton (*Gossypium barbadense*) constructed using genotyping-by sequencing based single nucleotide polymorphic markers and identification of fiber traits-related QTL in a recombinant inbred line population. *BMC Genom.* 2018;19:489.
47. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics.* 2009;25(14):1754–60.
48. Li H, Handsaker B, Wysoker A, et al. The sequence alignment/map format and SAMtools. *Bioinformatics.* 2009;25(16):2078–9.
49. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic acids research.* 2010;38(16):e164.
50. Pritchard JK, Wen W. Documentation for STRUCTURE software. Chicago: The University of Chicago Press; 2004.
51. Evanno G, Regaut S, Goudet J. Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Mol Ecol.* 2005;14:2611–20.
52. Hardy OJ, Vekemans X. SPAGEDi: a versatile computer program to analyse spatial genetic structure at the individual or population levels. *Mol Ecol Notes.* 2002;2:618–20.
53. Bradbury PJ, Zhang Z, Kroon DE, Casstevens RM, Ramdoss Y, Buckler ES. TASSEL: Software for association mapping of complex traits in diverse samples. *Bioinformatics.* 2007;23:2633–5.
54. Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB, Kresovich S, Buckler ES. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nat Genet.* 2006;38:203–8.
55. Hu G, Yu S. Extraction of high-quality total RNA in cotton leaf with improved CTAB Method. *Cotton Science.* 2007;19:69–70.
56. Xu P, Gao J, Cao Z, Chee PW, Guo Q, Xu Z, Paterson AH, Zhang X, Shen X. Fine mapping and candidate gene analysis of *qFL-chr1*, a fiber length QTL in cotton. *Theor Appl Genet.* 2017;130:1309–19.
57. Gao X, Britt RC Jr, Shan L, He P. Agrobacterium mediated virus-induced gene silencing assay in cotton. *Journal of Visualized Experiments.* 2011;54:e2938.

## Figures




**Figure 1**

Population structure analysis of 217 cotton cultivars. a Estimated LnP(K) of possible clusters (K) from 1 to 10. b Delta K based on the rate of change of LnP(K) between successive K. c Population structure of 217 Upland cotton accessions based on STRUCTURE when K=3




**Figure 2**

Manhattan and Q–Q plots for salt tolerance related traits in 2016. The horizontal dotted lines of the Manhattan plots with black color represent the genome-wide significance threshold of 0.0001




**Figure 3**

Manhattan and Q-Q plots for salt tolerance related traits in 2017. The horizontal dotted lines of the Manhattan plots with black color represent the genome-wide significance threshold of 0.0001




**Figure 4**

Box plots for the phenotypic values of the association for RPH on chromosome A13. \*\* indicate statistical significance at the 0.01 probability level C: C haplotype G: G haplotype




**Figure 5**

Expression profiles of three promising genes under NaCl stress at three time points in Miscott7913-83. a: Expression profiles of GH\_A13G0171 b: Expression profiles of GH\_A10G1620 c: Expression profiles of GH\_A08G0488 M: Miscott7913-83; Y: Leaves; G: Roots The samples were collected at 0 h, 12 h and 72 h under salt stress. The cotton actin gene was used as an internal reference. The data are mean  $\pm$  SE of three biological replicates. \* and \*\* indicate statistical significance at the 0.05 and 0.01 probability level, respectively.



**Figure 6**

Expression profiles of three promising genes under NaCl stress at three time points in Su 12. a: Expression profiles of GH\_A13G0171 b: Expression profiles of GH\_A10G1620 c: Expression profiles of GH\_A08G0488 S: Su 12; Y: Leaves; G: Roots The samples were collected at 0 h, 12 h and 72 h under salt stress. The cotton actin gene was used as an internal reference. The data are mean  $\pm$  SE of three biological replicates.\* and \*\* indicate statistical significance at the 0.05 and 0.01 probability level, respectively.



**Figure 7**

Functional characterization of GH\_A13\_G0171 VIGS in Miscot2111 a: Transcript levels of GH\_A13G0171 in pTRV2::00 and pTRV2:: GH\_A13G0171 inoculated plants. The cotton actin gene was used as an internal reference. The data are mean  $\pm$  SE of three biological replicates. \*\* indicate statistical significance at the 0.01 probability level, respectively. b: Phenotype identification of salt tolerance of the VIGS line and its control in seedling stage

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile6.xlsx](#)
- [Additionalfile5.tif](#)
- [Additionalfile4.xlsx](#)
- [Additionalfile3.tif](#)
- [Additionalfile2.xlsx](#)
- [Additionalfile1.xlsx](#)