

Maternal Autonomy and High-Risk Pregnancy in Bangladesh: The Mediating Influences of Childbearing Practices and Antenatal Care

Sumaiya Abedin (✉ su.abedin@gmail.com)

University of Rajshahi <https://orcid.org/0000-0003-4675-0029>

Dharma Arunachalam

Monash University

Research article

Keywords: Autonomy, antenatal care, childbearing practices, high-risk pregnancy, mediating effect.

Posted Date: July 2nd, 2020

DOI: <https://doi.org/10.21203/rs.2.15977/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 22nd, 2020. See the published version at <https://doi.org/10.1186/s12884-020-03260-9>.

Abstract

Background Maternal, infant and neonatal mortality rates are high in Bangladesh. Certain childbearing practices and poor utilisation of antenatal care services make Bangladeshi women more vulnerable to experience poor health during pregnancy and adverse pregnancy outcomes. Also, women in Bangladesh remain considerably subordinate to men in almost all aspects of their lives, from education and work opportunities to healthcare utilisation. This study investigates the severity of health complications during pregnancy in relation to women's autonomy, and how childbearing practices and utilisation of antenatal care mediate this relationship.

Methods Data from the most recent Bangladesh Demographic and Health Survey (BDHS) is used in this study. Multinomial regression models (MLRM) are employed to examine the relationship between the outcome variable - high risk pregnancy, and explanatory variables - women's autonomy, childbearing practices and use of antenatal care.

Results In Bangladesh, about 41.5% of women experienced high-risk pregnancies involving multiple health complications. Findings showed that women's autonomy in decision-making, freedom of movement and economic autonomy were significantly associated with high-risk pregnancies. However, women's autonomy in physical mobility in particular did so only through the mediating factors of maternal childbearing practices and antenatal care. Specifically, both early and delayed childbearing and shorter birth interval increased the likelihood of high-risk (multiple complications) pregnancies by about 30% and 31% respectively, high parity increased the risk by 23% and use of antenatal care decreased it by 46%.

Conclusion The Women's decision-making autonomy, freedom of movement and economic autonomy had significant effects on high-risk pregnancies. However, the effects were mediated by both maternal childbearing practices and use of antenatal care in a limited way. Policies and programmes aimed at improving pregnancy outcomes need to focus on all three sets of factors: women's autonomy, childbearing practices and use of antenatal care.

1. Background

Maternal and perinatal deaths as well as poor pregnancy outcomes is recognised as one of the major public health problems in Bangladesh. The common immediate causes of adverse pregnancy outcomes are mainly understood in terms of women's physiological health during pregnancy and childbirth. Although pregnancy is a universal part of female physiology and biology, this event is shaped by the surrounding social environment [1]. A pregnancy becomes high-risk when either the mother or the newborn (or both) have a significant risk of morbidity or mortality [2-3]. Studies on pregnancy health and childbirth are mainly concerned with physiological aspects, and only pay limited attention to maternal fertility behaviour, healthcare practices, and women's socio-economic background. This study extends

this knowledge by examining the relationship between maternal autonomy and the severity of pregnancy health complications followed by high-risk pregnancies among Bangladeshi women.

As highlighted in previous studies, women with low levels of education, low income and living in rural areas tend to enjoy low level of autonomy in reproductive health matters which increase the risk of poor pregnancy outcomes [4-9]. The identified risk factors for developing pregnancy complications were also found to be associated with certain childbearing practices which contribute to poor outcomes. A large number of studies have identified the adverse effects of adolescent pregnancies which increased the chance of experiencing several health complications during pregnancy: anaemia, hypertension or pre-eclampsia, gestational diabetes, placental problems, preterm labour and postpartum haemorrhage (10-14). In Bangladesh, most studies arrived at similar conclusions: adolescent pregnancies hold an increased risk of pregnancy and delivery complications [15-17].

Also, women who had given birth to a large number of children (more than three births) and had them in shorter birth intervals of less than 18 months were found to be more vulnerable to experience several pregnancy complications: pre-eclampsia, pregnancy infections, miscarriage, stillbirth and low birth weight [16, 18-22]. Additionally, access to reproductive healthcare services remained a key determinant of a sound pregnancy and healthy outcome [17, 23]. In this context, a large number of studies from developing countries including Bangladesh identified women's autonomy as a significant factor in the utilisation of antenatal care services [9, 24-27]. Although the associations between pregnancy health status, and maternal childbearing practices as well utilization of maternal healthcare have been widely recognized, the association between maternal autonomy and pregnancy health complications followed by high-risk pregnancies has not yet received similar attention. This study therefore investigates the association between women's autonomy and high-risk pregnancies, and also examines the mediating role of women's childbearing practices and antenatal care utilization in this relationship. The assumed key relations between autonomy and high-risk pregnancies through the pathways of maternal childbearing practices and antenatal care are shown in the diagram in Figure-1.

2. Data And Methodology

The present study is based on data from the *Verbal Autopsy Questionnaire* of Bangladesh Demographic and Health Survey 2011 (BDHS), which was the most recent survey that used the WHO *Verbal Autopsy Questionnaire*. Through this questionnaire, information on several health complications during pregnancy and delivery was collected from ever-married women aged 10-49 years for their most recent pregnancy in the five years preceding the survey. In the BDHS, the participants were recruited using a two-stage stratified cluster sample of households yielding a response rate of 98%. Details of data collection and management procedures are described elsewhere in 2011 BDHS [28]. The analysis reported here is based on a sample of 327 women for whom information was available for all the variables used in this paper.

Outcome variable

The outcome variable in this study is *high-risk pregnancy* which was derived by combining a number of health complications during pregnancy that a woman may have experienced: high blood pressure, heart disease, gestational diabetes, convulsion, vaginal bleeding, puffy face, blunted vision, severe headache, high fever, long labour, shortness of breath and excessive bleeding during labour. Each complication was assigned a score of '0' (did not experience that complication during pregnancy) or '1' (had that complication during pregnancy). The scores for all complications during pregnancy were combined to produce a single measure of pregnancy risk. This index may have values ranging from zero (no complication during pregnancy) to a maximum of 12 complications. Based on the univariate distribution of this variable, this variable was collapsed, for analytical purposes, into a three-category outcome variable: no risk (no complication), low risk (one-two complications), and high risk (three or more complications).

Predictor and mediating variables

The primary predictor variable in this study is comprised of three autonomy variables: *decision-making autonomy* (assessed through questions relating to decisions about household purchases, about their own healthcare, their children's healthcare, contraception, and the number and timing of births), *autonomy in physical mobility or freedom of movement* (assessed through questions related to the ability of women to visit friends and family, and access to healthcare services without any restriction), and *economic autonomy* (assessed through the ability to spend money when necessary, including for healthcare purposes). Responses to all the items were combined to produce a total score for each type of autonomy. Each total score was then collapsed into three categories: no autonomy (respondent can't decide at all, either husband or other family member decides about their lives); partial autonomy (respondent has the ability of make joint decisions with husband or other family members); and high autonomy (respondent has the ability to make the decision on their own). As there were relatively few women who reported 'partial autonomy', we combined the 'partial' and 'high' into a single 'autonomous' category.

In this study, maternal childbearing practices and uptake of antenatal care services were hypothesized to mediate the associations between maternal autonomy and levels of pregnancy risk. Childbearing practices included three variables: timing of the recent birth (low risk: on-time childbearing, 20 to 34 years; and high risk: early or delayed childbearing, below 20 years or above 34 years), parity (low-risk: 1-3 births; and high-risk: more than three births) and birth interval (low-risk: 18 months and more; and high-risk: less than 18 months). Antenatal care uptake was categorised as: women had insufficient antenatal care (0-2 visits) and had sufficient antenatal care (at least three visits as recommended by the WHO).

Statistical analysis

The responses obtained from the study population were analysed using descriptive statistics and cross-tabulations. To investigate the influences of maternal autonomy on high-risk pregnancy multinomial logistic regression models (MLRM) were fitted to the categorical response variable – pregnancy risk, and the odds of low (one or two complications) and high (three or more complications) risks versus no risk were estimated. We fitted two regression models to the data. First, high-risk pregnancy of women is

examined with only autonomy variables included in the model, and then the variables maternal childbearing practices and antenatal care were added to the model to assess their roles in influencing the relationship between autonomy and high-risk pregnancy.

3. Findings Of The Study

Prevalence of pregnancy complications

Table-1 presents information about the most recent live births for which mothers experienced any pregnancy complication. This table shows that over one in ten women (11.5%) had pregnancy-related diabetics. About 9.1% and 6.1% women (i.e. about one in seven women) had experienced two major complications (high blood pressure and convulsion, respectively) that were life-threatening for both the mother and newborn. A significant proportion of women experienced some minor health complications during their pregnancy such as blurred vision (30.3%), severe headache (36.7%), shortness of breath (21.2%) and puffy face (17%). About 28.8% of women had experienced labour lasting more than 12 hours, whereas about 19% women experienced excessive bleeding during labour. When all the complications were combined, it is clear that the vast majority (80%) had experienced at least one complication during pregnancy and childbirth. While about 38.6% of women had experienced one or two complications during pregnancy (low risk), a similar percentage (41.5%) of women were reported to have experienced at least three health complications during pregnancy (high risk).

Table 1: Distribution of health complications during pregnancy among Bangladesh women, BDHS 2011 (n=327)

Pregnancy complications	Percentage of women
High blood pressure	9.1
Gestational diabetes	11.5
Heart disease	4.6
Convulsion	6.1
Vaginal discharge	4.5
Puffy face	17.0
Blurred vision	30.3
Severe headache	36.7
High fever	37.1
Long labour	28.8
Excessive bleeding	19.3
Shortness of breath	21.2
<u>Pregnancy risk</u>	
No risk: No complication	19.9
Low risk: One to Two complications	38.6
High risk: More than two complications	41.5
All	100.0

Table 2 provides information on the three components of autonomy and the four factors, namely maternal age, parity, birth interval and use of antenatal care, that were hypothesised to mediate the relationship between women's autonomy and the experience of high-risk pregnancy. It is interesting that most Bangladeshi women tended to enjoy at least some degree of autonomy in decision making (61.5%) and freedom of mobility (54.5%). But they seem to enjoy very little economic autonomy: about 86% of women included in the study did not report any degree of autonomy in economic matters. In terms of the mediating factors, the vast majority had their recent birth at the appropriate age (neither early, nor late), have had fewer than four births, and the average interval between births was at least 18 months. Table 2 further shows that one half of the women did not receive sufficient antenatal care.

Table 2: Women's autonomy, maternal childbearing practices and antenatal care among Bangladeshi women, BDHS 2011 (n=327)

Autonomy and mediating factors	Percentage of women
Autonomy of women	
<i>Decision-making autonomy</i>	
No autonomy	38.5
Autonomous	61.5
<i>Autonomy in physical mobility</i>	
No autonomy	45.6
Autonomous	54.4
<i>Economic autonomy</i>	
No autonomy	86.2
Autonomous	13.8
-	
Maternal childbearing practices	
<i>Maternal age</i>	
Low-risk (20 to 34 years)	65.5
High-risk (<20 years and/or >34 years)	34.5
<i>Parity</i>	
Low-risk (1-3 births)	75.8
High-risk (> 3 births)	24.2
<i>Birth interval</i>	
Low-risk (> 18 months)	57.5
High-risk (< 18 months)	42.5
-	
Antenatal care visits	
No sufficient (< 3 ANC visits)	49.8
Sufficient (\geq 3 ANC visits)	50.2

High-risk pregnancy and associated factors

We first examine the relationship between women's autonomy, mediating factors and high-risk pregnancy through conventional cross-tabulations. We then present the results from the regression analysis. Percentage distribution in Table 3 shows that, of the three dimensions of autonomy, only economic autonomy (ability to spend money when necessary) had a clear association with pregnancy risk although it was not statistically significant ($p=0.11$). Among those who had the ability to decide on using the available economic resources (e.g. for health care), only 33% reported having experienced multiple complications during their last pregnancy compared to 50% among those who reported that they did not enjoy any autonomy in economic matters. The association was weak ($\text{chi-square}=4.36$, $p=0.11$) which may be related to the fact that there was only a relatively small number of respondents who had reported some level of economic autonomy (13.8%).

In contrast to autonomy variables, all the mediating variables had a strong and statistically significant relationship with pregnancy risks. Early or late childbearing, higher parity and shorter birth intervals increased the risk of experiencing multiple complications (high risk) during pregnancy (Table 3). On the other hand, as would be expected, sufficient utilisation of antenatal health services, decreased the risk of multiple complications.

Table 3: Percentage distribution of women by pregnancy risk, and women's autonomy, maternal childbearing practices and antenatal care of Bangladesh women ($n=327$)

Characteristics of respondents	Pregnancy risk (%)			
	No risk	Low risk	High risk	Chi square (p-value)
Autonomy of women				
<i>Decision-making autonomy</i>				2.874
No autonomy	20.6	29.4	50.0	(0.238)
Autonomous	19.4	34.3	46.3	
-				
<i>Autonomy in physical mobility</i>				4.920
No autonomy	24.8	28.2	47.0	(0.085)
Autonomous	15.7	36.0	48.3	
-				
<i>Economic autonomy</i>				4.355
No autonomy	19.1	30.9	50.0	(0.113)
Autonomous	24.4	42.2	33.4	
Maternal childbearing practices				
<i>Maternal age</i>				
Low-risk (20-34 years)				12.463
High-risk (< 20 years and/or > 34 years)	20.9	32.0	47.1	(0.002)
	17.6	33.7	49.0	
-				
<i>Parity</i>				8.864
Low-risk (1-3 births)	21.8	32.7	45.5	(0.011)
High-risk (> 3 births)	14.0	31.6	54.4	
-				
<i>Birth interval</i>				9.034
Low-risk (> 18 months)	18.5	30.5	51.0	(0.012)
High-risk (< 18 months)	21.6	36.2	42.2	
Antenatal care				
No sufficient (< 3 ANC visits)	18.4	23.9	57.7	14.342
Sufficient (\geq 3 ANC visits)	21.3	40.9	37.8	(0.001)

Low risk: One-Two pregnancy complications

High-risk: More than two pregnancy complications

The estimated odds ratios (OR) from two regression models are presented in Table 4. Model 1 included only the autonomy variables and Model 2 includes both autonomy and mediating variables. In model 1, all three autonomy variables had a strong (OR=0.63 for economic autonomy) to moderately strong (OR=0.79 for decision making) association with high risk (multiple complications) pregnancy. When the mediating variables were added in Model 2, women's autonomy in physical mobility was no more important. This shows that the influence of only one of the three autonomy variables seems to be mediated by factors related to childbearing and the use of antenatal care. However, all mediating variable were still important. Early and late age at childbirth increased the odds of high risk pregnancies by 30%, shorter birth interval and higher parity increased it by 31% and 23% respectively, and sufficient uptake of antenatal care services decreased the odds by almost 46%. The results confirmed that all but one autonomy variable had an independent effect on pregnancy risks, and that pregnancy risks are an outcome of the combined influences of respondents' childbearing practices, use of antenatal care and women's autonomy in decision making and economic autonomy. The role of women's autonomy in pregnancy complications was mediated only in a limited way by childbearing factors and use of antenatal care.

Table 4: Estimated odds ratios (multinomial logistic regression models) of the effects of women's autonomy, maternal childbearing practices, and antenatal care on pregnancy risk of Bangladesh women

Variables	Model 1		Model 2	
	Low-risk vs No risk	High-risk vs No risk	Low-risk vs No risk	High-risk vs No risk
Autonomy of women				
Decision-making autonomy				
No autonomy@	-	-	-	-
Autonomous	0.82	0.79*	0.85	0.81*
-				
Autonomy in physical mobility				
No autonomy@	-	-	-	-
Autonomous	0.86*	0.78*	0.71	1.08
-				
Economic autonomy				
No autonomy@	-	-	-	-
Autonomous	0.55	0.63*	0.53	0.49*
Maternal childbearing practices				
Maternal age				
Low-risk (20-34 years)@			1.21*	1.30*
High-risk (<20 and/or >34 years)				
-				
Parity				
Low-risk (1-3 births)@			-	-
High-risk (> 3 births)			1.24*	1.23*
-				
Birth interval				
Low-risk (> 18 months)@			-	-
High-risk (< 18 months)			1.23*	1.31*

Antenatal care				
Insufficient (< 3 ANC visits)@			-	-
Sufficient (≥ 3 ANC visits)			0.68*	0.54*

Note: * - $p < .05$; @ - reference category

4. Discussion

Pregnancy is a complex and long-term biological transition in a woman's life with many associated health complications. Most pregnancies that result in natural vaginal births involve minor complications, such as nausea and vomiting, headache, blurred vision, puffy face or shortness of breath. On the other hand, major health problems during pregnancy include high blood pressure, convulsion, gestational diabetes, or pregnancy infections. When women experience several of these complications, they are deemed to be carrying a high-risk pregnancy, which contribute to poor outcomes for mother and baby, requiring special medical attention. However, the concept of high-risk pregnancy has not received the same level of attention in the literature as childbearing in general. The present study therefore examined high-risk pregnancies in Bangladesh from a socio-demographic viewpoint.

The results of this study indicate that a significant proportion of women in Bangladesh have experienced pregnancy health complications, both minor and major. Also, a large number of women experienced two or more complications that were life-threatening. This phenomenon indicates an entrenched pattern of high-risk pregnancies in Bangladesh, making women vulnerable to adverse maternal and birth outcomes. The findings on the relationship between women's autonomy, childbearing practices and pregnancy risks illustrate that all three dimensions of women's autonomy, when no other variables were included in the analysis (see Table 4, Model 1), had significant influences on high-risk pregnancies. Earlier studies identified some indirect influences of autonomy indicators on pregnancy health status. It was found that freedom to go outside the home for healthcare purposes resulted in better maternal outcomes [9, 26, 29]. Independence in mobility can enable women to gather relevant information about pregnancy health, complications and remedy measures, thus contributed to healthy outcomes. In addition, high level of decision-making power about contraception, having children at a favourable time, spacing and number of children, as well as decision about own and child's healthcare also contributed to better pregnancy outcomes. In this context, decision-making autonomy, women's freedom of movement and ability to spend money when necessary to access maternal healthcare facilities reduce the prevalence of several health complications during pregnancy by ensuring higher levels of antenatal care utilization, which in turn decreases the chances of high-risk pregnancy. In this way, all three dimensions of maternal autonomy were found to be associated with sound pregnancy health.

In this study, childbearing practices and the use of antenatal care services appeared to have strong effects on the severity of childbearing complications that are followed by high-risk pregnancies. Adolescent pregnancy is a common phenomenon worldwide, but is more prevalent in developing

countries, including Bangladesh. This study found that teenage (and late) pregnancies are more likely to result in several life-threatening complications. The reasons for this might be that as teenage mothers they are more likely than older mothers to be less educated as well as less autonomous, and lacking in awareness and experience regarding the danger signs of pregnancy complications, so they are less likely to receive early prenatal and antenatal care. At the same time, they are more likely to experience several health complications during pregnancy because of the biological effects of early childbearing. A large number of earlier studies have also focused on this phenomenon [14-16, 30-32]. Previous research, as is the case in this study, also showed that the grand multiparous (more than three children) women were more likely to have caesarean deliveries, preterm births and stillbirths (19, 33). Findings of this study confirm this: high birth order, high-risk birth interval and adolescent or delayed childbearing increase the risk of multiple complications during pregnancy, thereby leading to high-risk pregnancy outcomes.

The present study found that short birth intervals of less than 18 months had significant effect on high-risk pregnancies, which is similar to findings other studies that investigated the adverse effects of short birth intervals on maternal health complications (20, 28, 35). Women with very short birth intervals are more likely to have anaemia, gestational diabetics and high-blood pressure, as well as an increased risk premature and low birthweight babies. In addition to this, maternal health complications could occur due to the adverse biological effects of high parity and short birth intervals. Generally, in high order births (more than three) women experience multiple and life-threatening complications. Grand multiparous women are older and less likely to have accessed antenatal care, which results in an increased risk of maternal complications and poor neonatal outcomes (16, 18, 34). The results also showed a significant relationship between high-risk pregnancies and maternal healthcare utilisation. Only one-half of women in the study had received sufficient antenatal care during pregnancy. Women who received sufficient antenatal care and who received treatment for health complications were more likely to experience sound pregnancy health and reduced chances of high-risk pregnancies.

5. Conclusion

The main focus of this research was to investigate the associations between women's autonomy and high-risk pregnancy, and as discussed above, all three indicators of autonomy appeared to have significant effects on high-risk pregnancy (Model 1). However, once the effects of childbearing practices and antenatal care were accounted for, the effects of autonomy in physical mobility became insignificant. This phenomenon indicates that women's freedom of movement influences high-risk pregnancies through maternal age, parity, birth interval as well as uptake of antenatal care. Thus, both childbearing practices and antenatal care mediate the association between maternal autonomy in physical mobility and high-risk pregnancies. In other words, the effects of autonomy are mediate only in a limited way through the mediating factors. To reduce the prevalence of high-risk pregnancies in Bangladesh, attention needs to be given to increase the level of autonomy of women, in particular their autonomy in decision making and economic matters. It is equally important to pay attention to women's age at birth, number of children, intervals between births and access to antenatal care services so as to improve mothers' reproductive health and pregnancy outcomes for both mother and child.

6. Abbreviations

ANC: Antenatal care

BDHS: Bangladesh Demographic and Health Survey

WHO: World Health Organization

7. Declarations

Ethics statement

We obtained permission from the MEASURE DHS to download and analyse the 2011 BDHS data-set for our study purpose. The data were originally collected by the Macro, Calverton, USA. The 2011 BDHS data collection procedures were approved by the ORC Macro-institutional review board. The protocol of the survey was reviewed and approved by the National Ethics Review Committee of the Bangladesh Ministry of Health and Family Welfare.

Consent for publication

Not applicable for this study.

Availability of data

The dataset generated and analysed for the present study is freely available at <http://dhsprogram.com/data/>

Competing interests

None of the authors have any competing interests.

Funding

The authors have no support or funding to report for the present study.

Authors' contributions

SA and DA contributed to the conceptualization and planning of this study. SA performed the statistical analyses and drafted the manuscript. DA revised and edited the final version of this manuscript. Both the authors read and approved the submission.

Acknowledgements

We would like to thank MEASURE DHS for providing us with the nationally representative dataset. We would also like to extend our thanks to the participants of the BDHS 2011.

References

1. Davis, K. and J. Blake 1956. "Social structure and fertility: An analytic framework". *Economic Development and Cultural Change* 211-235.
2. Gilbert, E. S. 2010. *Manual of High Risk Pregnancy and Delivery*. MO: Mosby.
3. Enkin, M., et al. 2000. *A Guide to Effective Care in Pregnancy and Childbirth*. Oxford University Press, Oxford.
4. Kabeer, N. 2011. "Between affiliation and autonomy: navigating pathways of women's empowerment and gender justice in rural Bangladesh". *Development and Change* 42(2): 499-528.
5. Rahman, M. M. and S. Abedin 2010. "Factors affecting neonatal mortality in Bangladesh". *Journal of Health Management* 12(2): 137-152.
6. Salihu, H. M., et al. 2005. "Perinatal mortality associated with abruption placenta in singletons and multiples". *American Journal of Obstetrics and Gynaecology* 193 (1): 198-203.
7. Savitz, D. A., et al. 2004. "Poverty, education, race, and pregnancy outcome". *Ethnicity & Disease* 14(3): 322-329.
8. Choudhury, N., et al. 2012. "Beliefs and practices during pregnancy and childbirth in urban slums of Dhaka, Bangladesh". *BMC Public Health* 12(1): 791.
9. Mumtaz, Z. and S. Salway 2005. "'I never go anywhere': extricating the links between women's mobility and uptake of reproductive health services in Pakistan". *Social Science & Medicine* 60(8): 1751-1765.
10. Sayem, A. and A. T. Nury 2011. "Factors associated with teenage marital pregnancy among Bangladeshi women". *Reproductive Health* 8(1): 16.
11. Liou, J. D., JJ Hsu, LM Lo, SF Chen, and TH Hung. 2010. "Advanced maternal age and adverse perinatal outcomes in an Asian population". *European Journal of Obstetrics & Gynaecology and Reproductive Biology* 148 (1): 21-26.
12. Muganyizi, P. S. and H. L. Kidanto 2009. "Impact of change in maternal age composition on the incidence of Caesarean section and low birth weight: analysis of delivery records at a tertiary hospital in Tanzania, 1999–2005." *BMC Pregnancy and Childbirth* 9(1): 30.
13. Luke, B. and Brown, M. B. 2007. "Elevated risks of pregnancy complications and adverse outcomes with increasing maternal age". *Human Reproduction* 22(5): 1264-1272.
14. Cunningham, A. J. 2001. "What's so bad about teenage pregnancy?" *Journal of Family Planning and Reproductive Health Care* 27(1): 36-41.
15. Gausia, K., et al. 2012. "Obstetric complications and psychological well-being: experiences of Bangladeshi women during pregnancy and childbirth". *Journal of Health, Population, and Nutrition* 30(2): 172.
16. Khan, R., et al. 2012. "An Examination of Women Experiencing Obstetric Complications Requiring Emergency Care: Perceptions and Socio-cultural Consequences of Caesarean Sections in Bangladesh". *Journal of Health, Population, and Nutrition* 30(2): 159.

17. Sikder, S. S., et al. 2011. "Accounts of severe acute obstetric complications in Rural Bangladesh". *BMC Pregnancy and Childbirth* 11(1): 76.
18. Teguate, I., et al. 2012. "Maternal and neonatal outcomes of grand multiparas over two decades in Mali". *Acta Obstetrics and gynaecology Scandinavia* 91(5): 580-586.
19. Shaikh, K., et al. 2011. "The association between parity, infant gender, higher level of paternal education and preterm birth in Pakistan: a cohort study". *BMC Pregnancy and Childbirth* 11(1): 88.
20. DaVanzo, J., et al. 2008. "The effects of pregnancy spacing on infant and child mortality in Matlab, Bangladesh: How they vary by the type of pregnancy outcome that began the interval". *Population Studies* 62(2): 131-154.
21. Royce, R. A. 2006. "Birth spacing—the long and short of it". *JAMA: The Journal of the American Medical Association* 295(15): 1837-1838.
22. Conde-Agudelo, A., et al. 2006. "Birth spacing and risk of adverse perinatal outcomes". *JAMA: the Journal of the American Medical Association* 295(15): 1809-1823.
23. Pervin, J., et al. 2012. "Association of antenatal care with facility delivery and perinatal survival—a population-based study in Bangladesh". *BMC Pregnancy and Childbirth* 12(1): 111.
24. Woldemicael, G. and E. Y. Tenkorang 2010. "Women's autonomy and maternal health-seeking behaviour in Ethiopia". *Maternal and Child Health Journal* 14(6): 988-998.
25. Mistry, R., and O Galal, M Lu. 2009. "Women's autonomy and pregnancy care in rural India: A contextual analysis". *Social Science and Medicine* 69(6): 926-933.
26. Mumtaz, Z. and S. Salway 2007. "Gender, pregnancy and the uptake of antenatal care services in Pakistan". *Sociology of Health & Illness* 29(1): 1-26.
27. Bloom, S. S., et al. 2001. "Dimensions of women's autonomy and the influence on maternal health care utilization in a north Indian city". *Demography* 38(1): 67-78.
28. National Institute of Population Research and Training (NIPORT), Mitra and Associates, ICF International. Bangladesh Demographic and Health Survey 2011. Dhaka and Rockvill.
29. Razzaque, A., et al. 2005. "Pregnancy spacing and maternal morbidity in Matlab, Bangladesh". *International Journal of Gynaecology & Obstetrics* 89: S41-S49.
30. Singh, S. and J. E. Darroch 2000. "Adolescent pregnancy and childbearing: levels and trends in developed countries". *Family Planning Perspectives* 32(1): 14-23.
31. Kim, J. M., L Labrique, KP West, M rashid. 2012. "Maternal morbidity in early pregnancy in rural northern Bangladesh". *International Journal of Gynaecology & Obstetrics*.
32. Lumbiganon, P., et al. 2010. "Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007–08". *The Lancet* 375(9713): 490-499.
33. Geidam, A., et al. 2011. "Pregnancy outcome among grand multiparous women at the University of Maiduguri Teaching Hospital: A case control study". *Journal of Obstetrics & Gynaecology* 31(5): 404-408.

34. Islam, M. A., et al. 2006. "Complications during pregnancy, delivery, and postnatal stages and place of delivery in rural Bangladesh". *Health Care for Women International* 27(9): 807-821.
35. Rutstein, S. O. 2005. "Effects of preceding birth intervals on neonatal, infant and under-five years' mortality and nutritional status in developing countries: evidence from the demographic and health surveys". *International Journal of Gynaecology and Obstetrics* 89(1): 7.

Figures




Figure 1

Relationship between autonomy and high-risk pregnancies through the pathways of childbearing practices and antenatal care