

Coronally Advanced Flap With Connective Tissue Graft For Treating Orthodontic-Associated Miller Class III Gingival Recession of The Lower Incisors: A One-Year Retrospective Study

Evgeny Weinberg (✉ evgenywein@gmail.com)

Tel Aviv University

Roni Kolerman

Tel Aviv University

Omer Cohen

Tel Aviv University

Alon Sebaoun

Tel Aviv University

Gil Slutzkey

Tel Aviv University

Research Article

Keywords: Gingival recession, Miller Class III, mandibular incisors, post-orthodontic

Posted Date: July 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-663623/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Clinical Medicine on January 1st, 2022. See the published version at <https://doi.org/10.3390/jcm11010235>.

Abstract

Aims

To assess the clinical outcome (after one year) of coronally advanced flap combined with connective tissue graft for the treatment of orthodontic-associated Miller Class III gingival recession of the lower incisors.

Materials and methods

This study included 15 patients who had undergone orthodontic treatment prior to development of recession. Measurements of recession depth, recession width, probing depth, and width of keratinized tissue were performed clinically immediately before surgery and after one year. In addition, digital measurements of recession depth, recession width, and root coverage esthetic score were performed on intraoral photographs.

Results

Significant reduction was observed for probing depth, recession depth, and recession width at one year, with significant increase in width of keratinized tissue. Mean root coverage was $83 \pm 24\%$ for recession depth, while complete root coverage was achieved in 10 out of 21 recessions (48%). The average root coverage esthetic score at 12 months was 7.1 ± 2.6 . An interaction was found between initial recession depth and mean root coverage.

Conclusions

Within the limitations of this study, our results confirm that combination of coronally advanced flap and connective tissue graft is effective in reducing post-orthodontic Miller Class III recessions of the mandibular incisors with a concomitant increase in width of keratinized tissue.

Clinical Relevance

These findings suggest that combination of coronally advanced flap with connective tissue graft may significantly improve post-orthodontic Miller Class III recessions of the mandibular incisors, even when the correction of the tooth malposition is unattainable.

Introduction

Gingival recession (GR) is the displacement of the soft tissue margin apical to the cemento-enamel junction (CEJ), which leads to root surface exposure to the oral environment [1]. Untreated buccal GR is associated with thermal and tactile sensitivity, esthetic complaints, and potential complications in maintaining optimal oral hygiene [2]. Moreover, untreated GR has a tendency for further apical displacement over time despite high patient self-care motivation, particularly when the keratinized tissue is reduced or absent [3].

A growing body of evidence relates orthodontic treatment to the development of GR [2, 4–6]. Experimental studies suggest that movement of the root beyond the envelope of the alveolar process causes stretching and thinning of the gingiva, which increases the risk of GR [7, 8], particularly in patients with reduced gingival thickness [5, 9]. Mandibular incisors seem to be the most vulnerable teeth to the development of post-orthodontic GR [5, 10, 11], due to a tooth-arch relationship that results in labially prominent teeth covered with a thin or nonexistent labial plate of bone and relatively thin marginal gingiva [12]. In addition, mandibular lingual fixed retention may be associated with a higher prevalence of GR, especially in proclined lower incisors [10, 13]. Another possible cause of post-orthodontic GR in mandibular incisors may be tooth malposition due to an unwanted force extracted by distortion and activation of the retainer's wire [14].

If malposition of teeth with roots positioned outside the alveolar envelope is the supposed etiology for GR, then orthodontic treatment should be considered with or without periodontal therapy [15]. However, orthodontic retreatment can be performed only when there is enough lingual bone support, assessed by computerized tomography (CBCT) [16]. Moreover, patients often refuse orthodontic correction of roots positioned outside the alveolar envelope, since this type of orthodontic movement requires reinsertion of a fixed orthodontic appliance in the lower arch for several months. In these cases, surgical root coverage can be considered despite the tooth malposition.

Although numerous surgical approaches for root coverage have been reported, the coronally advanced flap (CAF) together with connective tissue graft (CTG) is still considered the gold standard in Miller Class I or II buccal recessions, in terms of clinical outcomes [17–19]. However, limited data is available regarding the Miller Class III recessions [20], particularly in orthodontic-associated GRs of the lower incisors. In such cases, the interdental periodontal support loss is mild to moderate, soft tissue loss in the interdental area is present, or a malpositioning of the teeth prevents the attempt of full root coverage [21].

To help address this lacuna, the objective of this retrospective study was to assess the clinical outcome of CAF combined with CTG for the treatment of orthodontic-associated Miller Class III gingival recession of the lower incisors, one-year post-treatment.

Materials And Methods

The study design was approved by the Institutional Ethics Committee of Tel Aviv University (IEC No. 0002616-1). Data were collected by retrospective evaluation of all consecutive patients that met the inclusion criteria.

The patients were treated by one experienced periodontist (EW) between January 2018 and December 2019. All patients signed an informed consent concerning the planned treatment and received detailed information about alternative options.

The study included 15 patients (2 males, 13 females), aged 19–29 years (23.87 ± 3.2), in whom 21 sites were treated. The mean period between completion of orthodontic treatment and the surgical root coverage was seven years (ranging from 1 to 11).

All patients presented a buccal Miller Class III gingival recession of the lower incisors, had undergone orthodontic treatment prior to development of recession, and had complete clinical records and intraoral photographs. Other inclusion criteria were patient age ≥ 18 years, non-smoking status, and absence of any systemic disease that may affect periodontal treatment.

Exclusion criteria consisted of medically compromised patients, pregnant women, smokers, history of periodontal disease, previous mucogingival procedures including frenectomy, and/or poor hygiene levels (full-mouth plaque score more than 20%). Patients with non-identifiable cemento-enamel junction and those who underwent orthodontic correction of malposed incisors prior to surgery were excluded as well.

Initial therapy

Prior to the surgery, all patients underwent oral hygiene instruction, including atraumatic toothbrushing using modified Bass technique, as well as full-mouth supra- and subgingival scaling and professional tooth cleaning with the use of a rubber cup and low abrasive polishing paste. Surgical treatment of the recession defects was not scheduled until the patient demonstrated an adequate standard of supragingival plaque control.

Surgical Procedures

The surgical technique for gingival recession coverage was the trapezoidal-type of CAF [22], fully covering a CTG obtained by means of de-epithelialization of a free gingival graft (FGG) [23] (Fig. 1). The recipient site was prepared using a #15C blade by a variable (split-full-split) thickness trapezoidal flap extending to the alveolar mucosa that was elevated, and the roots were exposed. To give the CAF a vertical dimension and close adaptation of the alveolar bone and root surfaces, the labial submucosal tissue (LST) was isolated and removed by means of two incisions: one deep, to detach LST from the periosteum; and one superficial, to separate the LST from the inner surface of the alveolar mucosa [24]. FGG was harvested from the canine to first molar area of the palate and de-epithelialized, using a #15C blade extraorally. After mechanical treatment of the exposed root surface with Mini-Five™ Gracey curette (Hu Friedy, Chicago, IL, USA), the CTG was positioned at the level of the CEJ and sutured with single stiches using 7 – 0 resorbable suture (Vicryl®, Ethicon; Johnson & Johnson, Somerville, NJ, USA) to the de-epithelialized papilla. The flap was then positioned at least 1 mm coronally to the CEJ [25] and closed using the same 7 – 0 resorbable suture for single stiches on the lateral release incision and a nonresorbable 6 – 0 suture material (Prolene, Ethicon) for a coronal sling suture around the teeth.

All patients were administered with antibiotic therapy (875 mg of amoxicillin/clavulanate potassium twice daily or 150 mg of clindamycin four times a day for penicillin-sensitive patients) for one week. Anti-inflammatory analgesics (ibuprofen) were prescribed for five days. Patients were also instructed to avoid brushing the teeth in the treated area but to rinse with chlorhexidine solution (0.2%) twice daily for one minute. Sutures were removed after 14 days. Patients were recalled for control visits once a week in the first month, then monthly up to one year post-surgery.

Postoperative healing was uneventful in all treated cases and at all follow-up visits (Fig. 1).

Outcome measurements

The following variables were measured: recession depth (RD), recession width at the CEJ (RW), probing depth (PD), and width of keratinized tissue (KTW). The clinical measurements were performed by the operating periodontist (EW), at the mid-buccal aspect of the root, immediately before surgery (baseline) and at the one-year follow-up visit, with a 1 mm calibrated periodontal probe (PCP-UNC 15, Hu-Friedy). In addition, digital measurements of RD and RW were performed by an independent evaluator (GS), using ImageJ software (<https://imagej.net/>, ImageJ, RRID:SCR_003070) on magnified intraoral images (X10) photographed at the baseline and at the one-year follow-up visit, using a DSLR camera. To facilitate the subsequent analysis, the photographs were centered to the contact region of the lower central incisors at the midline. The digital measurements were calibrated to the true value using the same periodontal probe photographed aligned parallel to the buccal surface of the right central mandibular incisor. Digital measurements were repeated after a minimum of 14 days for all patients and recessions, enabling calculation of the method's error.

The esthetic outcome was evaluated using the root coverage esthetic score (RES) system [26] on intraoral photographs, with the same magnification as for the clinical measurements, by one operator (GS). The maximum esthetic score was 10.

Statistical Analysis

The inter-examiner reproducibility of the clinical and digital measurements of RD and RW, as well as intra-examiner reproducibility of all the digital measurements, was assessed by interclass correlation coefficient. Both descriptive statistics and analysis of variance (ANOVA) were performed to compare the baseline and 12-month outcomes. ANOVA with repeated measures was used to find interactions between the changes from baseline to 12 months in RD, RW, KTW (Δ RD, Δ RW, Δ KTW respectively), and percentage of root coverage (%RC), and the different baseline parameters. A correction by the Benjamini-Hochberg procedure was applied for multiple correlations between the different parameters. The analysis was performed using SPSS version 25.0 (IBM, Armonk, NY, USA).

Results

Significant reduction was observed for PD, RD, and RW at one-year post-treatment, compared to the baseline measurements, with significant increase in KTW (Table 1).

Table 1
Mean and SD at baseline and at 12 months of all evaluated parameters

	Baseline	12 months	p value
	Mean ± SD	Mean ± SD	Baseline Vs. 12 months
PD (mm)	2.3 ± 1.3	1.0 ± 0.5	0.0001
RD (mm)	4.2 ± 1.7	0.7 ± 0.8	0.0001
RW (mm)	3.8 ± 0.8	1.7 ± 1.7	0.0001
KTW (mm)	0.7 ± 0.9	2.6 ± 0.8	0.0001

p < 0.05 indicates statistically significant differences.

PD, probing depth; RD, recession depth; RW, recession width; KTW, keratinized tissue width.

Mean root coverage was $83 \pm 24\%$ for the recession depth, while complete root coverage was achieved in 10 out of 21 recessions (48%) (Table 2). The average root coverage esthetic score at 12 months was 7.1 ± 2.6 (Table 3).

Table 2
Percentage of root coverage and complete root coverage at 12 months

	12 months
	Mean ± SD
Percentage of root coverage (%RC)	$83 \pm 24\%$
Complete root coverage (CRC)	10/21 (48%)

Table 3
Root coverage esthetic score (RES) at 12 months

	Mean \pm SD
GM	4.4 \pm 1.8 (out of 6)
MTC	1.0 \pm 0.2 (out of 1)
STT	0.8 \pm 0.4 (out of 1)
MGJ	0.7 \pm 0.5 (out of 1)
Gingival color	0.4 \pm 0.5 (out of 1)
RES	7.1 \pm 2.6 (out of 10)
GM = gingival margin level; MTC = marginal tissue contour	
STT = soft tissue texture; MGJ = mucogingival junction	

An interaction was found for ΔRD (mm; Dependent variable) to RD and KW at baseline (independent variables). However, after linear regression and Benjamini-Hochberg correction for multiple comparisons, only RD at baseline had an interaction with ΔRD . When %RC, ΔRW , and ΔKTW were the dependent variables, no interactions were found.

The inter- and intra-examiner measure of agreement was Cohen's Kappa = 1, for both clinical and photographic measurements, which indicates a high level of internal consistency.

Discussion

The purpose of the present retrospective study was to evaluate the clinical and esthetic outcomes of the CAF with CTG in the treatment of post-orthodontic Miller Class III gingival recession of the lower incisors. The study included 15 patients who had undergone orthodontic treatment prior to development of recession, and for whom orthodontic correction was not feasible since the incisors were in malposition due to patient objection to corrective procedures or given the absence of lingual bone support as revealed by CBCT.

Most of the patients had completed the orthodontic treatment as young adults (between the ages of 14 and 18 years); however, accurate data could not be obtained regarding the exact starting time of the recession and the changes over time. All the patients reported fast progression of the recession, which caused them great concern and encouraged them to seek treatment urgently.

In this study, we classified gingival recessions using the Miller classification [21], probably the most commonly employed system for this purpose. Cairo and colleagues had suggested another classification system of gingival recessions, based on the interproximal clinical attachment level [27]. According to this

classification, all the treated Miller Class III recessions in this study were RT2. However, in Cairo and colleagues' classification, the malposition of the root is not considered a prognostic criterion, although it is a limiting factor for the amount of root coverage achieved at the buccal site after surgery [14]. This may be associated with the blood supply provided by interproximal soft tissue to the buccal flap/graft during the healing process. Therefore, in cases of post-orthodontic lower incisors recessions, the Miller classification, which takes into consideration the tooth position, is more appropriate.

Applying CAF combined with the CTG technique in Miller Class I or II buccal recession-type defects affecting the lower incisors, Zucchelli and colleagues reported up to 98% mean root coverage with 88% complete root coverage (CRC), provided the labial submucosal tissue was removed from the inner surface of the alveolar mucosa [24]. However, the scholarship on Miller Class III recessions is quite limited, mostly focused on the comparison of techniques or the evaluation of results obtained by new procedures.

In the present evaluation, mean root coverage for Miller Class III orthodontic-associated gingival recessions was found to be $83 \pm 24\%$, while CRC was recorded in 48% of the sites at one-year compared to baseline measurements. RD, RW, and PD were significantly reduced; however, only RD at baseline had demonstrated interaction with Δ RD. Interestingly, no interaction was found for %RC, RW, and KTW, pointing to the RD as the main prognostic factor for the root coverage.

A case series study evaluating CAF with CTG for the treatment of Miller Class III gingival recessions in mandibular central incisors found 86% of mean root coverage with 43% of CRC [28]. Interestingly, an additional study by the same authors using CTG in combination with tunnel technique showed only 74% of mean root coverage and 14% of CRC for Miller Class III defects in mandibular incisors [29]. The investigators explained these differences by facts that minimal efforts were made in coronally advancing the flap with the tunnel technique and that most of the root coverage occurred through vascularization of the exposed graft.

A recent study comparing clinical outcomes of CTG with and without enamel matrix derivative in the treatment of Miller Class III defects on mandibular anterior teeth reported 78% and 73% mean root coverage and 22% and 18% of CRC, respectively, 12-months post-treatment [30]. Another recent study presented results of modified tunnel double papilla procedure for root coverage of post-orthodontic Miller Class III recessions with either the removal or not of the bonded lingual retainer (BLR) [14]. An improvement of 43% was found without the removal of the BLR, compared to 87% once the BLR was removed prior to surgery; however, only a small number of cases was evaluated in each group.

Regarding the mean root coverage ($83 \pm 24\%$), our results are in line with the findings from the aforementioned studies; however, CRC was slightly higher (48%) in our evaluation. Although CRC was reached in nearly half of the sites only and consequently, given a final mean root coverage esthetic score of 7.1 ± 2.6 , this parameter may not be considered the most important treatment outcome; this is particularly true regarding non-esthetic areas such as anterior mandible, which is known to have worse results than other regions in the oral cavity (Harris, 1994; Nart et al., 2012). Evidently, a maintainable

situation was provided in all cases, by creating an adequate zone of keratinized tissue ($KTW_{12 \text{ months}} = 2.6 \pm 0.8 \text{ mm}$), thus preventing frenum and muscle pull and allowing easier brushing in this area.

The mean esthetic score in the present study was 7.1 ± 2.6 . In a recent meta-analysis, Cairo and colleagues compared esthetic results in different root coverage procedures [31]. The mean esthetic score in the different studies included in the meta-analysis for procedures with a CTG ranged between 7.5 and 9.45. Our results are somewhat lower than their results. However, we evaluated only Miller Class III recessions; since the RES is largely sensitive to the amount of root coverage [26] and Miller Class III recessions are considered highly challenging for CRC [20], the present esthetic score result is satisfactory.

An interesting finding of the present study is the statistically significant PD reduction (2.3 ± 1.3 to 1.0 ± 0.5), aligning with a previous report [24]. This favorable outcome may be due to the baseline PD measurements at the buccal aspect of lower incisors that are greater compared to the mean buccal PD reported in the literature for gingival recession [32]. A possible explanation involves the presence of deep bone dehiscence and a shallow vestibulum depth. In this clinical scenario, a buccal PD apical to the gingival recession forms more frequently, once a deep gingival defect reaches the vestibular fornx. The clinically significant PD and RD reductions, alongside the increase in KTW, represent a meaningful improvement in the periodontal conditions, which could be beneficial for the prognosis of the affected tooth and could justify the treatment of deep gingival recession affecting lower incisors even in the absence of the traditional esthetic or dentine hypersensitivity indications.

The major limitations of this study are its limited sample and lack of blinded examination of some of the clinical parameters. However, photographic examination by an independent evaluator and a high level of internal consistency partially compensate for these drawbacks.

Within the limitations of this study, our results confirm that combination of CAF and CTG may significantly improve post-orthodontic Miller Class III recessions of the mandibular incisors, even when correction of the tooth malposition is unattainable.

Declarations

Compliance with Ethical Standards

Conflict of Interest: Evgeny Weinberg declares that he has no conflict of interest. Roni Kolerman declares that he has no conflict of interest. Omer Cohen declares that he has no conflict of interest. Alon Sebaoun declares that he has no conflict of interest. Gil Slutzkey declares that he has no conflict of interest.

Funding: This research received no external funding.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards."

Informed consent: Informed consent was obtained from all individual participants included in the study.

References

1. Löe, H., Ånerud, Å. & Boysen, H. The Natural History of Periodontal Disease in Man: Prevalence, Severity, and Extent of Gingival Recession. *Journal of Periodontology*, **63**, 489–495 <https://doi.org/10.1902/jop.1992.63.6.489> (1992).
2. Zucchelli, G. & Mounssif, I. (2015) Periodontal plastic surgery. *Periodontology 2000* **68**:333–368. doi: 10.1111/prd.12059
3. Chambrone, L. & Tatakis, D. N. Long-Term Outcomes of Untreated Buccal Gingival Recessions: A Systematic Review and Meta-Analysis. *Journal of Periodontology*, **87**, 796–808 <https://doi.org/10.1902/jop.2016.150625> (2016).
4. Ji, J-J. *et al.* Prevalence of gingival recession after orthodontic treatment of infraversion and open bite. *Journal of Orofacial Orthopedics / Fortschritte der Kieferorthopädie*, **80**, 1–8 <https://doi.org/10.1007/s00056-018-0159-8> (2019).
5. Melsen, B. & Allais, D. Factors of importance for the development of dehiscences during labial movement of mandibular incisors: A retrospective study of adult orthodontic patients. *American Journal of Orthodontics and Dentofacial Orthopedics*, **127**, 552–561 <https://doi.org/10.1016/j.ajodo.2003.12.026> (2005).
6. Vasconcelos, G., Kjellsen, K., Preus, H., Vandevska-Radunovic, V. & Hansen, B. F. Prevalence and severity of vestibular recession in mandibular incisors after orthodontic treatment. *Angle Orthod*, **82**, 42–47 <https://doi.org/10.2319/021411-108.1> (2012).
7. Wennstrom, J. L., Lindhe, J., Sinclair, F. & Thilander, B. Some periodontal tissue reactions to orthodontic tooth movement in monkeys. *Journal of Clinical Periodontology*, **14**, 121–129 <https://doi.org/10.1111/j.1600-051x.1987.tb00954.x> (1987).
8. Steiner, G. G., Pearson, J. K. & Ainamo, J. (1981) Changes of the Marginal Periodontium as a Result of Labial Tooth Movement in Monkeys. *52*:314–320. doi: 10.1902/jop.1981.52.6.314
9. Yared, K. F. G., Zenobio, E. G. & Pacheco, W. (2006) Periodontal status of mandibular central incisors after orthodontic proclination in adults. *American Journal of Orthodontics and Dentofacial Orthopedics* **130**:6.e1-6.e8. doi: 10.1016/j.ajodo.2006.01.015
10. Renkema, A. M. *et al.* Gingival labial recessions in orthodontically treated and untreated individuals: a case - control study. *Journal of Clinical Periodontology*, **40**, 631–637 <https://doi.org/10.1111/jcpe.12105> (2013).
11. Bin Bahar, B. S. K., Alkhaldy, S. R., Kaklamanos, E. G. & Athanasiou, A. E. Do orthodontic patients develop more gingival recession in anterior teeth compared to untreated individuals? A systematic review of controlled studies. *Int. Orthod*, **18**, 1–9 <https://doi.org/10.1016/j.ortho.2019.08.025> (2020).
12. Dorfman, H. S. Mucogingival changes resulting from mandibular incisor tooth movement. *American Journal of Orthodontics*, **74**, 286–297 [https://doi.org/10.1016/0002-9416\(78\)90204-x](https://doi.org/10.1016/0002-9416(78)90204-x) (1978).

13. Pandis, N., Vlahopoulos, K., Madianos, P. & Eliades, T. Long-term periodontal status of patients with mandibular lingual fixed retention. *The European Journal of Orthodontics*, **29**, 471–476 <https://doi.org/10.1093/ejo/cjm042> (2007).
14. Beitzlitz, I., Barzilay, V., Rayyan, F., Sebaoun, A. & Sarig, R. Post-Orthodontic Lower Incisors Recessions: Combined Periodontic and Orthodontic Approach. *International Journal of Environmental Research and Public Health*, **17**, 8060 <https://doi.org/10.3390/ijerph17218060> (2020).
15. Laursen, M. G., Rylev, M. & Melsen, B. The role of orthodontics in the repair of gingival recessions. *American Journal of Orthodontics and Dentofacial Orthopedics*, **157**, 29–34 <https://doi.org/10.1016/j.ajodo.2019.01.023> (2020).
16. Chaushu, S. & Stabholz, A. (2013) Orthodontic-associated localized gingival recession of lower incisors: who should treat it? *Compend Contin Educ Dent* 34 Spec No:6–12; quiz 13
17. Rocuzzo, M., Bunino, M., Needleman, I. & Sanz, M. Periodontal plastic surgery for treatment of localized gingival recessions: a systematic review. *Journal of Clinical Periodontology*, **29**, 178–194 <https://doi.org/10.1034/j.1600-051x.29.s3.11.x> (2002).
18. Chambrone, L. *et al.* Root-Coverage Procedures for the Treatment of Localized Recession-Type Defects: A Cochrane Systematic Review. *Journal of Periodontology*, **81**, 452–478 <https://doi.org/10.1902/jop.2010.090540> (2010).
19. Sebaoun, A., Nemcovsky, C. E., Beitzlitz, I. & Moses, O. Modified Tunnel Double Papilla Procedure for Root Coverage in the Anterior Mandible. *Int J Periodontics Restorative Dent*, **39**, 503–509 <https://doi.org/10.11607/prd.3784> (2019).
20. Tatakis, D. N. *et al.* Periodontal Soft Tissue Root Coverage Procedures: A Consensus Report From the AAP Regeneration Workshop. *Journal of Periodontology*, **86**, S52–S55 <https://doi.org/10.1902/jop.2015.140376> (2015).
21. Miller, P. D. Jr A classification of marginal tissue recession. *Int J Periodontics Restorative Dent*, **5**, 8–13 (1985).
22. De Sanctis, M. & Zucchelli, G. Coronally advanced flap: a modified surgical approach for isolated recession-type defects: Three-year results. *Journal of Clinical Periodontology*, **34**, 262–268 <https://doi.org/10.1111/j.1600-051x.2006.01039.x> (2007).
23. Zucchelli, G. *et al.* Patient morbidity and root coverage outcome after subepithelial connective tissue and de-epithelialized grafts: a comparative randomized-controlled clinical trial. *Journal of Clinical Periodontology*:no-no, <https://doi.org/10.1111/j.1600-051x.2010.01550.x> (2010).
24. Zucchelli, G., Marzadori, M., Mounssif, I., Mazzotti, C. & Stefanini, M. Coronally advanced flap + connective tissue graft techniques for the treatment of deep gingival recession in the lower incisors. A controlled randomized clinical trial. *J Clin Periodontol*, **41**, 806–813 <https://doi.org/10.1111/jcpe.12269> (2014).
25. Pini Prato, G. P. *et al.* Coronally Advanced Flap: The Post-Surgical Position of the Gingival Margin Is an Important Factor for Achieving Complete Root Coverage. *Journal of Periodontology*, **76**, 713–722

- <https://doi.org/10.1902/jop.2005.76.5.713> (2005).
26. Cairo, F., Rotundo, R., Miller, P. D. & Pini Prato, G. P. Root Coverage Esthetic Score: A System to Evaluate the Esthetic Outcome of the Treatment of Gingival Recession Through Evaluation of Clinical Cases. *Journal of Periodontology*, **80**, 705–710 <https://doi.org/10.1902/jop.2009.080565> (2009).
27. Cairo, F., Nieri, M., Cincinelli, S., Mervelt, J. & Pagliaro, U. The interproximal clinical attachment level to classify gingival recessions and predict root coverage outcomes: an explorative and reliability study. *J Clin Periodontol*, **38**, 661–666 <https://doi.org/10.1111/j.1600-051X.2011.01732.x> (2011).
28. Nart, J. *et al.* Subepithelial connective tissue graft in combination with a coronally advanced flap for the treatment of Miller Class II and III gingival recessions in mandibular incisors: a case series. *Int J Periodontics Restorative Dent*, **32**, 647–654 (2012).
29. Nart, J. & Valles, C. Subepithelial Connective Tissue Graft in Combination with a Tunnel Technique for the Treatment of Miller Class II and III Gingival Recessions in Mandibular Incisors: Clinical and Esthetic Results. *Int J Periodontics Restorative Dent*, **36**, 591–598 <https://doi.org/10.11607/prd.2748> (2016).
30. Mercado, F., Hamlet, S. & Ivanovski, S. Subepithelial connective tissue graft with or without enamel matrix derivative for the treatment of multiple Class III-IV recessions in lower anterior teeth: A 3-year randomized clinical trial. *Journal of Periodontology*, **91**, 473–483 <https://doi.org/10.1002/jper.19-0058> (2020).
31. Cairo, F. *et al.* Aesthetic-And patient-related outcomes following root coverage procedures: A systematic review and network meta-analysis. *Journal of Clinical Periodontology*, **47**, 1403–1415 <https://doi.org/10.1111/jcpe.13346> (2020).
32. Cairo, F., Pagliaro, U. & Nieri, M. Treatment of gingival recession with coronally advanced flap procedures: a systematic review. *Journal of Clinical Periodontology*, **35**, 136–162 <https://doi.org/10.1111/j.1600-051x.2008.01267.x> (2008).

Figures



Figure 1

(A) Pre-op; (B) Flap elevation; (C) Fixation of CTG at CEJ level; (D) Closure of CAF above the CEJ; (E) 2 weeks follow-up; (F) 1 month follow-up; (G) 3 months follow-up; (H) 12 months follow-up.