Depression, Anxiety and Stress in Women with Breech Pregnancy - "Be not Afraid": A Prospective Study

Madeleine Schauer

Charite Universitätsmedizin Berlin

Elisabetta Latartara

Universita Cattolica del Sacro Cuore

Maria Alonso-Espias

La Paz University Hospital: Hospital Universitario La Paz

Emma Rossetti

University of Udine: Università degli Studi di Udine

Pimrapat Gebert

Charite Universitätsmedizin Berlin

Wolfgang Henrich

Charite University Hospital Berlin: Charite Universitätsmedizin Berlin

Larry Hinkson (✉ larry.hinkson@charite.de)

Charité University Hospital <https://orcid.org/0000-0002-9170-6345>

Research article

Keywords: Breech, Pregnancy, Depression, Anxiety, Stress, Anxiety Disorder, Depression Anxiety Stress Scale

Posted Date: June 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-664800/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Breech presentation at term is associated with higher perinatal complications through the increased incidence of cesarean section and complications by vaginal breech birth. The decision-making process between breech delivery, trial of external cephalic version or cesarean section may lead to maternal depression, stress and anxiety. Mental disorder in itself is associated with poorer maternal and fetal outcomes, such as preterm birth, small for gestational age infants, lower birth weight and increased risk for postnatal depression.

This study aims to evaluate the level of psychological distress for women with breech presentation compared to cephalic presentation. We hypothesized, that women with breech presentation have higher levels of depression, stress and anxiety than other pregnant women. Secondary objectives were to analyze potential demographic risk factors and comorbidity of psychological distress in breech pregnancy.

Methods: The breech study group was formed by 379 women with breech presentation. A sample of 128 women with cephalic presentation was recruited during routine clinical care. Depression, anxiety and stress symptoms were ascertained by means of the Depression-Anxiety-Stress-Score (DASS)-21 questionnaire. Categorical data was analyzed with Chi-square or exact test, continuous data with unpaired t-test or Mann-Whitney-U-test. Demographic risk factors were identified using a binary logistic regression model.

Results: Prevalence of psychological distress among women with breech was not higher compared to those of other pregnant women. Symptomatic depression, anxiety and stress affected 5.8%, 14.5% and 11.9% of women with breech, respectively. Decreasing age was identified as a risk factor for anxiety ($p = 0.006$). Multiparity increased risk for depression ($p = 0.001$), for anxiety ($p = 0.026$) and for perinatal stress ($p = 0.010$). More than 80% of women with depressive symptoms had comorbidities of psychological distress.

Conclusions: Breech presentation compared to cephalic presentation was not associated with higher levels of psychological distress. However, breech pregnancies are affected by symptoms of potential mental disorder. Multiparous women and younger women may need additional support and would benefit from a standardized screening tool for the assessment of perinatal psychological distress.

Clinical Trial Registration: Ethical approval (EA2/241/18) was granted by the Ethics Commission of the Charité University Hospital on the 23.01.2019 (ClinicalTrials.gov Identifier: NCT03827226)

Background

Approximately 2–4% of pregnancies are breech at term. [1, 2] Breech in itself is associated with higher stillbirth rate, fetal growth restriction and oligohydramnios. [1] In up to 95% a cesarean birth is planned. [3–6] Elevated perinatal mortality, birth trauma, low APGAR scores and neonatal asphyxia are reported for

vaginal breech compared to elective cesarean birth. [3, 4, 7–13] Cesarean birth minimizes the fetal risk, but increases risk for the mother and future pregnancies. [7, 14, 15] One effective method to decrease cesarean birth for breech is the external cephalic version.

Confronted with a breech pregnancy, women may be overburdened with decision-making before delivery with an increasing level of depression, anxiety and stress expected. Psychological distress in breech pregnancy has however never been investigated.

Small for gestational age infants, lower birth weight and increased risk for preterm birth and postnatal depression have been observed in women with mental disorders during pregnancy. [16–21] Despite this, few units establish routine clinical screening for psychological distress. [22]

The prevalence of antenatal depression decreases from first to third trimester. [16, 23, 24] A cohort study found 13.5% of the participants at 32 weeks of gestation at risk for depression. [25] Other authors indicated a point prevalence between 8.5% and 11.1% for minor and major depression in the third trimester. [16, 24, 26]

Antenatal anxiety is more prevalent than depression. [27–29] Lee et al. revealed that 54% of the women experienced symptoms of anxiety in at least one trimester and 35.8% in the last trimester. [27]

Pregnancy anxiety as fear linked to the pregnancy itself (e.g. fears about oneself and baby's well-being, concerns about labor and childbirth), is found to be one of the most potent risk factors for adverse outcomes. [17, 18]

High-risk-pregnancy, increasing gestational age, younger maternal age and history of drinking are shown to be predictors for anxiety. [27, 30] Depression as well, is associated with a high-risk pregnancy, lower maternal education and social factors such as lack of social support and domestic violence. [30, 31]

The primary objective of the study was to evaluate the prevalence of symptoms of depression, anxiety and stress for women with breech compared to women with cephalic presentation. We hypothesized higher levels of psychological distress among women with breech.

Secondary objectives were to determine (1) potential risk factors for high levels of distress among breech pregnancies, (2) the influence of gravidity on distress and (3) the level of comorbidity between depression, anxiety and stress.

Methods

We conducted a prospective observational study between February 2019 and September 2020 in the obstetric clinic of Charité University Hospital Berlin.

Ethical approval (EA2/241/18) was granted by the Ethics Commission of the Charité University Hospital (ClinicalTrials.gov Identifier: NCT03827226). Written informed consent was provided by all women who

agreed to take part in the study.

We used a self-designed questionnaire to collect demographic data, including age, body mass index (BMI), gestational week, gravidity, parity, history of cesarean birth, spontaneous birth, miscarriage and pregnancy termination. Any pre-existing health-condition and gestational complication as well as smoking and consumption of alcohol or drugs were recorded. Clinical symptoms of depression, anxiety and stress were ascertained by means of the Depression Anxiety Stress Score (DASS)-21 questionnaire. [32]

The breech study group was recruited from pregnant women with breech presentation attending the consultant-led breech clinic. Breech was confirmed on sonographic examination from the 36th week of pregnancy. The consultation involved discussing the options of vaginal breech birth, attempting external cephalic version or planning a cesarean birth.

Recruitment of the control group was conducted by direct approach in the general obstetric outpatient clinic. Those patients presented for normal delivery planning in the third trimester of pregnancy with cephalic presentation.

All women had a singleton gestation, were at least 18 years old, able to sign the informed consent and had basic German or English language skills. Exclusion criteria were history of mental disorder, use of antidepressant medication or anxiolytics and any fetal anomalies.

The DASS-21 is a self-report questionnaire consisting of 21 questions for measuring depression, anxiety and stress as negative emotional states. Developed by P. F. Lovibond and S. H. Lovibond as DASS-42, this short form version is also a well validated screening tool.[32, 33]

The questionnaire consists of three 7-item subscales that measure depression, anxiety and stress. Patients estimate the degree of symptoms they have experienced over the last seven days in a four-point-Likert-scale (0–3 points), with higher values indicating greater distress. The total score of each scale can range from 0 to 21 points and is built by summing all of the corresponding items. A total level of distress can be ascertained, ranging from 0 to 63.

Results can be converted in percentile ranks, which divide the scale in five severity groups, namely normal (≤ 7), mild (8–17), moderate (18–26), severe (27–34) and extremely severe (≥ 35) [34]. Cut-off scores for symptomatic depression, anxiety and stress were defined as moderate level or greater, meaning scores of 7, 6 and 10, respectively.

The use of somatic items (e.g. fatigue, sleep disturbance, constipation and decreased appetite) in the assessment of depression is common. [35, 36] This can cause an overestimation of depressive symptoms in pregnancy. [37] Due to the absence of somatic items in DASS-21, it is more appropriate for screening in pregnancy.

Statistical analysis

To calculate power we assumed the effect size of stress was 0.4, a sample of 379 in breech and 128 in control group, yielding a power of 97.4% with a significant level of 0.05 (two-sided). [38]

Continuous variables were presented with mean and standard deviation (SD), or median and interquartile range (25th percentile, 75th percentile), depending on the distribution. Histogram and Shapiro-Wilk test were used to explore the normal distribution. Chi-square or exact test by Monte-Carlo-Method was calculated for categorical data, whereas continuous data was analyzed with unpaired t-test for normal distribution or Mann-Whitney-U test for non-normal distribution.

In order to determine influencing factors for presence of psychological distress (moderate level or greater), we analyzed several potential demographic risk factors. Univariate association between distress and demographic factors was carried out using Chi-square. Variables tested were: maternal age, BMI before pregnancy, history of breech, family history breech, pregnancy risk, gestational diabetes, pre-existing health condition, hypothyroidism, gravidity, parity, history of spontaneous birth, cesarean birth, miscarriage or pregnancy termination and fetal presentation. All variables that showed statistical significance and relevant factors (e.g. age, BMI) were included in a multiple logistic regression model. Presence or absence of depression, anxiety, stress and total distress were defined as dependent variables.

A p-value < 0.05 was assumed as statistically significant. Data was analyzed with SPSS software version 25.0 (SPSS Inc., Chicago, Illinois, USA).

Results

A total of 564 women enrolled in the study, of which 57 women were excluded (Fig. 1). Of the remaining 507 pregnancies 379 had breech and 128 cephalic presentation.

Baseline characteristics group are shown in Table 1. There was no significant difference in maternal age and BMI between both groups ($p = 0.070$, $p = 0.447$).

Table 2 shows median scores and prevalence of severity grades of the DASS-21 questionnaire. Clinical symptoms of moderate to severe depression symptoms were found in 5.8% of breech pregnancies. Symptomatic antenatal anxiety occurred in 14.5%. Symptomatic stress was found in 12%.

Prevalence of symptomatic psychological distress (moderate to extremely severe) is presented in Fig. 2.

A Mann-Whitney-U-Test showed with no significant differences for scores of depression, stress and total distress between the groups. The median anxiety score of the control study group was significantly higher ($p = 0.033$).

No significant results were found for differences in severity grades, neither for the recommended five grades (normal to extremely severe) nor for those defined by our team (normal and symptomatic).

In order to determine demographic confounders and the influence of the fetal presentation on the presence of symptoms of mental disorders, a multiple logistic regression model was performed (Table 3).

Gestational diabetes was found to increase the risk of stress by 2.42 times ($p = 0.051$). Preexisting health condition (hypothyroidism excluded) had a 1.78 times higher risk of anxiety ($p = 0.042$). Multiparity significantly increased risk for depression, stress and total distress.

After adjusting for age, BMI as well as gestational and medical history, fetal presentation was not a significant predictor of psychological distress in pregnancy. Breech pregnancies were 1.51 times more likely to be stressed during third trimester ($p = 0.279$), but had lower risk for depression and anxiety, however these results were not statistically significant.

A multiple logistic regression model was performed to determine risk factors for psychological distress with breech. Adjusted Odds Ratios (95% CI) are shown in Table 4.

Multiparity was found to be the most important risk factor for symptoms of mental disorder. Women who have given birth to at least two children had an eightfold higher risk for depression ($p = 0.001$), threefold higher risk for anxiety ($p = 0.026$) and fourfold higher risk for stress ($p = 0.010$) compared to nulliparous women. Being younger increased the risk for the presence of symptomatic anxiety ($p = 0.006$). Gestational diabetes was 2.83 times more likely to cause high total distress scores ($p = 0.050$) and 2.61 times increased risk for stress, although this association was not significant ($p = 0.088$).

Breech pregnancies were divided into four groups: (1) primigravida, (2) multigravida with either history of term delivery, (3) history of miscarriage or (4) history of delivery and miscarriage.

Multigravida compared to primigravida were significantly more stressed ($p = 0.046$) and had higher total scores ($p = 0.044$).

Another relation was found for primigravida vs. multigravida with history of term delivery. Multigravida had higher depression scores ($p = 0.007$) as well as stress scores ($p = 0.043$).

All other combinations, especially the comparison between history of delivery and history of miscarriage did not show significant results. Comorbidity with depression and anxiety occurred in 3.6%, depression and stress in 4.2% and anxiety and stress in 6.3% of breech cases.

Multimorbidity (symptoms of depression, anxiety and stress simultaneously) was found in 11 women, representing 50.0%, 20.0% and 24.4% of the depressive, anxious and stressed population, respectively (Fig. 3). Of those with symptomatic depression, 9.1% were comorbid for anxiety and 22.7% for stress, the other 18.2% had no comorbid distress. Of those women with symptoms of anxiety, more than half had no comorbid distress, another 3.6% rated themselves as depressive and 23.6% as stressed. Combinations with either depression or anxiety was found in 11.1% and 28.8% of the participants suffering from stress, whereas a third of the stressed women had no other psychological comorbidities.

Discussion

Main findings

The prevalence of symptoms for moderate to extremely severe depression, anxiety and stress in the breech pregnancy was 5.8%, 14.5% and 11.9%, respectively. There was no statistical difference compared to cephalic pregnancy.

Interpretation

Studies show that women with high-risk pregnancy have a higher risk of psychological distress in pregnancy. [30, 39] We used the well-established DASS-21 questionnaire to assess symptoms of depression, anxiety and stress in breech compared to cephalic pregnancy. [32, 33]

In the breech group symptoms of mild depression were present in 5% and moderate to extremely severe symptoms in 5.8%. One study by Barber et al. found 10.5% of pregnant women as mildly depressive and 21.5% as moderate or highly depressive. [38] In this study however, women with known mental disorders were not excluded, pregnancies with breech were not defined and an online survey was used. Other studies showed variable prevalence rates of 13.5% and 11.1% for depression in pregnancy with the Edinburgh Postnatal Depression Scale. [16, 25] These variations are likely because women with known mental disorders were not excluded and gestational ages were also not defined. A meta-analysis on prevalence and incidence of perinatal depression excluded studies based on self-report screens and found a point prevalence of 8.5% for minor and major depression. [26]

Mild and symptomatic anxiety was found in 11.9% and 14.5% of the breech study group. Other studies such as Barber et al and Lee et al. showed higher rates (31.2% and 35.8%) for symptomatic anxiety in pregnancy. [38] [27] However, these studies were limited as patients with known mental disorders were not excluded, gestational age and fetal presentation was not defined and different screening tools were used. A meta-analysis by Dennis et al. could identify the difference between the prevalence of anxiety over all trimesters, when ascertained by self-report symptoms (18.0%) or clinical diagnosis as a measurement (15.2%). [28] Prevalence in third trimester was determined as being 24.6% for self-report and 15.4% for clinical diagnosis of any anxiety disorder, similar to the incidence in our breech study group.

In our study, 9% of breech pregnancies suffered from mild stress and 11.9% from moderate to extremely severe stress. Compared to depression and anxiety, antenatal stress seems to be neglected in research as it does not offer a medical diagnosis of a mental disorder. Two studies on stress in pregnancy presented varying results. Woods et al. used a clinical screening protocol for psychosocial strain and found 78% had low to moderate and 6% high stress levels. [22] In comparison, Barber et al. found that 16.3% of pregnant women had moderate or higher levels of stress. [38]

We performed an analysis of prevalence rates of combined components of psychological disorders in the breech population. Comorbid depression and anxiety affected 3.6%, depression and stress 4.2% and anxiety and stress 6.3%. Of the women with symptoms of depression, we found more than 80% had high levels of other psychological comorbidities. Comorbidity of antenatal depression and anxiety has been investigated in numerous studies. [20, 27, 29, 40, 41] A meta-analysis showed in the third trimester that

comorbid anxiety and mild to severe depression occurred in 9.5% of all cases and moderate to severe depression in 6.6%. [41] There remains a paucity in the literature however on the role of mental stress in pregnancy, therefore the association with other comorbidities has been rarely described.

We also analyzed predictors of psychological disorder in women with breech. We found, that women with multiparity, gestational diabetes and decreasing age were at higher risk. In the literature there are conflicting results on the influence of parity. Fairbrother et al. found no significant difference between nulliparas and multiparas, Dipietro et al. found a higher prevalence among multiparas whereas Gillespie et al. found a higher prevalence in primiparas. [42–44]

We found that breech pregnancy beyond 36 weeks of pregnancy, even with the challenges of deciding on interventions did not have significantly higher levels of psychological distress symptoms, compared to cephalic pregnancies. In fact, women with cephalic presentation in the control group scored higher on the anxiety subscale. Presumably, this might be a result of the collocation of the control group with other obstetric complications (e.g. hypertension, gestational diabetes, oligohydramnios, previous cesarean birth, previous stillbirth) which are associated with higher rates of anxiety. [30] However, a logistic regression analysis of our study and control groups showed no influence of these comorbidities on the presence of psychological distress. Anxiety could also be affected by fear of childbirth in general. Laursen et al observed a significant association of fear of childbirth with depressive and anxious symptoms. [45] Rouhe et al. found that fear of childbirth affects primarily nulliparous women and women with a history of cesarean birth. [46] A proportion of women presenting in the breech clinic may perhaps be more desirous for the chance to deliver vaginally and may have less fear of childbirth.

Strengths and Limitations

This is a large prospective study performed on over 370 participants with breech. This is the first study looking at breech and the influence of perinatal psychological distress symptoms. Importantly, all women with pre-known mental disorders were excluded to minimize selection bias and a well validated screening tool was used.[33]

The DASS-21 self-report questionnaire is however time dependent, evaluating symptom over the past week. Negative emotional states at other time points may have influenced self-assessment scores. Nevertheless, questionnaires are not used for diagnosis, but rather as a screening tool. Socio-economic confounders such as relationship status were not ascertained. There is already extensive research done on social predictors of psychological distress and we chose to look primarily at clinical parameters.

Despite no increase in psychological distress symptoms in women with breech in general, we found that those associated with multiparity, gestational diabetes and decreasing age are at higher risk of developing a mental illness and therefore require additional support. Screening for mental disorders should be established in clinical routine to detect women who may be at high risk of mental illness. Hare et al recently showed that anxiety in pregnancy was associated with a 15% increased risk of postnatal depression and this can negatively influence mother-infant bonding.[47] Optimal antenatal care should

therefore include assessment of the mental health status of expecting mothers and screening for stress in particular. Furthermore, perinatal and delivery complications such as can occur with breech pregnancy can increase a child's risk for anxiety independent from the parental psychopathology.[48] Freed et al also found it useful to screen for anxiety in pregnancy especially in mothers with a known psychiatric disease such as bipolar disorder as this can influence psychopathology in offspring.[49]

Support should then be tailored to meet individual needs. We therefore recommend a standardized screening tool such as the DASS-21 questionnaire for the specific categories of perinatal psychological distress in order to make comparisons between future studies compatible. Indeed, prenatal screening for anxiety can be implemented into prediction models used to earlier identify mothers and offspring at risk. [50]

Conclusion

In women with breech pregnancies significant symptomatic depression, anxiety and stress symptoms were found in 5.8%, 14.5% and 11.9% respectively. Compared to cephalic pregnancy this was not higher. However, multiparity, gestational diabetes and decreasing maternal age were identified as potential factors for developing mental distress in breech pregnancies and require additional support.

Declarations

Ethics approval and consent to participate

Ethical approval (EA2/241/18) was granted by the Ethics Commission of the Charité University Hospital on the 23.01.2019 (ClinicalTrials.gov Identifier: NCT03827226). All study participants provided written consent.

Availability of data and materials

Data sharing is available on reasonable request from the corresponding author.

Disclosure of Competing interests

The authors report no conflict of interest.

Funding

There were no sources of financial support.

Authors' Contributions

MS and LH wrote the paper. MS, LH, EL, MA-E and ER collected data. PG,MS and LH designed and analyzed the study. WH designed and reviewed the manuscript. All authors contributed, reviewed and

approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Macharey G, Gissler M, Rahkonen L, Ulander VM, Vaisanen-Tommiska M, Nuutila M, Heinonen S. Breech presentation at term and associated obstetric risks factors-a nationwide population based cohort study. *Arch Gynecol Obstet*. 2017;295(4):833–8.
2. Noli SA, Baini I, Parazzini F, Mauri PA, Vignali M, Gerli S, Favilli A, Cipriani S. Preterm Birth, Low Gestational Age, Low Birth Weight, Parity, and Other Determinants of Breech Presentation: Results from a Large Retrospective Population-Based Study. *Biomed Res Int*. 2019;2019:9581439.
3. Ekeus C, Norman M, Aberg K, Winberg S, Stolt K, Aronsson A. Vaginal breech delivery at term and neonatal morbidity and mortality - a population-based cohort study in Sweden. *The journal of maternal-fetal neonatal medicine: the official journal of the European Association of Perinatal Medicine the Federation of Asia Oceania Perinatal Societies the International Society of Perinatal Obstet*. 2019;32(2):265–70.
4. Fonseca A, Silva R, Rato I, Neves AR, Peixoto C, Ferraz Z, Ramalho I, Carocha A, Felix N, Valdoleiros S, et al. Breech Presentation: Vaginal Versus Cesarean Delivery, Which Intervention Leads to the Best Outcomes? *Acta Med Port*. 2017;30(6):479–84.
5. Lee HC, El-Sayed YY, Gould JB. Population trends in cesarean delivery for breech presentation in the United States, 1997–2003. *Am J Obstet Gynecol*. 2008;199(1):59 e51–8.
6. Hehir MP, O'Connor HD, Kent EM, Fitzpatrick C, Boylan PC, Coulter-Smith S, Geary MP, Malone FD. Changes in vaginal breech delivery rates in a single large metropolitan area. *Am J Obstet Gynecol*. 2012;206(6):498 e491–4.
7. Villar J, Carroli G, Zavaleta N, Donner A, Wojdyla D, Faundes A, Velazco A, Bataglia V, Langer A, Narvaez A, et al. Maternal and neonatal individual risks and benefits associated with caesarean delivery: multicentre prospective study. *BMJ*. 2007;335(7628):1025.
8. Diro M, Puangsricharn A, Royer L, O'Sullivan MJ, Burkett G: **Singleton term breech deliveries in nulliparous and multiparous women: a 5-year experience at the University of Miami/ Jackson Memorial Hospital**. *Am J Obstet Gynecol* 1999, **181**(2):247–252.
9. Su M, McLeod L, Ross S, Willan A, Hannah WJ, Hutton E, Hewson S, Hannah ME. Term Breech Trial Collaborative G: **Factors associated with adverse perinatal outcome in the Term Breech Trial**. *Am J Obstet Gynecol*. 2003;189(3):740–5.
10. Goffinet F, Carayol M, Foidart JM, Alexander S, Uzan S, Subtil D, Breart G, Group PS. Is planned vaginal delivery for breech presentation at term still an option? Results of an observational prospective survey in France and Belgium. *Am J Obstet Gynecol*. 2006;194(4):1002–11.

11. Giuliani A, Scholl WM, Basver A, Tamussino KF. Mode of delivery and outcome of 699 term singleton breech deliveries at a single center. *Am J Obstet Gynecol*. 2002;187(6):1694–8.
12. Azria E, Le Meaux JP, Khoshnood B, Alexander S, Subtil D, Goffinet F, Group PS. Factors associated with adverse perinatal outcomes for term breech fetuses with planned vaginal delivery. *Am J Obstet Gynecol*. 2012;207(4):285 e281–9.
13. Hofmeyr GJ, Hannah M, Lawrie TA. **Planned caesarean section for term breech delivery.** *Cochrane Database Syst Rev* 2015(7):CD000166.
14. Villar J, Valladares E, Wojdyla D, Zavaleta N, Carroli G, Velazco A, Shah A, Campodonico L, Bataglia V, Faundes A, et al. Caesarean delivery rates and pregnancy outcomes: the 2005 WHO global survey on maternal and perinatal health in Latin America. *Lancet*. 2006;367(9525):1819–29.
15. Young CB, Liu S, Muraca GM, Sabr Y, Pressey T, Liston RM, Joseph KS, Canadian Perinatal Surveillance S. Mode of delivery after a previous cesarean birth, and associated maternal and neonatal morbidity. *CMAJ*. 2018;190(18):E556–64.
16. Felice E, Saliba J, Grech V, Cox J. Prevalence rates and psychosocial characteristics associated with depression in pregnancy and postpartum in Maltese women. *J Affect Disord*. 2004;82(2):297–301.
17. Dunkel Schetter C. Psychological science on pregnancy: stress processes, biopsychosocial models, and emerging research issues. *Annu Rev Psychol*. 2011;62:531–58.
18. Dunkel Schetter C, Tanner L. Anxiety, depression and stress in pregnancy: implications for mothers, children, research, and practice. *Curr Opin Psychiatry*. 2012;25(2):141–8.
19. Alder J, Fink N, Bitzer J, Hosli I, Holzgreve W: **Depression and anxiety during pregnancy: a risk factor for obstetric, fetal and neonatal outcome? A critical review of the literature.** *The journal of maternal-fetal & neonatal medicine: the official journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstet* 2007, **20**(3):189–209.
20. Field T, Diego M, Hernandez-Reif M, Figueiredo B, Deeds O, Ascencio A, Schanberg S, Kuhn C. Comorbid depression and anxiety effects on pregnancy and neonatal outcome. *Infant Behav Dev*. 2010;33(1):23–9.
21. Grigoriadis S, Graves L, Peer M, Mamisashvili L, Tomlinson G, Vigod SN, Dennis CL, Steiner M, Brown C, Cheung A, et al: **Maternal Anxiety During Pregnancy and the Association With Adverse Perinatal Outcomes: Systematic Review and Meta-Analysis.** *J Clin Psychiatry* 2018, 79(5).
22. Woods SM, Melville JL, Guo Y, Fan MY, Gavin A. Psychosocial stress during pregnancy. *Am J Obstet Gynecol*. 2010;202(1):61.e61-67.
23. Teixeira C, Figueiredo B, Conde A, Pacheco A, Costa R. Anxiety and depression during pregnancy in women and men. *J Affect Disord*. 2009;119(1–3):142–8.
24. Gaynes BN, Gavin N, Meltzer-Brody S, Lohr KN, Swinson T, Gartlehner G, Brody S, Miller WC. **Perinatal depression: prevalence, screening accuracy, and screening outcomes.** *Evid Rep Technol Assess (Summ)* 2005(119):1–8.

25. Evans J, Heron J, Francomb H, Oke S, Golding J. Cohort study of depressed mood during pregnancy and after childbirth. *BMJ*. 2001;323(7307):257–60.
26. Gavin NI, Gaynes BN, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T. Perinatal depression: a systematic review of prevalence and incidence. *Obstet Gynecol*. 2005;106(5 Pt 1):1071–83.
27. Lee AM, Lam SK, Sze Mun Lau SM, Chong CS, Chui HW, Fong DY. Prevalence, course, and risk factors for antenatal anxiety and depression. *Obstetrics gynecology*. 2007;110(5):1102–12.
28. Dennis CL, Falah-Hassani K, Shiri R. Prevalence of antenatal and postnatal anxiety: systematic review and meta-analysis. *Br J Psychiatry*. 2017;210(5):315–23.
29. Andersson L, Sundstrom-Poromaa I, Wulff M, Astrom M, Bixo M. Depression and anxiety during pregnancy and six months postpartum: a follow-up study. *Acta Obstet Gynecol Scand*. 2006;85(8):937–44.
30. Rezaee R, Framarzi M. Predictors of mental health during pregnancy. *Iran J Nurs Midwifery Res*. 2014;19(7 Suppl 1):45–50.
31. Lancaster CA, Gold KJ, Flynn HA, Yoo H, Marcus SM, Davis MM. Risk factors for depressive symptoms during pregnancy: a systematic review. *Am J Obstet Gynecol*. 2010;202(1):5–14.
32. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol*. 2005;44(Pt 2):227–39.
33. Lovibond SH, Lovibond PF: **Manual for the Depression Anxiety Stress Scales. (2nd Ed.)**. In. Sydney: Psychology Foundation; 1995.
34. Crawford JR, Garthwaite PH, Lawrie CJ, Henry JD, MacDonald MA, Sutherland J, Sinha P. A convenient method of obtaining percentile norms and accompanying interval estimates for self-report mood scales (DASS, DASS-21, HADS, PANAS, and sAD). *Br J Clin Psychol*. 2009;48(Pt 2):163–80.
35. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry*. 1961;4:561–71.
36. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry*. 1960;23(1):56–62.
37. Klein MH, Essex MJ. Pregnant or depressed? The effect of overlap between symptoms of depression and somatic complaints of pregnancy on rates of major depression in the second trimester. *Depression*. 1994;2(6):308–14.
38. Barber CC, Steadman J. Distress levels in pregnant and matched non-pregnant women. *Aust N Z J Obstet Gynaecol*. 2018;58(1):128–31.
39. Costa ECV, Castanheira E, Moreira L, Correia P, Ribeiro D, Graca Pereira M. **Predictors of emotional distress in pregnant women: the mediating role of relationship intimacy.** *J Ment Health* 2017:1–9.
40. Heron J, O'Connor TG, Evans J, Golding J, Glover V, Team AS. The course of anxiety and depression through pregnancy and the postpartum in a community sample. *J Affect Disord*. 2004;80(1):65–73.

41. Falah-Hassani K, Shiri R, Dennis CL. The prevalence of antenatal and postnatal co-morbid anxiety and depression: a meta-analysis. *Psychol Med*. 2017;47(12):2041–53.
42. Gillespie SL, Mitchell AM, Kowalsky JM, Christian LM. Maternal parity and perinatal cortisol adaptation: The role of pregnancy-specific distress and implications for postpartum mood. *Psychoneuroendocrinology*. 2018;97:86–93.
43. Fairbrother N, Janssen P, Antony MM, Tucker E, Young AH. Perinatal anxiety disorder prevalence and incidence. *J Affect Disord*. 2016;200:148–55.
44. Dipietro JA, Costigan KA, Sipsma HL. Continuity in self-report measures of maternal anxiety, stress, and depressive symptoms from pregnancy through two years postpartum. *J Psychosom Obstet Gynaecol*. 2008;29(2):115–24.
45. Laursen M, Hedegaard M, Johansen C, Danish National Birth C. Fear of childbirth: predictors and temporal changes among nulliparous women in the Danish National Birth Cohort. *BJOG: an international journal of obstetrics gynaecology*. 2008;115(3):354–60.
46. Rouhe H, Salmela-Aro K, Halmesmaki E, Saisto T. Fear of childbirth according to parity, gestational age, and obstetric history. *BJOG: an international journal of obstetrics gynaecology*. 2009;116(1):67–73.
47. Hare MM, Kroll-Desrosiers A, Deligiannidis KM. Peripartum depression: Does risk versus diagnostic status impact mother-infant bonding and perceived social support? *Depress Anxiety*. 2021;38(4):390–9.
48. Hirshfeld-Becker DR, Biederman J, Faraone SV, Robin JA, Friedman D, Rosenthal JM, Rosenbaum JF. Pregnancy complications associated with childhood anxiety disorders. *Depress Anxiety*. 2004;19(3):152–62.
49. Freed RD, Tompson MC, Otto MW, Nierenberg AA, Hirshfeld-Becker D, Wang CH, Henin A. Early risk factors for psychopathology in offspring of parents with bipolar disorder: the role of obstetric complications and maternal comorbid anxiety. *Depress Anxiety*. 2014;31(7):583–90.
50. Hochman E, Feldman B, Weizman A, Krivoy A, Gur S, Barzilay E, Gabay H, Levy J, Levinkron O, Lawrence G. Development and validation of a machine learning-based postpartum depression prediction model: A nationwide cohort study. *Depress Anxiety*. 2021;38(4):400–11.

Tables

Table 1: Baseline characteristics of breech and control study group.

Variable	Breech (n = 379)	Control (n = 128)	p-value
	N (%)	N (%)	
Maternal age			
Mean (SD)	32.66 (4.49)	33.49 (4.50)	0.070
< 20	1 (0.3)	1 (0.8)	0.451
20 – 29	81 (21.4)	21 (16.4)	
30 – 39	271 (71.4)	97 (75.8)	
≥ 40	26 (6.9)	9 (7.0)	
Body Mass Index (kg/m²)			
Mean (SD)	23.46 (4.11)	23.13 (4.79)	0.447
Underweight	19 (5.1)	11 (8.7)	0.410
Normal weight	250 (66.7)	84 (66.7)	
Overweight	79 (21.0)	21 (16.7)	
Obesity	27 (7.2)	10 (7.9)	
Gravidity			
Median (IQR)	1.00 (1.00, 2.00)	2.00 (1.00, 2.00)	0.009
Primigravida	231 (61.1)	63 (49.2)	0.018
Multigravida	147 (38.9)	65 (50.8)	
Parity			
0	283 (74.9)	74 (57.8)	0.001
1	72 (19.0)	38 (29.7)	
≥ 2	23 (6.1)	16 (12.5)	
History of miscarriage			
Spontaneous abortion	80 (21.2)	19 (15.0)	0.128
Pregnancy termination	13 (3.4)	12 (9.4)	0.007
History of delivery			
Vaginal	76 (20.1)	33 (25.8)	0.177
Cesarean birth	13 (3.4)	22 (17.3)	< 0.001
Gestational complications			
Diabetes	23 (6.1)	18 (14.2)	0.004
Others	13 (3.4)	17 (13.4)	< 0.001
Pre-existing health condition			
Hypothyroidism	46 (12.1)	24 (18.9)	0.056
Others	73 (19.3)	48 (37.8)	< 0.001

Table 2: Median (IQR) and prevalence of severity grades of the DASS-21 of breech and control study group.

	Breech = 379				Control = 128			
Variable	Depression		Anxiety		Stress		Total	
	B	C	B	C	B	C	B	C
Score								
Median	1.00	1.00	2.00	2.50	4.00	4.00	7.00	8.00
(IQR)	(0.00, 2.00)	(0.00, 2.75)	(1.00, 4.00)	(1.00, 5.00)	(2.00, 7.00)	(2.00, 7.00)	(4.00, 13.00)	(4.00, 13.75)
p-value	0.577		0.035		0.597		0.404	
Severity grades N (%)								
Normal	338 (89.2)	113 (88.3)	279 (73.6)	81 (63.3)	300 (79.1)	103 (80.4)	292 (77.0)	96 (75.0)
Mild	19 (5.0)	6 (4.7)	45 (11.9)	19 (14.8)	34 (9.0)	13 (10.2)	37 (9.8)	14 (10.9)
Moderate	14 (3.7)	7 (5.4)	38 (10.0)	16 (12.5)	27 (7.1)	10 (7.8)	44 (11.6)	16 (12.5)
Severe	5 (1.3)	1 (0.8)	10 (2.6)	6 (4.7)	12 (3.2)	1 (0.8)	3 (0.8)	1 (0.8)
Extremely severe	3 (0.8)	1 (0.8)	7 (1.9)	6 (4.7)	6 (1.6)	1 (0.8)	3 (0.8)	1 (0.8)
p-value	0.904		0.120		0.660		0.976	

Table 3: Binary logistic regression model for demographic confounders of distress.

Variables	All pregnancies (n = 507)							
	Depression		Anxiety		Stress		Total	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value
Maternal age (yr)	0.95 (0.90 – 1.08)	0.738	0.93 (0.88 – 0.99)	0.015	0.96 (0.90 – 1.03)	0.218	0.95 (0.89 – 1.01)	0.101
Body Mass Index								
Underweight	0.47 (0.05 – 4.04)	0.490	1.98 (0.77 – 5.07)	0.155	1.57 (0.49 – 5.06)	0.446	0.79 (0.22 – 2.89)	0.727
Normal weight	1.00	-	1.00	-	1.00	-	1.00	-
Overweight	1.39 (0.54 – 3.60)	0.501	1.29 (0.69 – 2.43)	0.422	0.76 (0.34 – 1.71)	0.513	0.87 (0.43 – 1.78)	0.702
Obesity	2.48 (0.80 – 7.70)	0.117	1.50 (0.64 – 3.49)	0.348	2.31 (0.96 – 5.56)	0.062	1.74 (0.73 – 4.12)	0.212
Parity								
0	1.00	-	1.00	-	1.00	-	1.00	-
1	1.70 (0.65 – 4.49)	0.283	1.10 (0.59 – 2.04)	0.764	1.52 (0.75 – 3.11)	0.249	1.46 (0.75 – 2.81)	0.264
≥ 2	7.68 (2.77 – 21.28)	<0.001	2.24 (0.97 – 5.19)	0.060	4.2 (1.71 – 10.32)	0.002	4.04 (1.72 – 9.47)	0.001
Gestational diabetes								
Yes vs. No	0.52 (0.11 – 2.50)	0.415	1.58 (0.72 – 3.49)	0.254	2.42 (1.00 – 5.91)	0.051	2.33 (1.03 – 5.26)	0.043
Gestational complications								
Yes vs. No	1.28 (0.27 – 6.14)	0.756	1.43 (0.56 – 3.65)	0.459	1.35 (0.42 – 4.34)	0.616	1.38 (0.48 – 3.98)	0.556
Hypothyroidism								
Yes vs. No	1.06 (0.34 – 3.27)	0.923	1.07 (0.52 – 2.20)	0.849	0.70 (0.26 – 1.86)	0.469	1.16 (0.53 – 2.54)	0.704
Other pre-existing health condition								
Yes vs. No	1.21 (0.48 – 3.01)	0.689	1.78 (1.02 – 3.10)	0.042	0.95 (0.46 – 1.94)	0.881	1.38 (0.74 – 2.60)	0.311
Fetal presentation								
Cephalic	1.00	-	1.00	-	1.00	-	1.00	-
Breech	0.90 (0.37 – 2.18)	0.816	0.70 (0.40 – 1.23)	0.215	1.51 (0.72 – 3.17)	0.279	1.21 (0.63 – 2.34)	0.561

Table 4: Multiple logistic regression model for risk factors of distress in breech pregnancy

Variables	Depression		Anxiety		Stress		Total	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value
Maternal age	1.01 (0.91 – 1.13)	0.824	0.91 (0.85 – 0.97)	0.006	0.95 (0.88 – 1.03)	0.212	0.95 (0.89 – 1.02)	0.186
Body Mass Index								
Underweight	0.64 (0.07 – 6.17)	0.701	1.86 (0.55 – 6.32)	0.322	1.55 (0.40 – 6.10)	0.529	1.22 (0.31 – 4.76)	0.777
Normal weight	1.00	-	1.00	-	1.00	-	1.00	-
Overweight	1.00 (0.30 – 3.33)	0.996	1.41 (0.67 – 2.96)	0.367	0.69 (0.28 – 1.75)	0.438	0.82 (0.35 – 1.88)	0.631
Obesity	2.71 (0.73 – 10.04)	0.136	2.30 (0.87 – 6.08)	0.095	2.20 (0.79 – 6.10)	0.131	1.91 (0.69 – 5.29)	0.211
Parity								
0	1.00	-	1.00	-	1.00	-	1.00	-
1	1.35 (0.41 – 4.48)	0.625	1.16 (0.54 – 2.52)	0.705	1.55 (0.68 – 3.50)	0.296	1.24 (0.56 – 2.76)	0.597
≥ 2	8.61 (2.50 – 29.57)	0.001	3.35 (1.15 – 9.74)	0.026	4.27 (1.42 – 12.86)	0.010	4.33 (1.53 – 12.27)	0.006
Gestational diabetes								
Yes vs. No	0.37 (0.04 – 3.72)	0.398	2.08 (0.75 – 5.76)	0.157	2.61 (0.87 – 7.85)	0.088	2.83 (1.00 – 7.99)	0.050
Gestational complications								
Yes vs. No	1.48 (0.17 – 12.85)	0.725	1.21 (0.25 – 6.01)	0.814	1.66 (0.34 – 8.13)	0.533	1.35 (0.27 – 6.66)	0.713
Hypothyroidism								
Yes vs. No	0.65 (0.14 – 3.12)	0.589	0.74 (0.27 – 2.05)	0.564	0.52 (0.15 – 1.83)	0.307	0.83 (0.30 – 2.31)	0.727
Other pre-existing health condition								
Yes vs. No	1.27 (0.42 – 3.89)	0.675	1.75 (0.85 – 3.60)	0.132	1.24 (0.55 – 2.81)	0.607	1.43 (0.67 – 3.06)	0.361

Figures

Figure 1

Patient recruitment process.

Figure 2

Prevalence of symptomatic distress (moderate to extremely severe) measured by the DASS-21 for breech and control study group.

Figure 3

Psychological comorbidities of women with breech presentation. Subgroup analysis of women with symptoms of depression (A), anxiety (B) and stress (C). They had either no comorbidity (green: only one

scale of the DASS-21 above cut-off), one comorbidity (yellow: two scales above cut-off) or multimorbidity (red: all three scales above cut-off).