

Source Attribution of Salmonellosis by Time and Geography in New South Wales, Australia

Angus McLure (✉ angus.mclure@anu.edu.au)

National Centre for Epidemiology and Population Health, Australian National University

Craig Shadbolt

New South Wales Department of Primary Industries

Patricia M. Desmarchelier

Food Safety Principles

Martyn D. Kirk

National Centre for Epidemiology and Population Health, Australian National University

Kathryn Glass

National Centre for Epidemiology and Population Health, Australian National University

Research Article

Keywords: Salmonella infections, source attribution, foodborne disease, gastroenteritis, disease reservoir, zoonosis, Bayesian analysis

Posted Date: August 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-666044/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on January 4th, 2022. See the published version at <https://doi.org/10.1186/s12879-021-06950-7>.

Abstract

Background

Salmonella is major cause of zoonotic illness around the world, arising from direct or indirect contact with a range of animal reservoirs. In the Australian state of New South Wales (NSW), salmonellosis is believed to be primarily foodborne, but the relative contribution of animal reservoirs is unknown.

Methods

The analysis included 4,543 serotyped isolates from animal reservoirs and 30,073 serotyped isolates from domestically acquired human cases in NSW between January 2008 and August 2019. We used a Bayesian source attribution methodology to estimate the proportion of foodborne *Salmonella* infections attributable to broiler chickens, layer chickens, ruminants, pigs, and an unknown or unsampled source. Additional analyses included covariates for four time periods and five levels of rurality.

Results

A single serotype, *S. Typhimurium*, accounted for 65-75% of included cases during 2008-2014 but <50% during 2017-2019. Attribution to layer chickens was highest during 2008-2010 (48.7%, 95%CrI: 24.2-70.3%) but halved by 2017-2019 (23.1%, 95%CrI: 5.7-38.9%) and was lower in the rural and remote populations than in the majority urban population. The proportion of cases attributed to the unsampled source was 11.3% (95%CrI: 1.2%–22.1%) overall, but higher in rural and remote populations. The proportion of cases attributed to pork increased from approximately 20% in 2009-2016 to approximately 40% in 2017-2019 coinciding with a rise in cases due to *S. ser 4,5,12:i:-*.

Conclusion

Layer chickens were likely the primary reservoir of domestically acquired *Salmonella* infections in NSW circa 2010, but attribution to the source declined contemporaneously with increased vaccination of layer flocks and tighter food safety regulations for the handling of eggs.

Background

Salmonellosis is a common cause of foodborne illness and hospitalisation across Australia. Annual notification rates for salmonellosis in Australia's most populous state, New South Wales (NSW), ranged between 38.1 and 57.9 per 100,000 population during 2009–2019 [1]. However, for every notified case of salmonellosis, there are an estimated seven cases that go unreported [2]. In 2015, there were an estimated 64 thousand annual episodes of foodborne salmonellosis in Australia that resulted in three thousand hospitalisations, and thirteen deaths. The estimated annual societal cost of this illness was AUD 105 million [3–5].

Salmonella can be found in all major livestock and poultry species and a wide array of wild animals; however, the proportion of human *Salmonella* infections attributable to each of these sources is population- and geography-dependant, and often unknown. For instance, in the Netherlands, the primary sources were layers and pigs [6], in Italy and Denmark the primary sources were pigs [7, 8], while in Sweden over 80% of cases were attributed to travel [9]. Source attribution analyses in the Australian states of South Australia and Queensland both identified chickens (broilers and layers) to be the primary sources of illness [10, 11], but also highlighted attribution to nuts or environmental exposures in sub-tropical Queensland [10, 12]. Estimates of the proportion of infections arising from each source can highlight areas needing intervention and inform biosecurity and food safety policy across the entire food supply chain. For instance, source attribution studies in New Zealand in the early 2000s demonstrated that > 50% of *Campylobacter* infections were attributable to poultry meat, prompting targeted improvements to the food safety standards and practices around poultry. These improvements were followed by a 54% decline in incidence. Further source attribution analyses demonstrated the decline was due primarily to reductions in cases attributed to poultry [13]. In NSW, a broad range of interventions were put in place to prevent foodborne salmonellosis following the release of the new NSW Food Safety Strategy 2015–2021. Though incidence has declined in this period [1], the proportion of foodborne salmonellosis attributable to different animal reservoirs before and after this period was unknown.

Salmonella enterica subsp. *enterica*, which accounts for nearly all human cases of salmonellosis, can be classified into over 2500 serotypes [14]. Some serotypes are adapted to specific host species, such as *S. Typhi*, which has humans as the primary host and *S. Gallinarum*, which has poultry as the primary host [15]. Other serotypes can be found in one or more animal hosts and can cause human infections e.g. *S. Typhimurium*. *Salmonella* strains can also be differentiated using a range of typing schemes, including phage typing, multi-locus sequence typing (MLST), multiple-locus variable-number tandem repeat analysis (MLVA), whole genome sequencing and other techniques [14].

Several methods have been developed to attribute illness to specific sources [16, 17] Some rely on linking individual cases or outbreak cases to putative sources through epidemiological investigations, including case-control studies. However, these approaches are labour-intensive and often inconclusive due to small sample sizes. A suitable approach for sporadic cases that does not require food questionnaires is to compare the distribution of *Salmonella* types in isolates from cases to the distribution of types in putative sources. This approach estimates the overall proportion of cases attributable to each source. Hald and colleagues adopted a Bayesian source attribution framework [7], and their model and its variants have since been applied widely [10–12, 18–20]. A related approach accounts for genetic relatedness of strains using the asymmetric island model [20, 21]. Recent advances allow these models to be adjusted for covariates to identify trends or differences by subgroup [20].

We conducted a source attribution analysis for *Salmonella* infections acquired in New South Wales, Australia between January 2008 and August 2019 using variants of the Hald approach. We adjusted these models to account for trends over time and considered differences by age-group, gender, and rurality.

Methods

Human data

In NSW, treating pathology laboratories are required under public health legislation to report all *Salmonella* infections to the health department. We used deidentified data for 40,837 human cases notified between January 2008 and August 2019. The available case data included: typing information for the *Salmonella* isolates, patient gender, five-year age group, location, and travel history. Typing methods included serotyping, MLVA typing, and phage typing; however, not all typing methods were performed for every isolate. Location was encoded at the Statistical Area 2 (SA2) level as defined in the Australian Statistical Geography Standard. As we were primarily interested in cases acquired in NSW, we removed all cases recorded as being acquired outside NSW. However, travel history was known only for a small minority of cases, and before May 2010, cases that were known not to have travelled and cases with unknown travel history were recorded identically. Therefore, we also excluded all cases due to serotypes deemed to be *travel-associated*. A serotype was deemed to be travel-associated if more than half of cases after May 2010 with a known travel history were believed to be acquired outside NSW.

Non-human data

Data for non-human sources were collated from the National Enteric Pathogens Surveillance System (NEPSS) and the New South Wales Food Authority (NSWFA). The two datasets included non-human *Salmonella* isolates sampled from animals, animal products, farm environments (e.g. animal pens and barns), food, and the natural environment. Isolates considered as from potential animal sources were those derived from the faeces, carcass, or enclosures of an animal or from an animal product. Isolates that were not serotyped or were from a travel-associated serotype were excluded. We also excluded isolates without a recorded date or with a date before 2008.

Non-human animal isolates were categorised by reservoir animal source where possible from the recorded details. For example, all isolates from pig carcasses; pig faeces; cooked, cured, or raw pig meats; pig offal; pig intestinal contents; and pig farm environments (e.g. bedding) were categorised as *porcine*. Broiler and layer chickens were treated as separate reservoirs. Chicken meat products for human consumption and environmental samples from broiler farm premises and chicken meat processors were categorised as *broiler*, while isolates from eggs, egg products (e.g. mayonnaise) and egg farm premises were categorised as *layer*. Isolates that could not be definitively assigned to an animal reservoir were excluded, e.g. feather meal which might have been derived from broilers or layers, and fresh produce which might have been contaminated directly or indirectly by any wild or domestic animal. We excluded isolates from food products with multiple animal origins (e.g. 'egg and bacon roll') and isolates with ambiguous

descriptions (e.g. 'meat'). Where possible, putative sources were combined to form categories with at least 100 isolates. Putative sources that could not be reasonably combined into broader categories with at least 100 isolates were excluded.

As source attribution analysis requires the same typing scheme to be applied to both source and human data, and serotyping was the most consistently and completely applied method, analyses were conducted using serotyped data, and isolates without serotyping information were removed from both the human and non-human datasets.

Source attribution modelling framework

We generalised existing Bayesian source attribution methods [19, 20] to estimate changing attribution proportions over time, include covariates for the cases, and adjust for differences between types.

The proportion (θ_{ijst}) of cases in subpopulation s during period t that were due to type i from source j was modelled as:

$$\theta_{ijst} \propto a_{jst} w_j r_{ijt} q_i$$

with constraints $\sum_{i,j} \theta_{ijst} = 1$ and $\sum_i r_{ijt} = 1$ and where a_{jst} was the ability of source j during period t to act as a reservoir of infection for group s , w_j was a weight for the relative exposure of humans to contamination from source j , r_{ijt} was the relative prevalence of type i in source j during period t , and q_i was the relative ability of subtype i to lead to human infection. In each group s and period t the proportion of infections of all types attributed to a source, ξ_{jst} was:

$$\xi_{jst} = \sum_i \theta_{ijst} \propto a_{jst} w_j \sum_i r_{ijt} q_i$$

while the proportion of cases due to each type, μ_{ist} was:

$$\mu_{ist} = \sum_j \theta_{ijst} \propto q_i \sum_j a_{jst} w_j r_{ijt}$$

The estimation of these parameters occurred in two steps. The distribution of types in each source and period (r_{ijt}) were estimated first, and all other parameters were then estimated repeatedly with draws from the posterior distribution of r_{ijt} . In the first step, the number of isolates of each type observed in each source (X_{ijt}) were modelled with multinomial distributions, in one of two ways. For sources with many isolates in every period, the relative frequency of types were modelled independently for each period based only on the data collected in that period, i.e. $X_{.jt} \sim \text{Multinomial}(r_{.jt})$. With a unit Dirichlet prior this resulted in a Dirichlet posterior distribution: $p(r_{.jt} | X) \sim \text{Dirichlet}(1 + X_{.jt})$. For sources with too few samples in each period, the data across the whole study was used for every time period resulting in posterior estimates: $p(r_{.jt} | X) \sim \text{Dirichlet}(1 + \sum_{\tau} X_{.j\tau})$.

In estimating the remaining parameters, the efficiency of each type (q_i) and the exposure weights (w_j) were assumed to remain constant over time but source efficiencies (a_{jst}) were allowed to vary over time and by subgroup of cases. This was modelled as:

$$a_{jst} = \exp\left(\tau_{tj} + \sum_n F_{sn} \beta_{nj}\right).$$

where F was a matrix defining a linear predictor based on categorical, ordinal, or continuous covariates for each subgroup (i.e. category of a covariate) s of the cases; β was a matrix of parameters for each source j ; and τ defined temporal differences in source efficiency by time t and source j . A reference group was assigned to each covariate, and the associated parameters fixed to 0, while the remaining parameters are given unit normal priors.

The number of human cases in subpopulation s during period t that were due to pathogen type i , were modelled as independent multinomial variables for each period t and subgroup s , i.e. $Y_{.st} \sim \text{Multinomial}(\mu_{.st})$. The q_i were constrained with a hierarchical log-normal prior:

$$p(q_i | \sigma) \sim \text{logNormal}(0, \sigma^2) p(\sigma) \sim \text{Uniform}(0, 5).$$

For predominantly foodborne infections the exposure weights w_j can be approximated by the relative level of exposure to contaminated food products derived from each source. However, as this is not measured directly, we modelled this as $w_j = M_j k_j$, where M_j was the per capita consumption of food derived from source j , and k_j was the prevalence of the pathogen in food derived from source j , which was estimated from surveys of animal food products. For each source j , we modelled the number of total tests (N_j) and positive tests (P_j) $P_j \tilde{Binomial}(N_j, k_j)$, with an uninformative uniform prior on prevalence, i.e. $p(k_j) \tilde{Beta}(1, 1)$.

Our model framework was extended to include an 'unsampled source' by including an additional source j^* with no observed samples, i.e. $X_{jj^*t} = P_{j^*} = N_{j^*} = 0$.

Source attribution models

As a base case we considered a model with no covariates and no temporal variation, which was equivalent to the Modified Hald model applied to the whole study period [19]. As a sensitivity analysis we compared this to a model where all the type efficiency terms q_j were fixed at one, which was equivalent to the Dirichlet model of Liao et al. [20] with the addition of exposure weights. We considered each model with and without an 'unsampled source'.

We then considered models with combinations of covariates (age, rurality, and gender) with and without temporal variation. In models adjusting for age, we used age categories: 0–4, 5–19, 20–34, 35–64, and 65 and over. The rurality of each case was determined by matching the Australian Bureau of Statistics 2011 Statistical Area 2 (SA2) code provided for the case to one of the five rurality zones (Major Cities, Inner Regional, Outer Regional, Remote, and Very Remote) defined in the 2011 Australian Statistical Geography Remoteness Structure. However, as rurality was defined at the more granular SA1 level and each SA2 is built from SA1s, a few SA2s contained regions with different categories of rurality. Cases associated with these SA2s were assigned the rurality zone closest to the average rurality of the constituent SA1s. In our main analyses, rurality was modelled as an ordinal variable. In a sensitivity analysis we modelled rurality as a nominal categorical variable, combining the five categories down to three categories ('Major Cities', 'Regional' and 'Remote') to increase the number of cases in each category. The small number of cases that had a missing value for gender, age, or location were excluded only from analyses involving the missing covariate.

In models with temporal variation, we considered four three-year periods: 2008–2010, 2011–2013, 2014–2016, and 2017–2019. As there were > 200 isolates per period from broilers and > 450 isolates per period from layers, the relative frequency of serotypes was estimated independently for each period for broilers and layers. As there were relatively few isolates for ruminants and pigs in at least one period (< 30 for pigs and < 20 for ruminants), data across all study years was used to estimate the relative frequency of serotypes in ruminants and pigs in every period.

To improve model convergence, we excluded serotypes that rarely caused disease in humans (i.e. <10 human cases across the study period) from the analysis.

Exposure by source

Table 1 provides a summary of prevalence assumptions with references, while Table 2 summarises assumptions on food consumption per capita. The datasets used to determine the number of samples positive for each pathogen type (X_{ijt}) did not include data on the number of total tests, so the prevalence of *Salmonella* in food products was estimated from separate surveys, with an adjustment to reduce the number of positive samples by the fraction of cases from excluded serotypes from each source. As age, sex, location, and time-specific consumption and prevalence data were not available for all food sources, exposure weights and prevalence on animal products were assumed fixed across all time periods and the whole population. We used published national data on *Salmonella* from the *E. coli* and *Salmonella* monitoring (ESAM) program from the early 2000s to inform prevalence in ruminants and pork [22, 23], as publicly available contemporary surveys in NSW had insufficient samples to determine prevalence. The relative exposure of the human population to ruminants, pigs, and broilers were based on the apparent consumption of meats from these sources published by the Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES) [24]. For eggs, we assumed consumption of one egg was equivalent to 200g of meat (as in our previous work [11]) and used consumption estimates from a 2018 Australian Eggs annual report [25]. In models with an 'unsampled' source we adopted the conservative assumption that exposure to the unsampled source was equal to the source or group of sources with the highest consumption.

Table 1
Salmonella prevalence assumptions by source with references.

	Prevalence (95% CI)	Prevalence adjusted for types rare in cases (95% CI)	Reference
Chicken meat	48.4% (42.0–54.8)	38.6% (32.5–45.0)	NSW-specific data in Table 4 [49]
Chicken eggs	1.76% (0.70–3.59)	1.44% (0.66–2.72)	Following prior assumptions [11]
Pigs	1.88% (1.57–2.22)	1.14% (0.90–1.41)	National ESAM [†] data [22]
Ruminants	0.38% (0.33–0.43)	0.37% (0.32–0.42)	National ESAM [†] data [23]
† <i>E. coli</i> and <i>Salmonella</i> monitoring program			

Table 2

Relative exposure to potential sources of *Salmonella*, measured by mean consumption of meat and animal products per person per year in Australia.

	Relative Exposure (kg/person/year or equivalent)	Reference
Chicken meat	47	ABARES [†] [24]
Chicken eggs	49 [‡]	Australian eggs [25]
Pork	28	ABARES [†] [24]
Ruminants	34	ABARES [†] [24]
† Australian Bureau of Agricultural and Resource Economics and Sciences		
‡ Mean consumption was approximately 245 eggs per person per year in the 2017-18 financial year. The relative exposure was calculated assuming that one egg is equivalent to 200g of meat.		

Software

All analyses were conducted in the R programming environment [26]. Bayesian inference was performed using the No U-turn Algorithm with the Stan programming language [27] and the rstan R package [28]. Data cleaning and manipulation was done using the plyr [29], dplyr [30], and tidyr [31] R packages. Data visualisations were made with the ggplot2 R package [32].

Results

Data

After excluding isolates from humans with no serotype information (3,807 isolates, 9.3%), isolates from cases recorded as having travelled outside NSW (2,381 isolates, 5.8%), isolates of *S. Paratyphi* B bv Java and from serotypes that were commonly travel associated (4,089 isolates, 10.0%), and isolates of serotypes that were rare in humans (487 isolates, 1.2%), a total of 30,073 human isolates remained for source attribution analysis (Table 3). Most of the cases associated with these isolates were reported in major cities, with 51% of cases in females. After isolates from some animals were grouped and others were omitted (e.g. wildlife, horses, ducks, quail, and turkeys), the final source groups were broilers (N = 1,396), layers (N = 2,321), pigs (391), and ruminants (cattle, sheep, and goats; N = 435) (Table 4). The number of non-human isolates in our dataset was lowest in 2017–2019.

Table 3

Characteristics of *Salmonella* cases reported between January 2008 and August 2019 in New South Wales after excluding cases without known serotype (N = 3,807) travel associated cases (N = 6,470) and rare serotypes (N = 487).

	<i>S. Typhimurium</i> (N = 18,802)	Other serotypes (N = 11,271)	All serotypes (N = 30,073)	p value
Gender				0.010
Missing	47	22	69	
Female	9,680 (51.6%)	5,633 (50.1%)	15,313 (51.0%)	
Male	9,075 (48.4%)	5,616 (49.9%)	14,691 (49.0%)	
Rurality				< 0.001
Missing	142	48	190	
Major Cities	14,106 (75.6%)	7,521 (67.0%)	21,627 (72.4%)	
Inner Regional	3,434 (18.4%)	2,784 (24.8%)	6,218 (20.8%)	
Outer Regional	1,070 (5.7%)	844 (7.5%)	1,914 (6.4%)	
Remote	30 (0.2%)	48 (0.4%)	78 (0.3%)	
Very Remote	20 (0.1%)	26 (0.2%)	46 (0.2%)	
Age group				< 0.001
Missing	17	6	23	
00–04	4,191 (22.3%)	3,045 (27.0%)	7,236 (24.1%)	
05–19	4,186 (22.3%)	1,563 (13.9%)	5,749 (19.1%)	
20–39	4,986 (26.5%)	2,287 (20.3%)	7,273 (24.2%)	
40–64	3,434 (18.3%)	2,445 (21.7%)	5,879 (19.6%)	
65+	1,988 (10.6%)	1,925 (17.1%)	3,913 (13.0%)	
Year group				< 0.001
2008-10	4,780 (25.4%)	2,059 (18.3%)	6,839 (22.7%)	
2011-13	5,441 (28.9%)	2,407 (21.4%)	7,848 (26.1%)	
2014-16	5,861 (31.2%)	3,457 (30.7%)	9,318 (31.0%)	
2017-19	2,720 (14.5%)	3,348 (29.7%)	6,068 (20.2%)	

Table 4

The number of serotyped *Salmonella* isolates from humans, selected non-human sources sampled in NSW between January 2008 and June 2019. These counts exclude *S. Paratyphi B* by Java, types that were rare in humans (less than 10 cases over the period), and travel-associated types (types where more than half of cases with known travel history were acquired outside NSW). A selection of the most common excluded sources are displayed here to illustrate the sparsity of data over time.

Source	Year												Total
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
Human cases	1,759	2,101	2,979	2,864	2,275	2,709	3,416	2,818	3,084	2,489	2,063	1,516	30,073
Ruminants	58	60	49	87	60	33	25	42	10	6	2	3	435
Cattle	50	53	40	70	53	31	22	38	9	6	2	3	377
Ruminants – Other ¹	8	7	9	17	7	2	3	4	1	-	-	-	58
Broilers ²	152	99	98	76	117	37	61	360	185	84	56	71	1,396
Layers ²	288	160	155	270	204	251	225	158	109	136	128	237	2,321
Pigs	20	13	42	47	45	79	23	48	51	5	14	4	391
Total Included Non-Human ³	518	332	344	480	426	400	334	608	355	231	200	315	4,543
Wildlife ⁴	12	16	1	3	22	13	5	5	3	3	-	-	83
Poultry – Other ⁵	13	7	1	9	1	4	11	1	1	-	-	-	48
Horses	2	3	4	7	1	0	4	8	3	-	-	-	32
¹ Ruminant-other includes sheep and goats													
² Other isolates from poultry that could not be linked to either broilers or layers are not included in this table													
³ The included sources are ruminants, broilers, layer, and pigs													
⁴ Wildlife consists of a diverse group of non-captive non-domesticated animals													
⁵ Poultry-other consists of ducks, turkey and quail													

Type distribution in sources and humans

S. Typhimurium was the most common serotype in cases and every major source group except for broilers (Figure S1, Table S1). In broilers, *S. Typhimurium* was only the third most common serotype (10.1%, 141/1396) after *S. Infantis* (30.4%, 425/1396) and *S. Sofia* (20.6%, 287/1396) (Figure S1). The proportion of human cases due to *S. Typhimurium* was largest in major cities (65.2%), but this proportion declined with increasing rurality (Inner Regional 55.2%; Outer Regional 55.9%; Remote 38.5%; Very Remote 43%) (Table 3). Conversely, cases due to serotypes such as *S. Chester* and *S. Saintpaul* were less common in major cities than in regional and remote NSW. The proportion of human cases due to *S. Typhimurium* declined from 65–75% during 2008–2013 to < 50% during 2017–2019. The proportion of cases due to some serotypes rose over the same period, e.g. *S. Wangata* (2008–2010: 1.0%; 2017–2019: 9.2%), and *S. enterica* ser. 4,5,12:i:- (2008–2010: 0.0%; 2017–2019: 5.5%). The proportion of layer isolates that were serotyped as *S. Typhimurium* also declined over the same period (2008–2010: 41.6%, 251/603; 2017–2019: 17.4%, 87/501).

Subtyping

After restricting the analysis to serotyped isolates, only a small minority of non-*S. Typhimurium* isolates were also typed using phage or MLVA typing (human: 12.6%; reservoirs: 4.5%). While a larger proportion of *S. Typhimurium* isolates were typed using a least one additional method (human: 95.4%; reservoirs: 73.1%), this additional typing method was not consistent over time or between cases and reservoirs. Nearly all *S. Typhimurium* isolates from humans were phage typed prior to 2010, but < 7% were phage typed after 2010 (Table S1). Overall, fewer *S. Typhimurium* isolates from humans were phage typed (25.1%) than in the major source groups (58.2–84.4%). In contrast, while 93% of *S. Typhimurium* isolates from cases were MLVA-typed, less than 35% of *S. Typhimurium* isolates in each major source group were MLVA-typed and coverage varied by year and source. We were therefore unable to use phage typing or MLVA data in our source attribution analyses.

Source attribution without covariates

Table 5 summarises the attribution proportion for models without covariates or temporal variation. All models indicated layers as the leading source of illness. If we assumed all serotypes to be equally efficient at causing disease in humans, very few human cases were attributed to either broilers or pigs: 0.02% and 1.8% respectively. When allowing for differences between serotypes, a greater proportion of cases were attributed to broilers and pigs: 13.1% and 17.8% respectively. Equal- q and variable- q models with adjustments for unsampled sources attributed 5.8% and 11.3% to the unsampled source respectively and attributed fewer cases to ruminants and pigs than the models without adjustment for an unsampled source. The HaldDP model of Miller et al. [18] failed to converge with our dataset and is not shown.

Table 5

The percentage of human isolates attributed to ruminants, broilers, layers, pigs, and other unsampled sources with 95% credible intervals for models without covariates or temporal variation. In the equal- q model all *Salmonella* serotypes are assumed to be equally efficient in their ability to cause infection in humans, while in the variable- q model (equivalent to the modified Hald model [19]) serotypes are allowed to differ in their efficiency.

	Ruminants	Broilers	Layers	Pigs	Unsampled
Equal- q					
without unsampled source	41.3 (21.0–58.7)	0.02 (0.0–0.05)	56.9 (39.1–74.2)	1.8 (0.0–8.3)	-
with unsampled source	36.0 (16.9–55.5)	0.01 (0.0–0.03)	57.7 (39.7–74.0)	0.4 (0.0–3.1)	5.8 (0.0–10.7)
Variable- q					
without unsampled source	24.7 (0.0–55.8)	13.1 (0.0–44.9)	44.3 (9.8–76.7)	17.8 (0.0–39.1)	-
with unsampled source	10.5 (0.0–34.8)	18.2 (0.0–66.1)	48.4 (5.4–79.6)	12.1 (0.0–31.8)	11.3 (1.2–22.1)

Source attribution with covariates

In a temporal model without covariates, the proportion attributed to layers declined from 55% (95% CrI: 23–71%) in 2008–2010 to 34% (95% CrI: 16–51%) in 2014–2016 and 30% (95% CrI: 18–42%) in 2017–2019, with an increase in the proportion attributed to pigs in 2017–2019 and increase in attribution to ruminants in 2014–2016 (Fig. 1). When rurality was included in the model, a higher proportion of cases were attributed to layers in major cities, with this proportion declining with increasing rurality (Fig. 2). A similar rural-urban gradient was observed in a non-temporal model, although the proportion attributed to broiler chickens in that model (Figure S2) was substantially higher than all other models, potentially due to confounding with year ($p < 0.001$ for association between year and rurality).

There was weaker evidence of a rural-urban gradient in attribution to ruminants, with a higher proportion of infections attributed to ruminants in rural areas (Figs. 2 and S2), which disappeared in models with an unsampled source (Figures S3 and S4). In a sensitivity analysis where rurality was modelled as categorical variable with three levels ('Major Cities', 'Regional' and 'Remote') all findings remained qualitatively similar (results not shown).

A non-temporal univariable regression analysis found an association between attribution proportion and sex, with 5.7% (95% CrI: 1.3%–10.1%) fewer male cases attributed to layer chickens. Though a qualitatively similar sex difference was also observed in a

model including time and sex, and a model with age and sex, the sex differences were not statistically significant within any time period or age group. There were no consistent associations between age group and attribution proportions.

Estimates of the type-specific parameters, q , varied 1000-fold between the most efficient and least efficient serotypes. For instance, in the model including rurality, year-group, and an unsampled source, *S. Typhimurium* and *S. Abortusovis* were estimated to be 22.3 (95% CrI: 14.0–38.0) and 0.19 (95% CrI: 0.07–0.46) times as efficient as *S. Infantis*, while *S. Sofia* was estimated about as efficient as *S. Infantis* (q ratio 0.5; 95% CrI: 0.2–1.7) (Figure S5).

Discussion

We estimate that approximately half of all notified cases of non-typhoidal salmonellosis acquired in New South Wales during 2008–2019 were due to direct or indirect transmission from layer chickens. However, the proportion of cases attributed to layers declined from approximately 50% in 2009–2011 to approximately 25% in 2017–2019. The proportion of cases attributed to pork increased from approximately 20% in 2009–2016 to approximately 40% in 2017–2019, coinciding with a rise in cases due to *Salmonella enterica* ser 4,5,12:i:-, which has been found to be a persistent coloniser of pigs in Australia [33]. The proportion of cases attributed to layers was lower amongst rural-dwelling than urban-dwelling populations, similar to the rural-urban gradient found in a source attribution analysis of *Campylobacter* infections in New Zealand [20, 34].

The potential of zoonotic organism to lead to a notified case of foodborne disease depends on many factors including the organism's ability to survive transport, storage, and food preparation and its virulence in humans. Some source attribution modelling approaches (e.g. those applied to *Campylobacter* infections [20, 21]) have assumed that all types are equally able to lead to notified human cases. However, this assumption is inappropriate for *Salmonella*, which is known to have both virulent (e.g. *S. Typhimurium*) and avirulent (e.g. *S. Sofia* [35]) serotypes. In our study, *S. Typhimurium* was more prevalent in cases (63%) than in broilers (10%), layers (32%), pigs (23%), or ruminants (35%), while *S. Infantis* was less prevalent in cases (3%) than in broilers (30%), layers, (19%) or pigs (7%), indicating that *S. Typhimurium* is a more efficient foodborne pathogen than *S. Infantis*. Consequently, models that assumed all serotypes were equally efficient attributed < 0.1% of cases to broilers (in which *S. Sofia* and *S. Infantis* were most common). However, this is inconsistent with the common implication of chicken meat in outbreak investigations and case control studies (e.g. [36]). Although models that account for differences between serotypes led to source attribution estimates with much wider confidence intervals, we believe they are more appropriate for modelling source attribution of *Salmonella*.

The decline in attribution to layers from 2014 onwards may be the result of interventions including vaccination of layers against *S. Typhimurium*. The NSW Food Safety Strategy 2015–2021, released in 2014, including a specified 30% reduction target for foodborne salmonellosis [37]. Prior to 2014, many reported foodborne outbreaks in NSW were linked to poor handling of eggs and hygiene at retail food service (e.g. [38, 39]). Several measures were implemented to address this, including mandatory education and food safety training on the risks of raw egg use, improved cleaning and sanitising for retail businesses, guidelines for the safe use of eggs which made it an offence in NSW to prepare raw egg foods without adequate processing, and training and guidance for local council Environmental Health Officers to focus on areas of highest food safety risk during inspections. Since 2014, large egg producers across NSW have begun largescale vaccination of layer flocks against *S. Typhimurium*; estimates based on sales of vaccines indicate approximately 75% of commercial layer flocks were vaccinated in 2017–2019 (Dr. Christopher Morrow, personal communication, April 2021). The introduction of all these measures coincided with a marked decline in *S. Typhimurium* and total *Salmonella* incidence in NSW [40].

The models that included an 'unsampled' source attempted to quantify the plausible proportion of cases that could be attributed to any animal reservoirs other than domestic chickens, ruminants, and pigs. Since this approach necessarily drew inferences without information about of the relative abundance of different serotypes in the 'unsampled' animal reservoirs, these models need to be interpreted cautiously. In our study, the increasing trend in attribution to 'unsampled' sources is associated with the increasing incidence of serotypes that were absent or rare in the major food animals included in our analyses, e.g. *S. Wangata*, *S. Birkenhead*, *S. Waycross*, and *S. Hvittingfoss*. *S. Wangata* was not found in ruminants or pigs in our dataset and was very rare in broilers and layers (< 0.6%) but accounted for 9.0% of cases in 2017–2019 up from only 1.0% in 2008–2010. Similarly, though *S. Birkenhead* was only represented by a single isolate from layer chickens in 2019, the serotype accounted for 5.7% of cases in 2017–2019, up from 3.3% in 2008–2010. While *S. Wangata* is associated with contact with a range of wild animals in NSW [41–43], the animal reservoirs of *S. Birkenhead*, *S. Waycross* and *S. Hvittingfoss* are unclear. A source attribution analysis of *Salmonella* infections in the neighbouring

state of Queensland linked *S. Birkenhead*, *S. Waycross*, and *S. Hvittingfoss* to nuts (primarily locally grown macadamias) [10] suggesting environmental reservoirs of exposure. In north-eastern NSW, *S. Birkenhead* infection has been associated with not usually washing or peeling fruit and vegetables before eating raw. *S. Hvittingfoss* has caused outbreaks linked to contaminated cantaloupes in Australia with closely related isolates identified in bar-tailed godwits in north-western Australia, suggesting a common exposure to an unidentified animal or environmental reservoir [44]. *S. Hvittingfoss* has also been found in reptiles [45, 46], migratory ducks [47], and feral pigs [48] elsewhere in Australia. Consequently, a substantial part of the attribution to ‘unsampled’ sources in our model may be due to transmission from wild or feral animals through direct contact or indirect contamination of water, food, or the environment.

The present study has a number of strengths. The large dataset of *Salmonella* cases and a substantial dataset of isolates from major food animals, collected from across the state and over a long time period, have allowed us to identify the leading source of *Salmonella* infections, attribute separately to egg-laying and meat chickens, and identify temporal and rural-urban gradients in attribution. Furthermore, these key model outcomes were qualitatively robust across multiple model comparisons. In the source attribution framework we adopted, attribution is to animal reservoirs rather than routes or vehicles of transmission. For instance, our estimate of the proportion of cases attributed to layers includes infections that may have been acquired through direct contact with layer chickens, consumption of contaminated eggs, contact with manure, or consumption of water or food contaminated by layer chicken faeces at any point between farm and table. Attribution to animal reservoirs is simultaneously a strength – since all possible transmission routes are captured – and a limitation – since the relative contribution of different routes cannot be quantified.

Our study has a number of limitations. No high-resolution molecular subtyping scheme was consistently applied to *Salmonella* isolates from humans and animals over the whole study period, forcing us to restrict our analyses to the level of serotype. Because the majority of human cases were due to a single serotype, *S. Typhimurium*, this restriction substantially weakened inferences, leading to wide confidence intervals for many attribution estimates. Though the dataset of reported human cases provided more than adequate sample size, it is estimated that as few as one in eight *Salmonella* cases are reported [2]. As unreported cases are likely to have milder symptoms than reported cases, more virulent serotypes of *Salmonella* are likely to be over-represented in our dataset and the attribution proportions are therefore likely indicative of moderate and severe cases. Travel history was not available for all cases in our dataset, and though we took steps to remove cases not acquired in NSW from the study we may have removed some locally acquired cases in doing so. The isolates from animal reservoirs were not collected in a single sampling frame. The majority of non-human *Salmonella* sampling by the NSW Food Authority occurred through investigation of foodborne disease outbreaks and was often sporadic in nature. Some targeted survey work was also undertaken, but this was biased towards testing of layer flocks, or opportunistic *Salmonella* testing through poultry meat verification work focused on *Campylobacter*. To date, NSWFA have not conducted comparable ongoing surveys or studies focused on ruminants, pigs, or other potential reservoirs. The remaining isolates from non-human sources collected by the NEPSS included any isolate submitted to *Salmonella* typing laboratories in Australia and included public, private, and academic sampling efforts. For isolates from non-human sources, we only considered those collected in NSW; however, NSW residents consume food produced across Australia. Though lamb and mutton account for approximately 8% of meat consumption in Australia [24], very few *Salmonella* isolates from sheep isolates were available, forcing us to combine sheep, cattle, and goats into a single category for source attribution purposes. However, the relatively low consumption of sheep meat and the low prevalence of *Salmonella* in sheep [23] suggest that sheep are unlikely to be a major source of *Salmonella* infections in NSW. We only had sufficient samples to estimate the relative frequency of serotypes over time in poultry which may have influenced the estimated trend in source attribution. Finally, we had only limited samples from wild, feral, and companion animals, which may have been especially important for the accurate attribution of rural cases. Though attribution to ‘unsampled’ sources includes these reservoirs, we are unable to estimate the contribution different species or groups of species.

Conclusions

Layer chickens were the primary reservoir for human salmonellosis in NSW during 2009–2019; however, the importance of layers was less in rural populations and declined after 2015, concurrent with changes to food safety regulations and egg industry practices. The apparent increase in attribution to pigs and unsampled sources in 2017–19 warrants further investigation. Our study highlights the need for a high-resolution typing method to be consistently applied to *Salmonella* isolates collected from humans, food, animals, and the environment. Using a consistent approach would allow for more precise estimates for source attribution

analyses and assist in identifying specific sources in outbreak investigations. Furthermore, ongoing routine surveillance of *Salmonella* in food and food animals collected in a consistent sampling frame would improve the sample size and data quality for prevalence and source attribution estimates and allow these estimates to be updated regularly to monitor trends. Moreover, the mobility of people and food products between Australian states and territories calls for a nationally unified approach to the surveillance of *Salmonella*.

Abbreviations

NSW

New South Wales; CrI:Credible Interval; AUD:Australian Dollars; MLST:multi-locus sequence typing; MLVA:multiple-locus variable-number tandem repeat analysis; SA1:Australian Bureau of Statistics 2011 Statistical Area 1; SA2:Australian Bureau of Statistics 2011 Statistical Area 2; NEPSS:National Enteric Pathogens Surveillance System; NSWFA:New South Wales Food Authority; ESAM:*E. coli* and *Salmonella* Monitoring Program; ABARES:Australian Bureau of Agricultural and Resource Economics and Sciences

Declarations

Ethics approval and consent to participate

Ethics approval was provided by the Australian National University Human Research Ethics Committee (Protocol: 2019/470)

Consent for publication

Not applicable

Availability of data and material

The data used in this study were provided by New South Wales Department of Primary Industries, National Enteric Pathogens Surveillance Scheme, and Health Protection New South Wales; however, restrictions apply to the availability of these data, which were used under agreement for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of the relevant third parties.

Competing interests

The authors declare that they have no competing interests

Funding

This study was primarily funded by the New South Wales Department of Primary Industries, which also collected a subset of the non-human data and employs C.S. A.M. is supported through an ARC Discovery Project grant DP180100246 and M.D.K. is supported by an Australian National Health and Medical Research Council Fellowship (GNT1145997).

Authors' contributions

C.S., M.D.K., K.G., and conceived the study. A.M., C.S., and K.G. designed the study. A.M., P.M.D., and C.S. collated the data. A.M. wrote the R code, performed the analysis, and drafted the manuscript. All authors contributed to the interpretation of the results, revised the manuscript, and approved the final manuscript.

Acknowledgements

We would like to thank Health Protection New South Wales, Alison Imlay from the New South Wales Department of Primary Industries (NSWDPI) and the Mathilda Wilmot and Courtney Lane National Enteric Pathogen Surveillance System (NEPSS) hosted at the University of Melbourne (UoM) for providing access to the datasets used in this study. In particular we would like to thank Keira Glasgow and Neil Franklin (Health Protection NSW); Alison Imlay and Themy Saputra (NSWDPI); and Mathilda Wilmot and Courtney Lane (NEPSS/UoM). We would like to acknowledge Declan O'Donovan for assistance with data cleaning, and Dr Thea King from the New South Wales Department of Primary Industries for suggestions around *Salmonella* enterica ser 4,5,12:i:-in pigs and potential reservoirs of *S. Hvitvingfoss* and *S. Birkenhead*.

References

1. **National Notifiable Diseases Surveillance System.** Australian Government Department of Health. <http://www9.health.gov.au/cda/source/cda-index.cfm>. 2020/11/05.
2. Hall G, Yohannes K, Raupach J, Becker N, Kirk M: **Estimating Community Incidence of *Salmonella*, *Campylobacter*, and Shiga Toxin-producing *Escherichia coli* Infections, Australia.** *Emerging Infectious Diseases* 2008, **14**(10):1601–1609.
3. Kirk M, Ford L, Glass K, Hall G: **Foodborne Illness, Australia, Circa 2000 and Circa 2010.** *Emerging Infectious Diseases* 2014, **20**(11):1857–1864.
4. Ford L, Haywood P, Kirk MD, Lancsar E, Williamson DA, Glass K: **Cost of *Salmonella* Infections in Australia, 2015.** *Journal of Food Protection* 2019, **82**(9):1607–1614.
5. Ford L, Kirk M, Glass K, Hall G: **Sequelae of Foodborne Illness Caused by 5 Pathogens, Australia, Circa 2010.** *Emerging Infectious Diseases* 2014, **20**(11):1865–1871.
6. Mughini-Gras L, Enserink R, Friesema I, Heck M, Van Duynhoven Y, Van Pelt W: **Risk Factors for Human Salmonellosis Originating from Pigs, Cattle, Broiler Chickens and Egg Laying Hens: A Combined Case-Control and Source Attribution Analysis.** *PLoS ONE* 2014, **9**(2):e87933.
7. Hald T, Vose D, Wegener HC, Koupeev T: **A Bayesian Approach to Quantify the Contribution of Animal-Food Sources to Human Salmonellosis.** *Risk Analysis* 2004, **24**(1):255–269.
8. Mughini-Gras L, Barrucci F, Smid JH, Graziani C, Luzzi I, Ricci A, Barco L, Rosmini R, Havelaar AH, Van Pelt W *et al.*: **Attribution of human *Salmonella* infections to animal and food sources in Italy (2002–2010): adaptations of the Dutch and modified Hald source attribution models.** *Epidemiology and Infection* 2014, **142**(5):1070–1082.
9. Wahlström H, Andersson Y, Plym-Forshell L, Pires SM: **Source attribution of human *Salmonella* cases in Sweden.** *Epidemiology and Infection* 2011, **139**(8):1246–1253.
10. Fearnley EJ, Lal A, Bates J, Stafford R, Kirk MD, Glass K: ***Salmonella* source attribution in a subtropical state of Australia: capturing environmental reservoirs of infection.** *Epidemiology and Infection* 2018, **146**(15):1903–1908.
11. Glass K, Fearnley E, Hocking H, Raupach J, Veitch M, Ford L, Kirk MD: **Bayesian Source Attribution of Salmonellosis in South Australia.** *Risk Analysis* 2016, **36**(3):561–570.
12. Munck N, Smith J, Bates J, Glass K, Hald T, Kirk MD: **Source Attribution of *Salmonella* in Macadamia Nuts to Animal and Environmental Reservoirs in Queensland, Australia.** *Foodborne Pathogens and Disease* 2020, **17**(5):357–364.
13. Sears A, Baker MG, Wilson N, Marshall J, Muellner P, Campbell DM, Lake RJ, French NP: **Marked Campylobacteriosis Decline after Interventions Aimed at Poultry, New Zealand.** *Emerging Infectious Diseases* 2011, **17**(6):1007–1015.
14. Wattiau P, Boland C, Bertrand S: **Methodologies for *Salmonella enterica* subsp. *enterica* Subtyping: Gold Standards and Alternatives.** *Applied and Environmental Microbiology* 2011, **77**(22):7877–7885.
15. Thomson NR, Clayton DJ, Windhorst D, Vernikos G, Davidson S, Churcher C, Quail MA, Stevens M, Jones MA, Watson M *et al.*: **Comparative genome analysis of *Salmonella* Enteritidis PT4 and *Salmonella* Gallinarum 287/91 provides insights into evolutionary and host adaptation pathways.** *Genome Research* 2008, **18**(10):1624–1637.
16. Mughini-Gras L, Kooh P, Augustin J-C, David J, Fravallo P, Guillier L, Jourdan-Da-Silva N, Thébault A, Sanaa M, Watier L: **Source Attribution of Foodborne Diseases: Potentialities, Hurdles, and Future Expectations.** *Frontiers in Microbiology* 2018, **9**.
17. Pires SM, Vieira AR, Hald T, Cole D: **Source Attribution of Human Salmonellosis: An Overview of Methods and Estimates.** *Foodborne Pathogens and Disease* 2014, **11**(9):667–676.

18. Miller P, Marshall J, French N, Jewell C: **sourceR: Classification and source attribution of infectious agents among heterogeneous populations.** *PLOS Computational Biology* 2017, **13**(5):e1005564.
19. Mullner P, Jones G, Noble A, Spencer SEF, Hathaway S, French NP: **Source Attribution of Food-Borne Zoonoses in New Zealand: A Modified Hald Model.** *Risk Analysis* 2009, **29**(7):970–984.
20. Liao S-J, Marshall J, Hazelton ML, French NP: **Extending statistical models for source attribution of zoonotic diseases: a study of campylobacteriosis.** *Journal of The Royal Society Interface* 2019, **16**(150):20180534.
21. Wilson DJ, Gabriel E, Leatherbarrow AJH, Cheesbrough J, Gee S, Bolton E, Fox A, Fearnhead P, Hart CA, Diggle PJ: **Tracing the Source of Campylobacteriosis.** *PLoS Genetics* 2008, **4**(9):e1000203.
22. Hamilton DR, Smith P, Pointon A: **National *Salmonella* and *E. coli* Monitoring (ESAM) data from Australian pig carcasses from 2000 to 2006.** In: *Seventh international symposium on the epidemiology and control of foodborne pathogens in pork: 2007; Verona, Italy.* Iowa State University Digital Press; 2007: 129–132.
23. Jordan D, Morris S: **Analysis of ESAM data.** Sydney: Meat & Livestock Australia Limited; 2006.
24. **Agricultural commodity statistics.** Australian Bureau of Agricultural and Resource Economics and Sciences. <https://www.agriculture.gov.au/abares/research-topics/agricultural-outlook/data>. 2020/03/07.
25. **Australian eggs annual report 2017-18:** Australian Eggs; 2018.
26. R Core Team: **R: A Language and Environment for Statistical Computing** Vienna, Austria: R Foundation for Statistical Computing; 2020. <https://www.R-project.org/>
27. Stan Development Team: **Stan Modeling Language Users Guide and Reference Manual, 2.25.** 2020.
28. Stan Development Team: **RStan: the R interface to Stan.** 2020.
29. Wickham H: **The Split-Apply-Combine Strategy for Data Analysis.** *Journal of Statistical Software* 2011, **40**(1):1–29.
30. Wickham H, François R, Henry L, Müller K: **dplyr: A Grammar of Data Manipulation;** 2020. <https://CRAN.R-project.org/package=dplyr>
31. Wickham H: **tidyr: Tidy Messy Data;** 2020. <https://CRAN.R-project.org/package=tidyr>
32. Wickham H: **ggplot2: Elegant Graphics for Data Analysis:** Springer-Verlag New York; 2016.
33. Weaver T, Valcanis M, Mercoulia K, Sait M, Tuke J, Kiermeier A, Hogg G, Pointon A, Hamilton D, Billman-Jacobe H: **Longitudinal study of *Salmonella* 1,4,[5],12:i:- shedding in five Australian pig herds.** *Preventive Veterinary Medicine* 2017, **136**:19–28.
34. Lake RJ, Campbell DM, Hathaway SC, Ashmore E, Cressey PJ, Horn BJ, Pirikahu S, Sherwood JM, Baker MG, Shoemack P *et al.*: **Source attributed case-control study of campylobacteriosis in New Zealand.** *International Journal of Infectious Diseases* 2020.
35. Gan E, Baird FJ, Coloe PJ, Smooker PM: **Phenotypic and molecular characterization of *Salmonella enterica* serovar Sofia, an avirulent species in Australian poultry.** *Microbiology* 2011, **157**(4):1056–1065.
36. McPherson ME, Fielding JE, Telfer B, Stephens N, Combs BG, Rice BA, Fitzsimmons GJ, Gregory JE: **A multi-jurisdiction outbreak of *Salmonella* Typhimurium phage type 135 associated with purchasing chicken meat from a supermarket chain.** *Communicable Diseases Intelligence* 2006, **30**(4):449–455.
37. New South Wales Food Authority: **New South Wales Government food safety strategy 2015–2021: Safe food from paddock to plate.** Sydney: New South Wales Food Authority; 2014.
38. Ward K, Franklin N, Furlong C, Polkinghorne BG, Flint J: **New South Wales OzFoodNet Annual Report: 2013.** Sydney: Health Protection NSW; 2014.
39. New South Wales Health Communicable Diseases Branch: **New South Wales OzFoodNet Annual Report: 2012.** Sydney: Health Protection NSW; 2013.
40. New South Wales Health Communicable Diseases Branch: **New South Wales OzFoodNet Surveillance Report: 2018.** Sydney: Health Protection NSW; 2019.
41. Collins J, Simpson KMJ, Bell G, Durrheim DN, Hill-Cawthorne GA, Hope K, Howard P, Kohlenberg T, Lawrence K, Lilly K *et al.*: **A One Health investigation of *Salmonella enterica* serovar Wangata in north-eastern New South Wales, Australia, 2016–2017.** *Epidemiology and Infection* 2019, **147**.
42. Simpson KMJ, Hill-Cawthorne GA, Ward MP, Mor SM: **Diversity of *Salmonella* serotypes from humans, food, domestic animals and wildlife in New South Wales, Australia.** *BMC Infectious Diseases* 2018, **18**(1).

43. Simpson KMJ, Mor SM, Ward MP, Walsh MG: **Divergent geography of *Salmonella* Wangata and *Salmonella* Typhimurium epidemiology in New South Wales, Australia.** *One Health* 2019, **7**:100092.
44. Smith HG, Bean DC, Hawkey J, Clarke RH, Loyn R, Larkins J-A, Hassell C, Valcanis M, Pitchers W, Greenhill AR: ***Salmonella* enterica Serovar Hvittingfoss in Bar-Tailed Godwits (*Limosa lapponica*) from Roebuck Bay, Northwestern Australia.** *Applied and Environmental Microbiology* 2020, **86**(19).
45. Scheelings TF, Lightfoot D, Holz P: **Prevalence of *Salmonella* in Australian reptiles.** *Journal of Wildlife Diseases* 2011, **47**(1):1–11.
46. Iveson JB, Mackay-Scollay EM, Bamford V: ***Salmonella* and *Arizona* in reptiles and man in Western Australia.** *Journal of Hygiene* 1969, **67**(1):135–145.
47. Hoque MA, Burgess GW, Greenhil AR, Hedlefs R, Skerratt LF: **Causes of Morbidity and Mortality of Wild Aquatic Birds at Billabong Sanctuary, Townsville, North Queensland, Australia.** *Avian Diseases* 2012, **56**(1):249–256.
48. Ward MP, Cowled BD, Galea F, Garner MG, Laffan SW, Marsh I, Negus K, Sarre SD, Woolnough AP: ***Salmonella* infection in a remote, isolated wild pig population.** *Veterinary Microbiology* 2013, **162**(2–4):921–929.
49. **Baseline survey on the prevalence and concentration of *Salmonella* and *Campylobacter* in chicken meat on-farm and at primary processing;** 2010.

Figures




Figure 1

(A) Attribution proportion with 95% credible intervals for each of the major source groups for three-year periods and (B) change over time in the attribution proportion (in percentage points) with 2008-2010 as the reference. Dots indicate posterior mean, while dark and faint lines indicate 80% and 95% credible intervals respectively. The dashed horizontal line indicates no difference.




Figure 2

(A) Attribution proportion for each of the major source groups for cases residing in different rurality zones over time. (B) The difference in attribution proportion by rurality with residents of major cities as the reference. Dots indicate posterior mean, while dark and faint lines indicate 80% and 95% credible intervals respectively. The dashed horizontal line indicates no difference.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [NSWSalmonellaSourceAttributionSuppMaterialsBMC.docx](#)