

# Clinical Implications of Oral Assessment in Elderly Patients with Acute Heart Failure: A Single-center, Retrospective Study

Yusuke Uemura (✉ [yusuke0307@kosei.anjo.aichi.jp](mailto:yusuke0307@kosei.anjo.aichi.jp))

Anjo Kosei Hospital <https://orcid.org/0000-0002-2262-1452>

**Rei Shibata**

Nagoya University Graduate School of Medicine

**Haruna Ishikawa**

Anjo Kosei Hospital

**Ayumi Nagahori**

Anjo Kosei Hospital

**Yuta Katsumi**

Anjo Kosei Hospital

**Kenji Takemoto**

Anjo Kosei Hospital

**Shinji Ishikawa**

Anjo Kosei Hospital

**Toyoaki Murohara**

Nagoya University Graduate School of Medicine

**Masato Watarai**

Anjo Kosei Hospital

---

## Research article

**Keywords:** oral assessment, acute heart failure, elderly patients, activities of daily living

**Posted Date:** November 4th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-66811/v2>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

Oral health problems are common and are associated with various geriatric conditions in older adults. The importance of oral health has not been fully highlighted in the assessment and management of patients with heart failure. Here, we investigated the clinical implications of oral assessment in elderly patients with acute heart failure.

We evaluated oral health using the revised oral assessment guide in 77 patients aged 65 years or older who were admitted to hospital for acute heart failure. Poor oral health (defined as a revised oral assessment guide score  $\geq 9$ ) was identified in 66.2% of the patients. Patients with poor oral health had high prevalence of decreased physical function, undernutrition, and cognitive impairment. A reduction in the Barthel Index, as an indicator of activities of daily living during hospitalization, was significant in the enrolled patients. The Barthel Index decreased more in patients with poor oral health than those with normal oral health. Furthermore, the revised oral assessment guide score on admission was found to be the only independent predictor of changes in the Barthel Index during hospitalization in the multivariate regression analyses.

In conclusion, oral assessment using the revised oral assessment guide during hospitalization could provide useful information for the management of elderly heart failure patients.

## Introduction

The incidence and prevalence of heart failure (HF) increase with age [1, 2, 3, 4]. HF is the most common cause of hospitalization in patients aged 65 years or older. Hospitalization for HF is a significant public health problem, especially in elderly patients, because it is associated with higher rates of mortality, repeated re-hospitalization, and decline in physical activity, compared to that in younger patients [5, 6, 7, 8].

Elderly HF patients generally have complex comorbidity profiles, including not only comorbid diseases but also geriatric conditions [3, 9, 10]. These complex and diverse profiles are associated with poor prognosis of HF and increase the burden on health care services after discharge [11, 12]. Thus, a multidisciplinary approach, as well as medical treatment, is becoming important for the management of elderly patients with HF.

Oral health problems, like dry mouth, periodontal disease, dental caries, and inappropriate dentures, are common health conditions in older adults [13, 14]. Previous studies have demonstrated the association between poor oral health and various geriatric conditions, such as frailty, undernutrition, and cognitive impairment [15, 16, 17, 18]. However, the importance of oral health care has not been fully highlighted in the assessment and management of HF patients. In the present study, we investigated the clinical implications of oral assessment in elderly patients with acute HF.

## Materials And Methods

### Study population

Patients admitted to Anjo Kosei Hospital for the treatment of HF between October 2018 and March 2019 were reviewed. All patients were diagnosed with HF using the Framingham criteria [19]. During hospitalization, standardized cardiac inpatient rehabilitation was provided for all the patients according to standardized cardiac rehabilitation program from the Japanese Association of Cardiac Rehabilitation [20]. The study was conducted in accordance with the Declaration of Helsinki and approved by the Anjo Kosei Hospital ethics committee. Because of its retrospective nature, informed consent was deemed unnecessary according to the national regulation issued by the Japanese Ministry of Health, Labour and Welfare. However, the present study was carried out by the opt-out method of our hospital website.

A medical history was obtained to document past medical history, medications, and co-morbid disease. Hypertension was defined as systolic blood pressure (BP)  $\geq$  140 mmHg or diastolic BP  $\geq$  90 mmHg on repeated measurements, or receipt of antihypertensive treatment. Diabetes mellitus was defined as having a blood hemoglobin A1c  $\geq$  6.5%, 2-hour value  $\geq$  200 mg/dL ( $\geq$  11.1 mmol/L) on a 75 g oral glucose tolerance test, and/or taking glucose-modulating medication according to the diagnostic criteria of the Japan Diabetes Society [21].

### **Oral assessment**

Oral health was assessed by a certified dysphagia nurse using the revised oral assessment guide (ROAG) [22]. ROAG includes eight categories: voice, lips, mucous membranes, tongue, gums, teeth/dentures, saliva, and swallowing. Each category was described and rated as healthy (score 1) to severe (score 3). The total score ranged from eight, representing a normal oral health, to twenty-four, which represents severe oral health problems. The ROAG score was obtained within 1 week of hospitalization when the patient's respiratory state was stabilized without the need for oxygen. In the present study, patients with ROAG scores of 8 points were regarded to have a normal oral health and those with higher scores were regarded as having poor oral health (mild to moderate oral problems: 9-12 points, severe oral problems: 13-24 points).

### **Assessments for geriatric conditions**

Physical functional status was evaluated using the Barthel Index (BI), handgrip, and 10-meter normal/usual gait speed. The BI was obtained by ward nurses at admission and at discharge, as previously reported [8]. Changes in the BI were calculated as the difference between the BI on admission and the BI on discharge. Handgrip and 10-meter gait speed were evaluated by physical therapists before discharge.

Registered dietitians assessed nutritional status. Nutritional status was screened using the controlling nutritional status (CONUT) score and the geriatric nutritional risk index (GNRI) [23, 24]. Laboratory data at admission and body mass index (BMI) at the first measurement within 72 hours of hospitalization were used for calculation of the scores. Dietary energy intake was assessed by the proportion of nutritional intake from food compared to the predicted calorie requirement. Nutritional intake was calculated based on

the food intake for 3 days around the day of oral assessment. The predicted calorie requirement was defined as the total energy expenditure estimated from the Harris-Benedict equation [25].

Pharmacists assessed cognitive function using the mini-mental state examination (MMSE) before discharge [26].

## **Biomarker analysis and echocardiography**

Blood samples were obtained at the time of hospital admission. Complete blood counts were performed utilizing a Sysmex XE-5000 analyzer (Sysmex, Kobe, Japan). Plasma BNP was measured with the AIA-2000 enzymatic immunoassay analyzer (TOSOH, Tokyo, Japan). Other biomarkers were measured using a LABOSPECT 008 autoanalyzer (Hitachi Co., Tokyo, Japan). Estimated glomerular filtration rate (eGFR) was calculated by the Modification of Diet in Renal Disease formula [27]. Echocardiographic examination was performed by an experienced sonographer using Vivid E9 with XD clear (GE Healthcare, Tokyo, Japan). The images were recorded in a console and analyzed offline. Left ventricular ejection fraction was calculated using the modified Simpson's rule.

## **Statistical analysis**

All analyses were performed using PASW Statistics 21 software (SPSS Inc., Chicago, IL, USA). Continuous variables were presented as the mean  $\pm$  standard deviation or median (interquartile range). We assessed the normal distribution of continuous variables by the Shapiro-Wilk test. One-way analysis of variance (ANOVA) or Kruskal-Wallis test was used to determine the mean or median differences in variables between the groups. Categorical variables were presented as the count and/or percentage. Categorical variables were presented as the count and/or percentage and the chi-square test or Fisher's exact test was used for the group comparisons. Univariate correlations between changes in the BI during hospitalization and other variables were investigated using the Pearson's rank correlation test, and then a multiple linear regression analysis was performed. Age, gender, BI at admission, and variables with  $P < 0.05$  in the univariate analyses were incorporated into the multivariable model. In all analyses,  $P < 0.05$  was considered statistically significant.

## **Results**

### **Baseline characteristics**

Among a total of 171 eligible patients, 38 patients younger than 65 years or those with in-hospital death were excluded. We also excluded 56 patients whose ROAG score could not be obtained. Finally, the study included 77 patients. The mean age of overall patients was  $80.0 \pm 9.1$  years, and 58.4% of patients were men. The mean ROAG score was  $9.9 \pm 2.2$ . Poor oral health (ROAG score  $\geq 9$ ) was identified in 66.2% of the enrolled patients. Details of ROAG evaluations are shown in Table 1.

**Table 1. Baseline characteristics of enrolled patients**

	All patients (n = 77)	Normal oral health (n = 26)	Poor oral health (n = 51)	P- value
ROAG score	9.9 ± 2.2	8	10.9 ± 2.1	0.001
Age (years)	80.0 ± 9.1	73.8 ± 8.3	83.1 ± 7.9	0.001
Male sex	45 (58.4%)	19 (73.1%)	26 (51.0%)	0.063
Body mass index (kg/m <sup>2</sup> )	22.2 ± 3.9	23.1 ± 4.6	21.7 ± 3.5	0.159
History of admission due to HF	35 (45.5%)	14 (53.8%)	21 (41.2%)	0.291
History of CAD	26 (33.8%)	8 (30.8%)	18 (35.3%)	0.691
History of stroke	11 (14.3%)	3 (11.5%)	8 (15.7%)	0.623
Hypertension	65 (84.4%)	21 (80.8%)	44 (86.3%)	0.529
Diabetes mellitus	29 (37.7%)	7 (26.9%)	22 (43.1%)	0.165
Atrial fibrillation	36 (46.8%)	13 (50.0%)	23 (45.1%)	0.683
Hemoglobin (g/dL)	11.5 ± 2.3	12.1 ± 2.7	11.1 ± 2.0	0.072
CRP (mg/dL)	0.84 (0.17-2.54)	0.25 (0.13-0.86)	1.24 (0.28-4.19)	0.009
Albumin (g/dL)	3.4 ± 0.5	3.6 ± 0.5	3.3 ± 0.5	0.014
Creatinine (mg/dL)	1.60 ± 1.02	1.69 ± 1.20	1.56 ± 0.92	0.580
eGFR (mL/min/1.73 m <sup>2</sup> )	40.0 ± 20.6	42.5 ± 22.7	38.7 ± 19.5	0.446
Sodium (mEq/L)	139.9 ± 3.5	139.0 ± 3.7	140.3 ± 3.4	0.128
BNP (pg/mL)	805.0 (502.4-1304.8)	992.0 (415.5-1687.2)	732.8 (531.9-1101.0)	0.598
LV ejection fraction (%)	44.2 ± 17.4	39.7 ± 16.6	46.7 ± 17.5	0.106
Medications on admission				
ACE inhibitors/ARBs	31 (40.3%)	12 (46.1%)	19 (37.3%)	0.451
Beta-blockers	41 (53.2%)	16 (61.5%)	25 (49.0%)	0.298
Calcium channel blockers	32 (41.6%)	11 (42.3%)	21 (41.2%)	0.360
Loop diuretics	45 (58.4%)	15 (57.7%)	30 (58.8%)	0.924
MRAs	13 (27.5%)	6 (23.1%)	7 (13.7%)	0.300
Tolvaptans	14 (27.5%)	8 (30.8%)	14 (27.5%)	0.761

ta are presented as mean ± standard deviation (SD), median (interquartile range [IQR]), or n (%). Bold type indicates statistical significance. ROAG, revised oral assessment guide; HF, heart failure; CAD, coronary artery disease; CRP, C-reactive protein; eGFR, estimated glomerular filtration rate; BNP, brain natriuretic peptide; LV, left ventricular; ACE, angiotensin-converting enzyme; ARBs, angiotensin receptor blockers; MRA, mineralocorticoid receptor antagonist.

Patients were divided into three groups based on the ROAG score: a normal oral health group (ROAG score = 8, n = 26), a poor oral health group with mild to moderate problems (ROAG score = 9-12, n = 40), and a poor oral health group with severe problems (ROAG score = 13-24, n = 11) (Table 2). Patients with poor oral health were older and had lower albumin levels. C-reactive protein and sodium levels were higher in patients with poor oral health than in those with normal oral health. History of stroke and concomitant respiratory infection at admission were highly prevalent in patients with severe oral problems. There were no significant differences in gender, BMI, history of HF, etiology of coronary arterial disease (CAD), prevalence of current smoking, hypertension, diabetes and atrial fibrillation, renal function, hemoglobin, sodium, BNP level, left

ventricular ejection fraction, the use of non-invasive positive pressure ventilation or inotropes for the management of HF, length of bed rest, and the use of medications on admission (Table 2).

**Table 2. Details of ROAG evaluation of the enrolled patients**

Items	Evaluation (score)		
	Normal (1)	Minor problems (2)	Severe problems (3)
voice	47	27	3
lips	41	36	0
mucous membranes	75	2	0
tongue	60	17	0
gums	73	4	0
teeth/dentures	62	13	2
saliva	58	11	8
swallowing	65	10	2

Data are presented as number of patients. ROAG, revised oral assessment guide.

### Association between oral health and geriatric assessments on admission

We examined the association between oral health and common geriatric assessments on admission. Barthel Index, GNRI, dietary energy intake, and MMSE were significantly lower in the poor oral health group than in the normal oral health group. The COUNT score was significantly higher in patients with poor oral health than those with normal oral health (Table 3). There were no significant differences between the groups in handgrip and gait speed for the 10-meter walk. Use of dentures was also comparable between 3 groups. Thus, patients with poor oral health showed a higher prevalence of undernutrition and cognitive impairment.

**Table 3. Geriatric assessment**

	Normal oral health (n = 26)	Poor oral health (n = 51)	P-value
Barthel Index at admission	96.0 ± 9.8	76.6 ± 30.8	0.004
Handgrip (kg)	20.2 ± 8.7	14.5 ± 6.6	0.040
Gait speed for 10-meter walk (sec)	14.8 ± 6.2	19.5 ± 7.5	0.069
CONUT score	4.3 ± 2.9	5.7 ± 2.8	0.045
GNRI	98.1 ± 14.3	87.0 ± 9.9	< 0.001
Dietary energy intake (%)	89.6 ± 18.7	67.5 ± 32.3	0.002
MMSE	27.4 ± 2.4	25.2 ± 3.4	0.047

Data are presented as mean ± standard deviation (SD). Bold type indicates statistical significance. CONUT, controlling nutritional status; GNRI, geriatric nutritional risk index; MMSE, mini-mental state examination.

### Association between ROAG score and activities of daily living (ADL) preservation

Efforts to preserve patients' ability to perform ADL during hospitalization are important in the management of acute heart failure. The BI has been reported as a common tool to evaluate the ability to perform ADL. Here, we investigated the changes in the BI during hospitalization. The actual BI values on admission and discharge, and the BI changes in each group are shown in Table 4. Reduction in the BI during hospitalization was significant in the enrolled patients in both the normal oral health group and poor oral health group ( $P < 0.01$ ). Of note, the BI decreased more in patients with poor oral health than those with normal oral health ( $P < 0.01$ ). Hospital stay was significantly longer in patients with severe oral problems (Table 4).

**Table 4. Changes in the Barthel index during hospitalization**

	All patients (n = 77)	Normal oral health (n = 26)	Poor oral health (n = 51)	P-value
Barthel Index at admission	82.9 ± 27.4	96.0 ± 9.8	76.6 ± 30.8	0.004
Barthel Index at discharge	77.4 ± 32.3	93.5 ± 12.8	67.9 ± 36.4	0.001
Changes in Barthel Index	-7.4 ± 18.1	-1.2 ± 6.6	-11.0 ± 21.5	0.036

Data are presented as mean ± standard deviation (SD). Bold type indicates statistical significance.

Finally, to determine factors associated with ADL preservation during hospitalization in elderly patients with acute HF, we compared the clinical parameters on admission that were associated with the change in the BI during hospitalization using univariate and multivariate regression analyses. Changes in the BI during hospitalization were significantly correlated with age, hemoglobin levels, ROAG score, GNRI, and dietary energy intake on admission in the univariate regression analysis. Of those, ROAG score on admission was the only independent predictor of changes in the BI during hospitalization in the multivariate regression analyses (Table 5).

**Table 5. Independent predictors of change in the Barthel index based on linear regression analyses.**

	Univariate		multivariate	
	$\beta$	P-value	$\beta$	P-value
Age	-0.357	0.004	-0.175	0.219
Male sex	-0.150	0.233		
Body mass index	0.128	0.327		
History of admission due to HF	0.102	0.420		
Atrial fibrillation	0.080	0.527		
Hemoglobin	0.249	0.046	0.060	0.629
Log CRP	-0.183	0.165		
Albumin	0.114	0.365		
eGFR	0.062	0.622		
Sodium	-0.034	0.787		
Log BNP	0.047	0.710		
ROAG score	-0.421	< 0.001	-0.307	0.042
Barthel Index at admission	0.080	0.524		
Handgrip	0.313	0.087		
Gait speed for 10-meter walk	-0.282	0.124		
CONUT score	-0.210	0.098		
GNRI	0.276	0.034	0.106	0.431
Dietary energy intake	0.329	0.008	0.003	0.979
MMSE	0.154	0.810		

Bold type indicates statistical significance. HF, heart failure; CRP, C-reactive protein; eGFR, estimated glomerular filtration rate; BNP, brain natriuretic peptide; ROAG, revised oral assessment guide; CONUT, controlling nutritional status; GNRI, geriatric nutritional risk index; MMSE, mini-mental state examination.

## Discussion

The major findings of this study were as follows: 1) poor oral health, assessed using the ROAG score, is relatively common in elderly patients with acute HF, 2) patients with poor oral health had significantly more geriatric conditions than those with normal oral health, and 3) the ROAG score was independently correlated with changes in the BI during hospitalization in elderly patients with acute HF.

Oral health problems have been reported as common health conditions in older adults and hospitalized patients [17, 18, 28, 29]. Several points are known about the relationship between oral health problems and heart failure. Chronic inflammation caused by periodontal diseases is a risk factor for cardiac and cerebrovascular diseases, which are the most major comorbidities of HF [30, 31]. General fatigue, dyspnea, delirium, and sleep disturbance accompanied by decompensated HF might contribute to reducing adherence to oral hygiene. In addition, dry mouth is often a consequence of polypharmacy, particularly as a side effect of cardiovascular agents (angiotensin-converting enzyme inhibitors, beta-blockers, and diuretics) [32, 33].

However, oral health has been underrecognized in the assessment and management of patients with HF. Our data suggests the importance of oral assessment in the multidisciplinary management of elderly HF patients because poor oral health is highly prevalent and significantly associated with decline in physical function during hospitalization.

We have previously demonstrated that hospitalization for HF was significantly correlated with decreased BI as an assessment of ADL, and a decreased BI during hospitalization was associated with worse clinical outcomes<sup>8</sup>. It has also been reported that a decline in ADL due to acute HF is an independent risk factor of hospitalization for HF and mortality [34, 35]. Therefore, it is important to identify predictors of ADL decline during hospitalization in patients with HF. It has been demonstrated in elderly patients and patients with HF that age and the nutritional index are associated with ADL decline during hospitalization [34, 36]. Our present study involving elderly patients with HF has shown that the ROAG score serves as a good predictor of changes in ADL, and its predictive ability is comparable to age, nutritional indices, and various other parameters. Therefore, oral assessment of patients with HF using the ROAG score to screen for poor oral health may allow us to provide comprehensive care and rehabilitation to patients with poor oral health at an earlier stage. This may help prevent ADL decline during hospitalization for HF, leading to a better prognosis.

In the present study, nutritional status, assessed by CONUT and GNRI, was worse in the poor oral health group than in the normal oral health group. Furthermore, patients with poor oral health had lower dietary energy intake. Previous studies have demonstrated that poor oral health is associated with periodontal disease, dental caries, hyposalivation, and tooth loss or edentulousness, which pose risks of chewing difficulties, decreased masticatory function, and dysphagia [37, 38]. These oral problems may induce a preference for soft and easily chewable food and a need for changes in food texture to prevent aspiration and choking, leading to poor nutritional intake and undernutrition, and finally to sarcopenia, frailty, and decreased physical function [17, 39, 40]. Although undernutrition would be a major contributor, the association between poor oral health and changes in BI might be influenced by multiple factors. Besides undernutrition, higher prevalence of cognitive impairment among patients with poor oral health might also be associated with physical deconditioning caused by prolonged bed rest or inactivity. Furthermore, a decline in physical and cognitive function would, in turn, reduce self-care ability, including the ability to take care of one's oral health.

In the acute setting of decompensated HF, nurses have a pivotal role in oral assessment and care, because of the small number of dentists and oral hygienists. Several oral health assessment tools have been developed for non-dental health care professionals [22, 41, 42]. Among these indices, the ROAG is not only a simple and comprehensive assessment but it also has favorable validity and reliability [43]. Oral assessment in patients with acute HF might have several influences on multidisciplinary disease management. First, patients who require consultation with in-hospital or regional dental health care professionals are screened. Second, nurses receive feedback on oral care from their patients. Third, oral care is one of the fundamental self-care activities after discharge, and thus, the information that an oral assessment provides is useful for educating patients on dental health compliance. Fourth, several management strategies could be considered for patients with poor oral health. Nutritionists and dietitians could help to increase the dietary intake of the patients by changing meal content and food texture. Doctors and pharmacists could choose orally

disintegrating tablets for the ease of taking medicines. Although the efficacy of multidisciplinary interventions for patients with poor oral health has not been fully elucidated, we believe that oral assessment could provide useful information for the multidisciplinary management of elderly patients with HF.

The present study had several limitations. First, this was a single-center, retrospective study, and the sample size was relatively small. Second, the large number of eligible patients were excluded due to the lack of ROAG score. Because this is a pilot study to investigate the clinical implications of the oral assessment in patients with HF, oral health was assessed by only one certified dysphagia nurse to avoid the interobserver variation. Future prospective studies are necessary with larger patient populations. Third, the results could not be generalized to the routine assessment by ward nurses.

## **Conclusions**

Poor oral health, as assessed by the ROAG, is highly prevalent and oral assessment using the ROAG predicts a decline in physical function during hospitalization in elderly patients with acute HF. Thus, oral assessment during hospitalization could provide useful information for the management of elderly HF patients.

## **Declarations**

### **Data Availability**

The data that support the findings of this study are available from the corresponding author upon reasonable request.

### **Conflicts of interest**

The author(s) declare(s) that there is no conflict of interest regarding the publication of this article.

### **Funding statement**

None

### **Acknowledgements**

We are grateful to the members of the Multidisciplinary Team for Heart Failure at Anjo Kosei Hospital for their help in the acquisition of clinical data.

## **References**

1. Benjamin EJ, Virani SS, Callaway CW, Chamberlain AM, Chang AR, Cheng S, et al. Heart Disease and Stroke Statistics-2018 Update: A Report From the American Heart Association. *Circulation* 2018; 137: e67-e492, doi:10.1161/CIR.0000000000000558.
2. Shimokawa H, Miura M, Nochioka K, Sakata Y. Heart failure as a general pandemic in Asia. *Eur J Heart Fail* 2015; 17: 884-892.
3. Conrad N, Judge A, Tran J, Mohseni H, Hedgecott D, Crespillo AP, et al. Temporal trends and patterns in heart failure incidence: a population-based study of 4 million individuals. *Lancet* 2018; 391: 572-580.
4. Bui AL, Horwich TB, Fonarow GC. Epidemiology and risk profile of heart failure. *Nat Rev Cardiol* 2011; 8: 30-41.
5. Lawson CA, Zaccardi F, Squire I, Ling S, Davies MJ, Lam CSP, et al. 20-year trends in cause-specific heart failure outcomes by sex, socioeconomic status, and place of diagnosis: a population-based study. *Lancet Public Health* 2019; 4: e406-e420, doi:10.1016/S2468-2667(19)30108-2.
6. Hamaguchi S, Kinugawa S, Goto D, Tsuchihashi-Makaya M, Yokota T, Yamada S, et al. Predictors of long-term adverse outcomes in elderly patients over 80 years hospitalized with heart failure. - A report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J* 2011; 75: 2403-2410.
7. Chioncel O, Lainscak M, Seferovic PM, Anker SD, Crespo-Leiro MG, Harjola VP, et al. Epidemiology and one-year outcomes in patients with chronic heart failure and preserved, mid-range and reduced ejection fraction: an analysis of the ESC Heart Failure Long-Term Registry. *Eur J Heart Fail* 2017; 19: 1574-1585.
8. Uemura Y, Shibata R, Takemoto K, Koyasu M, Ishikawa S, Murohara T, et al. Prognostic Impact of the Preservation of Activities of Daily Living on Post-Discharge Outcomes in Patients With Acute Heart Failure. *Circ J* 2018; 82: 2793-2799.
9. Afilalo J, Alexander KP, Mack MJ, Maurer MS, Green P, Allen LA, et al. Frailty assessment in the cardiovascular care of older adults. *J Am Coll Cardiol* 2014; 63: 747-762.
10. Henkel DM, Redfield MM, Weston SA, Gerber Y, Roger VL. Death in heart failure: a community perspective. *Circ Heart Fail* 2008; 1: 91-97.
11. Chaudhry SI, Wang Y, Gill TM, Krumholz HM. Geriatric conditions and subsequent mortality in older patients with heart failure. *J Am Coll Cardiol* 2010; 55: 309-316.
12. van Deursen VM, Urso R, Laroche C, Damman K, Dahlstrom U, Tavazzi L, et al. Co-morbidities in patients with heart failure: an analysis of the European Heart Failure Pilot Survey. *Eur J Heart Fail* 2014; 16: 103-111.
13. Marcenes W, Kassebaum NJ, Bernabe E, Flaxman A, Naghavi M, Lopez A, et al. Global burden of oral conditions in 1990-2010: a systematic analysis. *J Dent Res* 2013; 92: 592-597.

14. Kassebaum NJ, Smith AGC, Bernabe E, Fleming TD, Reynolds AE, Vos T, et al. Global, Regional, and National Prevalence, Incidence, and Disability-Adjusted Life Years for Oral Conditions for 195 Countries, 1990-2015: A Systematic Analysis for the Global Burden of Diseases, Injuries, and Risk Factors. *J Dent Res* 2017; 96: 380-387.
15. Hasegawa Y, Sakuramoto A, Sugita H, Hasegawa K, Horii N, Sawada T, et al. Relationship between oral environment and frailty among older adults dwelling in a rural Japanese community: a cross-sectional observational study. *BMC Oral Health* 2019; 19: 23, doi:10.1186/s12903-019-0714-8.
16. Ramsay SE, Papachristou E, Watt RG, Tsakos G, Lennon LT, Papacosta AO, et al. Influence of Poor Oral Health on Physical Frailty: A Population-Based Cohort Study of Older British Men. *J Am Geriatr Soc* 2018; 66: 473-479.
17. Azzolino D, Passarelli PC, De Angelis P, Piccirillo GB, D'Addona A, Cesari M. Poor Oral Health as a Determinant of Malnutrition and Sarcopenia. *Nutrients* 2019; 11, doi:10.3390/nu11122898.
18. Kossioni AE. The Association of Poor Oral Health Parameters with Malnutrition in Older Adults: A Review Considering the Potential Implications for Cognitive Impairment. *Nutrients* 2018; 10, doi:10.3390/nu10111709.
19. McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 1971; 285: 1441-1446.
20. Izawa H, Yoshida T, Ikegame T, Izawa KP, Ito Y, Okamura H, et al. Standard Cardiac Rehabilitation Program for Heart Failure. *Circ J* 2019; 83: 2394-2398.
21. Committee of the Japan Diabetes Society on the Diagnostic Criteria of Diabetes M, Seino Y, Nanjo K, Tajima N, Kadowaki T, Kashiwagi A, et al. Report of the committee on the classification and diagnostic criteria of diabetes mellitus. *J Diabetes Investig* 2010; 1: 212-228.
22. Andersson P, Hallberg IR, Renvert S. Inter-rater reliability of an oral assessment guide for elderly patients residing in a rehabilitation ward. *Spec Care Dentist* 2002; 22: 181-186.
23. Ignacio de Ulibarri J, Gonzalez-Madrono A, de Villar NG, Gonzalez P, Gonzalez B, Mancha A, et al. CONUT: a tool for controlling nutritional status. First validation in a hospital population. *Nutr Hosp* 2005; 20: 38-45.
24. Bouillanne O, Morineau G, Dupont C, Coulombel I, Vincent JP, Nicolis I, et al. Geriatric Nutritional Risk Index: a new index for evaluating at-risk elderly medical patients. *Am J Clin Nutr* 2005; 82: 777-783.
25. Harris JA, Benedict FG. A Biometric Study of Human Basal Metabolism. *Proc Natl Acad Sci U S A* 1918; 4: 370-373.
26. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; 12: 189-198.

27. Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis* 2009; 53: 982-992.
28. Shiraishi A, Yoshimura Y, Wakabayashi H, Tsuji Y, Shimazu S, Jeong S. Impaired oral health status on admission is associated with poor clinical outcomes in post-acute inpatients: A prospective cohort study. *Clin Nutr* 2019; 38: 2677-2683.
29. Hanne K, Ingelise T, Linda C, Ulrich PP. Oral status and the need for oral health care among patients hospitalised with acute medical conditions. *J Clin Nurs* 2012; 21: 2851-2859.
30. Humphrey LL, Fu R, Buckley DI, Freeman M, Helfand M. Periodontal disease and coronary heart disease incidence: a systematic review and meta-analysis. *J Gen Intern Med* 2008; 23: 2079-2086.
31. Sen S, Giamberardino LD, Moss K, Morelli T, Rosamond WD, Gottesman RF, et al. Periodontal Disease, Regular Dental Care Use, and Incident Ischemic Stroke. *Stroke* 2018; 49: 355-362.
32. Smidt D, Torpet LA, Nauntofte B, Heegaard KM, Pedersen AM. Associations between oral and ocular dryness, labial and whole salivary flow rates, systemic diseases and medications in a sample of older people. *Community Dent Oral Epidemiol* 2011; 39: 276-288.
33. Kakudate N, Muramatsu T, Endoh M, Satomura K, Koseki T, Sato Y, et al. Factors associated with dry mouth in dependent Japanese elderly. *Gerodontology* 2014; 31: 11-18.
34. Takabayashi K, Kitaguchi S, Iwatsu K, Morikami Y, Ichinohe T, Yamamoto T, et al. A decline in activities of daily living due to acute heart failure is an independent risk factor of hospitalization for heart failure and mortality. *J Cardiol* 2019; 73: 522-529.
35. Katano S, Yano T, Ohori K, Kouzu H, Nagaoka R, Honma S, et al. Barthel Index Score Predicts Mortality in Elderly Heart Failure – A Goal of Comprehensive Cardiac Rehabilitation–. *Circ J* 2021; doi: 10.1253/circj.CJ-21-0584.
36. Hsu YH, Chou MY, Chu CS, Liao MC, Wang YC, Lin YT, et al. Predictive Effect of Malnutrition on Long-Term Clinical Outcomes among Older Men: A Prospectively Observational Cohort Study. *J Nutr Health Aging* 2019; 23: 876-882.
37. Gil-Montoya JA, de Mello AL, Barrios R, Gonzalez-Moles MA, Bravo M. Oral health in the elderly patient and its impact on general well-being: a nonsystematic review. *Clin Interv Aging* 2015; 10: 461-467.
38. Razak PA, Richard KM, Thankachan RP, Hafiz KA, Kumar KN, Sameer KM. Geriatric oral health: a review article. *J Int Oral Health* 2014; 6: 110-116.
39. Castrejon-Perez RC, Borges-Yanez SA, Gutierrez-Robledo LM, Avila-Funes JA. Oral health conditions and frailty in Mexican community-dwelling elderly: a cross sectional analysis. *BMC Public Health* 2012; 12: 773, doi:10.1186/1471-2458-12-773.

40. Zenthofer A, Rammelsberg P, Cabrera T, Hassel AJ. Increasing dependency of older people in nursing homes is associated with need for dental treatments. *Neuropsychiatr Dis Treat* 2014; 10: 2285-2290.
41. Chalmers JM, King PL, Spencer AJ, Wright FA, Carter KD. The oral health assessment tool—validity and reliability. *Aust Dent J* 2005; 50: 191-199.
42. Fjeld KG, Eide H, Mowe M, Hove LH, Willumsen T. Dental hygiene registration: development, and reliability and validity testing of an assessment scale designed for nurses in institutions. *J Clin Nurs* 2017; 26: 1845-1853.
43. Everaars B, Weening-Verbree LF, Jerkovic-Cosic K, Schoonmade L, Bleijenberg N, de Wit NJ, et al. Measurement properties of oral health assessments for non-dental healthcare professionals in older people: a systematic review. *BMC Geriatr* 2020; 20: 4, doi:10.1186/s12877-019-1349-y.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [BMCOAGTableS1.docx](#)