Large mid-upper arm circumference is associated with metabolic syndrome in middle-aged and elderly individuals: a community-based study

Jie Shi

Shanghai Jiaotong University School of Medicine Xinhua Hospital Chongming Branch

Zhen Yang (✉ yangzhen@xinhumed.com.cn)

Shanghai Jiaotong University School of Medicine Xinhua Hospital Chongming Branch

<https://orcid.org/0000-0003-0225-7099>

Yixin Niu

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Weiwei Zhang

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Xiaoyong Li

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Hongmei Zhang

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Ning Lin

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Hongxia Gu

Shanghai Jiaotong University School of Medicine Xinhua Hospital Chongming Branch

Jie Wen

Huashan Hospital Fudan University

Guang Ning

Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital

Li Qin

Shanghai Jiaotong University School of Medicine Xinhua Hospital Chongming Branch

Qing Su

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Research article

Keywords: metabolic syndrome, mid-upper arm circumference, central obesity, subcutaneous fat

Posted Date: April 2nd, 2020

DOI: <https://doi.org/10.21203/rs.2.16072/v3>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Endocrine Disorders on June 3rd, 2020. See the published version at <https://doi.org/10.1186/s12902-020-00559-8>.

Abstract

Background: The mid-upper arm circumference (MUAC) is a proxy for subcutaneous fat in the upper body and is a reliable screening measure for identifying individuals with abnormal local fat distribution. The purpose of this study was to evaluate the association between MUAC and metabolic syndrome (MetS) in middle-aged and elderly individuals.

Methods: We measured the MUAC in a cross-sectional sample with a total of 9787 subjects aged 40 years and older. The measurement of MUAC is performed on the right arm using a non-elastic tape held midway between the acromion and the olecranon processes in duplicate, with the arm hanging loosely at the side of the body. The MetS was defined according to the Joint Statement of the International Diabetes Federation Task Force on Epidemiology and Prevention.

Results: MUAC was positively correlated with waist circumference ($r=0.437$, $P<0.001$), BMI ($r=0.334$, $P<0.001$), fasting insulin ($r=0.348$, $P<0.001$), HOMA-IR ($r=0.134$, $P<0.001$), triglycerides ($r=0.138$, $P<0.001$), SBP ($r=0.124$, $P<0.001$), and DBP ($r=0.123$, $P<0.001$), and inversely correlated with adiponectin ($r=-0.147$, $P<0.001$) and HDL-cholesterol ($r=-0.176$, $P<0.001$) after adjusting for age and gender. Compared with the lowest quartile group, the odds ratios were substantially higher for MetS (OR 1.77; 95% CI 1.51-2.09, P for trend <0.001) in the highest MUAC quartile group after adjustment for potential cofounder.

Conclusion: Large mid-upper arm circumference is significantly associated with metabolic syndrome in middle-aged and elderly individuals.

Background

Metabolic syndrome (MetS), which contains a cluster of metabolic abnormalities including central obesity, hypertension, dysglycemia, and dyslipidemia that together culminate in the increased risk of cardiovascular disease (CVD) and diabetes, is a major global public health problem[1]. Of note, the prevalence of MetS is up to 30% in middle-aged and elderly people in China[2, 3]. Given the continuous increase of the aging population in China, high prevalence of MetS is a concerning observation that should be put on the agenda.

It is well documented that regional differences in fat accumulation exist imbalances in fatty acid storage and/or free fatty acid (FFA) release between fat depots[4]. In addition, the upper-body subcutaneous fat depot is lipolytic ally more active than lower-body subcutaneous fat[5, 6]. Mounting irresistible evidence suggests that upper-body subcutaneous adipose, usually correlated with increased visceral adipose, is associated with a metabolic disorder in different body mass indexes[7, 8]. Furthermore, previous studies have also demonstrated that systemic FFA concentration is primarily determined by upper-body subcutaneous fat[5, 9], indicating that upper arm adipose depot may play a pivotal role in risk factor pathogenesis. On the one hand, the transfer of excess FFA from systemic adipose tissue lipolysis to liver tissue will result in liver lipid accumulation. On the other hand, elevated FFA levels also provoke VLDL-

triglyceride production, insulin resistance, and inflammation, further increases the risk of metabolic disorders[10]. Furthermore, it has been demonstrated that dysregulation of fatty acid disposition, with ectopic lipid accumulation in non-adipose cells, is a major factor contributing to the development of MetS[11].

More recently, several studies have reported that mid-upper arm circumference (MUAC), a feasible and valid screening tool for identifying subjects with abnormal distribution of upper-body subcutaneous fat, was positively associated with higher risks of cardiometabolic disorders and subclinical atherosclerosis independent of waist circumference and BMI[12]. However, currently, the study on the association of MUAC with MetS among Chinese individuals is scarce. Thus, the purpose of this study was to evaluate the relationship between MUAC and MetS among Chinese middle-aged and elderly individuals.

Methods

Study population

The study is a part of the Risk Evaluation of cAncers in Chinese diabeTic Individuals: A lONgitudinal (REACTION) study, a community-based cross-sectional study conducted among 259,657 Chinese individuals aged 40 years and older[13]. REACTION study performed in 25 communities across mainland China, from 2011 to 2012. The study design and methods have been described previously in detail[13-15]. The data presented in this article are based on the baseline survey of subsamples from the Chongming District, Shanghai, China. A total of 9930 eligible subjects participated in the study. After excluding individuals with missing data about MUAC ($n = 143$), 9787 subjects (3156 men and 6631 women) were eventually included in the present analysis. The study protocol was approved by the Ethics Committee of Xinhua Hospital Affiliated to Shanghai Jiaotong University School of Medicine. Written informed consent was obtained from all participants.

Data collection

A standardized questionnaire was applied by trained physicians to collect essential information, including sex, age, lifestyle factors, educational attainment, physical activity, and previous medical history. Anthropometric measurements were collected by certified physicians using standard protocols in duplicate base on the National Health and Nutrition Examination Survey (NHANES) Anthropometry Procedures Manual[16]. The measurement of MUAC is performed on the right arm using a non-elastic tape held midway between the acromion and the olecranon processes, with the arm hanging loosely at the side of the body. Waist circumference was measured with a non-elastic tape held midway between the lower rib margin and the iliac crest at the end of a gentle expiration. Blood pressure (BP) was measured with an automated electronic device (OMRON Model1 Plus; Omron Company, Kyoto, Japan). Overweight was defined as body mass index (BMI) $\geq 24 \text{ kg/m}^2$, central obesity was defined as waist circumference $\geq 85 \text{ cm}$ for men and $\geq 80 \text{ cm}$ for women.

Biochemical measurements

Peripheral venous blood samples were collected after an overnight fast for at least 10 hours. The plasma glucose level was measured by the glucose oxidase method (ADVIA-1650 Chemistry System, Bayer, Leverkusen, Germany). Total cholesterol, triglycerides, low-density lipoprotein (LDL) cholesterol, and high-density lipoprotein (HDL) cholesterol were assessed by an automatic analyzer (Hitachi 7080; Tokyo, Japan). Hemoglobin A1c was measured with HPLC method (BIO-RAD, D10, CA). Fasting insulin was determined by RIA (Linco Research, St.Charles, MO). Insulin resistance was evaluated by the homeostasis model of assessment for insulin resistance (HOMA-IR).

Definition of MetS

The MetS was defined based on the Joint Statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity[1] as presenting at least three of the following five components: (1) waist circumference ≥ 85 cm in men or ≥ 80 cm in women, (2) triglycerides ≥ 1.7 mmol/L, (3) HDL cholesterol < 1.0 mmol/L in men or < 1.3 mmol/L in women, (4) systolic blood pressure (SBP) ≥ 130 and/or diastolic blood pressure (DBP) ≥ 85 mm Hg or current use of antihypertensive medications, (5) fasting plasma glucose ≥ 5.6 mmol/L, previous diagnosis of type 2 diabetes or use of antidiabetic agents.

Statistical analysis

Normally distributed data were reported as means $\pm$ SD, whereas variables with a skewed distribution were expressed as median (interquartile range) and log-transformed to approximate normality before analysis. Categorical variables were depicted by frequency and percentage. The subjects were divided into four groups according to the MUAC quartiles. For comparisons between groups, we performed one-way ANOVA for continuous variables with normal distributions and nonparametric tests for continuous variables with highly skewed distributions. The Chi-squared test was applied to compare categorical variables. Correlation coefficients between MUAC and metabolic features were calculated by partial correlation analysis after adjusted age and gender. Multivariate logistic regression analyses were used to test odds ratios (ORs) and 95% confidence intervals (CIs) of Mets for each MUAC quartile compared with the lowest quartile (Q1) group. Model 1 was adjusted for age and gender; model 2 was adjusted for the variables in model 1 plus smoking, alcohol drinking, educational attainment, physical activity, and self-reported CVD; model 3 was adjusted for the variables in model 2 plus CRP, adiponectin, and HOMA-IR; model 4 was adjusted for the variables in model 3 plus BMI. Data management and statistical analysis were performed with SPSS (version 23.0). $P < 0.05$ was considered statistically significant.

Results

Characteristics of participants according to MUAC quartiles

The mean of MUAC was 29.24 cm for males and 28.41 cm for females ($P < 0.001$), respectively. The individuals were divided into four groups based on the quartiles of MUAC. When analyzed by quartiles of

MUAC levels, as summarized in Table 1, the subjects with larger MUAC were more likely to be smokers, alcohol drinkers, and comorbidities including hypertension, hyperlipidemia, diabetes (all $P < 0.05$). With regard to metabolic parameters, the individuals in the higher MUAC quartiles showed higher levels of SBP, DBP, BMI, waist circumference or waist-to-hip ratio, fasting glucose, insulin, HOMA-IR, CRP, triglycerides (all $P < 0.001$), and LDL cholesterol ($P = 0.003$). Conversely, the subjects with larger MUAC displayed lower adiponectin and HDL cholesterol levels (both $P < 0.001$). However, elevated MUAC exhibited no association with the levels of total cholesterol in this study.

Association between MUAC and MetS

Partial correlation analysis showed that MUAC has the strongest correlation with waist circumference after adjusting for age and gender (Table 2). Additionally, MUAC was also strongly correlated with BMI and insulin.

As presented in Figure 1, the prevalence of MetS, overweight, and central obesity raised sharply from the small MUAC group to the large MUAC group both in men and women (Fig. 1). In addition, women seem to have a higher prevalence of MetS and central obesity compared with men. Furthermore, with the accumulation of MetS components, MUAC was gradually increased both in males and females (Fig. 2). It is worth noting that as the components of MetS increased one by one, the relationship between MUAC and MetS was statistically significant, both in subjects with MetS (number of MetS components ≥ 3) and the subjects without MetS (number of MetS components < 3). On this ground, MUAC may be directly or indirectly related to each MetS component.

As is shown in Table 3, the ORs for MetS and its components were significantly higher with increasing MUAC quartiles. In the largest MUAC quartile, the ORs (95% CIs) were 5.71 (4.97-6.56) for MetS, 14.70 (12.49-17.31) for central obesity, 2.23 (1.90-2.62) for elevated triglycerides, 2.11 (1.86-2.40) for elevated blood pressure, 2.06 (1.79-2.37) for reduced HDL cholesterol, 1.54 (1.35-1.76) for elevated fasting glucose after adjusting for age, gender, smoking, alcohol consumption, educational attainment, physical activity, CRP, adiponectin, and self-reported CVD (referencing to 1.00) (Model 2, all P for trend < 0.001). Further adjustment for HOMA-IR, CRP, and adiponectin, the OR was 3.24 (Model 3, 95% CI 2.79-3.76, P for trend < 0.001) for MetS in the highest MUAC quartile. Moreover, the correlation between MUAC and MetS was still statistically significant by additional adjustment for BMI (Model 4, OR 1.77; 95% CI 1.51-2.09; P for trend < 0.001).

Discussion

We observed a significant positive association between MUAC and MetS in a large-scale population study. Furthermore, we found that MUAC is strongly correlated with waist circumference, lipid parameters, blood pressure, and insulin, which indicated that abnormal local fat depot is a potential screening index for identifying metabolic disorders.

BMI and waist circumference are common screening measures for identifying individuals with abnormal distribution of body fat. Nevertheless, BMI cannot provide accurate information about the local distribution of body fat and it is difficult to obtain height and weight for patients who cannot stand. As for waist circumference, the deficiency of daily application lies in the big difference between preprandial and postprandial measurements. In view of the above reasons, MUAC began to show diagnostic value for assessing nutritional status. Compared to other anthropometric measurements, MUAC is not only easier to obtain, but also has other advantages such as being more accurate, convenient, and low-cost. Small MUAC has shown excellent performance in assessing malnutrition and predicting mortality both in children[17] and older individuals[18, 19]. More recently, large MUAC has been recognized as a valid tool for detecting overweight and obesity in children and adolescents[20, 21]. However, the study about whether MUAC is associated with obesity-related metabolic abnormality, such as MetS, is scarce. Currently, we found that MUAC, as a proxy of upper-body subcutaneous adipose, was positively associated with MetS. Moreover, consistent with previous studies[12], we found large MUAC also tightly correlated with central obesity, elevated blood pressure, and low HDL cholesterol.

The abnormal accumulation of fat affects adipose tissue metabolic capacities, endocrine, and immune function and leads to altered production of lipid mediators, adipokines, pro- or anti-inflammatory cytokines, and impaired signaling pathways that contribute to obesity-related metabolic abnormality[22]. Obesity increases the flux of fatty acids from adipose tissue to peripheral tissues[23]. The increased FFAs derived from adipose tissue is predominantly mediated by the resistance of adipose tissue to the anti-lipolytic effects of insulin[24]. There is compelling evidence that abnormally increased visceral fat is a maker of excessive systemic FFAs release[25], but it is worth noting that it is upper-body subcutaneous fat released the majority of FFAs[26]. Large MUAC means excessive subcutaneous fat accumulation, which contributes a greater portion of the fatty acids released into the circulation. Circulating FFAs is a crucial mediator in the development of metabolic disorders[27]. Elevated plasma FFAs induce insulin resistance, inflammation, and increase the synthesis and ectopic deposition of triglycerides[28-31]. Concomitantly, excessive FFAs also affect glucose metabolism by inhibiting glucose uptake, oxidation, glycogen synthesis, and increasing output hepatic glucose[32]. Additionally, increased FFAs can trigger oxidative stress which is an early instigator of MetS[33], and endoplasmic reticulum stress which intersects with many different inflammatory and stress signaling pathways by unfolded protein response[34-36]. The excess FFAs release derived from excess accumulation of arm subcutaneous adipose might be a potential mechanism to partly explain the correlation between MUAC and MetS.

Moreover, the adipose tissue is not only a depot of excess energy but also a highly active metabolic endocrine organ that secretes numerous biologically active molecules, which are collectively termed adipokines[37]. When adipose tissue expands, the capacity of adipocytes to function as endocrine cells and secrete various adipokines is altered in individuals with obesity and MetS[33, 38]. These abnormal levels of adipokine are linked to insulin resistance, impaired triglyceride storage and increased fatty acids in circulation[39]. Furthermore, as fat accumulation, substantial infiltration of immune cells occurs, and there is a specific crown-like disposition of macrophages around single necrotic adipocytes in obese people[40] and subjects with MetS[38]. Subsequently, proinflammatory pathways were activated, and

certain proinflammatory cytokines and chemokines were overflowed that result in low-grade inflammation and insulin resistance[22, 38]. In line with previous study[12], our findings also observed that MUAC is positively correlated with CRP. Overall, adipose dysfunctions, inflammation, and stress linking mid-upper arm obesity to insulin resistance and MetS.

To our knowledge, this is the first study to evaluate the association between MUAC and Mets among large-scale middle-aged and older people. The major strength of this study is the analysis based on a large sample. Potential covariates were strictly controlled in the analysis, so as to eliminated the possibility of residual confounding effects.

This study has several limitations. For one thing, we did not measure the tissue composition of the mid-upper arm. Due to the limitations of epidemiological screening conditions, we could not quantify adipose accumulation by more accurate radiographic measures. Therefore, the amount and size of subcutaneous adipocyte and muscle fat are not clear. For another, due to the present study is a cross-sectional analysis, we cannot draw the causality from our findings. Additionally, it is still unclear whether our findings in middle-aged/older Chinese subjects can be generalized to younger populations or individuals of other ethnicities.

In brief, our study observed that MUAC is positively associated with MetS even after adjustment for potential covariates. These findings provide a novel insight into the association between upper-body obesity and MetS, and a potential screening tool for identifying individuals with MetS. Further investigation of the underlying pathophysiological mechanism is needed to explain the positive association of mid-upper arm subcutaneous fat with MetS.

Conclusion

In summary, in this population-based study, there is a significantly positive association between MUAC and MetS in Chinese middle-aged and elderly individuals. Our results indicated that MUAC in the routine clinical measurement is necessary to screen for metabolic disorders, including dyslipidemia, dysglycemia, and elevated blood pressure.

Declarations

Acknowledgements

The authors thank all subjects who participated in the study and hospital staffs for their contribution in sample and data collection.

Funding

This work was supported by the National Natural Science Foundation of China (81670743 to QS, 81370953 to ZY), Shanghai Health System Outstanding Young Talents Training Program (XYQ2013098

to ZY), Shanghai Health and Family Planning Commission (21740173 to LQ). The funding body had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

The study is a part of the Risk Evaluation of cAncers in Chinese diabeTic Individuals: A lONgitudinal (REACTION) study. All data generated and analysed during this study are included in REACTION study. The data are held in a secure and confidential database which can only be assessed by members of the REACTION group. The REACTION has a website <http://www.rjh.com.cn/pages/neifenmike/REACTION/index.shtml>.

Author information

Affiliations

Department of Endocrinology, Xinhua Hospital Chongming Branch, Shanghai Jiaotong University School of Medicine, 25 Nanmen Road, Shanghai, China

Jie Shi, Zhen Yang, Yixin Niu, Hongxia Gu, Li Qin & Qing Su

Department of Endocrinology, Xinhua Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

Jie Shi, Zhen Yang, Yixin Niu, Weiwei Zhang, Xiaoyong Li, Hongmei Zhang, Ning Lin, Li Qin & Qing Su

Institute of Endocrinology and Diabetes, Department of Endocrinology and Metabolism, Huashan Hospital, Fudan University, Shanghai, China

Jie Wen

Shanghai Institute of Endocrinology and Metabolism, Department of Endocrine and Metabolic Diseases, Shanghai Clinical Center for Endocrine and Metabolic Diseases, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China

Guang Ning

Authors Contributions

ZY, LQ, and QS designed the study. YN, WZ, XL, HZ, NL, HG, JW, and GN recruited the subjects, processed samples, and contributed to acquisition of data. ZY, YN, WZ, NL and JS analyzed the data. JS wrote the manuscript. ZY revised the manuscript. All authors have read and approved the final manuscript.

Corresponding authors

Correspondence to Zhen Yang, Li Qin or Qing Su.

Ethics declarations

Ethics approval and consent to participate

The study protocol was approved by the Ethics Committee of Xinhua Hospital Affiliated to Shanghai Jiaotong University School of Medicine. Written informed consent was obtained from all participants.

Consent for publication

All authors have read the paper and agree that it can be published.

Competing interests

The authors declare that they have no competing interests.

Abbreviations

MUAC: mid-upper arm circumference

MetS: metabolic syndrome

FFAs: free fatty acids

BMI: body mass index

HDL: high-density lipoprotein

LDL: low-density lipoprotein

CVD: cardiovascular disease

T2D: type 2 diabetes

HOMA-IR: homeostasis model of assessment for insulin resistance

SBP: systolic blood pressure

DBP: diastolic blood pressure

OR: odds ratios

CI: confidence interval

CRP: C-reactive protein

References

1. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, Fruchart JC, James WP, Loria CM, Smith SC, Jr.: **Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity.** *Circulation* 2009, **120**(16):1640-1645.
2. Gu D, Reynolds K, Wu X, Chen J, Duan X, Reynolds RF, Whelton PK, He J: **Prevalence of the metabolic syndrome and overweight among adults in China.** *Lancet* 2005, **365**(9468):1398-1405.
3. He Y, Jiang B, Wang J, Feng K, Chang Q, Fan L, Li X, Hu FB: **Prevalence of the metabolic syndrome and its relation to cardiovascular disease in an elderly Chinese population.** *J Am Coll Cardiol* 2006, **47**(8):1588-1594.
4. Koutsari C, Ali AH, Mundi MS, Jensen MD: **Storage of circulating free fatty acid in adipose tissue of postabsorptive humans: quantitative measures and implications for body fat distribution.** *Diabetes* 2011, **60**(8):2032-2040.
5. Nielsen S, Guo Z, Johnson CM, Hensrud DD, Jensen MD: **Splanchnic lipolysis in human obesity.** *J Clin Invest* 2004, **113**(11):1582-1588.
6. Basu A, Basu R, Shah P, Vella A, Rizza RA, Jensen MD: **Systemic and regional free fatty acid metabolism in type 2 diabetes.** *Am J Physiol Endocrinol Metab* 2001, **280**(6):E1000-E1006.
7. Liang J, Teng F, Li Y, Liu X, Zou C, Wang Y, Li H, Qi L: **neck circumference and insulin resistance in Chinese adults: the Cardiometabolic Risk in Chinese (CRC) Study.** *Diabetes care* 2013, **36**(9):e145-e146.
8. Yang G-R, Yuan S-Y, Fu H-J, Wan G, Zhu L-X, Bu X-L, Zhang J-D, Du X-P, Li Y-L, Ji Y *et al*: **Neck circumference positively related with central obesity, overweight, and metabolic syndrome in Chinese subjects with type 2 diabetes: Beijing Community Diabetes Study 4.** *Diabetes care* 2010, **33**(11):2465-2467.
9. Jensen MD: **Role of body fat distribution and the metabolic complications of obesity.** *J Clin Endocrinol Metab* 2008, **93**(11 Suppl 1):S57-S63.
10. Boden G: **Fatty acid-induced inflammation and insulin resistance in skeletal muscle and liver.** *Curr Diab Rep* 2006, **6**(3):177-181.
11. Berk PD: **Regulatable fatty acid transport mechanisms are central to the pathophysiology of obesity, fatty liver, and metabolic syndrome.** *Hepatology* 2008, **48**(5):1362-1376.
12. Hou Y, Jia X, Xuan L, Zhu W, Deng C, Wang L, Zhao Z, Li M, Lu J, Xu Y *et al*: **Association between mid-upper arm circumference and cardiometabolic risk in Chinese population: a cross-sectional study.** *BMJ Open* 2019, **9**(9):e028904.

13. Ning G: **Risk Evaluation of cAncers in Chinese diabeTic Individuals: a IONgitudinal (REACTION) study.** *J Diabetes* 2012, **4**(2):172-173.
14. Yang Z, Yan C, Liu G, Niu Y, Zhang W, Lu S, Li X, Zhang H, Ning G, Fan J *et al*: **Plasma selenium levels and nonalcoholic fatty liver disease in Chinese adults: a cross-sectional analysis.** *Sci Rep* 2016, **6**:37288.
15. Qin L, Yang Z, Gu H, Lu S, Shi Q, Xing Y, Li X, Li R, Ning G, Su Q: **Association between serum uric acid levels and cardiovascular disease in middle-aged and elderly Chinese individuals.** *BMC Cardiovasc Disord* 2014, **14**:26.
16. <Anthropometry Procedures Manual .pdf>.
17. Berkley J, Mwangi I, Griffiths K, Ahmed I, Mithwani S, English M, Newton C, Maitland K: **Assessment of severe malnutrition among hospitalized children in rural Kenya: comparison of weight for height and mid upper arm circumference.** *Jama* 2005, **294**(5):591-597.
18. Schaap LA, Quirke T, Wijnhoven HAH, Visser M: **Changes in body mass index and mid-upper arm circumference in relation to all-cause mortality in older adults.** *Clin Nutr* 2018, **37**(6 Pt A):2252-2259.
19. Wijnhoven HA, van Bokhorst-de van der Schueren MA, Heymans MW, de Vet HC, Kruizenga HM, Twisk JW, Visser M: **Low mid-upper arm circumference, calf circumference, and body mass index and mortality in older persons.** *J Gerontol A Biol Sci Med Sci* 2010, **65**(10):1107-1114.
20. Talma H, van Dommelen P, Schweizer JJ, Bakker B, Kist-van Holthe JE, Chinapaw JMM, Hirasing RA: **Is mid-upper arm circumference in Dutch children useful in identifying obesity?** *Arch Dis Child* 2019, **104**(2):159-165.
21. Chaput JP, Katzmarzyk PT, Barnes JD, Fogelholm M, Hu G, Kuriyan R, Kurpad A, Lambert EV, Maher C, Maia J *et al*: **Mid-upper arm circumference as a screening tool for identifying children with obesity: a 12-country study.** *Pediatr Obes* 2017, **12**(6):439-445.
22. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr.: **Obesity is associated with macrophage accumulation in adipose tissue.** *The Journal of clinical investigation* 2003, **112**(12):1796-1808.
23. Arner P, Langin D: **Lipolysis in lipid turnover, cancer cachexia, and obesity-induced insulin resistance.** *Trends Endocrinol Metab* 2014, **25**(5):255-262.
24. Saltiel AR, Kahn CR: **Insulin signalling and the regulation of glucose and lipid metabolism.** *Nature* 2001, **414**(6865):799-806.
25. Guo Z, Hensrud DD, Johnson CM, Jensen MD: **Regional postprandial fatty acid metabolism in different obesity phenotypes.** *Diabetes* 1999, **48**(8):1586-1592.
26. Martin ML, Jensen MD: **Effects of body fat distribution on regional lipolysis in obesity.** *The Journal of clinical investigation* 1991, **88**(2):609-613.
27. Boden G: **Obesity, insulin resistance and free fatty acids.** *Curr Opin Endocrinol Diabetes Obes* 2011, **18**(2):139-143.

28. Kim F, Pham M, Luttrell I, Bannerman DD, Tupper J, Thaler J, Hawn TR, Raines EW, Schwartz MW: **Toll-like receptor-4 mediates vascular inflammation and insulin resistance in diet-induced obesity.** *Circ Res* 2007, **100**(11):1589-1596.
29. Shi H, Kokoeva MV, Inouye K, Tzamelis I, Yin H, Flier JS: **TLR4 links innate immunity and fatty acid-induced insulin resistance.** *The Journal of clinical investigation* 2006, **116**(11):3015-3025.
30. Morino K, Petersen KF, Shulman GI: **Molecular mechanisms of insulin resistance in humans and their potential links with mitochondrial dysfunction.** *Diabetes* 2006, **55** Suppl 2:S9-s15.
31. Hotamisligil GS: **Inflammation, metaflammation and immunometabolic disorders.** *Nature* 2017, **542**(7640):177-185.
32. Bergman RN, Ader M: **Free fatty acids and pathogenesis of type 2 diabetes mellitus.** *Trends Endocrinol Metab* 2000, **11**(9):351-356.
33. Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, Nakayama O, Makishima M, Matsuda M, Shimomura I: **Increased oxidative stress in obesity and its impact on metabolic syndrome.** *The Journal of clinical investigation* 2004, **114**(12):1752-1761.
34. Guo W, Wong S, Xie W, Lei T, Luo Z: **Palmitate modulates intracellular signaling, induces endoplasmic reticulum stress, and causes apoptosis in mouse 3T3-L1 and rat primary preadipocytes.** *Am J Physiol Endocrinol Metab* 2007, **293**(2):E576-586.
35. Ozcan U, Cao Q, Yilmaz E, Lee AH, Iwakoshi NN, Ozdelen E, Tuncman G, Gorgun C, Glimcher LH, Hotamisligil GS: **Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes.** *Science* 2004, **306**(5695):457-461.
36. Hotamisligil GS: **Endoplasmic reticulum stress and the inflammatory basis of metabolic disease.** *Cell* 2010, **140**(6):900-917.
37. Kershaw EE, Flier JS: **Adipose tissue as an endocrine organ.** *J Clin Endocrinol Metab* 2004, **89**(6):2548-2556.
38. Bremer AA, Devaraj S, Afify A, Jialal I: **Adipose tissue dysregulation in patients with metabolic syndrome.** *The Journal of clinical endocrinology and metabolism* 2011, **96**(11):E1782-1788.
39. Guilherme A, Virbasius JV, Puri V, Czech MP: **Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes.** *Nat Rev Mol Cell Biol* 2008, **9**(5):367-377.
40. Cancello R, Henegar C, Viguerie N, Taleb S, Poitou C, Rouault C, Coupaye M, Pelloux V, Hugol D, Bouillot JL *et al*: **Reduction of macrophage infiltration and chemoattractant gene expression changes in white adipose tissue of morbidly obese subjects after surgery-induced weight loss.** *Diabetes* 2005, **54**(8):2277-2286.

Tables

Table 1 Characteristics of study participants according to MUAC quartiles

Characteristics	Q1 (n=2524)	Q2 (n=2409)	Q3 (n=2427)	Q4 (n=2427)	P value for trend
	≤26.70	26.71-28.60	28.61-30.50	≥30.51	
MUAC (cm)	24.93±1.53	27.78±0.56	29.61±0.53	32.53±1.72	<0.001
Age (years) ^b	55.59±8.21	56.06±7.81	56.53±7.62	56.47±7.59	<0.001
Male (%) ^b	24.0	29.1	37.1	39.1	<0.001
Smoking (yes, %)	14.3	17.2	18.1	19.9	<0.001
Alcohol (yes, %)	18.1	24.5	26.0	28.1	<0.001
Educational attainment (%)					
0-6 years	20.7	21.7	21.8	25.4	0.003
7-9 years	50.0	49.1	48.6	48.2	
≥10 years	29.3	29.2	29.6	26.4	
Physical activity (%)					
Low	70.8	72.5	71.6	72.2	0.281
Moderate	21.4	19.6	21.2	19.7	
High	7.8	7.9	7.2	8.1	
SBP (mm Hg)	126.96±19.00	129.72±18.42	132.23±17.76	134.55±18.94	<0.001
DBP (mm Hg)	79.02±9.51	80.29±9.71	81.13±9.50	82.50±9.40	<0.001
BMI (kg/m ²)	22.79±2.90	24.47±2.48	25.46±2.61	26.94±3.08	<0.001
Waist circumference (cm)	78.38±11.41	82.94±8.49	86.54±8.75	91.11±8.77	<0.001
Waist to hip ratio	0.86±0.08	0.87±0.09	0.89±0.17	0.90±0.07	<0.001
Fasting glucose (mmol/L)	6.20±1.82	6.23±1.73	6.33±1.65	6.38±1.52	<0.001
2-h glucose (mmol/L)	8.30±4.01	8.61±3.94	8.91±3.79	9.18±3.73	<0.001
Insulin (mU/L) ^c	6.25(3.90-7.20)	7.18(4.50-8.50)	7.75(5.20-9.20)	9.40(6.10-11.50)	<0.001
HOMA-IR ^c	1.70(0.98-1.99)	2.00(1.22-2.45)	2.22(1.38-2.68)	2.70(1.65-3.38)	<0.001
CRP (ug/ml)	4.79±5.74	4.94±5.85	5.13±6.12	5.24±6.27	<0.001
Adiponectin (ug/ml)	12.34(7.92-15.67)	10.93(6.72-13.15)	8.17(5.24-11.33)	6.29(4.78-9.75)	<0.001
Triglycerides (mmol/L) ^c	1.44(0.83-1.66)	1.67(0.94-1.94)	1.83(1.03-2.13)	1.96(1.12-2.34)	<0.001
Total cholesterol (mmol/L)	4.64±1.05	4.62±1.03	4.65±1.04	4.67±1.03	0.291
LDL cholesterol (mmol/L)	2.58±0.78	2.59±0.76	2.63±0.77	2.65±0.76	0.003
HDL cholesterol (mmol/L)	1.32±0.34	1.23±0.32	1.19±0.30	1.16±0.28	<0.001
Comorbidities (%)					
Hypertension	20.8	25.0	31.4	36.9	<0.001
Hyperlipidemia	2.8	3.4	5.8	6.2	<0.001
Diabetes	8.8	9.3	10.8	10.8	0.036
CVD ^d	2.6	3.9	3.2	4.1	0.014

^a Data are means ± SD, median (interquartile range), or percentage; P value was calculated after adjustment for age and gender. ^b Not adjusted for itself. ^c These variables were log transformed before analysis. ^d Self-reported CVD including stroke and coronary heart disease.

Table 2 Partial correlation coefficients among MUAC and other clinical parameters

Variable	MUAC	
	r	P
BMI	0.334	<0.001
Waist circumference	0.437	<0.001
SBP	0.124	<0.001
DBP	0.123	<0.001
Fasting plasma glucose	0.045	0.043
2-h post-loading glucose	0.074	<0.001
Insulin	0.348	<0.001
HOMA-IR	0.134	<0.001
CRP	0.110	0.008
Adiponectin	-0.147	<0.001
HDL cholesterol	-0.176	<0.001
LDL cholesterol	0.024	0.016
Triglycerides	0.138	<0.001

All correlation coefficients were calculated after adjustment for age and gender.

Table 3 Adjusted ORs (95% CI) of MetS and its components according to the quartiles of MUAC

	ORs (95% CI)				P value for trend
	Q1	Q2	Q3	Q4	
MetS					
Model 1 ^a	1.00	1.79(1.60-2.02)	3.28(2.91-3.70)	5.77(5.08-6.54)	<0.001
Model 2 ^b	1.00	1.77(1.56-2.01)	3.25(2.85-3.70)	5.71(4.97-6.56)	<0.001
Model 3 ^c	1.00	1.55(1.35-1.78)	2.49(2.17-2.86)	3.24(2.79-3.76)	<0.001
Model 4 ^d	1.00	1.23(1.07-1.42)	1.68(1.45-1.95)	1.77(1.51-2.09)	<0.001
Central obesity					
Model 1 ^a	1.00	2.50(2.22-2.81)	5.67(5.00-6.43)	15.31(13.17-17.81)	<0.001
Model 2 ^b	1.00	2.47(2.18-2.81)	5.52(4.82-6.31)	14.70(12.49-17.31)	<0.001
Model 3 ^c	1.00	2.29(2.00-2.61)	4.69(4.08-5.40)	10.57(8.93-12.52)	<0.001
Model 4 ^d	1.00	1.41(1.20-1.65)	2.03(1.71-2.40)	3.46(2.81-4.25)	<0.001
Elevated blood pressure					
Model 1 ^a	1.00	1.27(1.13-1.43)	1.66(1.48-1.87)	2.15(1.91-2.42)	<0.001
Model 2 ^b	1.00	1.26(1.11-1.43)	1.64(1.45-1.86)	2.11(1.86-2.40)	<0.001
Model 3 ^c	1.00	1.18(1.03-1.34)	1.45 (1.27-1.65)	1.65 (1.44-1.88)	<0.001
Model 4 ^d	1.00	1.04(0.91-1.18)	1.16(1.02-1.33)	1.17(1.01-1.35)	<0.001
Elevated triglycerides					
Model 1 ^a	1.00	1.44(1.23-1.69)	1.81(1.55-2.11)	2.34(2.02-2.72)	<0.001
Model 2 ^b	1.00	1.39(1.17-1.65)	1.81(1.54-2.14)	2.23(1.90-2.62)	<0.001
Model 3 ^c	1.00	1.26(1.06-1.50)	1.49(1.26-1.77)	1.52(1.28-1.80)	<0.001
Model 4 ^d	1.00	1.21(1.02-1.45)	1.36(1.14-1.64)	1.40(1.17-1.67)	<0.001
Reduced HDL cholesterol					
Model 1 ^a	1.00	1.45(1.27-1.66)	1.70(1.49-1.93)	2.18(1.91-2.48)	<0.001
Model 2 ^b	1.00	1.36(1.18-1.58)	1.68(1.45-1.94)	2.06(1.79-2.37)	<0.001
Model 3 ^c	1.00	1.29(1.11-1.50)	1.50(1.30-1.74)	1.68(1.45-1.95)	<0.001
Model 4 ^d	1.00	1.23(1.06-1.43)	1.38(1.19-1.61)	1.46(1.24-1.71)	<0.001
Elevated fasting glucose					
Model 1 ^a	1.00	1.28(1.13-1.45)	1.48(1.32-1.72)	1.51(1.33-1.73)	<0.001
Model 2 ^b	1.00	1.28(1.12-1.47)	1.51(1.04-1.34)	1.54(1.35-1.76)	<0.001
Model 3 ^c	1.00	1.23(1.08-1.41)	1.41(1.23-1.61)	1.38(1.20-1.57)	<0.001
Model 4 ^d	1.00	1.14(0.99-1.32)	1.15(1.00-1.32)	1.25(1.09-1.44)	<0.001

^a Model 1 was adjusted for age and gender. ^b Model 2 was further adjusted for smoking, alcohol drinking, educational attainment, physical activity, and self-reported CVD. ^c Model 3 was further adjusted for CRP, adiponectin, and HOMA-IR. ^d Model 4 was further adjusted for BMI. Q, quartile. Q1 is the reference group. Multivariate logistic regression models were used to estimate the odds ratios (ORs) with corresponding 95% confidence intervals (CIs) for MetS and its components. The final model is statistically significant ($\chi^2 = 34.927$, $P < 0.001$).

Figures

Figure 1

Prevalence of the MetS, overweight, and central obesity by quartiles of MUAC among 9787 participants. The quartiles of MUAC were calculated by gender respectively. In male, the cutoffs of MUAC were ≤ 27.40 , 27.41-29.25, 29.26-31.10, and ≥ 31.11 cm; in female, the cutoffs were ≤ 26.45 , 26.46-28.40, 28.41-30.30, and ≥ 30.31 cm. From 1st to 4th quartile (Q1 to Q4), the prevalence of MetS were 32.0%; 45.6%; 59.3%; 72.4% ($P = 0.007$) in men and 35.4%; 49.4%; 64.5%; 75.1% ($P < 0.001$) in women, respectively. The prevalence of overweight was 30.2%; 48.2%; 66.7%; 83.6% in men, and 24.9%; 43.3%; 67.0%; 83.3% in women, both $P < 0.001$. The prevalence of central obesity was 29.5%; 51.6%; 70.4%; 84.7% in men, and 33.0%; 54.5%; 72.3%; 88.4% in women, both $P < 0.001$. MUAC, mid-upper arm circumference; MetS, metabolic syndrome; Q, quartile.

Figure 2

MUAC according to the number of MetS components. Data are shown as the means $\pm$ SEM. The mean values of MUAC for those with none to five components were 27.14, 28.01, 28.63, 29.70, 30.21, 30.71 cm for male, and 26.62, 27.00, 27.83, 28.79, 29.43, 29.8 cm for female, respectively. *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$; ****: $P < 0.0001$. MUAC, mid-upper arm circumference; MetS, metabolic syndrome.