

Rapid assessments to differentiate dementia using plasma biomarkers in primary care: a case control study

Ping-Huang Tsai (✉ phtsai@nycu.edu.tw)

National Yang-Ming University: National Yang Ming Chiao Tung University - Yangming Campus

<https://orcid.org/0000-0002-7236-0211>

Li-Cho Hsu

National Yang Ming Chiao Tung University Hospital

Hsuan-Ming Tsao

National Yang Ming Chiao Tung University Hospital

Shieh-Yueh Yang

MagQu Co Ltd.

Kuen-Lin Chen

National Chung Hsing University

Research

Keywords: Dementia plasma biomarker panel, Dementia, Alzheimer's disease, amyloid-beta protein, Tau protein

Posted Date: July 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-673269/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

The aging of society has increased the incidence of dementia, which is more common among older individuals. Older individuals are typically cared for by primary care providers in hospitals. However, more than 60% of patients with early-stage dementia are unrecognized in primary care. Several groups have developed dementia screening tools for primary care purposes. In this work, assessments based on plasma biomarkers for differentiating among various types of dementia were developed for primary care applications.

Methods

Forty-six patients with very mild dementia (VMD) or mild cognitive impairment (MCI), fifty patients with Alzheimer's disease (AD), and four patients with non-AD dementia were enrolled. Plasma amyloid-beta 1–40 ($A\beta_{1-40}$), $A\beta_{1-42}$, total Tau (T-Tau), and phosphorylated Tau (p-Tau181) were assayed using immunomagnetic reduction for each subject.

Results

The results show that non-AD dementia can be discriminated from other forms of dementia using plasma $A\beta_{1-40}$, with a cutoff value of 50.03 pg/ml resulting in an area under the curve (AUC) of 0.794. The plasma $A\beta_{1-42}$ -to- $A\beta_{1-40}$ ratio can serve as an index for discriminating AD from VMD and MCI, with a cutoff value of 0.3015 resulting in an AUC of 0.674.

Conclusions

A biomarker panel measuring the levels of plasma $A\beta_{1-40}$ and the ratio of $A\beta_{1-42}$ to $A\beta_{1-40}$ could potentially assist primary care practitioners in evaluating whether a patient suffers from non-AD dementia, AD, or VMD and MCI.

Trial registration

The research was approved by the IRB of NYCUH (IRB No.2017A033 and No. 2018A004).

Background

The prevalence of dementia among older people has been increasing. In a 2013 national survey of dementia prevalence in Taiwan, the prevalence of dementia among individuals older than 65 years was 8% [1]. Dementia represents an essential public health issue. In general, older individuals routinely visit Primary Care (PC) to receive treatment rather than the Neurological Division, particularly for their first neurological screenings. The appropriate screening and identification of dementia among older individuals in the PC setting are critical to ensure the timely initiation of management, treatment, or

intervention strategies for patients with mild (MCI) or subjective cognitive impairment. However, screening for cognitive impairment or dementia is not commonly performed by PC practitioners. Hypertension, hyperlipidemia, or type II diabetes are the common diseases treated by PC practitioners [2–4], whereas assessments of cognitive impairment are not regularly performed in PC settings. Symptoms of cognitive impairment may not be apparent during a routine PC visit, and cognitive impairment is traditionally diagnosed in a PC setting based on clinical suspicion in response to the patient's symptoms or caregivers' concerns [5]. From 1992 to 2006, the reported accuracy of mild dementia diagnoses among PC providers was 14–69% [6–13]. The missed or delayed diagnosis of dementia occurs frequently.

In the early 2000s, PC physicians utilized cognitive tests or evaluated patients' functional abilities to assess mild dementia. Cruz-Orduña et al. reported that the sensitivity of detection based on patient-reported or suspected cognitive impairment over a 49-year period increased to 75% with the use of the Mini-Mental State Examination (MMSE) or the Functional Activities Questionnaire (FAQ) [14].

Chronic conditions have been demonstrated to contribute to dementia prediction [15–17]. Tsai et al. integrated various risk factors, such as age, sex, body mass index, education, stroke, diabetes, hypertension, hyperlipidemia, head trauma, and depression, with the evaluation of cognitive tests for the assessment of dementia in PC settings [18], which resulted in a model capable of differentiating patients with dementia from normal controls with a sensitivity greater than 90%.

Significant progress in enhancing the sensitivity of dementia diagnoses in PC settings has identified additional diagnostic needs, such as the identification of information that would allow PC physicians to recognize which type of dementia a patient is presenting to ensure the timely application of appropriate management or treatment strategies.

Although the efficacy of monoclonal antibodies for Alzheimer's disease (AD) treatment remains uncertain, an increasing number of studies have explored the diagnosis of AD using biomarkers. The assessment of plasma biomarkers represents a convenient and cheap diagnostic method compared with the current gold standard of positron emission tomography (PET), which may allow for plasma biomarkers to be tested annually. Several papers reported by neurologists demonstrated the feasibility of assaying plasma biomarkers for identifying AD [19–22], Parkinson's disease [23–25], or frontotemporal dementia [26]. These results suggest the possibility of developing assessments that use plasma biomarkers to differentiate among various types of dementia in PC settings. In this work, 100 patients with various types of dementia were recruited. Plasma amyloid-beta 1–40 ($A\beta_{1-40}$), amyloid-beta 1–42 ($A\beta_{1-42}$), total Tau (T-Tau), and phosphorylated Tau (p-Tau181) were assayed using immunomagnetic reduction for each subject. Rapid assessments for the differentiation of dementia types using plasma biomarkers were developed for the PC setting.

Methods And Materials

Recruitment of subjects

All patients aged 50 years or older were recruited from neurological clinics at National Yang Ming Chiao Tung University Hospital (NYCUH) (former National Yang-Ming University Hospital). All participants received cognitive assessments. Dementia was diagnosed according to the National Institute on Aging and Alzheimer's Association (NIA-AA) criteria [27]. Trained research assistants also administered the Chinese version of the MMSE [28], which features a total score of 30. The Clinical Dementia Rating (CDR) was also used to determine the severity of dementia after separate semi-structured interviews with the patient and a knowledgeable informant were conducted by a neurologist or psychologist. The CDR scores are categorized as follows: 0 for normal, 0.5 for MCI or very mild dementia (VMD) [29, 30], 1 for mild dementia, 2 for moderate dementia, and 3 for severe dementia [31]. Individuals with VMD were classified according to the presentation of mild impairment in two or more cognitive domains and a slight decline in daily function; cognitive deficits sufficient to interfere with independence in daily life, community affairs, or at-home hobbies; or based on the outcome of the CDR. MCI was diagnosed according to the NIA-AA-recommended criteria, defined as a change in cognition with impairment in one or more cognitive domains but no evidence of impairment in social or occupational functions as assessed by the CDR, activities of daily living (ADL) [32], and instrumental activities of daily living (IADL) [33]. The final diagnosis was reviewed after one year of follow-up.

The research was approved by the Institutional Review Board (IRB) of NYCUH (IRB No.2017A033 and No. 2018A004). Written informed consent was obtained from all participants in the study. All patients with dementia and their proxies provided written informed consent. Informed consent could only be signed directly by patients with MCI because the IRB agreed that patients with MCI, who suffered from cognitive impairments in one or more cognitive domains but presented with no evidence of impairment in social or occupational functions, were capable of understanding the study procedures and non-invasive assessments being performed in this study. We also explained this informed consent requirement to any proxies who accompanied MCI patients.

All participants received medical, neurological, neuropsychological, and psychiatric assessments and blood examinations. The neurological assessments performed for each participant included a cerebral computed tomography scan to exclude intracranial pathologies (i.e., brain tumors or stroke) that may have contributed to cognitive decline.

Plasma preparation

A 9-ml EDTA tube (455036, Greiner) was used to draw blood. No fasting was required for the blood draw. The tube was gently inverted 10 times immediately after the blood draw. A swing-out (bucket) rotor was used to centrifuge the blood at 15–25°C at 1500–2500 × *g* for 15 minutes. Then, 1 ml plasma (supernatant) was transferred to a fresh 1.5-ml Eppendorf tube. All aliquoted plasma samples were stored at –80°C within 4.5 hours after the blood draw and prior to performing biomarker assays.

Assays of plasma biomarkers

The frozen human plasma sample was moved from -80°C to wet ice and room temperature. To assaying $\text{A}\beta_{1-40}$, T-Tau, and p-Tau181 (Tau phosphorylated at threonine 181), 40 μl plasma was mixed with 80 μl of the respective reagent (MF-AB0-0060, MF-TAU-0060, and MF-PT1-0060; MagQu). For assaying $\text{A}\beta_{1-42}$, 60 μl plasma was mixed with 60 μl reagent (MF-AB2-0060, MagQu). For each batch of measurements, calibrators (CA-DEX-0060, CA-DEX-0080, MagQu) and control solutions were used. An immunomagnetic reduction (XacPro-S361, MagQu) analyzer was utilized. For each sample, duplicate measurements were performed for each biomarker. The averaged concentration of the duplicate measurements was reported.

Statistical analysis

Continuous variables for each measurement are presented as the mean \pm standard deviation. SPSS (version 22.0) for Windows (SPSS Inc., Chicago, IL, USA) was used to perform statistical analyses. Baseline demographic characteristics, including age, MMSE score, and CDR-Sum of Boxes (SOB) scores, were coded as continuous variables and compared using a T-test to determine p -values. Receiver operating characteristic curves were analyzed for each plasma biomarker to explore cutoff values, sensitivity, specificity, and area under the curve (AUC) for the differentiation of various types of dementia. All statistical tests were two-tailed, and significance levels were established at p -values of less than 0.05.

Results

A total of 46 patients with VMD or MCI, referred to as VMD + MCI; 50 patients with AD; and 4 patients with non-AD dementia were enrolled. In the non-AD group, one patient had dementia with Lewy bodies, one patient had depression, one patient was folic acid deficient, and one patient had VMD with depression. Among the 46 VMD + MCI patients, 11 patients had VMD, and 35 patients had MCI. Notably, the 11 VMD patients progressed to AD within 3 years following their assessment in this study, indicating that VMD in these patients likely represented the early stages of AD. The demographic information of all enrolled subjects is listed in Table 1. Women comprised 50.0%, 50.0%, and 65.2% of the non-AD, AD, and VMD + MCI groups, respectively. The mean ages were 68.5 ± 12.4 years in the non-AD group, 78.4 ± 8.19 years in the AD group, and 72.2 ± 8.6 years in the VMD + MCI group. No significant differences in age were observed among the non-AD, AD, and VMD + MCI groups ($p > 0.05$). The CDR-SOB scores were 2.00 ± 1.78 , 3.77 ± 2.65 , 1.95 ± 1.27 for the non-AD, AD, and VMD + MCI groups, respectively. Patients with AD showed significantly higher scores on the CDR-SOB than non-AD and VMD + MCI patients ($p < 0.01$). The MMSE scores were 24.3 ± 5.7 , 20.0 ± 5.5 , and 23.9 ± 4.3 for the non-AD, AD, and VMD + MCI, respectively. AD shows significantly lower scores of MMSE than non-AD and VMD + MCI ($p < 0.01$).

Table 1
Demographic information of the enrolled subjects.

Group	Non-AD	AD	VMD + MCI		
			VMD	MCI	Combined
n (female%)	4 (50%)	50 (50.0%)	11 (63.6%)	35 (65.7%)	46 (65.2%)
Age (years)	68.5 ± 12.4	78.4 ± 8.19	74.6 ± 5.7	71.5 ± 9.3	72.2 ± 8.6
CDR-SOB	2.00 ± 1.78	3.77 ± 2.65	2.55 ± 1.06	1.76 ± 1.28	1.95 ± 1.27
MMSE	24.3 ± 5.7	20.0 ± 5.5	23.3 ± 4.2	24.1 ± 4.4	23.9 ± 4.3
Aβ ₁₋₄₀ (pg/ml)	48.99 ± 1.27	51.92 ± 4.16	54.74 ± 4.16	54.24 ± 4.20	54.36 ± 4.15
Aβ ₁₋₄₂ (pg/ml)	15.65 ± 1.06	16.41 ± 1.27	15.71 ± 0.97	15.93 ± 1.00	15.88 ± 0.99
T-Tau (pg/ml)	19.97 ± 2.29	21.58 ± 4.04	19.17 ± 2.18	19.73 ± 2.57	19.60 ± 2.47
p-Tau181 (pg/ml)	3.12 ± 0.48	3.41 ± 0.73	3.01 ± 0.54	3.19 ± 0.53	3.14 ± 0.53
AD: Alzheimer's disease; VMD: very mild dementia; MCI: mild cognitive impairment; CDR-SOB: clinical dementia ranking-sum of boxes; MMSE: Mini-Mental State Examination.; Aβ: amyloid-beta					

The measured Aβ₁₋₄₀ levels in the plasma were 48.99 ± 1.27 pg/ml for the non-AD group, 51.92 ± 4.16 pg/ml for the AD group, and 54.36 ± 4.15 pg/ml for the VMD + MCI group. The plasma Aβ₁₋₄₀ level in the non-AD group was significantly lower than those in the AD and VMD + MCI groups ($p < 0.01$). The plasma Aβ₁₋₄₂ levels were 15.65 ± 1.06 pg/ml for the non-AD group, 16.41 ± 1.27 pg/ml for the AD group, and 15.88 ± 0.99 pg/ml for the VMD + MCI group. The AD group showed a significantly higher Aβ₁₋₄₂ level than that for the VMD + MCI group ($p < 0.05$), but not compared with the level for the non-AD ($p > 0.05$). The T-Tau levels in plasma were 19.98 ± 2.29 pg/ml for the non-AD group, 21.58 ± 4.04 pg/ml for the AD group, and 19.60 ± 2.47 pg/ml for the VMD + MCI. The AD group presented significantly higher levels of plasma T-Tau than those of the non-AD and VMD + MCI groups ($p < 0.05$). The measured p-Tau181 levels in plasma for the non-AD (3.12 ± 0.48 pg/ml), AD (3.41 ± 0.73 pg/ml), and VMD + MCI (3.14 ± 0.53 pg/ml) groups are listed in Table 1. No significant differences in plasma p-Tau181 levels were observed among the non-AD, AD, and VMD + MCI groups ($p > 0.05$).

The VMD + MCI group included 11 VMD patients and 35 MCI patients. The VMD (74.6 ± 5.7 years) and MCI (71.5 ± 9.3 years) patients were aged matched. Except for the CDR-SOB scores ($p < 0.05$), no significant differences were observed between the VMD and MCI patients, including MMSE scores and plasma Aβ₁₋₄₀, Aβ₁₋₄₂, T-Tau and p-Tau181 levels ($p > 0.05$).

Discussion

The goal of this study was to develop a rapid assessment for the differentiation of different types of dementia using plasma biomarkers for the PC setting. As listed in Table 1, the non-AD group showed relatively lower levels of plasma $A\beta_{1-40}$ compared with those in the AD and VMD + MCI groups, suggesting that plasma $A\beta_{1-40}$ level might represent a promising index for discriminating non-AD patients from other groups. Through ROC curve analysis, the plasma $A\beta_{1-40}$ level the cutoff value for discriminating non-AD from AD and VMD + MCI was found to be 50.03 pg/ml. The corresponding clinical sensitivity and specificity were 75% and 100%. The AUC was 0.794.

An ROC curve analysis was performed using individual plasma biomarkers to differentiate AD from VMD + MCI, and the results are listed in Table 2. In addition, several groups have proposed that the combinations of plasma biomarkers might represent a more adequate index for discriminating among various dementia types or severities [19, 34]. The results of the ROC curve analysis using combined plasma biomarkers to differentiate AD from VMD + MCI are listed in Table 2.

Table 2
ROC curve analysis for differentiating AD from VMD + MCI using individual or combined plasma biomarkers.

Biomarker	Cutoff value	Sensitivity (95% CI)	Specificity (95% CI)	AUC	<i>p</i> -value
Tau	19.5 pg/ml	0.600 (0.452–0.736)	0.544 (0.390–0.691)	0.635	< 0.05
p-Tau181	3.26 pg/ml	0.540 (0.393–0.682)	0.587 (0.432–0.730)	0.600	> 0.05
$A\beta_{1-42}$	16.15 pg/ml	0.620 (0.472–0.754)	0.544 (0.390–0.691)	0.622	< 0.05
$A\beta_{1-42}$ -to- $A\beta_{1-40}$	0.3015	0.640 (0.492–0.771)	0.674 (0.520–0.805)	0.674	< 0.05
$A\beta_{1-42} \times$ Tau	303.5 pg ² /ml ²	0.640 (0.492–0.771)	0.544 (0.390–0.691)	0.631	< 0.05
$A\beta_{1-42} \times$ Tau/ $A\beta_{1-40}$	5.722 pg/ml	0.652 (0.472–0.734)	0.587 (0.432–0.730)	0.651	< 0.05

ROC: receiver operating characteristic; AD: Alzheimer's disease; VMD: very mild dementia; MCI: mild cognitive impairment; AUC: area under the curve; CI: confidence interval; $A\beta$: amyloid-beta

Among these biomarkers, the p-Tau181 level did not significantly differentiate AD from VMD + MCI ($p > 0.05$). However, published papers have demonstrated that plasma p-Tau181 represents a promising index

for differentiating AD patients from those with MCI due to AD, who comprised the VMD group in this study [35, 36]. Through a careful inspection of the data collected here, the p -value for the comparison of plasma p-Tau181 levels between AD (3.41 ± 0.73 pg/ml) and VMD (3.01 ± 0.54 pg/ml) patients was found to be less than 0.05, whereas the p -value was greater than 0.05 for the comparison of pTau181 levels between the AD and MCI (3.19 ± 0.53 pg/ml) groups, which is consistent with the results of previously published papers.

As listed in Table 2, the $A\beta_{1-42}$ -to- $A\beta_{1-40}$ ratio shows the highest AUC value (0.674), whereas $A\beta_{1-42}$ shows the lowest AUC value (0.622) for discriminating AD from VMD + MCI. This result suggests that the $A\beta_{1-42}$ -to- $A\beta_{1-40}$ ratio could represent the most effective index for differentiating AD from VMD + MCI. The cutoff value for the $A\beta_{1-42}$ -to- $A\beta_{1-40}$ ratio was established as 0.3015 to discriminate AD from VMD + MCI patients. The clinical sensitivity and specificity were 0.640 and 0.674, respectively.

The further discrimination of VMD from MCI patients included in the VMD + MCI group is not necessary for PC settings. Such discrimination should be conducted using neuropsychological tests performed by neuropsychologists or neurophysiologists. For example, as listed in Table 1, the VMD (3.08 ± 1.67) group showed a significantly higher CDR-SOB score than the MCI (1.76 ± 1.28 ; $p < 0.01$) group. An alternative method for discriminating between VMD and MCI is follow-up monitoring. All VMD patients progressed to AD within 3 years after the visit reported in this study.

A flow chart for rapid assessments using plasma biomarkers to differentiate among non-AD, AD, and VMD + MCI groups is illustrated in Fig. 1. First, plasma $A\beta_{1-40}$ levels were assayed. A measured plasma $A\beta_{1-40}$ level below 50.03 pg/ml would be highly indicative of non-AD. Otherwise, the subject is suspected of AD or VMD + MCI. Second, plasma $A\beta_{1-42}$ levels were assayed to obtain the ratio of $A\beta_{1-42}$ -to- $A\beta_{1-40}$, with a value above 0.3015 indicating AD and a value below 0.3015 indicating VMD or MCI.

The flow chart shown in Fig. 1 could assist PC physicians to easily assess the dementia types observed in patients, allowing for the PC physician to provide patients with adequate disease management strategies or refer the patient to neurologist care. Delayed interventions and treatments could be prevented among patients who visit a PC practitioner instead of the Neurological Division.

Conclusion

A rapid assessment for discriminating non-AD, AD, and VMD + MCI patients was demonstrated using plasma biomarkers. The plasma $A\beta_{1-40}$ level can be used to discriminate non-AD, followed by the assessment of the plasma $A\beta_{1-42}$ -to- $A\beta_{1-40}$ ratio to differentiate AD from VMD + MCI. The cutoff values for the plasma $A\beta_{1-40}$ level and the $A\beta_{1-42}$ -to- $A\beta_{1-40}$ ratio are suggested. This type of biomarker panel has the potential to help doctors in PC settings evaluate the risks of suffering from non-AD dementia, AD, or VMD and MCI in each patient.

Limitation

This study is limited to a small case number. There were no standard cerebrospinal fluid biomarkers as standard, but it isn't easy in Taiwan due to culture.

Abbreviations

PC

Primary Care;

MCI

Mild cognitive impairment;

MMSE

Mini-Mental State Examination;

FAQ

Functional Activities Questionnaire;

AD

Alzheimer's disease;

PET

Positron emission tomography;

$A\beta_{1-40}$

Amyloid-beta 1–40;

$A\beta_{1-42}$

amyloid-beta 1–42;

T-Tau

total Tau;

p-Tau181

phosphorylated Tau;

NYCUH

National Yang Ming Chiao Tung University Hospital;

NIA-AA

National Institute on Aging and Alzheimer's Association;

CDR

Clinical Dementia Rating;

VMD

Very mild dementia;

ADL

Activities of daily living;

IADL

Instrumental activities of daily living;

IRB

Institutional Review Board;

CDR-SOB

CDR-Sum of Boxes SOB;
AUC
Area under the curve

Declarations

Ethics approval and consent to participate

The research was approved by the Institutional Review Board (IRB) of National Yang Ming Chiao Tung University Hospital (IRB No.2017A033 and No. 2018A004). Written informed consent was obtained from all participants in the study.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

The study was funded by National Yang Ming Chiao Tung University Hospital

Authors' contributions

Ping-Huang Tsai: Conception, design of the work, drafted the work, and substantively revised it

Li-Cho Hsu: Acquisition, analysis

Hsuan-Ming Tsao: Acquisition, analysis, Project Administration

Shieh-Yueh Yang: Interpretation of data, drafted the work

Kuen-Lin Chen: conception, design of the work

Acknowledgements

Laboratory Department, National Yang Ming University Hospital.

References

1. Sun Y, Lee HJ, Yang SC, Chen TF, Lin KN, Lin CC, et al. A nationwide survey of mild cognitive impairment and dementia, including very mild dementia, in Taiwan. *PLoS One*. 2014;9.
2. Wändell P, Carlsson AC, Wettermark B, Lord G, Cars T, Ljunggren G. Most common diseases diagnosed in primary care in Stockholm, Sweden, in 2011. *Fam Pract*. 2013;30.
3. Ornstein SM, Nietert PJ, Jenkins RG, Litvin CB. The prevalence of chronic diseases and multimorbidity in primary care practice: A PPRNet report. *J Am Board Fam Med*. 2013;26.
4. Finley CR, Chan DS, Garrison S, Korownyk C, Kolber MR, Campbell S, et al. What are the most common conditions in primary care? Systematic review. *Can Fam Physician*. 2018;64.
5. Brayne C, Fox C, Boustani M. Dementia screening in primary care: Is it time? *J. Am. Med. Assoc.* 2007.
6. Lagaay AM, Van der Meij JC, Hijmans W. Validation of medical history taking as part of a population based survey in subjects aged 85 and over. *Br Med J*. 1992;304.
7. Verhey FRJ, Jolles J, Ponds RWHM, Rozendaal N, Plugge LA, De Vet RCW, et al. Diagnosing dementia: a comparison between a monodisciplinary and a multidisciplinary approach. *J Neuropsychiatry Clin Neurosci*. 1993;5.
8. Eefsting JA, Boersma F, Van Den Brink W, Van Tilburg W. Differences in prevalence of dementia based on community survey and general practitioner recognition. *Psychol Med*. 1996;26.
9. O'Connor DW, Pollitt PA, Hyde JB, Brook CPB, Reiss BB, Roth M. Do general practitioners miss dementia in elderly patients? *Br Med J*. 1988;297.
10. Ólafsdóttir M, Skoog I, Marcusson J. Detection of dementia in primary care: The Linköping study. *Dement Geriatr Cogn Disord*. 2000;11.
11. Valcour VG, Masaki KH, Curb JD, Blanchette PL. The detection of dementia in the primary care setting. *Arch Intern Med*. 2000;160.
12. Löppönen M, Riihala I, Isoaho R, Vahlberg T, Kivelä SL. Diagnosing cognitive impairment and dementia in primary health care - A more active approach is needed. *Age Ageing*. 2003.
13. Borson S, Scanlan JM, Watanabe J, Tu SP, Lessig M. Improving identification of cognitive impairment in primary care. *Int J Geriatr Psychiatry*. 2006;21.
14. Cruz-Orduña I, Bellón JM, Torrero P, Aparicio E, Sanz A, Mula N, et al. Detecting MCI and dementia in primary care: Efficiency of the MMS, the FAQ and the IQCODE. *Fam Pract*. 2012;29.
15. Kivipelto M, Ngandu T, Laatikainen T, Winblad B, Soininen H, Tuomilehto J. Risk score for the prediction of dementia risk in 20 years among middle aged people: a longitudinal, population-based study. *Lancet Neurol*. 2006;5.
16. Chen JH, Lin KP, Chen YC. Risk factors for dementia. *J. Formos. Med. Assoc.* 2009.
17. Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C. Potential for primary prevention of Alzheimer's disease: An analysis of population-based data. *Lancet Neurol*. 2014;13.
18. Tsai PH, Liu JL, Lin KN, Chang CC, Pai MC, Wang WF, et al. Development and validation of a dementia screening tool for primary care in Taiwan: Brain Health Test. *PLoS One*. 2018;13.

19. Chiu MJ, Yang SY, Horng HE, Yang CC, Chen TF, Chieh JJ, et al. Combined plasma biomarkers for diagnosing mild cognition impairment and Alzheimer's disease. *ACS Chem Neurosci*. 2013;4.
20. Lue LF, Sabbagh MN, Chiu MJ, Jing N, Snyder NL, Schmitz C, et al. Plasma levels of A β 42 and Tau identified probable Alzheimer's dementia: Findings in two cohorts. *Front Aging Neurosci*. 2017;9.
21. Teunissen CE, Chiu MJ, Yang CC, Yang SY, Scheltens P, Zetterberg H, et al. Plasma Amyloid- β (A β 42) Correlates with Cerebrospinal Fluid A β 42 in Alzheimer's Disease. *J Alzheimers Dis*. 2018;62.
22. Jiao F, Yi F, Wang Y, Zhang S, Guo Y, Du W, et al. The Validation of Multifactor Model of Plasma A β 42 and Total-Tau in Combination With MoCA for Diagnosing Probable Alzheimer Disease. *Front Aging Neurosci*. 2020;12.
23. Lin CH, Yang SY, Horng HE, Yang CC, Chieh JJ, Chen HH, et al. Plasma α -synuclein predicts cognitive decline in Parkinson's disease. *J Neurol Neurosurg Psychiatry*. 2017;88.
24. Wang HL, Lu CS, Yeh TH, Shen YM, Weng YH, Huang YZ, et al. Combined assessment of serum alpha-synuclein and RAB35 is a better biomarker for Parkinson's disease. *J Clin Neurol*. 2019;15.
25. Lin WC, Lu CH, Chiu PY, Yang SY. Plasma Total α -Synuclein and Neurofilament Light Chain: Clinical Validation for Discriminating Parkinson's Disease from Normal Control. *Dement Geriatr Cogn Disord*. 2021;49.
26. Yang S-Y, Liu H-C, Lin C-Y, Chiu M-J, Chen T-F, C-HL and H-HC. Development of Assaying Plasma TDP-43 Utilizing Immunomagnetic Reduction. *Neurol Disord [Internet]*. 2020;8:1–8. Available from: <https://www.hilarispublisher.com/open-access/development-of-assaying-plasma-tdp43-utilizing-immunomagnetic-reduction.pdf>.
27. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement*. 2011;7.
28. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12.
29. Andersen K, Lolk A, Nielsen H, Andersen J, Olsen C, Kragh-Sørensen P. Prevalence of very mild to severe dementia in Denmark. *Acta Neurol Scand*. 1997;96.
30. Shyu YIL, Yip PK. Factor structure and explanatory variables of the mini-mental state examination (MMSE) for elderly persons in Taiwan. *J Formos Med Assoc*. 2001;100.
31. Hughes CP, Berg L, Danziger WL, Coben LA, Martin RL. A new clinical scale for the staging of dementia. *Br J Psychiatry*. 1982;140.
32. MAHONEY FI, BARTHEL DW. FUNCTIONAL EVALUATION: THE BARTHEL INDEX. *Md State Med J*. 1965;14.
33. Lawton MP, Brody EM. Assessment of older people: Self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969;9.

34. Lin CH, Chiu SI, Chen TF, Jang JSR, Chiu MJ. Classifications of neurodegenerative disorders using a multiplex blood biomarkers-based machine learning model. *Int J Mol Sci.* 2020;21.
35. Yang CC, Chiu MJ, Chen TF, Chang HL, Liu BH, Yang SY. Assay of plasma phosphorylated tau protein (threonine 181) and total tau protein in early-stage Alzheimer's disease. *J Alzheimer's Dis.* 2018;61.
36. Chiu M-J, Yang S-Y, Chen T-F, Lin C-H, Yang F-C, Chen W-P, et al. Synergistic Association between Plasma A β 1–42 and p-tau in Alzheimer's Disease but Not in Parkinson's Disease or Frontotemporal Dementia. *ACS Chem Neurosci.* 2021.

Figures




Figure 1

Flow chart showing the assessment of plasma biomarkers to differentiate non-Alzheimer's disease (AD), AD, and very mild dementia (VMD)+mild cognitive impairment (MCI).