Screening and Identification of Potential iNOS Inhibitors to Curtail Cervical Cancer Progression: An in-Silico Drug Repurposing Approach

PAVAN KUMAR POLEBOYINA (✉ pavanbiotech2012@gmail.com)

Osmania University Faculty of Science <https://orcid.org/0000-0003-0393-3917>

SMITA C PAWAR

Osmania University Faculty of Science

AKBAR PASHA

Osmania University

RAVINDER DONETI

Osmania University

SNEHA MALLESWARI POLEBOYINA

Andhra University

SHIVAJI BANOTHU

Osmania University

Deepthi pasumarthi

Osmania University

Annapurna SD

Osmania University

Kumbhakar Divya

Osmania University

Research Article

Keywords: NOS2 (iNOS), Nitric Oxide Synthase, Cervical Cancer, Drug repurposing, FDA approved anti-cancer drugs, Small Molecules

Posted Date: July 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-674491/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Applied Biochemistry and Biotechnology on October 27th, 2021. See the published version at <https://doi.org/10.1007/s12010-021-03718-2>.

Screening and identification of potential iNOS inhibitors to curtail cervical cancer progression: An in-silico drug repurposing approach

Pavan Kumar Poleboyina¹, Akbar Pasha¹, Ravinder Doneti¹, Sneha Malleswari Poleboyina², Shivaji. B¹, Deepti .P¹, Annapurna S D¹, Divya K¹, Smita C Pawar^{1*}

¹Department of Genetics & Biotechnology, University College of Science, Osmania University, Hyderabad-500007, Telangana, India. Email: pavanbiotech2012@gmail.com

²Department of Pharmaceutical Biotechnology, AU College of Pharmaceutical Sciences, Andhra University, Visakhapatnam-530003, Andhra Pradesh, India.

***Corresponding author email address: dr.smitapawar@osmania.ac.in**

ABSTRACT:

Cervical cancer is the second most common cause of cancer deaths in women worldwide and remains the main reason of mortality amongst women of reproductive age in developing countries. Nitric oxide is involved in several physiological functions inclusive of inflammatory and immune responses. However, the function of NO in tumor biology is debatable. The inducible NOS (iNOS/NOS2) isoform is the one responsible to maintain the levels of NO and it exhibits pleiotropic effects in various cancer with concentration-dependent pro- and anti-tumor effects. NOS2 triggers angiogenesis and endothelial cell migration in tumors by regulating the levels of vascular endothelial growth factor (VEGF). In drug discovery, drug repurposing involves investigations of approved drug candidates to treat various other diseases. In this study, we used FDA-approved anti-cancer drugs and small molecules to target iNOS and identify a potential selective iNOS inhibitor. The structures of ligands were geometrically optimized, and energy minimized using Hyperchem software. Molecular docking was performed using Molegro virtual docker and ligands were selected based on MolDock score, Rerank score, and H-bonding energy. In the study showed 4 compounds, Degarelix, Goserelin, Triptorelin pamoate, and venetoclax demonstrated excellent binding affinity to NOS2 protein. These compounds exhibited the lowest MolDock score, Rerank score, with better H-bonding energy to NOS2. Based on the results these ligands project to be promising potential NOS2 inhibitors to curtail cervical cancer progression

Keywords: NOS2 (iNOS), Nitric Oxide Synthase, Cervical Cancer, Drug repurposing, FDA approved anti-cancer drugs, Small Molecules

INTRODUCTION

Continual human papillomavirus (HPV) infection is the primary reason for cervical cancer and it has led to the development of prophylactic vaccines to prevent HPV infection and HPV assays.[1]. In World, cervical cancer (CC) ranks second in causing cancer deaths in women. Due to the increased pap smear test for early detection of CC, the average death rate has declined in developed countries[2]. The global prevalence of cervical cancer is about 510,000 new cases annually, with about 288,000 deaths globally[3]. In India, 122,844 women get diagnosed and 67,477 succumb to death from the disease of Cervical Cancer annually. In the Indian population, 15 years and older-aged women are at high risk of developing cancer and cervical cancer is the second most common cancer in women aged 15-44 years. By comparing the highest standardized incidence of cervical cancer in South Asia, India has the highest age at 22 than other countries i.e. 19.2 in Bangladesh, 13 in Sri Lanka, and 2.8 in Iran.

Nitric Oxide (NO) is one of the principal oxides of nitrogen that mediates a variety of actions such as neurotransmission, host defense, iron metabolism, and vasodilatation but increased NO production contribute to a variety of disorders including cancer[4]. NOS2(iNOS) is a member of a Nitric Oxide Synthase(NOS) family of enzymes and produces NO[5]. There are three isoforms of NOS in nature: endothelial (eNOS), neural (nNOS), and inducible(iNOS), the inducible NOS exhibits increased expression in several types of cancer but the prognostic role in cervical cancer still remains unclear[6].

In our previous study, we demonstrated that iNOS expression levels were reduced by using plant-derived diterpenoid andrographolide, in both in-silico and in-vitro[7]. In the present study we adopt an Insilco drug repurposing approach to identify a potential inhibitor for NOS2 by using FDA approved anti-cancer drugs and small molecules.

MATERIALS AND METHODS

The study involves retrieval of target protein in PDB format from RCSB Protein Data Bank[8] and retrieval of ligands, FDA approved anti-cancer drugs and small molecules from Pubchem database[9], Zinc database[10], and Drug bank[11]. Due to the presence of cofactors and peptide bonds in ligands, some structures were drawn and converted into pdb files through Hyperchem8 software[12] and ChemDraw Pro 12.0 software (RRID:SCR_016768)[13]. Molecular Docking was carried out by Molegro Virtual Docker (MVD2019_7_0_0)(RRID:SCR_000190)[14], data analyzed by Molegro Data modeler[15] and results were envisioned by Molegro Molecular Viewer[16].

Retrieval of target protein

3d structure of NOS2 (iNOS) Human inducible nitric oxide synthase with inhibitor protein (PDB ID: 4NOS) was retrieved from RCSB Protein Data Bank. Before docking, the unwanted hetero atoms, water molecules, and other ligand compounds were removed by MVD.

Retrieval of FDA approved anti-cancer drugs and small molecules

1,116 compounds were virtually screened and a total of 162 top hit lead optimized compounds were selected for further analysis. The 3d and 2d structures of ligands were retrieved from Pubchem database, zinc database, drug bank, and some structures were drawn in ChemDraw pro 12.0 and Hyperchem 8 package; they were pre-optimized using molecular mechanics force field procedure (MM⁺, AMBER) in Hyperchem8[17]. To get the conformers of FDA-approved anti-cancer drugs and small molecules with the lowest energy, the semi-empirical method AM-1 was applied to the molecular structures. To avoid the local stability, each molecular structure was geometrically optimized several times with different starting points using the Polak-Rebierre algorithm, till the root-mean-square gradient becomes equal to 0.01 and 0.001 kcal Å⁻¹ mol⁻¹[18]. Finally the energy minimizations of each ligand structure was performed. the first step and second step calculations i.e. single point calculations that are used to determine the total molecular energy of structure and geometric optimization calculations (MM⁺, AMBER force field) to determine the energy minimization algorithms that locate the flexible structures were done by using Hyperchem8[19].

Prediction of ligand binding active sites

The ligand-binding active sites in NOS2 (iNOS, PDB ID: 4NOS) protein for FDA-approved anti-cancer drugs and small molecule ligands are not well characterized. The amino acid residues responsible for active site formation in NOS2 protein were identified through the MVD cavity detection algorithm. This MVD detected 5 different active sites with high volume surface in NOS2 protein.

Protein preparation

NOS2 human inducible nitric oxide synthase with inhibitor protein (PDB ID: 4NOS) was selected for the molecular docking studies using MVD. All necessary H atom addition, valency checks were performed. MVD provided utilities and protein preparation i.e. protonation addition of mismatched amino acid residues was repaired, optimized and rebuilt by MVD. For docking studies it is important that the imported structures must be prepared accurately, the atom connectivity and bond orders are correct and the partial atomic charges are assigned. PDB files have a poor or missing assignment of explicit hydrogen's, and the PDB file format cannot accommodate bond order information. Then the repair, rebuilt and optimized protein was saved in *.Mol format. The final structure was visualized and analyzed with SPDBV 4.1.0 (RRID:SCR_013295)[20] window control panel and BIOVIA Discovery studio (BIOVIA_2021.DS2021Client) software (RRID:SCR_015651)[21].

Ligand preparation

The selected FDA-approved anti-cancer drugs and small molecules were downloaded from Pubchem, Zinc database, and drug bank. Some torsion and peptide bond containing candidates were drawn and minimized its energy by using Hyperchem8 software and imported to MVD workspace in *.Mol format. Before import, the small molecules and FDA-approved anti-cancer

drugs are subjected to series of steps to generate optimization and variation of the structure, as the energy minimized ligands have the capability to form better bonding with protein.

Docking studies

In the current study, a docking algorithm called MolDock has been used. It is based on a new hybrid search algorithm, called guided differential evolution. It combines differential evolutionary optimization technique with an active site prediction algorithm[22]. For the docking process GRID resolution was set to 0.30Å, and the cavities were set on the coordinate values X= 16.90, Y= 61.22 and Z=17.64. Other embedded parameters were used which consist of a maximum iteration of 1,500, simplex evolution of 300, Neighbour distance factor 1.00, and the highest population size of 50. For each ligand 10 independent runs were performed with the differential evolution algorithm, with every 10 independent runs of these docking returning 5 docking poses and selecting the top most pose for consideration. MVD shows higher docking accuracy than the other docking software(MVD: 87%, Glide: 82%, Surflex: 75%, FlexX: 58%)[23]. The scoring function used by MVD is called MolDock which is derived from the PLP scoring functions. The docking scoring function MolDock score energy, E_{score} is defined by (1), where E_{inter} is the ligand-receptor interaction energy and E_{intra} is the internal energy of the ligand. According to (2) equation, E_{inter} is calculated.

$$E_{score} = E_{inter} + E_{intra} \quad (1)$$

$$E_{inter} = \sum_{i=ligand} \sum_{j=protein} \left[E_{PLP}(r_{ij}) + 332.0 \frac{q_i q_j}{4r_{ij}^2} \right] \quad (2)$$

The E_{PLP} is a piecewise linear potential by using two different sets of parameters i.e. the steric and Hydrogen bonds[24]. According to (3) equation E_{intra} is calculates the energy involves in the pair of atoms of the ligand and torsional energy where h is the torsional angle of the bond. E_{clash} assigns a penalty of 1,000kcal/mol if the distance between two heavy atoms similar than 2.0Å.

$$E_{intra} = \sum_{i=ligand} \sum_{j=protein} [E_{PLP}(r_{ij})] + \sum_{flexible\ bond} A[1 - \cos(m\theta - \theta_0)] + E_{clash}. \quad (3)$$

Pharmacophore modeling

By using the LigandScout 4.4.5 software package(RRID:SCR_014889) the structure-based pharmacophore model was generated[25]. Herein the proposed algorithm studies and interprets

ligand-receptor interactions such as hydrogen bonds, hydrophobic regions, and charge transfer regions of the macromolecular environment from PDB files and allowing the automatic binding of the pharmacophore model. The exclusion volume spheres were added to the generated pharmacophore model which represents the inaccessible areas along with any potential ligand[26].

Drug-likeness and ADMET properties

The drug scan and ADMET properties were performed to determine the inhibitor has the fulfilled conditions as the drug candidate based on Lipinski's Rule of Five[27]. The pharmacokinetic properties, toxicity, and Bioavailability of selected ligands were carried out by using the Molinspiration cheminformatics tool[28], SwissADME tool[29], and pkCSM tool[30].

RESULTS

Target protein structural information

The NOS2 (iNOS) human inducible nitric oxide synthase protein(PDB ID: 4NOS) with a resolution of 2.25Å was retrieved from the RCSB protein data bank. The structural confirmations of protein were obtained by PDBsum[31] and obtained PROCHECK[32], ERRAT[33] validations by uploading its structure in structure validation server (SAVESv6.0)[34]. The obtained 3d cartoon, secondary structural analysis, and Ramachandran plot are shown in Fig.1

Fig.1 shows structural information of protein NOS2 (a) 3d cartoon structure (b) secondary structure of 3pty mapped obtained using PDBsum (c) and (d) Ramachandran plot

The NOS2 protein active sites were obtained by the MVD cavity detection algorithm the five detected cavities with their volume are illustrated in Fig.2

Fig.2 MVD generated cavities of protein NOS2

LIGAND STRUCTURE INFORMATION

Total 162 ligand structures of FDA-approved anti-cancer drugs and small molecules were used in docking to find out its inhibitory binding effect with the target protein. The drug compounds structural information and its flexibility confirmation with after single point and geometry optimization minimized energies are represented in Table 1

Docking Results validation

Herein we used MVD for docking studies to obtain MolDock score, Rerank score, Interaction energy, torsions, and H-bonding energy. To find out the protein-ligand interaction each compound undergoes 10runs with threshold energy 100 and is finally given 5 best poses. Out of 5 poses, we have selected the top 1 pose for each ligand having the lower energy along with higher MolDock score, lower rerank score, and best hydrogen bonding. The obtained energies of each ligand are shown in Table 2

Among all docked 162 ligands, 4 compounds showed harmonious inhibitory effect with NOS2 protein. Based on the average MolDock score -150kcal/mol to -160kcal/mol, we have selected above-average MolDock score obtained ligands for further studies. The top 4 compounds; Degarelix, Goserelin, Triptorelin pamoate, and Venetoclax revealed the lowest MolDock score on

NOS2 protein which is -226.10kcal/mol, -189.77kcal/mol, -185.73kcal/mol, and -168.21kcal/mol respectively.

Degarelix binds into the active site of NOS2 protein with MolDock score -226.10kcal/mol and its binding site consist of amino acid residues were Glu394C, Glu393C, Asn390C, Val386C, Glu387C, Glu91D, Pro474D, Ser92D, His477D, Glu441D, Lys445D, Ile204D, Asn449D, Glu448D, Met444D, Met94D, Asn89D, Lys88D, Thr95D, Arg86D. The docked amino acid residues against the ligand were Asn449D, Met94D, Ser92D, Pro474D, Val386C, Asn89D.

Goserelin binds into the active site of NOS2 protein with MolDock score -189.77kcal/mol and its binding site consist of amino acid residues were Val386C, Glu387C, Gly91D, Glu93D, Ser92D, Asn89D, Met94D, Phe96D, Glu448D, Met447D, Lys445D, Met444D, Glu478D, Pro474D, Gly441D. The docked amino acid residues against the ligand were Glu441D, Gly91D, Met444D, Ser92D.

Triptorelin pamoate binds into the active site of iNOS protein with MolDock score -185.73kcal/mol and its binding site consist of amino acid residues were Pro474D, Val386C, Glu441D, Lys445D, Met444D, Arg452D, Asn449D, Glu448D, Phe96D, Met94D, Gly478D, Asn89D, Ser92D. The docked amino acid residues against the ligand were Glu448D, Met444D, Ser92D.

Venetoclax binds into the active site of iNOS protein with MolDock score -168.21kcal/mol and its binding site consist of amino acid residues were Gly93D, Met94D, Asn89D, Ser92D, Glu448D, Lys445D, Met444D, Pro474D, Glu441D. The docked amino acid residues against the ligand were Ser92D, Glu448D. Fig.3 illustrates the top 4 ligands hydrogen bonding interactions with amino acid residues of NOS2 protein.

Fig.3: Hydrogen bond interactions of ligands with protein NOS2 (a) Degarelix (b) Goserelin (c)Triptorelin pamoate (d) Venetoclax

Fig.4 represents the key amino acid residues of NOS2 protein which forming steric interactions with ligands. Fig.5 and Fig. 6 show the ligand docked against the crystal protein structures of human inducible nitric oxide synthase and the amino acid residues around active & docked against NOS2 protein respectively (Table 3). Our validated results of pharmacophore models are illustrated in Fig.7

(a)

(b)

(c)

(d)

Fig.4 Steric interactions of ligands with NOS2 protein (a) Degarelix (b) Goserelin (c) Triptorelin pamoate (d) Venetoclax

Fig.5Ligand docked against the crystal protein structure of NOS2 (a) Degarelix (b) Goserelin (c) Triptorelin pamoate (d) Venetoclax

(a)

(b)

(c)

(d)

Fig.6The amino acid residues around active site and docked amino acids against NOS2 (a) Degarelix (b) Goserelin (c) Triptorelin pamoate (d) Venetoclax

Fig.7pharmacophore models (a) Degarelix (b) Goserelin (c) Triptorelin pamoate (d) Venetoclax

ADMET properties and Drug-likeness results

A rapid dissociation rate will occur with the weaker binding, the more negative binding energy analogue to the strong binding of selected compounds to the target NOS2 protein[35]. The lower binding energy exhibited compounds gives the strong binding with NOS2 protein, the selected compounds have shown the lower binding energies. According to the Rule of Five, a molecule might be no longer orally active if it violates or greater than the 4 rules[36].To obtain the pharmacophore properties ligands canonical smiles were uploadedto pkCSM, swissADME, and Molinspiration cheminformatics tools, and the results are illustrated in Table 4

DISCUSSION

The enzyme NOS2 also called inducible NOS or inflammatory NOS plays a critical role in inflammation and tissue damage in cancer cells[37]. Docking studies have proved to be an essential tool to discover and identify the potential target candidates for a particular protein[38]. In this study FDA-approved anti-cancer drugs and small molecules were used to identify the potential inhibitor for human inducible nitric oxide synthase protein. Among 162 docked compounds we have identified 4 compounds that showed better inhibitory effect on NOS2.

The first compound Degarelix sold under the brand name of Firmagon is a gonadotrophin-releasing hormone antagonist used in the first-line treatment of advanced prostate cancer[39]. It has shown the lowest MolDock score, Rerank score, and better H-bonding energy with NOS2 protein which is -226.10kcal/mol, -106.48kcal/mol, and -10.57kcal/mol respectively.

The second compound Goserelin sold under the brand name Zoladex and it reduces the oestrogen levels in premenopausal women by inhibiting the hypothalamic-pituitary-ovarian axis[40]. It is a luteinizing hormone-releasing hormone antagonist widely used in the treatment of Breast cancer[41]. It has shown the lowest MolDock score, Rerank score, and H-bonding energy with NOS2 protein which has -189.77kcal/mol, -92.09kcal/mol, and -7.7kcal/mol respectively.

The third selected compound Triptorelin pamoate sold under the brand name of Decapeptyl or Gonapeptyl and is a gonadotrophin-releasing hormone agonist. It is widely used in the palliative treatment of advanced prostate cancer[42], adjuvant treatment with the combination of tamoxifen-Triptorelin pamoate in the early stage of Breast cancer[43]. It has shown the lowest MolDock score -185.73kcal/mol, Rerank score -59.99kcal/mol, and H-bonding energy -10.24kcal/mol with NOS2 protein.

The fourth compound venetoclax sold under the brand name of Venclexta and it is an antineoplastic agent which blocks BCL-2 and promotes apoptosis in various cancer cells[44]. In docking it has shown the lowest MolDock score -168.21kcal/mol, Rerank score -64.67kcal/mol, and H-bonding energy -5.23kcal/mol with NOS2 protein. All these drug candidates are widely using in the treatment of various cancers. Thus this study identified the 4 best inhibitory compounds from FDA-approved anti-cancer drugs and small molecules to inhibit NOS2 protein in cervical cancer. However further in-vitro and in-vivo studies are warranted to validate the promising potential of these compounds to be used as NOS2 protein inhibitors to restrain the cervical cancer progression.

Findings

1. The key amino acids of NOS2 protein which are forming H-bonding interactions with ligands are Asn449D, Met94D, Ser92D, Pro474D, Val386C, Asn89D, Glu441D, Gly91D, Met444D, and Glu448D.
2. The common amino acid Ser92D is involved in all ligand interactions and forms H-bonding with ligands. So, from this we deduce that the amino acid Ser92D of NOS2 protein has the highest binding efficiency to form H-bonding with ligands.
3. The key amino acids which are involved in steric interactions are Val386C, Ser92D, Met94D, Met444D, Lys445D, Gln448D, Phe96D, Arg452D, Glu97D, Glu393C, Asn390C, Asn449D, and Ile204D.
4. Based on our in-silico analysis we conclude that Degarelix, Goserelin, Triptorelin pamoate, and Venetoclax exhibit the potential to effectively inhibit NOS2 expression in cervical cancer.

CONCLUSION

In summary, FDA-approved anti-cancer drugs and small molecules from the PubChem, zinc database, and drug bank were docked against NOS2 protein. Molecular docking studies revealed that Degarelix, Goserelin, Triptorelin pamoate, and Venetoclax drugs interacted with human inducible nitric oxide synthase (NOS2) protein. These ligands demonstrate an excellent inhibitory binding effect on NOS2 and are potential candidates to be used as drugs to reduce iNOS expression in cervical cancer tumor cells. These results open the way for further experimental confirmation as a drug repurposing approach for NOS2 inhibitor to combat cervical cancer.

DECLARATIONS

Data availability

Data supporting the productivity of this investigation are available from the corresponding author upon request.

Conflict of interests

The authors declare that there is no conflict of interest. No additional benefits will be received from a third party directly or indirectly by the authors.

Author's contributions

SCP & PKP conceptualized, designed interpreted data and edited manuscript; PKP, AP, DP, SDA, SB, SMP, DK designed and conducted the study. All authors have approved the manuscript in the current form.

Acknowledgments

Thankful to the staff of the Department of Genetics & Biotechnology, Osmania University for providing the Bioinformatics/software facilities.

Funding

This study was supported by grants of Dr. Smita C. Pawar received from the Science and Engineering Research Board(SERB), Govt. of India, reference Nos. SB/EMEQ-471/2014 and EEQ/2019/000569 dt 06-Jan-2020, and the fellowship was provided by DHR/HRD/Women Scientist/TypeV to SDA, by CFMT: RUSA 2.0 to DK, by CSIR-UGC to PKB & AP, by ICMR to SB. The facilities were provided by DSTPURSE-II.

Ethical approval

Not applicable

Consent to participate

Not applicable

Consent to publish

Not applicable

Table 1 Ligand structural information and after single point and geometry optimization energy minimization calculations

Name of the inhibitor	Pubchem CID	Molecular formula (Molecular weight g/mol)	STRUCTURE	After Single point Energy	After Geometry Optimization minimizing energy
Abarelix	16131215	<u>C</u> <u>72</u> <u>H</u> <u>95</u> <u>Cl</u> <u>N</u> <u>14</u> <u>O</u> <u>14</u> (1416.1)		461.92 Kcal/mol	29.72 Kcal/mol
Abemaciclib	46220502	<u>C</u> <u>27</u> <u>H</u> <u>32</u> <u>F</u> <u>2</u> <u>N</u> <u>8</u> (506.6)		117.59 Kcal/mol	39.88 Kcal/mol
Abiraterone acetate	9821849	<u>C</u> <u>26</u> <u>H</u> <u>33</u> <u>NO</u> <u>2</u> (391.5)		96.74 Kcal/mol	41.36 Kcal/mol
Acalabrutinib	71226662	<u>C</u> <u>26</u> <u>H</u> <u>23</u> <u>N</u> <u>7</u> <u>O</u> <u>2</u> (465.5)		602.20 Kcal/mol	39.20 Kcal/mol
Afatinib	10184653	<u>C</u> <u>24</u> <u>H</u> <u>25</u> <u>Cl</u> <u>F</u> <u>N</u> <u>5</u> <u>O</u> <u>3</u> (485.9)		116.72 Kcal/mol	32.17 Kcal/mol

Afatinib dimaleate	15606394	<u>C₃₂H₃₃ClFN₅O₁₁</u> (718.1)		116.71 Kcal/mol	32.17 Kcal/mol
Alectinib	49806720	<u>C₃₀H₃₄N₄O₂</u> (482.6)		357.92 Kcal/mol	46.84 Kcal/mol
Alimta	135916113	<u>C₂₀H₁₉N₅O₆⁻²</u> (425.4)		126.05 Kcal/mol	23.31 Kcal/mol
Alpelisib	56649450	<u>C₁₉H₂₂F₃N₅O₂S</u> (441.5)		145.89 Kcal/mol	38.96 Kcal/mol
Amifostine	2141	<u>C₅H₁₅N₂O₃PS</u> (214.23)		60.35 Kcal/mol	27.52 Kcal/mol
Anagrelide	135409400	<u>C₁₀H₇Cl₂N₃O</u> (256.08)		49.61 Kcal/mol	13.98 Kcal/mol

Anastrozole	2187	$\text{C}_{17}\text{H}_{19}\text{N}_5$ (293.4)		538.2 Kcal/mol	25.80 Kcal/mol
Andrographolide	5318517	$\text{C}_{20}\text{H}_{30}\text{O}_5$ (350.4)		Kcal/mol	Kcal/mol
Anexsia	1983	$\text{C}_8\text{H}_9\text{NO}_2$ or $\text{HOC}_6\text{H}_4\text{NHCOCH}_3$ (151.16)		Kcal/mol	Kcal/mol
Apalutamide	24872560	$\text{C}_{21}\text{H}_{15}\text{F}_4\text{N}_5\text{O}_2\text{S}$ (477.4)		419.5 Kcal/mol	61.71 Kcal/mol
Aprepitant	135413536	$\text{C}_{23}\text{H}_{21}\text{F}_7\text{N}_4\text{O}_3$ (534.4)		114.85 Kcal/mol	48.16 Kcal/mol
Aredia	60342	$\text{C}_3\text{H}_{19}\text{NNa}_2\text{O}_{12}\text{P}_2$ (369.11)		137.22 Kcal/mol	27.57 Kcal/mol
ofatumumab	6918251	$\text{C}_{12}\text{H}_{10}\text{N}_4\text{O}_2$ (242.23)		Kcal/mol	Kcal/mol

Axitinib	6450551	<u>C₂₂H₁₈N₄OS</u> (386.5)		117.47 Kcal/mol	30.09 Kcal/mol
Belinostat	6918638	<u>C₁₅H₁₄N₂O₄S</u> (318.3)		135.17 Kcal/mol	35.57 Kcal/mol
Bendamustine	65628	<u>C₁₆H₂₁Cl₂N₃O₂</u> (358.3)		72.08 Kcal/mol	25.61 Kcal/mol
Besponsa	135298753	<u>0635715014</u>		Kcal/mol	Kcal/mol
Binimetinib	10288191	<u>C₁₇H₁₅BrF₂N₄O₃</u> (441.2)		99.41 Kcal/mol	40.68 Kcal/mol
Bortezomib	387447	<u>C₁₉H₂₅N₄O₄</u> (384.2)		119.28 Kcal/mol	8.73 Kcal/mol

Bosutinib	5328940	<u>C₂₆H₂₉Cl₂N₅O₃</u> (530.4)		348.87 Kcal/mol	26.24 Kcal/mol
Brigatinib	68165256	<u>C₂₉H₃₉ClN₇O₂P</u> (584.1)		149.58 Kcal/mol	68.10 Kcal/mol
Bromfenac	60726	<u>C₁₅H₁₂BrNO₃</u> (334.16)		76.69 Kcal/mol	10.53 Kcal/mol
Cabazitaxel	9854073	<u>C₄₅H₅₇NO₁₄</u> (835.9)		356.35 Kcal/mol	112.50 Kcal/mol
Cabozantinib	25102847	<u>C₂₈H₂₄FN₃O₅</u> (501.5)		316.45 Kcal/mol	198.91 Kcal/mol
Capecitabine	60953	<u>C₁₅H₂₂FN₃O₆</u> (359.35)		82.56 Kcal/mol	21.28 Kcal/mol

Carfilzomib	11556711	<u>C₄₀H₅₇N₅O₇</u> (719.9)		384.57 Kcal/mol	217.77 Kcal/mol
Carmustine	2578	<u>C₅H₉Cl₂N₃O₂</u> (214.05)		29.95 Kcal/mol	4.98 Kcal/mol
Cedazuridine	25267009	<u>C₉H₁₄F₂N₂O₅</u> (268.21)		57.96 Kcal/mol	17.23 Kcal/mol
Cinacalcet	156419	<u>C₂₂H₂₂F₃N</u> (357.4)		83.97 Kcal/mol	27.52 Kcal/mol
Clofarabine	119182	<u>C₁₀H₁₁ClFN₅O₃</u> (303.68)		63.41 Kcal/mol	27.81 Kcal/mol
Cobimetinib	16222096	<u>C₂₁H₂₁F₃IN₃O₂</u> (531.3)		137.58 Kcal/mol	57.52 Kcal/mol

Copanlisib	135565596	<u>C₂₃H₂₈N₈O₄</u> (480.5)		120.79 Kcal/mol	31.25 Kcal/mol
Copiktra	137528190	<u>C₂₂H₁₉ClN₆O₂</u> (434.9)		112.23 Kcal/mol	64.89 Kcal/mol
Cisplatin	5702198	PtCl ₂ (NH ₃) ₂ or <u>Cl₂H₆N₂Pt</u>		Kcal/mol	Kcal/mol
Crizotinib	11626560	<u>C₂₁H₂₂Cl₂FN₅O</u> (450.3)		99.94 Kcal/mol	28.30 Kcal/mol
Dabrafenib	44462760	<u>C₂₃H₂₀F₃N₅O₂S₂</u> (519.6)		195.59 Kcal/mol	57.51 Kcal/mol
Dacomitinib	11511120	<u>C₂₄H₂₅ClFN₅O₂</u> (469.9)		112.70 Kcal/mol	47.95 Kcal/mol
Darolutamide	67171867	<u>C₁₉H₁₉ClN₆O₂</u> (398.8)		347.86 Kcal/mol	29.47 Kcal/mol

Dasatinib	3062316	<u>C₂₂H₂₆ClN₇O₂S</u> (488.0)		117.68 Kcal/mol	41.12 Kcal/mol
Decitabine	451668	<u>C₈H₁₂N₄O₄</u> (228.21)		46.97 Kcal/mol	14.03 Kcal/mol
Degarelix	16136245	<u>C₈₂H₁₀₃ClN₁₈O₁₆</u> (1632.3)		542.80 Kcal/mol	30.89 Kcal/mol
Docetaxel	148124	<u>C₄₃H₅₃NO₁₄</u> (807.9)		348.28 Kcal/mol	107.73 Kcal/mol
Doxorubicin	31703	<u>C₂₇H₂₉NO₁₁</u> (543.5)		129.23 Kcal/mol	67.89 Kcal/mol
Eloxatin 2d	9887054	<u>C₈H₁₄N₂O₄Pt</u> (397.3)		69.89 Kcal/mol	3.70 Kcal/mol

Eloxatin Region	9887054	<u>C₈H₁₄N₂O₄Pt</u> (397.3)		Kcal/mol	Kcal/mol
Enasidenib	89683805	<u>C₁₉H₁₇F₆N₇O</u> (473.4)		122.88 Kcal/mol	30.29 Kcal/mol
Encorafenib	50922675	<u>C₂₂H₂₇ClFN₇O₄S</u> (540.0)		171.10 Kcal/mol	50.87 Kcal/mol
Entrectinib	25141092	<u>C₃₁H₃₄F₂N₆O₂</u> (560.6)		128.70 Kcal/mol	63.25 Kcal/mol
Enzalutamide	15951529	<u>C₂₁H₁₆F₄N₄O₂S</u> (464.4)		378.92 Kcal/mol	25.65 Kcal/mol
Epirubicin	41867	<u>C₂₇H₂₉NO₁₁</u> (543.5)		128.01 Kcal/mol	74.27 Kcal/mol

Erdafitinib	67462786	<u>C₂₅H₃₀N₆O₂</u> (446.5)		109.94 Kcal/mol	37.97 Kcal/mol
Eribulin	11354606	<u>C₄₀H₅₉NO₁₁</u> (729.9)		Kcal/mol	Kcal/mol
Eribulin mesylate	17755248	<u>C₄₁H₆₃NO₁₄S</u> (826.0)		Kcal/mol	Kcal/mol
Erlotinib	176870	<u>C₂₂H₂₃N₃O₄</u> (393.4)		542.11 Kcal/mol	19.89 Kcal/mol
Everolimus	6442177	<u>C₅₃H₈₃NO₁₄</u> (958.2)		282.28 Kcal/mol	73.94 Kcal/mol
Exemestane	60198	<u>C₂₀H₂₄O₂</u> (296.4)		88.62 Kcal/mol	35.14 Kcal/mol

Fedratinib	16722836	<u>C₂₇H₃₆N₆O₃S</u> (524.7)		167.74 Kcal/mol	57.49 Kcal/mol
Fentanyl	3345	<u>C₂₂H₂₈N₂O</u> (336.5)		82.58 Kcal/mol	22.27 Kcal/mol
Fentanyl citrate	13810	<u>C₂₈H₃₆N₂O₈</u> (528.6)		200.98 Kcal/mol	76.46 Kcal/mol
Fentanyl citrate 2d				14.85 Kcal/mol	5.29 Kcal/mol
Fentanyl citrate 2d				14.85 Kcal/mol	5.29 Kcal/mol
Ferumoxides	6432052	<u>Fe₃O₄</u> (231.53)		-0.008 Kcal/mol	-0.0081 Kcal/mol
Flutamide	3397	<u>C₁₁H₁₁F₃N₂O₃</u> (276.21)		85.60 Kcal/mol	11.78 Kcal/mol
Fulvestrant	104741	<u>C₃₂H₄₇F₅O₃S</u> (606.8)		136.96 Kcal/mol	47.47 Kcal/mol

Fusilev	136047905	<u>C₂₀H₂₁CaN₇O₇</u> (511.5)		135.77 Kcal/mol	17.72 Kcal/mol
Gefitinib	<u>123631</u>	<u>C₂₂H₂₄ClFN₄O₃</u> (446.9)		86.06 Kcal/mol	25.85 Kcal/mol
Gilteritinib	49803313	<u>C₂₉H₄₄N₈O₃</u> (552.7)		112.40 Kcal/mol	41.56 Kcal/mol
Glasdegib	25166913	<u>C₂₁H₂₂N₆O</u> (374.4)		347.31 Kcal/mol	33.85 Kcal/mol
Goserelin	5311128	<u>C₅₉H₈₄N₁₈O₁₄</u> (1269.4)		446.54 Kcal/mol	58.84 Kcal/mol
Granisetron	3510	<u>C₁₈H₂₄N₄O</u> (312.4)		82.59 Kcal/mol	40.08 Kcal/mol
Ibritumomab	92002593	<u>C₂₄H₃₅N₅O₁₁</u> (569.6)		308.48 Kcal/mol	35.38 Kcal/mol

Ibrutinib	24821094	<u>C</u> <u>2</u> <u>5</u> <u>H</u> <u>2</u> <u>4</u> <u>N</u> <u>6</u> <u>O</u> <u>2</u> (440.5)		124.08 Kcal/mol	38.32 Kcal/mol
Idelalisib	11625818	<u>C</u> <u>2</u> <u>2</u> <u>H</u> <u>1</u> <u>8</u> <u>F</u> <u>N</u> <u>7</u> <u>O</u> (415.4)		114.28 Kcal/mol	30.34 Kcal/mol
Imatinib	5291	<u>C</u> <u>2</u> <u>9</u> <u>H</u> <u>3</u> <u>1</u> <u>N</u> <u>7</u> <u>O</u> (493.6)		124.04 Kcal/mol	31.10 Kcal/mol
Ingenol mebutate	6918670	<u>C</u> <u>2</u> <u>5</u> <u>H</u> <u>3</u> <u>4</u> <u>O</u> <u>6</u> (430.5)		316.31 Kcal/mol	234.29 Kcal/mol
Interferon alfa-2B	71306834	<u>C</u> <u>1</u> <u>6</u> <u>H</u> <u>1</u> <u>7</u> <u>Cl</u> <u>3</u> <u>I</u> <u>2</u> <u>N</u> <u>3</u> <u>Na</u> <u>O</u> <u>5</u> <u>S</u> (746.5)		73.67 Kcal/mol	26.49 Kcal/mol
Iodixanol	3724	<u>C</u> <u>3</u> <u>5</u> <u>H</u> <u>4</u> <u>4</u> <u>I</u> <u>6</u> <u>N</u> <u>6</u> <u>O</u> <u>1</u> <u>5</u> (1550.2)		208.03 Kcal/mol	34.12 Kcal/mol
Irinotecan	60838	<u>C</u> <u>3</u> <u>3</u> <u>H</u> <u>3</u> <u>8</u> <u>N</u> <u>4</u> <u>O</u> <u>6</u> (586.7)		144.23 Kcal/mol	45.13 Kcal/mol

Ivosidenib	71657455	<u>C₂₈H₂₂ClF₃N₆O₃</u> (583.0)		442.37 Kcal/mol	54.98 Kcal/mol
Ixabepilone	6445540	<u>C₂₇H₄₂N₂O₅S</u> (506.7)		288.12 Kcal/mol	205.05 Kcal/mol
Ixazomib	25183872	<u>C₁₄H₁₉BCl₂N₂O₄</u> (361.0)		97.88 Kcal/mol	8.19 Kcal/mol
Lapatinib	208908	<u>C₂₉H₂₆ClFN₄O₄S</u> (581.1)		182.83 Kcal/mol	52.24 Kcal/mol
Larotrectinib	<u>46188928</u>	<u>C₂₁H₂₂F₂N₆O₂</u> (428.4)		114.06 Kcal/mol	40.12 Kcal/mol
Lenalidomide	216326	<u>C₁₃H₁₃N₃O₃</u> (259.26)		77.02 Kcal/mol	16.88 Kcal/mol
Lenvatinib	9823820	<u>C₂₁H₁₉ClN₄O₄</u> (426.9)		287.33 Kcal/mol	195.54 Kcal/mol

Letrozole	3902	<u>C₁₇H₁₁N₅</u> (285.30)		583.27 Kcal/mol	17.03 Kcal/mol
Leuprolide acetate	657180	<u>C₆₁H₈₈N₁₆O₁₄</u> (1269.4)		432.43 Kcal/mol	55.15 Kcal/mol
Leuprorelin	657181	<u>C₅₉H₈₄N₁₆O₁₂</u> (1209.4)		414.72 Kcal/mol	55.26 Kcal/mol
Levoleucovorin	135398559	<u>C₂₀H₂₃N₇O₇</u> (473.4)		135.77 Kcal/mol	17.72 Kcal/mol
Lonsurf	9829639	<u>C₁₉H₂₃Cl₂F₃N₆O₇</u> (575.3)		142.01 Kcal/mol	24.02 Kcal/mol
Lorlatinib	71731823	<u>C₂₁H₁₉FN₆O₂</u> (406.4)		351.30 Kcal/mol	29.98 Kcal/mol
Lutathera	132274234	<u>C₆₅H₈₇LuN₁₄O₁₉S₂</u> (1609.5)		Kcal/mol	Kcal/mol

Magnesium silicate, hydrous	16211421	$\text{H}_2\text{Mg}_3\text{O}_{12}\text{Si}_4$ or $\text{Mg}_3(\text{OH})_2\text{Si}_4\text{O}_{10}$ or $\text{Mg}_3(\text{Si}_4\text{O}_{10})(\text{OH})_2$ (379.27)		1683.13 Kcal/mol	0.03 Kcal/mol
Marqibo	138911098	$\text{C}_{46}\text{H}_{58}\text{N}_4\text{O}_{14}\text{S}$ (923.0)		494.99 Kcal/mol	148.72 Kcal/mol
Mechlorethamine	4033	$\text{C}_5\text{H}_{11}\text{Cl}_2\text{N}$ (156.05)		16.04 Kcal/mol	9.36 Kcal/mol
Melphalan flufenamide	9935639	$\text{C}_{24}\text{H}_{30}\text{Cl}_2\text{FN}_3\text{O}_3$ (498.4)		88.32 Kcal/mol	8.47 Kcal/mol
Methoxsalen	4114	$\text{C}_{12}\text{H}_8\text{O}_4$ (216.19)		70.23 Kcal/mol	22.29 Kcal/mol
Midostaurin	9829523	$\text{C}_{35}\text{H}_{30}\text{N}_4\text{O}_4$ (570.6)		212.05 Kcal/mol	79.99 Kcal/mol
Miraluma	449763	$\text{C}_{36}\text{H}_{66}\text{N}_6\text{O}_6\text{Tc}$ (777.9)		158.66 Kcal/mol	10.87 Kcal/mol

Morphine sulfate	16051935	<u>C₃₄H₄₀N₂O₁₀S</u> (668.8)		67.01 Kcal/mol	32.34 Kcal/mol
Nelarabine	3011155	<u>C₁₁H₁₅N₅O₅</u> (297.27)		58.26 Kcal/mol	29.07 Kcal/mol
Neratinib	9915743	<u>C₃₀H₂₉ClN₆O₃</u> (557.0)		381.56 Kcal/mol	22.04 Kcal/mol
Neratinib maleate	67307512	<u>C₃₄H₃₃ClN₆O₇</u> (673.1)		381.56 Kcal/mol	22.04 Kcal/mol
Netupitant	6451149	<u>C₃₀H₃₂F₆N₄O</u> (578.6)		148.62 Kcal/mol	32.73 Kcal/mol
Nilotinib	644241	<u>C₂₈H₂₂F₃N₇O</u> (529.5)		158.65 Kcal/mol	29.82 Kcal/mol

Niraparib	24958200	<u>C₁₉H₂₀N₄O</u> (320.4)		85.27 Kcal/mol	31.15 Kcal/mol
Olaparib	23725625	<u>C₂₄H₂₃FN₄O₃</u> (434.5)		290.16 Kcal/mol	190.08 Kcal/mol
Omacetaxine mepesuccinate	285033	<u>C₂₉H₃₉NO₉</u> (545.6)		120.60 Kcal/mol	46.58 Kcal/mol
Ondansetron	4595	<u>C₁₈H₁₉N₃O</u> (293.4)		85.55 Kcal/mol	36.31 Kcal/mol
Osimertinib	71496458	<u>C₂₈H₃₃N₇O₂</u> (499.6)		137.22 Kcal/mol	41.54 Kcal/mol
Oxaliplatin	9887053	<u>C₈H₁₄N₂O₄Pt</u> (397.3)		215.23 Kcal/mol	3.68 Kcal/mol

Paclitaxel	36314	<u>C</u> <u>47</u> <u>H</u> <u>51</u> <u>N</u> <u>O</u> <u>14</u> (853.9)		Kcal/mol	Kcal/mol
Paclitaxel	36314	<u>C</u> <u>47</u> <u>H</u> <u>51</u> <u>N</u> <u>O</u> <u>14</u> (853.9)		377.83 Kcal/mol	104.81 Kcal/mol
Palbociclib	5330286	<u>C</u> <u>24</u> <u>H</u> <u>29</u> <u>N</u> <u>7</u> <u>O</u> <u>2</u> (447.5)		112.47 Kcal/mol	34.66 Kcal/mol
Palonosetron	6337614	<u>C</u> <u>19</u> <u>H</u> <u>24</u> <u>N</u> <u>2</u> <u>O</u> (296.4)		71.61 Kcal/mol	34.57 Kcal/mol
Panobinostat	6918837	<u>C</u> <u>21</u> <u>H</u> <u>23</u> <u>N</u> <u>3</u> <u>O</u> <u>2</u> (349.4)		91.83 Kcal/mol	26.32 Kcal/mol
Panobinostat lactate	23725423	<u>C</u> <u>24</u> <u>H</u> <u>29</u> <u>N</u> <u>3</u> <u>O</u> <u>5</u> (439.5)		91.83 Kcal/mol	26.32 Kcal/mol
Pazopanib	10113978	<u>C</u> <u>21</u> <u>H</u> <u>23</u> <u>N</u> <u>7</u> <u>O</u> <u>2</u> <u>S</u> (437.5)		166.35 Kcal/mol	57.22 Kcal/mol

Pefloxacin mesylate	119525	<u>C</u> <u>1</u> <u>8</u> <u>H</u> <u>2</u> <u>4</u> <u>F</u> <u>N</u> <u>3</u> <u>O</u> <u>6</u> <u>S</u> (429.5)		87.76 Kcal/mol	20.81 Kcal/mol
Pexidartinib hydrochloride	73053710	<u>C</u> <u>2</u> <u>0</u> <u>H</u> <u>1</u> <u>6</u> <u>Cl</u> <u>2</u> <u>F</u> <u>3</u> <u>N</u> <u>5</u> (454.3)		108.61 Kcal/mol	20.32 Kcal/mol
Pirfenidone	40632	<u>C</u> <u>1</u> <u>2</u> <u>H</u> <u>1</u> <u>1</u> <u>N</u> <u>O</u> (185.22)		Kcal/mol	Kcal/mol
Plerixafor	65015	<u>C</u> <u>2</u> <u>8</u> <u>H</u> <u>5</u> <u>4</u> <u>N</u> <u>8</u> (502.8)		80.82 Kcal/mol	52.15 Kcal/mol
Pomalidomide	134780	<u>C</u> <u>1</u> <u>3</u> <u>H</u> <u>1</u> <u>1</u> <u>N</u> <u>3</u> <u>O</u> (4273.24)		98.87 Kcal/mol	22.36 Kcal/mol
Ponatinib	24826799	<u>C</u> <u>2</u> <u>9</u> <u>H</u> <u>2</u> <u>7</u> <u>F</u> <u>3</u> <u>N</u> <u>6</u> <u>O</u> (532.6)		615.46 Kcal/mol	38.82 Kcal/mol
Ponatinib hydrochloride	46908927	<u>C</u> <u>2</u> <u>9</u> <u>H</u> <u>2</u> <u>8</u> <u>Cl</u> <u>F</u> <u>3</u> <u>N</u> <u>6</u> <u>O</u> (569.0)		615.46 Kcal/mol	38.82 Kcal/mol

Porfimer sodium	137321858	<u>C₃₄H₃₈N₄NaO₅</u> [±] (605.7)		187.59 Kcal/mol	60.68 Kcal/mol
Pralatrexate	148121	<u>C₂₃H₂₃N₇O₅</u> (477.5)		584.65 Kcal/mol	10.35 Kcal/mol
Pralsetinib	129073603	<u>C₂₇H₃₂FN₉O₂</u> (533.6)		140.80 Kcal/mol	41.37 Kcal/mol
Premarin	23667301	<u>C₁₈H₂₁NaO₅S</u> (372.4)		133.15 Kcal/mol	56.51 Kcal/mol
Pregabalin	5486971	<u>C₈H₁₇NO₂</u> (159.23)		Kcal/mol	Kcal/mol
Quadramet	6918027	<u>C₆H₂₀N₂O₁₂P₄Sm</u> (589.05)		231.74 Kcal/mol	99.44 Kcal/mol
Raloxifene	5035	<u>C₂₈H₂₇NO₄S</u> (473.6)		133.38 Kcal/mol	39.35 Kcal/mol

Raloxifene hydrochloride	54900	$\text{C}_{28}\text{H}_{28}\text{ClNO}_4\text{S}$ (510.0)		133.38 Kcal/mol	39.35 Kcal/mol
Regorafenib	11167602	$\text{C}_{21}\text{H}_{15}\text{ClF}_4\text{N}_4\text{O}_3$ (482.8)		139.04 Kcal/mol	22.51 Kcal/mol
Ribociclib	44631912	$\text{C}_{23}\text{H}_{30}\text{N}_8\text{O}$ (434.5)		Kcal/mol	Kcal/mol
Rolapitant	10311306	$\text{C}_{25}\text{H}_{26}\text{F}_6\text{N}_2\text{O}_2$ (500.5)		126.69 Kcal/mol	25.03 Kcal/mol
Romidepsin	5352062	$\text{C}_{24}\text{H}_{36}\text{N}_4\text{O}_6\text{S}_2$ (540.7)		151.38 Kcal/mol	23.03 Kcal/mol
Rucaparib	9931954	$\text{C}_{19}\text{H}_{18}\text{FN}_3\text{O}$ (323.4)		88.16 Kcal/mol	30.15 Kcal/mol
Sacituzumab govitecan	91668186	$\text{C}_{76}\text{H}_{104}\text{N}_{12}\text{O}_{24}\text{S}$ (1601.8)		407.54 Kcal/mol	81.69 Kcal/mol

Sancuso	5284566	<u>C₁₈H₂₄N₄O</u> (312.4)		82.59 Kcal/mol	40.08 Kcal/mol
Secretin	16129665	<u>C₁₃₀H₂₁₉N₄₃O₄₂</u> (3056.4)		1026.73 Kcal/mol	82.26 Kcal/mol
Selinexor	71481097	<u>C₁₇H₁₁F₆N₇O</u> (443.3)		143.49 Kcal/mol	20.49 Kcal/mol
Sonidegib	24775005	<u>C₂₆H₂₆F₃N₃O₃</u> (485.5)		130.03 Kcal/mol	30.42 Kcal/mol
Sorafenib	<u>216239</u>	<u>C₂₁H₁₆ClF₃N₄O₃</u> (464.8)		133.75 Kcal/mol	22.55 Kcal/mol
Sordinol	12454	<u>C₂₂H₂₅ClN₂OS</u> (401.0)		80.78 Kcal/mol	29.99 Kcal/mol
Sunitinib	5329102	<u>C₂₂H₂₇FN₄O₂</u> (398.5)		131.12 Kcal/mol	47.29 Kcal/mol

Talazoparib	135565082	<u>C₁₉H₁₄F₂N₆O</u> (380.4)		97.35 Kcal/mol	21.35 Kcal/mol
Tamoxifen	2733526	<u>C₂₆H₂₉NO</u> (371.5)		93.02 Kcal/mol	24.71 Kcal/mol
Technetium	<u>23957</u>	<u>Tc</u> (96.90636)		Kcal/mol	Kcal/mol
Telotristat ethyl	25181577	<u>C₂₇H₂₆ClF₃N₆O₃</u> (575.0)		140.85 Kcal/mol	25.19 Kcal/mol
Temozolomide	5394	<u>C₆H₆N₆O₂</u> (194.15)		64.76 Kcal/mol	19.33 Kcal/mol
Temsirolimus	6918289	<u>C₅₆H₈₇NO₁₆</u> (1030.3)		296.57 Kcal/mol	74.90 Kcal/mol
Tepotinib	25171648	<u>C₂₉H₂₈N₆O₂</u> (492.6)		365.8 Kcal/mol	21.47 Kcal/mol

Tivozanib	9911830	<u>C₂₂H₁₉ClN₄O₅</u> (454.9)		126.0 Kcal/mol	35.66 Kcal/mol
Tivozanib hydrochloride	11547978	<u>C₂₂H₂₂Cl₂N₄O₆</u> (509.3)		126.0 Kcal/mol	35.66 Kcal/mol
Topotecan	60700	<u>C₂₃H₂₃N₃O₅</u> (421.4)		111.58 Kcal/mol	28.14 Kcal/mol
Topotecan hydrochloride	60699	<u>C₂₃H₂₄ClN₃O₅</u> (457.9)		111.58 Kcal/mol	28.14 Kcal/mol
Trabectedin	108150	<u>C₃₉H₄₃N₃O₁₁S</u> (761.8)		Kcal/mol	Kcal/mol
Trametinib	11707110	<u>C₂₆H₂₃FIN₅O₄</u> (615.4)		328.5 Kcal/mol	199.13 Kcal/mol
Trilaciclib dihydrochloride	124081865	<u>C₂₄H₃₂Cl₂N₈O</u> (519.5)		106.6 Kcal/mol	44.28 Kcal/mol

Triptorelin pamoate 2d	25074469	<u>C</u> <u>8</u> <u>7</u> <u>H</u> <u>9</u> <u>8</u> <u>N</u> <u>1</u> <u>8</u> <u>O</u> <u>1</u> <u>9</u> (1699.8)		622.43 Kcal/mol	76.75 Kcal/mol
Tucatinib	51039094	<u>C</u> <u>2</u> <u>6</u> <u>H</u> <u>2</u> <u>4</u> <u>N</u> <u>8</u> <u>O</u> <u>2</u> (480.5)		132.9 Kcal/mol	40.68 Kcal/mol
Umbralisib	72950888	<u>C</u> <u>3</u> <u>1</u> <u>H</u> <u>2</u> <u>4</u> <u>F</u> <u>3</u> <u>N</u> <u>5</u> <u>O</u> <u>3</u> (571.5)		150.52 Kcal/mol	31.44 Kcal/mol
Uridine triacetate	20058	<u>C</u> <u>1</u> <u>5</u> <u>H</u> <u>1</u> <u>8</u> <u>N</u> <u>2</u> <u>O</u> <u>9</u> (370.31)		110.57 Kcal/mol	11.00 Kcal/mol
Valrubicin	454216	<u>C</u> <u>3</u> <u>4</u> <u>H</u> <u>3</u> <u>6</u> <u>F</u> <u>3</u> <u>N</u> <u>O</u> <u>1</u> <u>3</u> (723.6)		222.82 Kcal/mol	37.77 Kcal/mol
Uroxatral	25137863	<u>C</u> <u>1</u> <u>9</u> <u>H</u> <u>2</u> <u>8</u> <u>C</u> <u>l</u> <u>N</u> <u>5</u> <u>O</u> <u>4</u> (425.9)		Kcal/mol	Kcal/mol
Vandetanib	3081361	<u>C</u> <u>2</u> <u>2</u> <u>H</u> <u>2</u> <u>4</u> <u>Br</u> <u>F</u> <u>N</u> <u>4</u> <u>O</u> <u>2</u> (475.4)		84.68 Kcal/mol	24.63 Kcal/mol
Vemurafenib	42611257	<u>C</u> <u>2</u> <u>3</u> <u>H</u> <u>1</u> <u>8</u> <u>C</u> <u>l</u> <u>F</u> <u>2</u> <u>N</u> <u>3</u> <u>O</u> <u>3</u> <u>S</u> (489.9)		179.24 Kcal/mol	54.66 Kcal/mol

Venetoclax	49846579	<u>C</u> <u>4</u> <u>5</u> <u>H</u> <u>5</u> <u>0</u> <u>Cl</u> <u>N</u> <u>7</u> <u>O</u> <u>7</u> <u>S</u> (868.4)		372.50 Kcal/mol	80.60 Kcal/mol
Vismodegib	24776445	<u>C</u> <u>1</u> <u>9</u> <u>H</u> <u>1</u> <u>4</u> <u>Cl</u> <u>2</u> <u>N</u> <u>2</u> <u>O</u> <u>3</u> <u>S</u> (421.3)		148.64 Kcal/mol	43.01 Kcal/mol
Vyxeos	11422859	<u>C</u> <u>3</u> <u>6</u> <u>H</u> <u>4</u> <u>2</u> <u>N</u> <u>4</u> <u>O</u> <u>1</u> <u>5</u> (770.7)		225.20 Kcal/mol	51.24 Kcal/mol
Zanubrutinib	135565884	<u>C</u> <u>2</u> <u>7</u> <u>H</u> <u>2</u> <u>9</u> <u>N</u> <u>5</u> <u>O</u> <u>3</u> (471.5)		131.26 Kcal/mol	41.94 Kcal/mol
Zoledronic acid	68740	<u>C</u> <u>5</u> <u>H</u> <u>1</u> <u>0</u> <u>N</u> <u>2</u> <u>O</u> <u>7</u> <u>P</u> <u>2</u> (272.09)		140.04 Kcal/mol	56.99 Kcal/mol

Table 2 MolDock score, rerank score, interaction, torsions and hbond energy of the docked compounds with NOS2 protein.

Name of the inhibitor	MolDock Score Kcal/mol	Rerank Score Kcal/mol	Interaction energy Kcal/mol	Torsion angle Kcal/mol	HBond Kcal/mol

Abarelix	-157.81	-0.93	-88.92	38	-2.82
Abemaciclib	-110.70	-79.76	-114.47	7	-1.06
Abiraterone acetate	-88.80	-68.04	-95.37	3	-1.60
Acalabrutinib	-113.68	-77.93	-104.46	6	-2.20
Afatinib	-115.59	-82.64	-119.65	8	-4.12
Afatinib dimaleate	-113.95	-18.00	-100.96	8	-2.58
Alectinib	-112.83	--83.57	-119.19	4	-0.85
Alimta	-124.39	-81.05	-119.96	9	-5.10
Alpelisib	-99.15	-79.74	-102.08	6	-3.75

Amifostine	-60.77	-41.28	-58.29	7	-1.48
Anagrelide	-77.11	-59.47	-78.49	0	0
Anastrozole	-82.61	-44.79	-86.85	6	-5.05
Andrographo lide	-75.25	-58.00	-77.85	4	-8.86
Anexsia	-59.88	-48.51	-62.40	1	-2.80
Apalutamide	-98.34	-70.24	-88.52	5	-1.46
Aprepitant	-117.15	-74.06	-114.18	8	0
Aredia	-52.58	-39.92	-54.66	4	-4.20
Arzerra	-89.34	-71.87	-88.85	1	-2.5

Axitinib	-107.75	-76.35	-104.53	5	-0.18
Belinostat	-95.39	-62.95	-76.57	5	-0.45
Bendamustine	-101.90	-59.29	-99.32	9	-2.6
Besponsa	-167.05	-59.76	-97.25	13	-1.91
Benimetinib	-81.37	-45.09	-103.65	6	-3.76

Bortezomib	-97.34	-64.69	-97.64	9	-5.43
Bosutinib	-116.04	-58.63	-88.59	10	-1.47
Brigatinib	-78.77	-40.92	-96.56	8	-1.87
Bromfenac	-82.54	-67.10	-85.77	4	-0.09
Cabazitaxel	-130.75	-66.02	-104.44	16	-1.38
Cabozantinib	-107.09	-86.89	-124.84	8	-
Capecitabine	-91.76	20.55	-99.02	7	-6.55
Carfilzomib	-118.56	-58.62	-107.65	20	-3.05
Carmustine	-65.14	-51.94	-65.12	5	-2.96

Cedazuridine	-68.87	-60.84	-81.77	2	-7.41
Cisplatin	-21.76	-25.30	-21.76	0	-4.8
Cinacalcet	-91.56	-64.25	-87.26	7	0
Clofarabine	-86.66	-68.85	-86.68	2	-6.11
Cobimetinib	-114.111	-88.24	-117.28	4	-2.62
Copanlisib	-110.56	-51.89	-109.47	8	-0.23
Copiktra	-84.23	4.34	-88.58	4	-0.85
Crizotinib	-108.28	-78.07	-106.19	5	-0.03
Dabrafenib	-101.94	-60.11	-78.09	6	0

Dacomitinib	-87.37	-63.81	-96.69	8	-0.35
Darolutamide	-131.34	-83.17	-116.02	7	-5.67
Dasatinib	-111.44	-37.08	-110.99	6	-3.71
Decitabine	-80.81	-55.99	-84.11	2	-6.55
Degarelix	-226.10	-106.48	-226.47	42	-10.57
Docetaxel	-125.02	-72.22	-114.53	14	-4.38
Doxorubicin	-41.22	-48.19	-102.29	5	-8.92
Elliotts Region 10	-36.43	-30.21	-36.43	0	-5.63
Elliotts Region 20	-45.60	-43.73	-64.28	1	-9.11

Elliotts Region 5	-23.25	-20.59	-23.25	0	-3.05
Eloxatin Region	-35.65	-33.73	-46.47	0	-2.24
Eloxatinregion n 1	-39.77	-35.02	-43.60	1	-5.00

Enasidenib	-65.55	-62.06	-97.55	8	-0.04
Encorafenib	-137.86	-85.89	-111.07	10	-0.72
Entrectinib	-121.73	-65.00	-96.89	7	-0.53
Enzalutamide	-92.14	-53.70	-88.76	5	-2.5
Epirubicin	-48.75	-50.50	-90.22	5	-7.23
Erdafitinib	-107.92	-74.48	-102.36	9	-7.50
Erlotinib	-124.80	-88.19	-126.94	11	-0.97
Everolimus	-111.73	-75.61	-110.00	9	-6.11
Exemestane	-93.36	-69.43	-94.16	0	-1.63

Fedratinib	-117.43	-65.67	-119.91	11	-0.07
Fentanyl	-66.66	21.70	-83.94	6	0
Fentanyl citrate	-72.74	-56.61	-93.85	6	-2.5
Flutamide	-76.50	-63.66	-80.19	4	-5.4
Fulvestrant	-97.24	-59.85	-106.40	15	-3.8
Fusilev	-110.75	-72.73	-91.98	10	-4.9
Gefitinib	-111.97	-69.10	-102.30	8	-2.7
Gilteritinib	-124.01	-91.63	-124.53	9	-4.6
Glasdegib	-116.23	-30.53	-100.83	6	-0.57

Goserelin	-189.77	-92.09	-133.3	34	-7.7
Granisetron	-80.37	-55.51	-83.82	2	0
Ibritumomab	-114.25	-59.08	-92.89	21	-3.9
Ibrutinib	-107.41	-72.47	-106.97	5	-2.0
Idelalisib	-103.54	-75.62	-103.18	5	-0.4
Imatinib	-106.68	-34.06	-120.29	7	-2.5
Ingenol mebutate	-87.17	-50.38	-89.31	5	-6.7
Interferon alfa-2B	-108.07	-68.29	-92.12	8	0
Iodixanol	-144.14	-74.14	-94.80	22	-4.0

Irinotecan	-112.18	-75.20	-124.62	5	-3.9
Ivosidenib	-156.93	-73.68	-119.95	8	-0.3
Ixabepilone	-110.69	-64.43	-79.83	3	-3.6
Ixazomib	-95.99	-67.60	-88.00	7	-4.5
Lapatinib	-120.46	-43.73	-92.37	11	-3.9
Larotrectinib	-134.84	-76.76	-95.18	5	0
Lenalidomide	-77.62	-64.36	-87.35	1	-0.6
Lenvatinib	-108.85	-68.09	-100.74	8	-2.0
Letrozole	-90.19	-70.54	-94.06	5	0

Leuprolide acetate	-154.10	-54.06	-83.42	32	-7.3
Leuprorelin	-164.38	-62.48	-123.52	32	-6.0
Levoleucovorin	-115.66	-71.22	-94.11	10	-5.15
Lonsurf	-73.91	-64.55	-83.91	3	-5.35
Lorlatinib	-117.89	-76.58	-100.39	1	-1.73
Marqibo	-128.23	-61.32	-96.05	11	-0.56
Mechlorethamine	-45.91	-35.92	-47.06	4	0
Melphalan flufenamide	-117.04	-67.83	-103.58	14	-0.62
Methoxsalen	-78.31	-47.00	-80.89	1	-1.79

Midostaurin	-142.42	-91.15	-117.86	3	-0.47
Miraluma	-42.80	-38.82	-47.08	3	-3.16
Morphine sulfate	-80.13	-49.42	-86.61	0	-5.18
Nelarabine	-85.12	-66.53	-89.28	3	-6.47
Neratinib	-103.98	-77.61	-113.81	12	-0.64
Neratinib maleate	-117.95	-90.09	-117.32	12	-2.00
Netupitant	-109.24	-68.92	-112.85	7	-2.46
Nilotinib	-125.02	-82.46	-124.81	7	-2.50
Niraparib	-76.81	-64.25	-87.52	3	-2.38

Olaparib	-76.45	-58.21	-91.69	4	-1.24
Omacetaxine mepesuccinate	-101.87	-47.71	-85.96	11	-4.18
Ondansetron	-88.11	-62.67	-86.36	2	-1.80
Osimertinib	-80.97	-66.57	-83.53	10	-3.91
Oxaliplatin	-46.54	-40.41	-52.19	0	-1.83
Paclitaxel	-132.68	-60.71	-108.71	14	-0.94
Palbociclib	-101.88	-72.68	-96.65	4	-3.87
Palonosetron	-70.65	-63.12	-92.20	1	0
Panobinostat	-102.40	-68.77	-98.26	7	-5.15

Panobinostat lactate	-127.42	-56.35	-125.90	7	-2.11
Pazopanib	-128.95	-95.28	-115.36	5	-2.5
Pefloxacin mesylate	-91.76	-75.39	-97.96	3	-2.14
Pexidartinib hydrochloride	-114.10	-73.64	-108.80	6	0
Pirfenidone	-51.70	-35.69	-70.22	1	-0.29
Plerixafor	-128.58	-83.06	-108.01	4	-0.72
Pomalidomide	-84.84	-70.11	-91.58	1	-2.50
Ponatinib	-133.52	-99.85	-136.60	8	-2.79
Ponatinib hydrochloride	-113.27	-89.39	-123.75	8	-0.07

Porfimer sodium	-138.19	-84.70	-104.66	8	-4.07
Pralatrexate	-125.24	-70.01	-103.10	11	-2.34
Pralsetinib	-129.3	-77.32	-94.91	8	0
Premarin	-94.11	-53.64	-95.13	2	-1.79
Pregabalin	-62.75	-48.37	-61.69	5	-3.0
Quadramet	-83.41	-56.73	-82.60	11	-1.53
Raloxifene	-136.16	-86.90	-107.13	7	-1.68
Regorafenib	-126.41	-87.24	-114.04	8	-3.33
Rolapitant	-114.82	-74.44	-116.35	7	0

Romidepsin	-114.11	-70.00	-96.65	3	-0.93
Rucaparib	-94.88	-71.76	-96.57	3	0
Sancuso	-80.33	-55.64	-83.72	2	0
Sorafenib	-119.79	-81.70	-102.21	8	-5.1
Sordinol	-107.71	-21.19	-94.20	6	0
Sunitinib	-124.80	-83.98	-103.60	8	-0.8
Talazoparib	-116.04	-80.38	-110.84	2	-2.0
Tamoxifen	-106.25	-80.40	-99.08	9	0
Telotristat ethyl	-122.38	-37.67	-91.80	11	-3.7

Temozolomi de	-70.12	-61.23	-73.40	1	-3.2
Temsirolimu s	-100.64	-71.73	-109.37	11	-4.2
Tepotinib	-100.97	-29.96	-109.74	8	-1.6
Tivozanib	-132.09	-83.29	-100.45	8	-1.0
Trametinib	-96.02	-19.80	-105.67	5	-4.54
Trilaciclib dihydrochlori de	-133.21	-83.41	-130.22	3	-0.20
Triptorelin pamoate	-185.73	-59.99	-129.30	33	-2.72
Tucatinib	-121.69	-66.33	-96.52	5	-2.40
Umbralisib	-147.49	-87.26	-106.60	6	-0.68

Uridine triacetate	-116.59	-87.17	-115.36	8	-5.08
Uroxatral	--117.41	-55.64	-111.05	8	-1.13
Valrubicin	--115.98	-61.15	-115.51	12	-5.78
Vandetanib	-102.72	-77.22	-112.77	6	-3.16
Vemurafenib	-124.80	-69.76	-94.51	7	-0.31
Venetoclax	-168.21	-64.67	-109.12	12	-0.54
Vismodegib	-97.78	-73.73	-97.91	4	-1.51
Vyxeos	-73.33	-59.61	-78.45	2	-3.41
Zanubrutinib	-101.69	-74.84	-100.71	6	-0.81

Zoledronic acid	-62.61	-22.65	-63.54	4	-3.33
--------------------	--------	--------	--------	---	-------

Table 3 Amino acid residues around active site and docked against NOS2 protein

Name of the Inhibitor	Amino acid residues around active site	Ligand binding amino acids
Degarelix	Glu394C,Glu393C,Asn390C,Val386C,Glu387C,Glu91D,Pro474D,Ser92D,His477D,Glu441D,Lys445D,Ile204D,Asn449D,Glu448D,Met444D,Met94D,Asn89D,Lys88D,Thr95D,Arg86D	Asn449D,Met94D,Ser92D,Pro474D,Val386C,Asn89D
Gosarelin	Val386C,Glu387C,Gly91D,Glu93D,Ser92D,Asu89D,Met94D,Phe96D,Glu448D,Met447D,Lys445D,Met444D,Glu478D,Pro474D,Gly441D	Glu441D,Gly91D,Met444D,Ser92D
Triptorelin Pamoate	Pro474D,Val386C,Glu441D,Lys445D,Met444D,Arg452D,Asn449D,Glu448D,Phe96D,Met94D,Gly478D,Asn89D,Ser92D	Glu448D,Met444D,Ser92D
Venetoclax	Gly93D,Met94D,Asn89D,Ser92D,Glu448D,Lys445D,Met444D,Pro474D,Glu441D	Ser92D,Glu448D

Table 4 Pharmacophore kinetic and drug-likeness and ADMET properties of top selected ligands

Property	Model Name	Degarelix	Goserelin	Triptorelin pamoate	Venetoclax	Unit
Absorption	Water solubility	-2.89	-2.89	-2.89	-3.03	log mol/L
	Caco2 permeability	0.622	-1.50	-2.58	0.84	log Papp in 10–6 cm/s
	Intestinal absorption(Human)	0	0	0	100	%Absorbed
	Skin Permeability	-2.73	-2.73	-2.73	-2.73	log Kp
Distribution	VDss(Human)	-0.536	-0.24	-0.012	0.329	log l/kg
	Fraction unbound(Human)	0.276	0.36	0.379	0.169	Fu
	BBB permeability	-2.726	-3.43	-4.13	-1.74	log BB
	CNS permeability	-6.852	-7.31	-7.49	-3.11	logPS
Metabolism	CYP2D6 substrate	No	No	No	No	Yes/No
	CYP3A4 substrate	Yes	Yes	No	Yes	
	CYP1A2 inhibitor	No	No	No	No	
	CYP2C19 inhibitor	No	No	No	No	
	CYP2C9 inhibitor	No	No	No	No	
	CYP2D6 inhibitor	No	No	No	NO	
	CYP3A4 inhibitor	No	No	No	Yes	
Excretion	Total clearance	-0.897	0.414	-1.04	-0.09	log ml/min/kg
	Renal OCT2 substrate	No	No	No	No	Yes/No
Toxicity	AMES toxicity	No	No	Yes	No	
	Max. tolerated dose(Human)	0.438	0.438	0.438	0.278	log mg/kg/day
	hERG I inhibitor	No	No	No	No	Yes/No
	Oral rat acute toxicity(LD50)	2.482	2.482	2.482	2.60	Mol/kg

	Oral rat chronic toxicity(LOAEL)	10.202	7.479	10.343	1.924	log mg/kg_bw/day
	Hepatotoxicity	Yes	Yes	No	Yes	Yes/No
	Skin sensitization	No	No	No	No	
	T.pyriformis toxicity	0.285	0.285	0.285	0.285	log µg/L
	Minnow toxicity	6.928	10.099	7.236	-0.48	log µg/L
Drug likeness	Lipinski	No	No	No	No	Yes/No
Bioactivity Score	GPCR ligand	-4.01	-3.84	-3.87	-2.36	
	Ion channel modulator	-4.04	-3.92	-3.94	-3.42	
	Kinase inhibitor	-4.05	-3.93	-3.95	-3.00	
	Nuclear receptor ligand	-4.08	-3.98	-3.99	-3.33	
	Protease inhibitor	-3.98	-3.77	-3.82	-1.92	
	Enzyme inhibitor	-4.03	-3.88	-3.91	-2.87	

References

1. Arbyn, M.; Weiderpass, E.; Bruni, L.; de Sanjosé, S.; Saraiya, M.; Ferlay, J.; Bray, F. Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. *Lancet Glob. Heal.* **2020**, *8*, e191–e203, doi:10.1016/S2214-109X(19)30482-6.
2. Fontham, E.T.H.; Wolf, A.M.D.; Church, T.R.; Etzioni, R.; Flowers, C.R.; Herzig, A.; Guerra, C.E.; Oeffinger, K.C.; Shih, Y.T.; Walter, L.C.; et al. Cervical cancer screening for individuals at average risk: 2020 guideline update from the American Cancer Society. *CA. Cancer J. Clin.* **2020**, *70*, 321–346, doi:10.3322/caac.21628.
3. Kaarthigeyan, K. Cervical cancer in India and HPV vaccination. *Indian J. Med. Paediatr. Oncol.* **2012**, *33*, 7–12, doi:10.4103/0971-5851.96961.
4. Ambs, S.; Merriam, W.G.; Bennett, W.P.; Felley-Bosco, E.; Ogunfusika, M.O.; Oser, S.M.; Klein, S.; Shields, P.G.; Billiar, T.R.; Harris, C.C. Frequent nitric oxide synthase-2 expression in human colon adenomas: Implication for tumor angiogenesis and colon cancer progression. *Cancer Res.* **1998**, *58*, 334–341.
5. Hussain, S.P.; Trivers, G.E.; Hofseth, L.J.; He, P.; Shaikh, I.; Mechanic, L.E.; Doja, S.; Jiang, W.; Subleski, J.; Shorts, L.; et al. Nitric oxide, a mediator of inflammation, suppresses tumorigenesis. *Cancer Res.* **2004**, *64*, 6849–6853, doi:10.1158/0008-5472.CAN-04-2201.
6. Chen, H.H.W.; Su, W.C.; Chou, C.Y.; Guo, H.R.; Ho, S.Y.; Que, J.; Lee, W.Y. Increased expression of nitric oxide synthase and cyclooxygenase-2 is associated with poor survival in cervical cancer treated with radiotherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2005**, *63*, 1093–1100, doi:10.1016/j.ijrobp.2005.03.062.
7. Pasha, A.; Kumbhakar, D.V.; Doneti, R.; Kumar, K.; Dharmapuri, G.; Poleboyina, P.K.; S. K., H.; Basavaraju, P.; Pasumarthi, D.; S. D., A.; et al. Inhibition of Inducible Nitric Oxide Synthase (iNOS) by Andrographolide and in Vitro Evaluation of Its Antiproliferative and Proapoptotic Effects on Cervical Cancer. *Oxid. Med. Cell. Longev.* **2021**, *2021*, doi:10.1155/2021/6692628.
8. RCSB PDB: Homepage Available online: <https://www.rcsb.org/> (accessed on Jun 24, 2021).
9. PubChem Available online: <https://pubchem.ncbi.nlm.nih.gov/> (accessed on Jun 24, 2021).
10. Irwin, J.J.; Sterling, T.; Mysinger, M.M.; Bolstad, E.S.; Coleman, R.G. ZINC: A free tool to discover chemistry for biology. *J. Chem. Inf. Model.* **2012**, *52*, 1757–1768.
11. Wishart, D.S.; Feunang, Y.D.; Guo, A.C.; Lo, E.J.; Marcu, A.; Grant, J.R.; Sajed, T.; Johnson, D.; Li, C.; Sayeeda, Z.; et al. DrugBank 5.0: A major update to the DrugBank database for 2018. *Nucleic Acids Res.* **2018**, *46*, D1074–D1082, doi:10.1093/nar/gkx1037.

12. Hyper Available online: <https://hyper.com/> (accessed on Jun 24, 2021).
13. ChemDraw - PerkinElmer Available online: <https://perkinelmerinformatics.com/products/research/chemdraw/> (accessed on Jun 24, 2021).
14. Bitencourt-Ferreira, G.; de Azevedo, W.F. Molegro virtual docker for docking. In *Methods in Molecular Biology*; Humana Press Inc., 2019; Vol. 2053, pp. 149–167.
15. Molegro Data Modeller | Macs in Chemistry Available online: <https://www.macinchem.org/blog/files/671a3395f65b525e75347a60edf7935e-236.php> (accessed on Jun 24, 2021).
16. Molegro Molecular Viewer – Molexus Available online: <http://molexus.io/molegro-molecular-viewer/> (accessed on Jun 24, 2021).
17. Daoud, I.; Melkemi, N.; Salah, T.; Ghalem, S. Combined QSAR, molecular docking and molecular dynamics study on new Acetylcholinesterase and Butyrylcholinesterase inhibitors. **2018**, doi:10.1016/j.compbiolchem.2018.03.021.
18. Sadeghi, F.; Afkhami, A.; Madrakian, T.; Ghavami, R. Computational study to select the capable anthracycline derivatives through an overview of drug structure-specificity and cancer cell line-specificity. *Chem. Pap.* **2021**, 75, 523–538, doi:10.1007/s11696-020-01321-z.
19. Hocquet, A.; Langgrd, M. An evaluation of the MM+ force field. *J. Mol. Model.* **1998**, 4, 94–112, doi:10.1007/s008940050128.
20. Swiss PDB Viewer - Home Available online: <https://spdbv.vital-it.ch/> (accessed on Jun 24, 2021).
21. Balakumar, C.; Ramesh, M.; Tham, C.L.; Khathi, S.P.; Kozielski, F.; Srinivasulu, C.; Hampannavar, G.A.; Sayyad, N.; Soliman, M.E.; Karpoormath, R. Ligand- and structure-based in silico studies to identify kinesin spindle protein (KSP) inhibitors as potential anticancer agents. *J. Biomol. Struct. Dyn.* **2018**, 36, 3687–3704, doi:10.1080/07391102.2017.1396255.
22. Prasad, C.; Rao, A.V.; Rao, M.V. Computer aided design and molecular docking studies on a series of 1, 3-thiazolidine-2, 4-diones as new class of 5-lipoxygenase inhibitors. **2014**, 8, 858–863.
23. Kaushik, P.; Lal Khokra, S.; Rana, A.C.; Kaushik, D. Pharmacophore modeling and molecular docking studies on pinus roxburghii as a target for diabetes mellitus. *Adv. Bioinformatics* **2014**, 2014, doi:10.1155/2014/903246.
24. Yang, J.M.; Chen, C.C. GEMDOCK: A Generic Evolutionary Method for Molecular Docking. *Proteins Struct. Funct. Genet.* **2004**, 55, 288–304, doi:10.1002/prot.20035.
25. Wolber, G.; Langer, T. LigandScout: 3-D pharmacophores derived from protein-bound ligands and their use as virtual screening filters. *J. Chem. Inf. Model.* **2005**, 45, 160–169,

doi:10.1021/ci049885e.

26. Rella, M.; Rushworth, C.A.; Guy, J.L.; Turner, A.J.; Langer, T.; Jackson, R.M. Structure-based pharmacophore design and virtual screening for novel Angiotensin Converting Enzyme 2 inhibitors. *J. Chem. Inf. Model.* **2006**, *46*, 708–716, doi:10.1021/ci0503614.
27. Alhazmi, M.I. Hypothesis Volume 11(4) Molecular docking of selected phytochemicals with H1N1 Proteins. *Bioinformation* **2015**, *11*, 196.
28. Thangavelu, R.; Faisal, S. Prediction of molecular properties, bioactivity & docking of novel hydrazines. *Int. J. Trends Pharm. Life Sci.* **2016**, *2*, 757–768.
29. Yadav, A.R.; Mohite, S.K. *Anticancer Activity and In-Silico ADMET Analysis of Malvastrum Coromandelianum*;
30. Pires, D.E. V; Blundell, T.L.; Ascher, D.B.; 1ga, U.K. pkCSM: Predicting Small-Molecule Pharmacokinetic and Toxicity Properties Using Graph-Based Signatures. **2015**, doi:10.1021/acs.jmedchem.5b00104.
31. Laskowski, R.A.; Jabło nska, J.; Pravda, L.; Svobodov Va rekov, R.; Thornton, J.M. TOOLS FOR PROTEIN SCIENCE PDBsum: Structural summaries of PDB entries. **2017**, doi:10.1002/pro.3289.
32. Laskowski, R.A.; MacArthur, M.W.; Moss, D.S.; Thornton, J.M. PROCHECK: a program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.* **1993**, *26*, 283–291, doi:10.1107/s0021889892009944.
33. Colovos, C.; Yeates, T.O. Verification of protein structures: Patterns of nonbonded atomic interactions. *Protein Sci.* **1993**, *2*, 1511–1519, doi:10.1002/pro.5560020916.
34. SAVESv6.0 - Structure Validation Server Available online: <https://saves.mbi.ucla.edu/> (accessed on Jun 25, 2021).
35. Copeland, R.A. Conformational adaptation in drug-target interactions and residence time. *Future Med. Chem.* **2011**, *3*, 1491–1501.
36. Guan, L.; Yang, H.; Cai, Y.; Sun, L.; Di, P.; Li, W.; Liu, G.; Tang, Y. ADMET-score-a comprehensive scoring function for evaluation of chemical drug-likeness. *Medchemcomm* **2019**, *10*, 148–157, doi:10.1039/C8MD00472B.
37. Galea, E.; Feinstein, D.L. Regulation of the expression of the inflammatory nitric oxide synthase (NOS2) by cyclic AMP. **2018**, 2125–2137.
38. Sharma, V.; Sharma, P.C.; Kumar, V. In Silico Molecular Docking Analysis of Natural Pyridoacridines as Anticancer Agents. **2016**, 2016.
39. Shore, N.D. Experience with degarelix in the treatment of prostate cancer. *Ther. Adv. Urol.* **2013**, *5*, 11–24, doi:10.1177/1756287212461048.
40. Jonat, W. Goserelin (Zoladex TM) – its role in early breast cancer in pre- and perimenopausal women. **2001**, *85*, 1–5.

41. Matched, A.R.; Control, C.; Li, Q.; Zhu, Y.; Sun, J.; Li, F. Role of Goserelin in Combination with Endocrine Therapy for the Treatment of Advanced Breast Cancer in Premenopausal Women Positive for Hormone Receptor : **2013**, 28, doi:10.1089/cbr.2012.1436.
42. Aslam N, Nadeem K, Noreen R, J.A.C. Prostate Cancer Prostate Cancer. *Abeloff's Clin. Oncol. 5/e* **2015**, 8, 938–944.
43. Frampton, J.E. Triptorelin: A Review of its Use as an Adjuvant Anticancer Therapy in Early Breast Cancer. *Drugs* **2017**, 77, 2037–2048, doi:10.1007/s40265-017-0849-3.
44. Li, Q.; Cheng, L.; Shen, K.; Jin, H.; Li, H.; Cheng, Y.; Ma, X. Efficacy and safety of BcL-2 inhibitor venetoclax in hematological malignancy: A systematic review and meta-analysis of clinical trials. *Front. Pharmacol.* **2019**, 10, 1–12, doi:10.3389/fphar.2019.00697.