Outcomes of COVID-19 patients that needed emergency general surgery: predictors of mortality and postoperative complications

Katya Bozada-Gutiérrez

Hospital General Dr Manuel Gea Gonzalez

Mario Trejo-Avila (✉ mario.trejo.avila@gmail.com)

Hospital General Dr Manuel Gea Gonzalez <https://orcid.org/0000-0001-9249-3321>

Carlos Valenzuela-Salazar

Hospital General Dr Manuel Gea Gonzalez

Jesús Herrera-Esquivel

Hospital General Dr Manuel Gea Gonzalez

Mucio Moreno-Portillo

Hospital General Dr Manuel Gea Gonzalez

Research Article

Keywords: COVID-19, perioperative outcomes, COVID-19 mortality, surgery in COVID-19, SARS-CoV-2

Posted Date: July 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-676027/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Purpose

There is limited data about the perioperative outcomes of COVID-19 patients that needed emergency general surgery. The aims of the present study were to describe the perioperative outcomes of COVID-19 patients that underwent emergency general surgery and to determine possible predictors of mortality and postoperative complications.

Methods

A prospective study of positive COVID-19 patients that needed an emergency general surgery procedure at our center was performed.

Results

From March 2020 to February 2021, 44 patients were included. We found that patients with SARS-CoV-2 symptomatic disease have increased postoperative complications, higher ICU admissions, prolonged length of stay, and decreased 90-day survival as compared with asymptomatic COVID-19 patients. The 90-day survival probability of the entire cohort was 70.1% (60.3–79.9) and was significantly lower in patients with COVID-19 symptoms 63.4% (50.5–76.2). We found the following cut-off values for the prediction of mortality: ferritin ≥ 438.5 ng/mL (AUC = 0.908), CRP value ≥ 12.5 mg/dL (AUC = 0.715), leukocyte $\geq 13.8 \times 10^3/\mu\text{L}$ (AUC = 0.706), and albumin ≤ 2.78 g/dL (AUC = 0.704). Also, a cut-off value of CRP of ≥ 12.5 mg/dL yielded an accuracy of 82.9% for the prediction of postoperative complications ($p < 0.001$).

Conclusion

Patients with symptomatic COVID-19 that needed emergency surgery have increased postoperative complications, higher ICU admissions, prolonged length of stay, and decreased 90-day survival as compared with asymptomatic COVID-19 patients. Preoperative ferritin, CRP, leukocytes, and albumin could be used as predictors of mortality.

Introduction

A novel coronavirus (severe acute respiratory syndrome coronavirus 2, SARS-CoV-2) emerged in China, in December 2019, causing a new disease named “COVID-19” (coronavirus disease 2019) [1]. SARS-CoV-2 rapidly spread across the world and was declared a pandemic by the World Health Organization (WHO) on March 11, 2020 [1]. In Mexico, until March 31, 2021, a total of 2,455,219 suspected COVID-19 patients were registered, of which 1,993,629 were treated in Mexico City [2].

This pandemic originated unprecedented changes for health systems, hospitals and surgical departments. In the world, and also in Mexico, the non-essential surgical procedures or elective surgeries were cancelled and postponed. In our country, hospitals and institutions were converted to specialized units for attention to COVID-19 patients. Our hospital and surgical department maintained only emergency surgery, so patients with surgical emergencies with and without SARS-CoV-2 infection were treated.

Early reports suggested an increased risk of morbidity and mortality for COVID-19 patients who underwent surgical procedures [3-7]. Patients with acute abdomen or surgical emergencies plus concomitant COVID-19 started to present to the emergency department. The effects of SARS-CoV-2 infection in patients that needed emergency surgery are still poorly described and few evidence have been published [6, 8].

Recently published data suggested an overall postoperative 30-day mortality of 23.8%, with more than half of the patients developing postoperative pulmonary complications [6]. To note, emergency in comparison to elective surgery was associated with 30-day mortality (1.67 [1.06–2.63], $p=0.026$) [6]. Also, recent data suggested that surgery performed ≥ 7 weeks after SARS-CoV-2 diagnosis was associated with a similar mortality risk to baseline, so when possible surgery should be delayed for at least 7 weeks [9]. However, several diseases or abdominal emergencies required fast or early treatment.

Until now, there are limited published data describing the perioperative outcomes of patients that needed an emergency general surgery procedure with concomitant COVID-19. The aims of the present study were to describe the perioperative outcomes of patients with COVID-19 that underwent emergency general surgery and to determine possible predictors of mortality and postoperative complications.

Patients And Methods

Study population

We prospectively enrolled adult patients with confirmed COVID-19 diagnosis that required emergency general surgery procedures from March 2020 to February 2021. Patients were admitted to Hospital General Dr. Manuel Gea González in Mexico City, Mexico. COVID-19 was diagnosed according to the World Health Organization (WHO) COVID-19 case definitions [1, 9-11]. We enrolled symptomatic and asymptomatic SARS-CoV-2 positive patients (asymptomatic carriers) that required emergency surgery. We only included patients with the SARS-CoV-2 infection confirmed either by reverse-transcriptase polymerase chain reaction (RT-PCR) assay of a nasopharyngeal swab, a rapid antigenic test, or with a highly suspicious computed tomography (CT) findings (COVID-19 Reporting and Data

System, CO-RADS category ≥ 5 [10]). All patients had chest CT scans previous to surgery. Patients that developed COVID-19 after surgery, but during the same admission (<7 days after surgery), were included. We enrolled patients that underwent emergency general surgery procedures including: gastrointestinal or abdominal emergencies, and wound debridements for necrotizing soft tissue infections (NSTI).

We excluded patients that required thoracic, neurosurgical or head and neck surgical procedures, we also excluded patients with negative laboratory tests or CT scans with CO-RADS <4 . The study protocol was approved by the ethics and research committee of the hospital and conducted in accordance with the principles of the Declaration of Helsinki. Prospective informed consent was waived by competent authorities due to the use of anonymous data.

Data collection

Several variables were recorded including demographic parameters, preoperative quick Sequential Organ Failure Assessment (qSOFA) score [12], type of surgical procedure, preoperative laboratory studies, type of postoperative complications, data regarding intensive care, hospital length of stay (days), and 90-day mortality.

Demographic data included age (years), gender, body mass index (BMI), comorbidities, smoking status, American Society of Anesthesiology classification (ASA). We calculated the preoperative qSOFA score [12] and divided the patients in high risk (>2 point) or not high risk patients (0-1 points).

The type of emergency general surgery procedures included abdominal, gastrointestinal and wound debridements for necrotizing soft tissue infections (NSTI). Abdominal surgeries were divided in laparoscopic, open or converted. All surgical procedures were performed in a separated theater dedicated to COVID-19 patients. Preoperative laboratory data consisted of a complete blood count, C-reactive protein (CRP) expressed in mg/dL, lactate dehydrogenase (LDH) in IU/L, serum ferritin ng/mL, and albumin in g/dL. Postoperative complications were classified and presented according with the Clavien-Dindo classification [13]. The need for intensive care (ICU), vasopressors and invasive mechanical ventilation, timing of intensive care in relation to surgery (pre- or postoperative) as well as organ failures were recorded. The hospital length of stay (LOS) in days was registered. Survival was calculated at 90-days.

We presented the data of the entire cohort of patients and based on the presence or absence of SARS-CoV-2 symptoms (including pneumonia). Pneumonia was diagnosed by clinical symptoms (fever, dyspnea, cough) and a confirmatory CT scan imaging (consolidation, pleural effusion, etc). COVID-19 related symptoms included fever, cough, dyspnea, fatigue, myalgia and headache, among others [14, 15].

Statistical Analysis

All data was collected prospectively in a digital database. For the purpose of analysis we divided the patients in four groups: based on their survival status (survivors vs. non-survivors) and based on the presence or absence of postoperative complications (postoperative complicated vs. non-complicated). Comparisons of baseline preoperative and postoperative variables were performed. Categorical data were presented as totals (n) and proportions as percentages (%). Categorical data were compared using the Pearson's Chi-square test or Fisher's exact test. Univariate analysis was performed in order to identify possible risk factors for mortality and postoperative complications. All variables with a p value inferior to 0.10 in the univariate analysis were considered as potential risk factors and were entered into multivariate backward logistic regression analysis. The variables representing the lowest risk for each complication was considered to be the reference group (OR=1). Odds ratios (OR) and 95% confidence intervals (95% CI) were calculated for the risk of postoperative complications. Kaplan-Meier method was employed to perform survival analysis (90-day survival) and the differences were compared with the log-rank test. Cox proportional hazards (Hazard ratios, HR) regression model was employed to identify possible independent prognostic factors for 90-day mortality. To evaluate the discriminatory ability (accuracy) of each risk factor, receiver operating characteristic (ROC) curves were generated, and the areas under the curve (AUC) were calculated. The optimal cut-offs were determined by ROC analysis. The sensitivity, specificity, and positive and negative likelihood ratios (LR), of each cut-off value were calculated. A two-tailed p value <0.05 was considered to be statistically significant. All data were analyzed using SPSS statistic version 22.0 (IBM Corporation, Armonk, New York, NY).

Results

We prospectively collected data of 44 COVID-19 confirmed patients that needed an emergency general surgery procedure during the pandemic at our institution. The baseline characteristics of the included patients are presented in **Table 1 and 2**. Concerning the entire cohort, 52.3% were males, the mean age was 53.5 years, the mean BMI was 28.8, 43.2% were current smokers, 59.1% have comorbidities, and 63.6% were classified as High Risk qSOFA score (**Table 1**). Regarding COVID-19 timing of diagnosis, 30 (68.2%) were diagnosed preoperatively and 14 (31.8%) postoperatively.

The surgical procedures performed in this cohort were abdominal (n=39, 88.6%) and soft tissue debridements for NSTI (n= 5, 11.4%). Regarding the abdominal procedures, there were 11 appendectomies, 10 cholecystectomies, and 18 emergency laparotomies. The causes of emergency laparotomies were: perforated gastroduodenal ulcer (n=3), strangulated inguinal hernia (n=1), acute mesenteric ischemia (n=2), penetrating abdominal trauma (n=2), strangulated ovarian cyst (n=1), non-penetrating abdominal trauma (n=1), perforated diverticulitis (n=2), peritoneal dialysis catheter insertion or revision (n=5), and necrotizing pancreatitis (n=1). Abdominal surgeries were performed open (n=20, 51.3%), laparoscopic (n=16, 41%), and laparoscopic converted to open (n=3, 7.7%). Five patients

(11.4%) had necrotizing soft tissue infection that required surgical exploration and several surgical debridements.

Comparisons between COVID-19 patients with symptomatic vs. asymptomatic disease

Of the total of patients, 30 (68.1%) had SARS-CoV-2 associated symptoms (**Table 1**). Patients with symptomatic COVID-19 were significantly associated with older age, higher BMI, were current smokers, had higher ASA score, had more comorbidities (more diabetes and hypertension), and had higher preoperative qSOFA score. Regarding preoperative studies (**Table 3**), patients with symptomatic disease had lower hemoglobin and albumin levels, and had higher CRP, LDH and ferritin levels. Postoperative complications were more frequent in patients with symptoms, as well as more incidence of ARDS, septic shock and MOF. Patients with symptoms require more ICU admissions, mechanical ventilation, vasopressors, and had more prolonged LOS. Mortality was more frequent in patients with symptomatic disease (7.1 vs. 36.7%, $p=0.041$).

Comparisons between survivors vs. non-survivors

Of the 44 patients, 32 survived (72.7%) and the rest were died at 90 days ($n=12$, 27.3%). Patients baseline characteristics were similar between these two groups (**Table 2**). Only BMI was different, being more obese the group of patients that died ($BMI \geq 30$: 34.4 vs. 75%, $p=0.016$). Comorbidities as well as the preoperative qSOFA score were similar in both groups. Non-survivors had higher rates of pneumonia (59.4 vs. 91.7%, $p=0.041$). Regarding the preoperative laboratory studies, non-survivors had higher CRP, leucocyte count and ferritin, and lower albumin. The rate of postoperative complications was similar between both groups, however patients in the non-survivors group developed more ARDS, septic shock, and multi-organic failure. Also, non-survivors required more ICU admissions, invasive mechanical ventilation, and vasopressors.

Comparisons between patients with postoperative complications and patients without postoperative complications

A total of 23 (52.3%) patients developed at least one postoperative complication (**Table 2 and 4**). Postoperative complications were classified according to Clavien-Dindo and are presented in **Table 5**.

The majority of baseline and demographic data were similar between complicated and uncomplicated patients. Patients that developed postoperative complications had higher ASA classification, elevated preoperative CRP, and higher rate of pneumonia. Also, complicated patients were associated with higher incidence of ARDS and septic shock, requiring more mechanical ventilation, ICU admission, and vasopressors. The hospital LOS was shorter in patients without postoperative complications (21.8 vs. 38.8 days, $p=0.043$).

Predictors of mortality and postoperative complications

The overall mortality rate was 27.3% (12/44). The 90-day survival probability after surgery of the entire cohort was 70.1% (95% CI 60.3-79.9) (**Table 3 and 4**). The 90-day survival probability was significantly lower in patients with SARS-CoV-2 symptoms (63.4 [95% CI: 50.5-76.2]) in comparison with patients without symptoms (84.5 [95% CI: 74.3-94.3]) (**Figure 1**).

The 90-day survival probability was similar between patients with (65.3 [95% CI: 51.2-79.5]) and without (75.3 [95% CI: 62.2-88.4]) postoperative complications.

The univariate analysis revealed that higher BMI (HR=4.2 [1.14-15.66]), developing ARDS (HR=8.4 [1.08-65.17]), need of vasopressors (HR=10.4 [1.34-80.65]) and invasive ventilation (HR=8.41 [1.08-65.17]), as well as higher values of ferritin (HR=1.003 [1.002-1.005]), CRP (HR=1.05 [1.01-1.10]) and leukocytes (HR=1.097 [1.003-1.198]) were risk factors for decreased 90-day survival. Multivariate analysis identified high preoperative ferritin values as independent risk factor (HR= 1.003 (1.002-1.005); $p<0.001$) for decreased 90-day survival as shown in **Table 6**.

To assess the discriminatory capacity of each risk factor, ROC curves were constructed for survival status and postoperative complications at 90-days. The area under the ROC curve (AUC), sensitivity, specificity, and LR were calculated for mortality and postoperative complications, as displayed in **Table 7**.

Regarding the accuracy of prediction, we found the following cut-off values associated with decreased 90-day survival: ferritin ≥ 438.5 ng/mL (AUC= 0.908, $p<0.001$), CRP value ≥ 12.5 mg/dL (AUC=0.715, $p=0.030$), leukocyte count $\geq 13.8 \times 10^3/\mu\text{L}$ (AUC=0.706, $p=0.037$), and albumin ≤ 2.78 g/dL (AUC=0.704, $p=0.039$) (**Figure 2**).

Regarding the probability for postoperative complications, univariate and multivariate analysis revealed that higher levels of preoperative CRP (OR= 0.022 [0.013-0.032], $p<0.001$) was associated with increased

risk of postoperative complications. We found that a cut-off value of CRP of ≥ 12.5 mg/dL yielded an accuracy of 82.9% for the prediction of postoperative complications in COVID-19 patients ($p < 0.001$) (Figure 3).

Discussion

We found in the present study that symptomatic COVID-19 patients that needed an emergency general surgery procedure had worse postoperative outcomes, increased postoperative complications, higher ICU admissions, prolonged length of stay, and decreased 90-day survival probability as compared with asymptomatic COVID-19 patients. We also found that preoperative ferritin value of ≥ 438.5 ng/mL, CRP value of ≥ 12.5 mg/dL, total leukocyte count of $\geq 13.8 \times 10^3/\mu\text{L}$ and albumin of ≤ 2.78 g/dL, could be used as predictor factors for decreased 90-day survival with acceptable accuracy. Also, a preoperative CRP value of ≥ 12.5 mg/dL predicted the development of postoperative complications.

To date, there is limited data that describes outcomes of COVID-19 patients that underwent surgery. The firsts published studies regarding the impact of COVID-19 on surgical outcomes analyzed elective and emergency surgeries together and included several specialties together [4,8,7,16]. A series of 34 patients from China, at the beginning of the pandemic, with occult COVID-19 infections at the time of surgery reported the development of pneumonia in all patients, ARDS in 32%, 44.1% were admitted to the ICU and a 20.5% postoperative mortality rate [4]. However, this study included several types of surgeries including neurosurgery, orthopedics, ophthalmology and general surgery. In the cohort of Dogietto et al. [17], they described factors associated with surgical mortality and complications among patients with and without COVID-19 in Italy. They concluded that the surgical mortality and complications were higher in patients with COVID-19 (19.5%) compared with patients without COVID-19, where pulmonary (14.6%) and thrombotic complications (9.7%) are significantly associated with it. The meta-analysis conducted by Abate et al. [7] revealed a very high global rate of postoperative mortality among COVID-19 patients of 20% (95% CI: 15-26) and a postoperative ICU admission rate of 15% (95% CI: 10-21). However, this meta-analysis included patients of several specialties (mostly orthopedics), and included elective and emergency procedures.

There are some published studies regarding outcomes after emergency or acute surgery in COVID-19 patients. Seeliger et al. [8], described 13 patients with acute surgical condition and concomitant COVID-19. They concluded that co-infection did not result in more complications for emergency abdominal surgery, although an acute abdomen with severe COVID-19 had dismal prognosis. In a multi-center cohort study conducted by Carrier et al. [18], they analyzed 44 patients with COVID-19, with 31 surgeries (71%) being urgent. They described that the postoperative 30-day mortality in COVID-19 patients undergoing surgery was high (15.9%). This mortality was higher in patients with symptoms

(23.1%) compared to those without symptoms (5.6%), although not statistically significant. Fallani et al. [19], reported a higher rate of postoperative complications during the COVID-19 pandemic in patients with peritonitis. They found that the ASA score, severity of peritonitis, qSOFA score, diagnosis other than appendicitis, and COVID period resulted independent predictors of complications.

One strength of the present study is that we limited the procedures to general surgeries (mostly abdominal emergencies). Also, all our patients had confirmed COVID-19. We noted that patients with COVID-19 were at a substantially higher risk of postoperative morbidity and mortality. In our study, 52.3% of patients experienced at least one complication, the postoperative mortality rate was 27.3%, the mean hospital stay was 30.7 days, and 61.4% required ICU admission. It is important to mention that patients in our cohort have high rates of comorbidities (59.1%), almost the half of patients were classified as ASA III (43.2%), and the majority of patients (63.6%) have a high-risk preoperative qSOFA score. This could contribute to the high rate of surgical complications, need of ICU, prolonged LOS, and decreased survival rate.

The data analyzed in the present study suggests that patients with COVID-19 who needed emergent gastrointestinal or soft tissue surgery have an increased risk of postoperative morbidity and mortality. These findings have several important implications. First, it is necessary to rule out COVID-19 in all patients that are admitted with an acute surgical disorder, in order to inform the patients or relatives about the potential impact of COVID-19 on surgical complications, length of stay, need of ICU, and mortality. Secondly, we consider that laparoscopy can be safely performed in patients with diseases commonly treated by laparoscopy (like appendicitis and cholecystitis), with all the isolation and protective measures needed to avoid unnecessary infections to the health personnel. Thirdly, we found a high prevalence of comorbidities, obesity, and smoking, that could impact on the severity of COVID-19, thus affecting postoperative outcomes.

We would like to add that there is no scientific evidence concerning the risk of transmission of COVID-19 by laparoscopic surgery [20-23]. A recent systematic review of 22 articles [20] concluded that laparoscopic surgery can be used with the recommended precautions because of its benefits compared to open surgery. In our study, we performed 19 laparoscopic procedures. All these laparoscopic procedures were performed in a special theater and within an area dedicated to only SARS-CoV-2 positive patients. We employed a special filter for gas evacuation and all staff had personal protective equipment. We found that after following protocols recommended by scientific societies and our internal protocols [20-23], no staff members (surgeons, anesthesiologist and nurses) were infected with COVID-19 after laparoscopic surgeries.

There are some limitations to our study that need to be mentioned. The most important limitation is that this is a single-center observational study, which predisposes to all the biases inherent to the design. Another important limitation is the inherent problems with the criteria for diagnosing COVID-19. For example, CT-scan findings depend on the criteria of the radiologists [10]. Other limitation was that although a comprehensive review of all medical records was performed, we cannot exclude the possibility that some clinical parameters, including symptoms and comorbidities, were not correctly registered. However, despite these limitations we consider that the results of this study could help us to better understand the implications of SARS-CoV-2 in patients that needs emergency gastrointestinal and soft tissue surgery.

Conclusion

In conclusion, we found in our study that positive SARS-CoV-2 patients that needed an emergency general surgery procedure with concomitant COVID-19 related symptoms had worse postoperative outcomes, increased postoperative complications, higher ICU admissions, prolonged length of hospital stay, and decreased 90-day survival rate as compared with asymptomatic COVID-19 patients. Also, preoperative serum ferritin, C-reactive protein, leukocytes and albumin values could be used as accurate predictors of decreased 90-day survival.

Declarations

Availability of data and material

The data that support the findings of this study are available on request from the corresponding author.

Authors' Contributions

KBG and MTA: contributed equally to study conception and design, acquired data, performed the data analyses and wrote the manuscript, critical revision of the manuscript, and approved the final version.

CVS, JHE, MMP: study conception and design, acquisition of data, critical revision of the manuscript, and approved the final version.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declarations

Conflict of Interest The authors declare that they have no conflicts of interest.

Ethical approval: The study protocol was approved by the ethics and research committee of the hospital

Ethical statement: the study was conducted in accordance with the principles of the Declaration of Helsinki.

Consent to participate: Informed consent was waived according to the regulations of our institution for observational studies.

Acknowledgments

We would like to thank Adolfo Cuendis-Velázquez, Elisafat Arce-Liévano, Jessica Betancourt-Ferreya, Miguel Blas-Franco, Roberto Delano-Alonso, Ignacio Del Rio-Suárez, Eduardo Cárdenas-Lailson, and Sujey Romero-Loera for sharing their data about their surgical cases. Special thanks to Dr. Pablo Maravilla for his helpful advise. We also would like to thank all medical and nurse staff dedicated to COVID-19 surgical care.

References

1. World Health Organisation (WHO) Coronavirus disease (COVID-19) outbreak webpage. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
2. Antonio-Villa NE, Fernandez-Chirino L, Pisanty-Alatorre J, Mancilla-Galindo J, Kammar-García A, Vargas-Vázquez A, González-Díaz A, Fermín-Martínez CA, Márquez-Salinas A, Guerra EC, Bahena-López JP, Villanueva-Reza M, Márquez-Sánchez J, Jaramillo-Molina ME, Gutiérrez-Robledo LM, Bello-Chavolla OY (2021) Comprehensive evaluation of the impact of sociodemographic inequalities on adverse outcomes and excess mortality during the COVID-19 pandemic in Mexico City. Clin Infect Dis. [https://doi: 10.1093/cid/ciab577](https://doi.org/10.1093/cid/ciab577)
3. Vranis NM, Bekisz JM, Daar DA, Chiu ES, Wilson SC (2021) Clinical Outcomes of 2019 COVID-19 Positive Patients Who Underwent Surgery: A New York City Experience. J Surg Res 261:113-122. [https://doi: 10.1016/j.jss.2020.10.032](https://doi.org/10.1016/j.jss.2020.10.032).
4. Lei S, Jiang F, Su W, Chen C, Chen J, Mei W, Zhan LY, Jia Y, Zhang L, Liu D, Xia ZY, Xia Z (2020) Clinical characteristics and outcomes of patients undergoing surgeries during the incubation period of COVID-19

infection. *EClinicalMedicine* 21:100331. [https://doi: 10.1016/j.eclinm.2020.100331](https://doi.org/10.1016/j.eclinm.2020.100331)

5. Aminian A, Safari S, Razeghian-Jahromi A, Ghorbani M, Delaney CP (2020) COVID-19 outbreak and surgical practice: Unexpected fatality in perioperative period. *Ann Surg* 272:e27ee29. [https://doi:10.1097/SLA.0000000000003925](https://doi.org/10.1097/SLA.0000000000003925).
6. COVIDSurg Collaborative (2020) Mortality and pulmonary complications in patients undergoing surgery with perioperative SARS-CoV-2 infection: an international cohort study. *Lancet* 396(10243):27–38. [https://doi: 10.1016/S0140-6736\(20\)31182-X](https://doi.org/10.1016/S0140-6736(20)31182-X)
7. Abate S, Mantefardo B, Basu B (2020) Postoperative mortality among surgical patients with COVID-19: a systematic review and meta-analysis. *Patient Safety in Surgery* 14:37. <https://doi.org/10.1186/s13037-020-00262-6>
8. Seeliger B, Philouze G, Cherkaoui Z, Felli E, Mutter D, Pessaux P (2020) Acute abdomen in patients with SARS-CoV-2 infection or co-infection. *Langenbecks Arch Surg* 405(6):861-866. [https://doi: 10.1007/s00423-020-01948-2](https://doi.org/10.1007/s00423-020-01948-2)
9. COVIDSurg Collaborative; GlobalSurg Collaborative (2021) Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. *Anaesthesia* 76(6):748-758. [https://doi: 10.1111/anae.15458](https://doi.org/10.1111/anae.15458)
10. Prokop M, van Everdingen W, van Rees Vellinga T, Quarles van Ufford H, Stöger L, Beenen L, Geurts B, Gietema H, Krdzalic J, Schaefer-Prokop C, van Ginneken B, Brink M, COVID-19 Standardized Reporting Working Group of the Dutch Radiological Society (2020) CO-RADS: A Categorical CT Assessment Scheme for Patients Suspected of Having COVID-19-Definition and Evaluation. *Radiology* 296(2):E97-E104. [https://doi: 10.1148/radiol.2020201473](https://doi.org/10.1148/radiol.2020201473).
11. Gao Y, Xi H, Chen L (2020) Emergency Surgery in Suspected COVID-19 Patients With Acute Abdomen: Case Series and Perspectives. *Ann Surg* 272(1):e38-e39. [https://doi: 10.1097/SLA.0000000000003961](https://doi.org/10.1097/SLA.0000000000003961)
12. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, Hotchkiss RS, Levy MM, Marshall JC, Martin GS, Opal SM, Rubenfeld GD, van der Poll T, Vincent JL, Angus DC (2016) The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 315(8):801-10. [https://doi: 10.1001/jama.2016.0287](https://doi.org/10.1001/jama.2016.0287).
13. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240:205–213. [https://doi: 10.1097/01.sla.0000133083.54934.ae](https://doi.org/10.1097/01.sla.0000133083.54934.ae).
14. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW; the Northwell COVID-19 Research Consortium, Barnaby DP, Becker LB, Chelico JD, Cohen SL, Cookingham J, Coppa K, Diefenbach MA, Dominello AJ, Duer-Hefe J, Falzon L, Gitlin J, Hajizadeh N, Harvin TG, Hirschwerk DA,

- Kim EJ, Kozel ZM, Marrast LM, Mogavero JN, Osorio GA, Qiu M, Zanos TP (2020) Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* 323(20):2052-2059. [https://doi: 10.1001/jama.2020.6775](https://doi.org/10.1001/jama.2020.6775)
15. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, Du B, Li LJ, Zeng G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS; China Medical Treatment Expert Group for Covid-19 (2020) Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 382(18):1708-1720. [https://doi: 10.1056/NEJMoa2002032](https://doi.org/10.1056/NEJMoa2002032)
16. Pai E, Chopra S, Mandloi D, Upadhyay AK, Prem A, Pandey D (2020) Continuing surgical care in cancer patients during the nationwide lockdown in the COVID-19 pandemic—perioperative outcomes from a tertiary care cancer center in India. *J Surg Oncol* 122(5):1–6. [https://doi: 10.1002/jso.26134](https://doi.org/10.1002/jso.26134)
17. Doglietto F, Vezzoli M, Gheza F, Lussardi GL, Domenicucci M, Vecchiarelli L, Zanin L, Saraceno G, Signorini L, Panciani PP, Castelli F, Maroldi R, Rasulo FA, Benvenuti MR, Portolani N, Bonardelli S, Milano G, Casiraghi A, Calza S, Fontanella MM (2020) Factors associated with surgical mortality and complications among patients with and without coronavirus Disease 2019 (COVID-19) in Italy. *JAMA Surg* 155(8):1–14. [https://doi: 10.1001/jamasurg.2020.2713](https://doi.org/10.1001/jamasurg.2020.2713)
18. Carrier F, Amzallag E, Lecluyse V (2021) Postoperative outcomes in surgical COVID- 19 patients: a multicenter cohort study. *BMC Anesthesiology* 21:15. <https://doi.org/10.1186/s12871-021-01233-9>.
19. Fallani G, Lombardi R, Masetti M, Chisari M, Zanini N, Cattaneo GM, Filosa M, Zanzi F, Guerra E, Bonilauri S, Di Donato L, Garulli G, Lucchi A, Grassia M, Ugolini G, Pasini F, Vetrone G, Benini C, Nicosia S, Jovine E (2021) Urgent and emergency surgery for secondary peritonitis during the COVID-19 outbreak: an unseen burden of a healthcare crisis. *Updates Surg* 73(2):753-762. [https://doi: 10.1007/s13304-020-00943-y](https://doi.org/10.1007/s13304-020-00943-y)
20. El Boghdady M, Ewalds-Kvist BM (2021) Laparoscopic Surgery and the debate on its safety during COVID-19 pandemic: A systematic review of recommendations. *Surgeon* 19(2):e29-e39. [https://doi: 10.1016/j.surge.2020.07.005](https://doi.org/10.1016/j.surge.2020.07.005)
21. Di Saverio S, Khan M, Pata F, Ietto G, De Simone B, Zani E, Carcano G (2020) Laparoscopy at all costs? Not now during COVID-19 outbreak and not for acute care surgery and emergency colorectal surgery: A practical algorithm from a hub tertiary teaching hospital in Northern Lombardy, Italy. *J Trauma Acute Care Surg* 88(6):715-718. [https://doi: 10.1097/TA.0000000000002727](https://doi.org/10.1097/TA.0000000000002727)
22. Francis N, Dort J, Cho E, Feldman L, Keller D, Lim R, Mikami D, Phillips E, Spaniolas K, Tsuda S, Wasco K, Arulampalam T, Sheraz M, Morales S, Pietrabissa A, Asbun H, Pryor A (2020) SAGES and EAES recommendations for minimally invasive surgery during COVID-19 pandemic. *Surg Endosc* 34(6):2327-2331. [https://doi: 10.1007/s00464-020-07565-w](https://doi.org/10.1007/s00464-020-07565-w)

23. Schwarz L, Tuech JJ (2020) Is the use of laparoscopy in a COVID-19 epidemic free of risk? Br J Surg 107(7):e188. [https://doi: 10.1002/bjs.11649](https://doi.org/10.1002/bjs.11649)

Tables

Table 1. Characteristics of positive SARS-CoV-2 patients with and without symptomatic disease, that underwent emergency general surgery				
	All patients (N= 44)	Asymptomatic patients (n= 14)	Symptomatic patients (n= 30)	P value
Sex, n (%)				0.659
Female	21 (47.7)	6 (42.9)	15 (50)	
Male	23 (52.3)	8 (57.1)	15 (50)	
Age, mean (SD)	53.5 (17)	42.2 (18.8)	58.7 (2.4)	0.002
Age, n (%)				0.038
<59	28 (63.6)	12 (85.7)	16 (53.3)	
≥60	16 (36.4)	2 (14.3)	14 (46.7)	
BMI, mean (SD)	28.8 (6.8)	26.2 (7.08)	30 (6.48)	0.093
BMI, n (%)				0.029
<29.9	24 (54.5)	11 (78.6)	13 (43.3)	
≥30	20 (45.5)	3 (21.4)	17 (56.7)	
Smoking, n (%)				0.008
Yes	19 (43.2)	2 (14.3)	17 (56.7)	
No	25 (56.8)	12 (85.7)	13 (43.3)	
ASA classification				<0.001
I	5(11.4)	5 (35.7)	0	
II	20 (45.5)	8 (57.1)	12 (40)	
III	19 (43.2)	1 (7.1)	18 (60)	
Comorbidities, n (%)				0.005
Yes	26 (59.1)	4 (28.6)	22 (73.3)	
Comorbidities, n (%)				
Diabetes	16 (36.4)	2 (14.3)	14 (46.7)	0.038
Hypertension	25 (56.8)	4 (28.6)	21 (70)	0.01
COPD	3 (6.8)	0	3 (10)	0.220
CRD	9 (20.5)	2 (14.3)	9 (20.5)	0.488
Preoperative qSOFA score				0.009

Not High Risk (0-1)	16 (36.4)	9 (64.3)	7 (23.3)	
High Risk (>2)	28 (63.6)	5 (35.7)	23 (76.7)	
Index surgery, n (%)				
Appendectomy	11 (25)	6 (42.8)	5 (16.7)	0.061
Cholecystectomy	10 (22.7)	4 (28.6)	6 (20)	0.527
Emergency Laparotomy	13 (29.5)	3 (21.4)	10 (33.3)	0.420
PDCI	5 (11.4)	0	5 (16.7)	0.160
Soft tissue debridement	5 (11.4)	1 (7.1)	4 (13.3)	0.546
Modality of abdominal surgery, n (%)				
Laparoscopic	16 (41)	8 (61.5)	8 (30.8)	0.065
Open	20 (51.3)	5 (38.5)	15 (57.7)	0.320
Converted to open	3 (7.7)	0	3 (11.5)	0.537
BMI:body mass index; ASA: American Society of Anesthesiology; SOFA: : PDCI: peritoneal dialysis catheter insertion; CRD: chronic renal disease; COPD: chronic obstructive pulmonary disease.				

Table 2. Characteristics of positive SARS-CoV-2 patients that underwent emergency general surgery. Two comparisons based on 90-day mortality status and presence or absence of postoperative complications.

	90-Day Survivors (n= 32)	90-Day non survivors (n= 12)	P value	No Postoperative complications (n= 21)	Postoperative complications (n= 23)	P value
Sex, n (%)			0.853			0.989
Female	15 (46.9)	6 (50)		10 (47.6)	11 (47.8)	
Male	17 (53.1)	6 (50)		11 (52.4)	12 (52.2)	
Age, mean (SD)	52.1 (18.2)	56.5 (13.5)	0.121	49.6 (17.9)	57.1 (15.7)	0.171
Age, n (%)			0.798			0.098
<59	20 (62.5)	8 (66.7)		16 (76.2)	12 (52.2)	
≥60	12 (37.5)	4 (33.3)		5 (23.8)	11 (47.8)	
BMI, mean (SD)	28.0 (6.6)	30.9 (7.1)	0.216	28.4 (7.2)	29.1 (6.5)	0.755
BMI, n (%)			0.016			0.349
<29.9	21 (65.6)	3 (25)		13 (61.9)	11 (47.8)	
≥30	11 (34.4)	9 (75)		8 (38.1)	12 (52.2)	
Smoking, n (%)			0.214			0.062
Yes	12 (37.5)	7 (58.3)		6 (28.6)	13 (56.5)	
ASA classification			0.105			0.023
I	5 (15.6)	0		5 (23.8)	0	
II	16 (50)	4 (33.3)		10 (47.6)	10 (43.5)	
III	11 (34.4)	8 (66.7)		6 (28.6)	13 (56.5)	
Comorbidities, n (%)			0.189			0.802
Yes	17 (53.1)	9 (75)		12 (57.1)	14 (60.9)	
Comorbidities, n (%)						
Diabetes	11 (34.4)	5 (41.7)	0.654	7 (33.3)	9 (39.1)	0.690
Hypertension	17 (53.1)	8 (66.7)	0.419	11 (52.4)	14 (60.9)	0.570
COPD	2 (6.3)	1 (8.3)	0.807	0	3 (13)	0.086

CRD	6 (18.8)	3 (25)	0.647	6 (28.6)	3 (13)	0.202
Preoperative qSOFA score						0.690
Not High Risk (0-1)	13 (40.6)	3 (25)	0.337	7 (33.3)	9 (39.1)	
High Risk (>2)	19 (59.4)	9 (75)		14 (66.7)	14 (60.9)	
Index surgery, n (%)						
Appendectomy	9 (28.1)	2 (16.7)	0.434	6 (28.5)	5 (21.7)	0.601
Cholecystectomy	9 (28.1)	1 (8.3)	0.162	4 (19)	6 (26.1)	0.577
Emergency Laparotomy	8 (25)	5 (41.7)	0.080	4 (19)	9 (39.1)	0.144
PDCI	2 (6.3)	3 (25)	0.080	4 (19)	1 (4.3)	0.124
Soft tissue debridement	4 (12.5)	1 (8.3)	0.698	3 (14.3)	2 (8.7)	0.559
Modality of abdominal surgery, n (%)						
Laparoscopic	13 (46.4)	3 (27.3)	0.273	9 (50)	7 (33.3)	0.291
Open	13 (46.4)	7 (63.6)	0.333	9 (50)	11 (52.4)	1.000
Converted	2 (7.2)	1 (9)	0.837	0	3 (14.3)	0.234
BMI:body mass index; ASA: American Society of Anesthesiology; SOFA: : PDCI: peritoneal dialysis catheter insertion; CRD: chronic renal disease; COPD: chronic obstructive pulmonary disease.						

Table 3. Clinical evolution of COVID-19 patients with and without symptoms that underwent emergency general surgery

	All patients (n= 44)	Asymptomatic patients (n= 14)	Symptomatic Patients (n= 30)	P value
Preoperative studies, mean (SD)				
Hemoglobin	12.0 (2.9)	14.1 (2.5)	11.0 (2.5)	0.001
CRP	19.3 (13.5)	10.5 (9.1)	23.4 (13.2)	0.002
Platelets	272 (121.6)	286.8 (94.6)	265.0 (133.3)	0.586
Leucocytes	13.1 (5.7)	12.2 (5.3)	13.6 (6.0)	0.468
Lymphocytes	1.38 (1.1)	1.72 (1.0)	1.22 (1.0)	0.158
Neutrophils	13.5 (17.3)	8.8 (4.6)	15.6 (20.5)	0.232
Monocytes	1.73 (2.7)	2.82 (3.4)	1.22 (2.2)	0.069
LDH	251.8 (131.5)	192.7 (56.0)	279.4 (147.6)	0.040
Ferritin	708.3 (389.3)	509.9 (186.9)	800.9 (425.7)	0.019
Albumin	3.0 (0.9)	3.84 (0.7)	2.61 (0.6)	<0.001
Postoperative complications, n (%)				0.001
Yes	23 (52.3)	2 (14.3)	21 (70)	
Organ Failures, n (%)				<0.001
ARDS	27 (61.4)	1 (7.1)	26 (86.7)	0.092
ARF	10 (22.7)	1 (7.1)	9 (30)	0.001
Septic shock	15 (34.1)	0	15 (50)	0.026
MOF	13 (29.5)	1 (7.1)	12 (40)	
Mechanical ventilation, n (%)				<0.001
Not required	17 (38.6)	13 (92.9)	4 (13.3)	
Preoperative	16 (36.4)	0	16 (53.3)	
Postoperative	11 (25.0)	1 (7.1)	10 (33.3)	
ICU Admission, n (%)				<0.001
No	17 (38.6)	13 (92.9)	4 (13.3)	
Preoperative	15 (34.1)	0	15 (50)	

Postoperative	12 (27.3)	1 (7.1)	11 (36.7)	
ICU treatment, n (%)				<0.001
Invasive ventilation	27 (61.4)	1 (7.1)	26 (86.7)	<0.001
Vasopressors	25 (56.8)	1 (7.1)	24 (80)	
Hospital LOS, mean (SD)	30.7 (28.1)	9.6 (14.8)	40.6 (27.4)	<0.001
Mortality				
90-day mortality, n (%)	12 (27.3)	1 (7.1)	11 (36.7)	0.041
90-day survival probability (Kaplan-Meier) (95% CI)	70.1 (60.3-79.9)	84.5 (74.3-94.3)	63.4 (50.5-76.2)	<0.05
<p>SD: standard deviation; CRP: C-Reactive Protein; LDH: lactate deshydrogense; ARDS: Acute Respiratory Distress Syndrome , ARF: acute renal failure, MOF: multi organic failure; LOS: length of stay;</p> <p>GPS: Glasgow Prognostic Score; PI: Prognostic Index; NLR: neutrophil to lymphocyte ratio; PLR: platelet to lymphocyte ratio; LMR: Lymphocyte to monocyte ratio; PNI: prognostic nutritional index.</p>				

Table 4. Clinical evolution of COVID-19 patients that underwent emergency general surgery. Comparisons between survivors and non-survivors and complicated and non-complicated patients.

	90-Day Survivors (n= 32)	90-Day non survivors (n= 12)	P value	No Postoperative complications (n= 21)	Postoperative complications (n= 23)	P value
Preoperative studies, mean (SD)						
Hemoglobin	12.1 (3.1)	11.7 (2.5)	0.682	12.6 (3.3)	11.4 (2.4)	0.171
CRP			0.013	11.1 (9.1)	26.8 (12.4)	<0.001
Platelets	16.2 (11.4)	27.4 (15.4)	0.728	279.4 (101.5)	265.2 (139.5)	0.704
Leucocytes	275.9 (122.8)	261.4 (123.0)	0.045	11.4 (4.3)	14.7 (6.5)	0.063
Lymphocytes			0.807	1.48 (1.0)	1.29 (1.1)	0.556
Neutrophils	12.1 (5.2)	16 (6.3)	0.952	9.4 (4.6)	17.2 (23.2)	0.142
Monocytes	1.35 (0.9)	1.45 (1.3)	0.626	2.29 (3.4)	1.22 (1.7)	0.197
LDH		13.7 (7.0)	0.745	222.6 (120.5)	278.5 (138.1)	0.161
Ferritin	13.4 (20.0)		<0.001	614.8 (344.7)	278.5 (138.1)	0.129
Albumin	1.85 (2.7)	1.40 (2.8)	0.050	3.24 (1.0)	2.78 (0.7)	0.087
	247.8 (130.9)	262.5 (138.4)				
	536.0 (210.3)	1167.9 (388.8)				
	3.16 (0.9)	2.56 (0.7)				
Postoperative complications, n (%)			0.242	-	-	
Yes	15 (46.9)	8 (66.7)				
Symptomatic COVID-19			0.041	9 (30)	21 (70)	0.001
Yes	19 (59.4)	11 (91.7)				
Organ Failures, n (%)			0.011			
ARDS	16 (50)	11 (91.7)	0.066	7 (33.3)	20 (87)	<0.001
	5 (15.6)	5 (41.7)	0.038	4 (19)	6 (26.1)	0.578

ARF	8 (25)	7 (58.3)	<0.001	2 (9.5)	13 (56.5)	0.001
Septic shock	1 (3.1)	12 (100)		4 (19)	9 (39.1)	0.145
MOF						
Mechanical ventilation, n (%)			0.037			<0.001
Not required						
Preoperative	16 (50)	1 (8.3)		14 (66.7)	3 (13)	
Postoperative	9 (28.1)	7 (58.3)		6 (28.6)	10 (43.5)	
	7 (21.9)	4 (33.3)		1 (4.8)	10 (43.5)	
ICU Admission, n (%)			0.03			<0.001
No	16 (50)	1 (8.3)		14 (66.7)	3 (13)	
Preoperative	8 (25)	7 (58.3)		6 (28.6)	9 (39.1)	
Postoperative	8 (25)	4 (33.3)		1 (4.8)	11 (47.8)	
ICU treatment, n (%)						
Invasive ventilation	16 (50)	11 (91.7)	0.011	7 (33.3)	20 (87)	<0.001
Vasopressors	14 (43.8)	11 (91.7)	0.004	7 (33.3)	18 (78.3)	0.003
Hospital LOS, mean (SD)	26.7 (28.3)	41.5 (25.2)	0.121	21.8 (26.9)	38.8 (27.1)	0.043
Mortality	-	-	-			
90-day mortality, n (%)				4 (19)	8 (34.8)	0.242
90-day survival probability (Kaplan-Meier) (95% CI)				75.3 (62.2-88.4)	65.3 (51.2-79.5)	0.278
SD: standard deviation; CRP: C-Reactive Protein; LDH: lactate deshydrogense; ARDS: Acute Respiratory Distress Syndrome , ARF: acute renal failure, MOF: multi organic failure; LOS: length of stay;						
GPS: Glasgow Prognostic Score; PI: Prognostic Index; NLR: neutrophil to lymphocyte ratio; PLR: platelet to lymphocyte ratio; LMR: Lymphocyte to monocyte ratio; PNI: prognostic nutritional index.						

Table 5. Description of postoperative complications according with the Clavien-Dindo classification (N=44)	
Clavien-Dindo Classification	N (%)
Grade I, n (%)	20 (45.4)
Grade II, n (%)	
Blood transfusion	16 (36.3)
Total parenteral nutrition	17 (38.6)
Postoperative Ileus	12 (27.2)
Pneumonia	21 (47.7)
Delirium	11 (25)
Superficial surgical infection	12 (27.2)
Deep surgical infection	5 (11.3)
Post-chole biliary leak	1 (2.2)
Pancreatic fistula	1 (2.2)
Grade IIIa, n (%)	
Percutaneous peripancreatic collection drainage	1 (2.2)
Percutaneous drainage of intra-abdominal collection	2 (4.5)
Grade IIIb, n (%)	
Evisceration	1 (2.2)
Bleeding	2 (4.5)
Intestinal occlusion	1 (2.2)
Pancreatic necrosis	2 (4.5)
Negative pressure therapy	1 (2.2)
Appendicular stump leak	1 (2.2)
Anastomotic leak	1 (2.2)
Need of ERCP	4 (9.0)
Grade IVa, n (%)	20 (45.5)
Single organ dysfunction	
Grade IVb, n (%)	13 (29.5)
Multiorgan dysfunction	

Grade V, n (%)	
Mortality	12 (27.3)

Table 6. Logistic regression analysis of factors associated with postoperative mortality and development of postoperative complications in patients with COVID-19				
	Univariate analysis		Multivariate analysis	
Factors	Hazard Ratio (95% CI)	P value	Hazard Ratio (95% CI)	P value
Univariate and multivariate analysis for factors associated with 90-day survival (Cox regression)				
BMI, n (%)		0.030		
<29.9	Ref.			
≥30	4.23 (1.14-15.66)			
COVID-19 Pneumonia	6.03 (0.78-46.77)	0.085		
ARDS	8.41 (1.08-65.17)	0.042		
ARF	2.94 (0.93-9.34)	0.066		
Septic shock	2.87 (0.90-9.06)	0.072		
Invasive ventilation	8.41 (1.08-65.17)	0.042		
Vasopressors	10.40 (1.34-80.65)	0.025		
Cox regression analysis for laboratory parameters associated with 90-day survival				
Factors	Hazard Ratio (95% CI)	P value	Hazard Ratio (95% CI)	P value
Ferritin	1.003 (1.002-1.005)	<0.001	1.003 (1.002-1.005)	<0.001
CRP	1.05 (1.01-1.10)	0.016		
Leukocytes	1.097 (1.003-1.198)	0.042		
Albumin	0.511 (0.246-1.062)	0.072		
Linear regression analysis for laboratory parameters associated with postoperative complications				
Factors	Odds Ratio (95% CI)	P value	Odds Ratio (95% CI)	P value
CRP	0.022 (0.013-0.032)	<0.001	0.022 (0.013-0.032)	<0.001
Leukocytes	0.025 (-0.001-0.050)	0.063		
Albumin	-0.145 (-0.312-0.022)	0.087		
BMI: body mass index; ARDS: acute respiratory distress syndrome; ARF: acute renal failure; CRP: C-Reactive protein. For uni and multivariable logistic regression analysis odds ratio (OR) and 95% confidence interval (CI) are presented. Odds ratios of factors associated to 90-days survival. Only				

significant results are shown in the multivariate analysis. The reference category has an odds ratio of 1.00.

Table 7. ROC analysis for prediction of postoperative mortality and postoperative complications in COVID-19 patients that underwent emergency surgery

	AUC (95% CI)	Cut-off	Sensitivity (%)	Specificity (%)	LR +	LR -	P value
ROC analysis for prediction of postoperative mortality							
Ferritin	0.908 (0.785-1.000)	438.5	91.7	62.5	1.47	0.22	<0.001
CRP	0.715 (0.522-0.908)	12.5	83.3	62.5	1.40	0.41	0.030
Leukocytes	0.706 (0.528-0.884)	13.8	75	31.3	2.40	0.36	0.037
Albumin	0.704 (0.534-0.875)	2.78	66.7	37.5	1.78	0.53	0.039
ROC analysis for prediction of postoperative complications							
CRP	0.829 (0.702-0.956)	12.5	91.3	38.1	2.40	0.14	<0.001
ROC: Receiver Operator Curve; AUC: area under the curve; LR: likelihood ratios; CRP: C-Reactive Protein							

Figures

Figure 1

The 90-day survival probability after surgery was significantly lower in patients with SARS-CoV-2 symptoms in comparison with patients without symptoms (Log rank test < 0.05).

Figure 2

ROC curves for prediction of mortality: CRP, AUC=0.715; leukocytes, AUC=0.706; and Ferritin, AUC=0.908.

Figure 3

ROC curve for CRP prediction of postoperative complications, AUC=0.829.