A novel insight into nitrogen and auxin signaling in lateral root formation in tea plant [*Camellia sinensis* (L.) O. Kuntze]

Shunkai Hu(Former Corresponding Author)

Nanjing Agricultural University <https://orcid.org/0000-0002-9094-259X>

Mi Zhang

Nanjing Agricultural University

Yiqing Yang

Nanjing Agricultural University

Wei Xuan

Nanjing Agricultural University

Zhongwei Zou

Manitoba

Emmanuel Arkorful

Nanjing Agricultural University

Yi Chen

Nanjing Agricultural University

Qingping Ma

Nanjing Agricultural University

Anburaj Jeyaraj

Nanjing Agricultural University

Xuan Chen

Nanjing Agricultural University

Xinghui Li(New Corresponding Author) (✉ lxh@njau.edu.cn)

Research article

Keywords: Nitrogen; Auxin; Lateral roots; Signaling pathway; Transcriptome; *Camellia sinensis*

Posted Date: January 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.16108/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Plant Biology on May 24th, 2020. See the published version at <https://doi.org/10.1186/s12870-020-02448-7>.

Abstract

Background: Tea plant (*Camellia sinensis*) is one of the most popular non-alcoholic beverage worldwide. Lateral roots (LRs) of tea plant are the main organ used for tea plant to absorb soil moisture and nutrients. Lateral roots formation and development are tightly regulated by the nitrogen and auxin signaling pathway. In order to understand the function of auxin and nitrogen signaling in LRs formation and development, transcriptome analysis was applied to investigate the differentially expressed genes involved in lateral roots of tea plants treated with indole-3-butyric acid (IBA), N-1-naphthylphthalamic acid (NPA), low and high nitrogen concentration.

Results: A total of 296 common differentially expressed genes were mainly identified and annotated to four signaling pathways, such as nitrogen metabolism, plant hormone signal transduction, Glutathione metabolism and transcription factors. RNA-sequencing results revealed that majority of differentially expressed genes play important roles in nitrogen metabolism and hormonal signal transduction. Low nitrogen condition induced the biosynthesis of auxin and accumulation of transcripts, thereby regulating lateral roots formation. Furthermore, metabolism of cytokinin and ethylene biosynthesis were also involved in lateral roots development. Transcription factors like *MYB* genes also contributed to the lateral roots formation of tea plants through secondary cell wall biosynthesis. Reversed phase ultra performance liquid chromatography (RP-UPLC) results showed that the auxin concentration in lateral roots was increased, while the nitrogen level decreased. Thus, tea plant lateral roots formation could be induced by low nitrogen concentration via auxin biosynthesis and accumulation.

Conclusion: This study provides new insights into the mechanisms associated with nitrogen and auxin signaling pathways to regulate LRs formation and arises new clues for the efficient utilization of nitrogen in tea plant at the genetic level.

Background

Plant exploring the soil environments for water and nutrients absorption is highly dependent on the architecture of its root systems[1, 2]. The lateral roots (LRs) have high physiological activities, and allow plants to adapt to various nutrients, temperatures and soil environments[3, 4]. Due to the importance of plant growth and agricultural production, the mechanisms of LRs formation has been reported previously[5].

Nitrogen and auxin are the two major factors affecting plant LRs formation. Nitrogen is a necessary component for the synthesis of biological compounds such as DNA, amino acids, proteins and plant hormones[6]. Plant roots can absorb nitrogen as organic (amino acids and peptides) and inorganic (nitrate and ammonium nitrogen) forms[7]. The plasticity of lateral roots is sensitive to different nitrogen suppliers[8, 9]. It has been reported that over supplying of nitrate can inhibit LRs development, while lower nitrate level enhances the growth of LRs. On the contrary, LRs growth will be inhibited at supra-optimal N supply[10, 11]. A previous study showed that nitrate transporter genes (NRTs) are responsible for the high-

affinity NO_3^- transport system, which can able to induce LR growth[12]. Interestingly, Okamoto et al. (2003) found that the expression levels of *NRT1.1* and *NRT2.1* can be strongly induced by the treatment with high level of NO_3 [13]. According to Remans *et al.* (2006), *NRT1.1* can regulate the growth of primary and LR, *NRT2.1* plays a major role in the absorption of NO_3^- in *Arabidopsis*, and also determines the root architecture by controlling LR formation[14]. The *ANR1* (A NO_3^- -inducible *Arabidopsis* gene) was the first identified gene involved in signaling transduction, which associated with external NO_3 concentration affecting LR development[15]. The ammonia transporter gene *AtAmt1.1* has also been found to play a critical role in restructuring LR architecture under N-starvation[16]. Vidal *et al.* (2010) reported auxin- NO_3^- pathway by identifying a passive feedforward mechanism consisting of auxin receptor AUXIN SIGNALING F-BOX 3 (AFB3) and microRNA 393 (miRNA 393)[12]. Auxin biosynthesis gene *TAR2* (tryptophan aminotransferase related 2) are also reported to reprogram root architecture under low nitrogen conditions[17]. Therefore, nitrogen supplying has a strong correlation with auxin synthesis and metabolism. These findings have provided a clear basis for further exploration of the regulatory relationship between nitrogen metabolism-related genes and plant LR growth. However, most of these studies based this discipline have focused on the model crop *Arabidopsis* only, and rarely included woody plants.

Plant hormonal signals were participated in various stages of plant growth and development, including embryogenesis, seed germination, vegetative growth, fruit ripening and leaf senescence. Then, the exploration of plant hormones and their regulatory roles in growth and development of LR has been reported extensively. For example, abscisic acid (ABA) restrains LR development after emergence of LR primordium from primary root before excitation of LR meristem[18]. Auxins regulate LR formation with the stimulation of several endogenous and environmental signals[19]. Cytokinin also affects primary and LR initiation, organ differentiation depends on the ratio of cytokinin-to-auxin, and high cytokinin-to-auxin ratio inhibits root formation and vice versa[20]. Gibberellins in combination with other hormones (ABA, ethylene, and auxin) can control LR formation[21]. Plant hormonal involvement in LR formation is controlled by complex signal crosstalk. However, the potential molecular mechanisms of hormonal involvement in LR formation remain unclear.

Tea plant [*Camellia sinensis* (L.) O. Kuntze] is one of the most popular non-alcoholic natural beverage worldwide. It is a nitrogen-preferring perennial woody plant having well-developed LR. Nitrogen fertilizer is essential for tea leaf increasing and quality. LR are the main organs used for tea plant to absorb soil nutrients and water. Therefore, the LR development has a direct impact on yield and quality of tea plant. Developed LR can increase the absorption capacity of tea plant for soil nitrogen utilization efficiency. However, the molecular mechanism of nutrient conditions for LR formation in tea plants has not yet been explored. In addition, the metabolic relationship between nitrogen and auxin has not been well studied, and the effects of nitrogen and auxin relationship on LR formation are still not clear. The objectives of this study are to compare differentially expressed genes involved in the formation of LR and to determine the concentration of auxin in tea plant by using RNA sequencing and RP-UPLC, respectively, under the nitrogen-deficient, nitrogen-toxicity, IBA (Indole-3-butyric acid, promoting root

development), NPA (N -1-naphthylphthalamic acid, preventing root development) treatments. The mechanisms underlying LR phenotypic changes and their responses to different nitrogen concentrations in tea plants were investigated. The signaling pathways and regulatory networks of nitrogen and auxin in LR formation were proposed.

Results

Tea plant growth performance in response to diverse N levels and auxin

The tea seedlings showed normal leaves growth under various nitrogen treatments in hydroponic culture. All seedlings showed varied condition of growth after ten weeks, resulting in significant change with N treatment (Fig 1). Primary roots of seedlings were slimmer and longer with lower nitrogen supplement. The seedlings showed greener leaves, shortened length of lateral roots, and significant reduction in numbers of LR as N concentration was increased. The aboveground parts of seedlings treated with NN (no nitrogen), LN (low nitrogen) and HN (high nitrogen) were shorter than the control (Fig. 1), and with the increase of nitrogen concentration, the number and length of lateral roots decreased significantly (Fig.3). These results indicated that low nitrogen concentration would promote tea plants LR formation and development while high nitrogen concentration would inhibit the LR growth. In addition, it was observed that the development of LR of seedlings with hormone treatment was investigated. The results showed that the development of LR of seedlings with HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 8 weeks) were restored in the medium after eight weeks, which significantly increase the number and length of lateral roots (Fig. 2a, Fig. 3), while LR of seedlings with LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 4 weeks) were inhibited and decreased obviously as well as the color turned to brown relative to LN treatment after four weeks (Fig. 2c, Fig. 3). These findings suggested that the exogenous auxin could break the inhibition of the LR growth with HN treatment in tea plant. Therefore, since the auxin accumulation and production were inhibited by HN treatment, the LR growth would be restrained by exogenous auxin (IBA).

Effect of nitrogen and IBA supply on auxin and zeatin concentration in LR

There was a significant reduction of auxin concentration in LR treated with HN concentration (Fig. 4). This suggests that HN could inhibit production of auxin, while low nitrogen could induce production of auxin in LR. It also showed that concentration of auxin in LR decreases when exogenous auxin concentration increases. There was a positive correlation between the zeatin concentration and nitrogen concentration of LR. Therefore, zeatin concentration in LR was also increased when seedlings treated with HN concentration. Also, high exogenous auxin concentration decreased zeatin concentration in LR.

Transcriptomes data analysis

A total of 136.13Gb clean data with a minimum Q30 of 92.83% was obtained (Table 1). The clean reads of each sample were sequence-aligned with the designated reference genome, and the efficiency of the

alignment was over 76.06%. Sample correlation coefficients of data obtained revealed that the throughput and quality of sequencing was high enough for further analysis (Fig. 5).

Function annotation of the tea LR transcriptome

Assembled unigenes were annotated with NR, NT, Swiss-Port, KEGG, COG and GO databases. Annotation to NR database revealed that 12,552 NR annotated unigenes showed high identity to *Vitis vinifera*, which was more similar than other species (Fig. 6). The assembled genes associated with 'metabolic process', which the 'single-organism process' and 'cellular process' accounts the majority of GO classification. In cellular component, a total of 13,249, 13,180, and 9,876 unigenes were annotated in cell part, cell and organelle, respectively. In molecular function, a total of 17,987 unigenes were annotated in catalytic activity and 16,499 unigenes were annotated to binding. In biological process, a total of 22,439, 18,248, and 14,766 unigenes were annotated in metabolic process, cellular process and single-organismal process, respectively. COG functional classification analysis also revealed that most unigenes were annotated to 'amino acid transport and metabolism', 'cytoskeleton' and 'cell motility'.

GO and KEGG analysis of differentially expressed genes

A total of 7,784 DEGs were identified from the LR of seedlings treated by LN and CK, and among them, a total of 3,164 and 4,622 unigenes were up-regulated and down-regulated, respectively. A total of 5,432 unigenes (3,216 up-regulated and 2,219 down-regulated unigenes) were differentially expressed in LR of seedlings cultured with CK and HN. Also, a total of 7,756 DEGs (3,709 up-regulated and 4,047 down-regulated unigenes) were obtained from the LR of seedlings treated with LN and LN + NPA. There were 21,671 DEGs in the LR between the seedlings treated with HN and HN + IBA. Among them, there were 11,338 up-regulated and 10,333 down-regulated unigenes identified (Fig. 7).

Gene ontology cellular component analysis indicated that DEGs derived from the LR treated with LN and CK were almost involved in cell, cell part and organelle. Molecular function analysis revealed that most DEGs identified were involved in catalytic activity, binding and transporter activity. The DEGs identified in biological contributed into metabolic process, cellular process, and single-organism process (Fig. S1).

The DEGs derived from the LR of seedling treated with the HN and CK indicated that most of them were involved in cell part, cell and organelle for GO cellular component. Go molecular function analysis revealed most of DEGs were identified from catalytic activity, binding and transporter activity. GO biological process analysis showed that DEGs contributed into to metabolic process, cellular process and single-organism process (Fig. S2). GO cellular component analysis indicated that DEGs derived from the LR treated with LN and LN + NPA were almost involved in cell, cell part and organelle. Molecular function analysis revealed that most DEGs identified were involved in catalytic activity, binding and transporter activity. The DEGs identified in biological contributed into metabolic process, cellular process, and single-organism process (Fig. S3). The DEGs derived from the LR of seedling treated with the HN and HN + IBA indicated that most of them were involved in cell part, cell and organelle for GO cellular component. Go molecular function analysis revealed most of DEGs were identified from catalytic activity, binding and

transporter activity. GO biological process analysis showed that DEGs contributed into to metabolic process, cellular process and single-organism process (Fig. S4).

Among the DEGs identified from LR of seedlings between LN and CK treatments, 387 DEGs were assigned to 72 KEGG pathways. Of these pathways, DEGs involved in carbohydrate metabolism (76 genes) were the most abundant, followed by translation (65 genes), and 'folding, shorting and degradation' (42 genes) (Fig. S5). Among the DEGs identified from LR of seedlings between HN and CK treatments, 279 DEGs were assigned to 101 KEGG pathways. Of these, DEGs involved in carbohydrate metabolism (62 genes) were the most abundant, followed by translation (39 genes), and 'folding, shorting and degradation' (30 genes) (Fig. S6). In addition, a total of 499 DEGs were assigned to 116 KEGG pathways between the LR of seedlings treated with LN and LN + NPA. Of these, 101, 60, and 42 DEGs were involved in carbohydrate metabolism 'metabolism of other amino acids', and in amino acids metabolism, respectively (Fig. S7). Finally, 1606 DEGs were assigned to 132 KEGG pathways between LR of seedlings treated with HN and HN + IBA. The most abundant of these DEGs were involved in carbohydrate metabolism (274 genes), followed by translation (176 genes), amino acid metabolism (176 genes), lipid metabolism (171 genes), and 171 DEGs in 'folding, sorting and degradation' (Fig. S8).

DEGs related tonitrogen and auxintreatments

Finally, a total of 296 common DEGs were identified from the LR of seedlings in all treatments (Fig. 8a). KEGG enrichment analysis showed that these genes were involved in various metabolic pathways. Most of the DEGs were annotated in nitrogen metabolism, plant hormone signal transduction, Glutathione metabolism, and translation (Fig. 8b). In nitrogen metabolism, DEGs included high affinity nitrate transporter (*NRT*) genes, ammonium transporter (*AMT*) gene (Table 3), and nrt1/ptr family (*NPF*) protein genes (Table S2). In the plant hormone signal transduction, the auxin response factor 2 (*ARF2*) genes (Table 3), *Aux/IAA*, *GH3* (indole-3-acetic acid-amido synthetases) genes (Table S2) were identified in auxin metabolism. Adenylate isopentenyltransferase, cytokinin hydroxylases, and cytokinin dehydrogenase genes were involved in cytokinin metabolism, while ethylene biosynthetic process genes participate in ethylene metabolism (Table 3, Table S2). In transcription factors, the annotated genes encoding P-type R2R3 MYB protein (Table 3), MADS-box, NAC, and WRKY family were identified (Table S2). Glutathione S-transferases (*GSTs*) genes which regulate Glutathione metabolism (Table S2), and F-box/kelch genes which participate in F-box protein metabolism were also obtained from the KEGG analysis (Table 3).

Relative expression of differentially expressed genes

The reliability of RNA-Seq data was validated by qRT-PCR using eleven genes. These eleven genes were included in the regulation of nitrogen and auxin signaling pathways. Of these genes, three are connected with nitrogen metabolism; one is related to tryptophan aminotransferase; one is associated with auxin response factor; two are involved in cytokinin biosynthesis; two are connected to F-box/kelch protein; and one is related to glutathione metabolism (Table S1). Gene expression analysis revealed that qRT-PCR result was consistent with RNA-Seq analysis, suggesting that RNA-Seq data was reliable (Fig. 9).

Discussion

Nitrogen metabolism genes are responsible for various N levels and auxin

Nitrogen and auxin significantly affect LR development via N signaling, and regulate biosynthesis and transport of hormones such as ABA, GA and IAA. In previous studies, N-deficiency induced high affinity nitrate transporter *NRT1.1*, *NRT2.1*, *NRT2.4* and *NRT2.5* in roots of *Arabidopsis* plant[28]. It was also reported that *NRT2.1*, *NRT2.2*, *NRT2.4* and *NRT2.5* could synergistically confer to plants ability to adapt to low N condition[29]. *NRT1* and *NRT2* are not only for NO_3^- transportation but also for auxin transportation under low nitrogen condition. When plants are subjected to low nitrogen, *NRT1* and *NRT2* have high expression and induce auxin accumulation and transportation for regulating the LR formation[30]. Similarly, in this study, we found that two common DEGs (CSA011051; MSTRG.51865) *NRT2.4* and *NRT2.5* were identified from LR under all four treatments. *NRT2.4* and *NRT2.5* were up-regulated and down-regulated in the LR of seedlings under LN and LN + NPA treatments, respectively. And from the results of hormone determination, the N-deficiency could increase the content of auxin in lateral roots. Therefore, LN treatment could induce *NRTs* expression and thus increase auxin production and accumulation. And they could be regulated by nitrogen concentration and auxin treatment contributing to tea plant LR formation.

Auxin and nitrogen signaling could control LR development. Tryptophan (Trp) aminotransferase of *Arabidopsis*1 (TAA1) is an important enzyme which plays important function in Trp transformation in to indole-3-pyruvic acid (IPyA), an auxin biosynthesis pathway (IPyA pathway)[31]. Recent studies have shown that TAR regulates plant roots and shoots development: Tryptophan aminotransferase related genes (*TAR1-4*) were involved in IPyA pathway[32], while tryptophan aminotransferase related 2 (*TAR2*) gene was induced by N deficiency that can improve auxin biosynthesis in *Arabidopsis*, and increase IAA levels in LR development[33, 34]. *TAR2* is desired to accumulate auxin in LR under low nitrogen condition. In the present study, one tryptophan aminotransferase related 2 gene (CSA015778) was down-regulated under HN + IBA treatment, indicating that LN treatment could induce *TAR2* expression for auxin accumulation. However, under excessive exogenous auxin treatment, *TAR2* would be down-regulated to balance LR auxin concentration. The Tryptophan aminotransferase related 4 (*TAR4*) gene was up-regulated under low nitrogen condition in the shoot of *Arabidopsis*[33]. Similar observation was present in this study as *TAR4* (CSA001598) was up-regulated in LR under LN treatment. Therefore, we speculate that *TAR4* also participate in LR development under low nitrogen condition. The auxin concentration increased as nitrogen level decreased in LR of seedlings. In summary, the tea plant LR formation could be induced by low nitrogen concentration via auxin biosynthesis and accumulation.

Arabidopsis has six AMT-type ammonium transporters including *AMT1.1* to *AMT1.5* and *AMT2.1*[35]. Previous reports demonstrated that ammonium supply can induce the LR initiation and branching in *Arabidopsis*, this may be because that ammonium is complementary to nitrate in regulating LR development, which happens in AMT-dependent manners[36]. Ruan *et al.* (2016) reported that ammonia improves nitrate uptake in tea roots, however, the present study revealed that *AMT* genes expression is

diametrically opposite to *NRT* genes in LN and HN + IBA treatment[37]. Among the six AMT-type ammonium transporters identified in the model plant *Arabidopsis*, *AMT1.4* was expressed in the pollen[35], but *AMT1.4* (CSA018499) was identified to express in tea plant roots in the present study. It was down-regulated in LRs under LN and HN + IBA treatments, and up-regulated in LN + NPA treatment. We concluded that *AMT* genes were down-regulated under low nitrogen condition and auxin treatment, and their expressions may inhibit LRs formation. In addition, the expression analysis of *NRT*, *AMT*. and *NPF* family genes revealed that the synergistic relationship of auxin and nitrogen signaling on LR development does exist. These genes contribute to N utilization efficiency exploration and provide gene reference for selection of high nitrogen-efficient varieties of tea plants. This can also contribute to the exploration of regulation of LRs formation through regulating nitrogen and auxin signal pathway.

Plant hormone metabolism and signaling pathways

The plant hormone, auxin is critical for plant growth and development processes, and plays its regulatory role mainly by inducing expression of early auxin response genes including *Aux/IAA*, *GH3* and *SAUR*. *GH3* can combine free auxin with disparate acid; therefore, overexpression of *GH3* would result in expression of severe auxin-deficient phenotypes[38, 39]. In the present study, one *GH3* gene was up-regulated under LN treatment but downregulated under LN + NPA treatment, while the HN + IBA treatment regulated *GH3* genes. In addition, *Aux/IAA* proteins play the role of transcriptional repressors by heterodimerizing with auxin response factor (*ARF*) transcription factors, while *ARF* family acts as key regulator of root development[40, 41]. At low auxin concentration, *Aux/IAA* proteins could restrain transcriptional activation of *ARF* proteins; thereby, preventing response genes transcription of auxin. But higher auxin concentration could induce *ARF* genes expression and then promote LRs development[41]. Under LN treatment, the *Aux/IAA* proteins genes (CSA031541, MSTRG.7473) were down-regulated, while auxin response factor 2 (*ARF2*) gene (CSA011327) was up-regulated. This; therefore, shows that in LN condition, the LRs' auxin concentration would be increased, and then inhibit *Aux/IAA* genes expression and improve *ARF* genes expression to induce LRs development. Moreover, under LN + NPA treatment, the *ARF2* gene (CSA011327) was down-regulated, while *ARF2* gene (CSA012843) was up-regulated under HN + IBA treatment. This; therefore, indicates that low nitrogen and auxin treatments could improve *ARFs* expression and enhance LRs formation.

Cytokinin is important for plants proliferation, plants cell division, secondary metabolism, and regulation of plants shoot and roots development[42]. Other researchers have shown that some nitrogen signals are substituted by cytokinins as local and long-distance signal, and; thus, various genes were regulated by these plant hormones, including metabolism, growth development and nutrient absorption[43]. High cytokinin content is reported to improve shoot development while high auxin content enhances root formation[44]. Therefore, there exist an important signaling pathway among nitrogen, auxin and cytokinin in the regulation of plant development. The present study identified DEGs involved in cytokinin metabolism. Under LN treatment, one adenylate isopentenyltransferase (CAS006753) gene which induce cytokinin biosynthesis was down-regulated, while two cytokinin dehydrogenase genes which inhibit cytokinin biosynthesis were up-regulated. Under LN + NPA treatment, one cytokinin dehydrogenase gene

(CAS011288) was down-regulated, but expressed in LN treatment. Under HN + IBA treatment, cytokinin hydroxylases gene (CAS017731) which induce cytokinin biosynthesis was down-regulated, while 7 cytokinin dehydrogenase genes were up-regulated. RP-UPLC technique revealed that cytokinin concentration increased with increasing nitrogen concentration and decreased with exogenous auxin treatment. It suggests that LN treatment and auxin treatment could inhibit cytokinin biosynthesis, while high auxin condition induces LR formation in tea plants.

Ethylene is connected to plant's physiological and morphological responses to nitrogen deficiency, and nitrate transporters *NRT1.1* and *NRT2.1* are also sensitive to ethylene[45, 46]. Under low external nitrate concentration, *NRT2.1* induces and promotes ethylene biosynthesis and signaling activity[47]. Auxin and ethylene signaling pathways show specific regulation of plants growth and development, such as root elongation and root hair formation[48]. Studies also suggest that ethylene might stimulate localized auxin biosynthesis[49]. In the present study, ethylene biosynthesis varied with different levels of nitrogen treatment. Under LN, ethylene biosynthesis genes were up-regulated, but down-regulated under HN treatment. Under LN + NPA treatment, 3 ethylene biosynthesis genes were down-regulated while 14 genes were up-regulated under HN + IBA treatment. This; therefore, suggests that LN treatment improves ethylene biosynthesis; thereby, promoting auxin response genes expression and LR formation, while HN treatment down-regulates ethylene biosynthesis; thus, inhibits LR formation. Comparison between HN + IBA and LN + NPA treatments revealed that auxin could promote ethylene biosynthesis and stimulate LR formation. These results; therefore, clearly indicate that auxin and nitrogen could regulate tea plant LR formation through ethylene biosynthesis pathway.

Transcription Factors

Transcription factors (TFs) can control the expression of stress resistance genes[50]. Many TF families such as *NAC*, *MYB*, *MADS-box* and *WRKY* have been explored[51], and these families can regulate cell division and expansion, lateral root development and secondary cell wall biosynthesis. Several TFs have been expressed in plants exposed to N-deficient situations[52, 53].

R2R3-MYB is reported to be the most abundant MYB protein. MYBs46/83 is speculated to be the prime regulator of secondary cell wall biosynthesis, while *AtMYB58* specifically activate lignin biosynthesis, as regulated by *AtMYB46*[54]. Therefore, MYBs expression would thicken the secondary cell wall and inhibit cells division and elongation[55, 56]. In the present study, the P-type R2R3 MYB protein (*MYB83*) homologous gene was down-regulated under LN and HN + IBA treatment, but up-regulated with NPA treatment. This; therefore, indicates that LN treatment could inhibit *MYB* genes expression and restrain secondary cell wall biosynthesis, and; thus, regulate roots development. Similar inference can be made with auxin treatment. This also confirms the hypothesis that accumulation of auxin can be promoted under low nitrogen conditions in LR of tea plant.

NAC transcription factors are important for plant growth as they could regulate plants cell division, lateral root development and secondary cell wall biosynthesis[57, 58]. A few *NAC* genes have been identified as key and effective regulation factor in auxin signaling pathway which directly affect LR development[59,

60]. In the present study, 7 *NAC* DEGs were expressed under LN treatment; 43 *NAC* DEGs were expressed under HN + IBA treatment and most of them were up-regulated. Similarly, DEGs of *NAC* also up-regulated under LN + NPA treatment. Consistent with previous reports, the present study; therefore, observed that *NAC* genes could be induced by nitrogen treatment and auxin treatment to regulate tea plants LR formation; however, the detail signaling pathway still needs further exploration.

MADS-box TFs control plants root, flower and fruit development[61, 62]. Previous study has shown that *AGL21* is induced by N-deprivation, and auxin also promotes *AGL21*, while *AGL21* proteins interact with *ANR1* (*AGL44*) to mediate LR development[63, 64]. In the present study, 12 MADS-box protein genes were down-regulated under LN treatment. Under HN + IBA treatment, there are 19 DEGs. Also, *AGL* genes were up-regulated under LN + NPA treatment. Thus, it is tempting to suggest that in tea plant, the MADS-box protein genes expression might differ from that of *Arabidopsis*.

WRKY transcription factors could be involved in various plant developmental processes, such as biological and abiotic stresses, and seed germination and dormancy[65]. WRKY TF is a major player in plant's innate immune system. Beet cyst nematode is reported to regulate WRKY TF genes expression to enhance roots development in *Arabidopsis*[66]. In a previous study, WRKY TF families were induced under N-deficient condition[67]. In the current study, WRKY TF family genes were induced by various levels of nitrogen treatments (Table S2). This reveals that WRKY TF family genes could be significantly induced by auxin signaling, and; thus, take part in nitrogen metabolism under various nitrogen conditions. It can also be deduced that TFs play important roles in nitrogen and auxin network in tea plant LR formation. It is; therefore, important to explore the involvement of WRKY TF family gene via nitrogen and auxin signaling pathway in LR formation in tea plant.

Glutathione Metabolism

KEGG analysis revealed a significant change in glutathione metabolism in all treatments. GSH-dependent developmental pathway induces and sustains cell division during root development, and regulates auxin transport and evolution[67, 68]. Glutathione also acts as thiol/disulfide buffer. It can regulate the balance between GSH (reduced form) and GSSG (oxidized form) by GSH oxidation through reactive oxygen species, and GSSG reduction through glutathione reductase⁶⁸. Exogenous GSSG could not induce roots in normal conditions but promote root development under auxin treatment[69]; therefore, auxin and GSSG interaction would regulate plants roots development. It is reported that the reduction ratio of GSH/GSSG inhibits lateral roots in the presence of auxin[70]. Glutathione S-transferases (GSTs) transforms GSH to GSSG, while glutathione reductase (GR) induces the reduction of GSSG into GSH[71]. In the present study, 8 GSTs genes were up-regulated and 12 GST genes were down-regulated with LN treatment. Under HN condition, 14 GSTs genes were down-regulated, while 5 GSTs genes were up-regulated. IBA treatment up-regulated 5 glutathione reductase genes and 65 GSTs genes. Finally, 37 GSTs genes were up-regulated under NPA treatment. Nitrogen and auxin treatments had many DEGs in glutathione metabolism, both of them could affect GSH/GSSG ratio to regulate LR formation in tea plant; however, the specific adjustment mechanism is still vague and requires further research.

F-box Protein

F-box proteins are important ingredient of proteasome pathway and participate in cellular functions, such as auxin receptor (TIR1), which mediates transcriptional response to auxin in a F-box protein[72]. Many researches have shown that adventitious roots formation accompanies soluble and insoluble carbohydrates accumulation, and the *At1g23390* is shown to be related to such metabolism[73, 74]. When tea plants were treatment with IBA, a F-box/kelch gene similar to *AT1g23390* was identified and this suggests a complex regulatory network during adventitious roots development[70]. In the current study, *AT1g23390* (CAS036587) was up-regulated by HN + IBA treatment, and its homogenous gene was down-regulated by LN + NPA treatment but up-regulated by LN treatment. It; therefore, depicts that this gene could be induced by nitrogen treatment, and its function was similar with auxin treatment in root development. It also indicates that LN treatment could induce auxin production in tea plant LRs. Hence, the putative role of this gene might be to regulate LRs formation through N and auxin signaling pathway in tea plants.

Conclusions

In this study, transcriptome analysis of LRs of tea plant (*Camellia sinensis*) treated with low and high nitrogen concentration, indole-3-butyric acid (IBA) and N -1-naphthylphthalamic acid (NPA) was carried out to reveal the function of auxin and nitrogen in LRs formation and development. Nitrogen deficiency could induce the expression of *NRT* genes, thus increase the production and accumulation of auxin, promote lateral roots formation directly. In the process of cytokinin synthesis, nitrogen deficiency could restrain cytokinin production, and the auxin accumulation and synthesis could control cytokinin production by promoting cytokinin hydroxylase and cytokinin dehydrogenase to improve auxin/cytokinin ratio, and further improve LRs formation and development. Nitrogen deficiency and auxin accumulation and synthesis also regulate growth and development of lateral roots by promoting metabolic synthesis of ethylene. Auxin and N deficiency would inhibit secondary cell wall biosynthesis by inhibiting *MYB* genes expression to facilitate lateral roots development. Auxin and N deficiency also synergistically control the GSH/GSSG ratio by effecting Glutathione metabolism. TF families like *NAC*, *WRKY* and *MAD-box* genes were affected by nitrogen and auxin treatments, similar effect was observed in early auxin response genes including *Aux/IAA*, *GH3* (Fig. 10). The results in this study are meaningful in building an overall regulation network, but requires additional genetic and physiological data to realize this. This study would offer a foundation for further exploration into LRs formation, and also accelerate genomic studies on tea plant.

Methods

Plant materials

Seeds (*Camellia sinensis* cv. Fuding dabai) were surface-sterilized and pre-grown in a liquid medium for six weeks. Purebred fuding dabai tea seeds from Fujian Province Lianfeng Tea Co., Ltd

(<http://1016526.b2b.qth58.cn>). The solution contained 0.75 mM $(\text{NH}_4)_2\text{SO}_4$, 0.25 mM $\text{Ca}(\text{NO}_3)_2 \cdot 4(\text{H}_2\text{O})_3$, 0.05 mM KH_2PO_4 , 0.35 mM K_2SO_4 , 0.395 mM CaCl_2 , 0.21 mM MgSO_4 , 35.0 μM NaFeEDTA, 46.1 μM H_3BO_3 , 2.0 μM MnSO_4 , 0.3 μM CuSO_4 , 2.0 μM ZnSO_4 and 0.5 μM Na_2MoSO_4 . The basic nitrogen (N) nutrient solution was set into four levels, including a gradient N concentration of 0 mM (labelled as no nitrogen, NN), 0.25 mM (labelled as low nitrogen, LN), 1 mM (labelled as control, CK), and 2.5 mM (labelled as high nitrogen, HN). The control experiment (1 mM) was supplemented with 0.75 mM ammonium and 0.25 mM nitrate using $(\text{NH}_4)_2\text{SO}_4$ and $\text{Ca}(\text{NO}_3)_2 \cdot 4(\text{H}_2\text{O})$ respectively, which the best combination of N concentration for seedlings growth.

The seedlings were grown in a growth chamber at 28/25°C (day/night), 75% relative humidity, 16/8 h (light/darkness) photoperiod, and 300 $\mu\text{mol}^{-2}\text{s}^{-1}$ light intensity. The growth medium was replaced with the same fresh liquid medium every seven days. The seedlings have grown LRs post ten weeks of treatment, and then the 2.5 and 0.25 mM nitrogen treated seedlings were supplemented with 0.4 mM IBA and 20 μM NPA, respectively. After 24 h, the LRs treated with 0.25, 1, 2.5 mM nitrogen, 2.5 mM nitrogen + 0.4 mM IBA and 0.25 mM nitrogen + 20 μM NPA were harvested and frozen in liquid nitrogen for latter RNA extraction. To acquire the LRs from the tea seedlings, the seedlings were treated with 0.25 mM, 1 mM and 2.5 mM nitrogen respectively for 10 weeks, and then the tea seedlings cultured with the 2.5 mM nitrogen were further treatment with 10 μM IBA for eight weeks to harvest lateral roots. Finally, the four kinds of LRs obtained from the collected samples were used to measure hormones using RP-UPLC examination and phenotypic observation. The tea seedlings cultured with the 0.25 mM nitrogen were further treatment with 10 μM NPA for four weeks to be only used for phenotypic observation.

Lateral roots number and length analysis

The lateral roots of each tea plant were cut down and laid flat on a flat plate. The WinRHIZO root analysis system (Regent Instruments, Inc., Canada) was used to scan the lateral roots and record the total number and length of the lateral roots of each tea plant. Three biological replicates were performed for each sample.

RNA isolation, library construction and RNA sequencing

Total RNA was isolated from LRs of tea seedlings using plant RNA extraction kit with DNase (TIANDZ, Inc., Beijing, China) according to the manufacturer's protocol. The quantity and quality of the RNA samples were determined by using NanoDrop2000 Spectrophotometer (Thermo Fisher Scientific, USA), 1.2 % agarose gel electrophoresis, and Agilent 2100 Bioanalyzer (Agilent Technologies, Inc., Santa Clara, CA, USA). Then, RNA samples were sent to Genepioneer Biotech Corporation (Nanjing, China) for library construction and sequencing. Eighteen libraries were constructed and sequenced with PE150 using Illumina HiSeq platform. Each sample was sequenced with three biological replications. All sequenced data have been deposited to the NCBI Sequence Read Archive under the GenBank accession number SRA number (uploading).

Sequencing data analysis

Raw reads were cleaned by removing adaptor sequence, ambiguous reads ('N'×10), and low-quality reads (i.e., where more than 50% of bases in a read had a quality value $Q \leq 5$) using an in-house perl script. High-quality clean reads were mapped to the latest version (CAS institute of Botany, Kunming) of tea plant reference genome (http://www.plantkingdomgdb.com/tea_tree) using HISAT2 software[22, 23].

Functional annotation and pathway analysis

All mapped genes and unmapped genes obtained in this study were annotated using public databases including NR database (<http://www.ncbi.nlm.nih.gov/>), Swiss-Prot database (<http://www.expasy.ch/sprot>), COG (<http://www.ncbi.nlm.nih.gov/COG/>) and Pfam databases (<http://pfam.xfam.org/>) using BLASTX (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Gene Ontology (GO) functional classification for all annotated genes was analysed using WEGO software[24]. A Python script was used to retrieve Kyoto Encyclopedia of Genes and Genomes (KEGG) annotation from blast results. GO enrichment analysis of DEGs were performed using Singular Enrichment Analysis (SEA) method with $P < 0.01$ and $FDR < 0.05$ by agriGO. The hypergeometric Fisher exact test ($P < 0.01$) and Benjamini (FDR < 0.05) were performed to detect statistically significant enrichments of KEGG pathway. GO and KEGG enrichment analysis was performed using the whole tea tree transcriptome setting as background.

Pearson correlation analysis

According to the amount of gene expression in the sample, the correlation coefficient between samples can be calculated to distinguish the sample correlation. For the project of biological duplication, the success of biological duplication can be evaluated by the correlation of samples. Pearson's correlation coefficient R (Pearson's correlation coefficient) was used as the evaluation index of correlation among samples[25]. The closer "R" is to 1, the stronger the correlation between the two samples is. In this study, five samples were analysed for transcriptome, and each sample was repeated three times (T1-T15).

Differential expression analysis

Fragments per kb per million reads (FPKM) method was used to quantify the expression levels of transcripts. The DESeq2 package was used to identify DEGs[26]. The $FDR \leq 0.01$ and the absolute value of $\log_2$ ratio ≥ 1 were set as thresholds value for significant differential gene expression between two samples.

Reversed phase ultra performance liquid chromatography (RP-UPLC) analysis

The harvested tea samples were extracted by methanol for auxin concentration determination. Then the extracted samples were applied for RP-UPLC analysis and the chromatographic conditions were as follows: Hypersil ODS C18 column (250 mm X 4.0 mm, 5 μ m); methanol and ultrapure water (0.5% glacial acetic acid) as mobile phase, and gradient elution. The column temperature was 35 °C, injection volume was 15 μ L, flow rate was 1 mL/min, and detection wavelength was 254 nm. External standard calibration

curve method was employed for the quantitative analysis. For standard preparation and standard curve construction, standard IAA and ZT were dissolved in 50% chromatographic methanol. Peak area (Y) was used as linear regression curve for the mass (X, nmol) to obtain the linear range of the regression equation, standard curve, and detection limits. The concentration gradient was 12.5, 25, 50, 100 and 200 ng/mL. Each measurement was replicated three times.

Quantitative RT-PCR (qRT-PCR) analysis

Total RNA was extracted using plant RNA extraction kit with DNase (TIANDZ Inc., Beijing, China) according to the manufacturer's protocol. First-strand cDNA was synthesized from 2 µg of total RNA using Prime Script RT Reagent Kit (Takara, Japan). The qRT-PCR reaction was performed in a 96-well plates using Bio-Rad Real-time thermal cycler CFX96 with SYBR Premix ExTaq™ Kit (Takara, Dalian, China). The glyceraldehyde-3-phosphate dehydrogenase gene (*GAPDH*) and *ACTIN* of tea plant were used as a reference gene. The relative mRNA expression levels were calculated using the $2^{-\Delta\Delta C_t}$ method[27]. Three biological replicates were performed for each sample. Microsoft Excel 2016 and Sigmaplot 13.0 (Systat Software, Canda) were used for statistical analysis. The difference between the two samples was assessed using the student t-test method, $P < 0.05$ was considered significant statistically. The qPCR primers of differentially expressed genes were designed using NCBI primer-BLAST (<https://www.ncbi.nlm.nih.gov/tools/primer-blast/>). Information on genes and primer sequences used in this study are listed in Table S1.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data sustaining the results in this study are included in this article or its supplementary information files. Other datasets generated during this study are available upon reasonable request from the corresponding author (Xinghui Li).

Competing Interest

The authors declare no competing Interest

Funding

This work was supported financially by the China Earmarked Fund for Modern Agro-industry Technology Research System (CARS-19), Student Innovation Research and Entrepreneurship Training Program (201910307016Z) and Top-notch Academic Programs Project of Jiangsu Higher Education Institutions (PPZY2015B156).

Contributions

SH, WX and XL conceived the study. SH, YY, MM carried out bioinformatics data analysis. ZZ, EA and YC performed qPCR analysis and participated in the sequence alignment. SH and XL wrote the manuscript. QM, AJ, XC critically reviewed the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Not Applicable.

Corresponding authors

Correspondence to Xinghui Li.

Abbreviations

LRs: Lateral roots

indole-3-butyric acid: IBA

N-1-naphthylphthalamic acid: NPA

Eversed phase ultra performance liquid chromatography: RP-UPLC

NN: no nitrogen

LN: low nitrogen

HN: high nitrogen

Nitrate transporter genes: NRTs

AUXIN SIGNALING F-BOX 3: AFB3

Tryptophan aminotransferase related 2: TAR2

Abscisic acid: ABA

Ammonium transporter: AMT

Auxin response factor 2: ARF2

Glutathione S-transferases: GSTs

Tryptophan: Trp

Tryptophan aminotransferase of Arabidopsis1: TAA1

Tryptophan aminotransferase related 4: TAR4)

Transcription factors: TFs

NAM, ATAF1/2 and CUC2: NAC

References

1. Hodge A, Berta G, Doussan C, Merchan F, Crespi M: **Plant root growth, architecture and function.** *Plant Soil* 2009, **321**(1-2):153-187.
2. Varney GT, Canny MJ: **Rates of Water-Uptake into the Mature Root-System of Maize Plants.** *New Phytol* 1993, **123**(4):775-786.
3. Seiler GJ: **Influence of temperature on primary and lateral root growth of sunflower seedlings.** *Environ Exp Bot* 1998, **40**(2):135-146.
4. Li HB, Xia M, Wu P: **Effect of phosphorus deficiency stress on rice lateral root growth and nutrient absorption.** *Acta Bot Sin* 2001, **43**(11):1154-1160.
5. Khan MA, Gemenet DC, Villordon A: **Root System Architecture and Abiotic Stress Tolerance: Current Knowledge in Root and Tuber Crops.** *Front Plant Sci* 2016, **7**:1584.
6. Crawford NM, Forde BG: **Molecular and developmental biology of inorganic nitrogen nutrition.** *Arabidopsis Book* 2002, **1**:e0011.
7. Gruffman L, Jamtgard S, Nasholm T: **Plant nitrogen status and co-occurrence of organic and inorganic nitrogen sources influence root uptake by Scots pine seedlings.** *Tree Physiol* 2014, **34**(2):205-213.
8. Lynch JP: **Steep, cheap and deep: an ideotype to optimize water and N acquisition by maize root systems.** *Ann Bot-London* 2013, **112**(2):347-357.
9. Duan LN, Dietrich D, Ng CH, Chan PMY, Bhalerao R, Bennett MJ, Dinneny JR: **Endodermal ABA Signaling Promotes Lateral Root Quiescence during Salt Stress in Arabidopsis Seedlings.** *Plant Cell* 2013, **25**(1):324-341.
10. Guo YF, Chen FJ, Zhang FS, Mi GH: **Auxin transport from shoot to root is involved in the response of lateral root growth to localized supply of nitrate in maize.** *Plant Sci* 2005, **169**(5):894-900.
11. Gruber BD, Giehl RFH, Friedel S, von Wiren N: **Plasticity of the Arabidopsis Root System under Nutrient Deficiencies.** *Plant Physiol* 2013, **163**(1):161-179.
12. Vidal EA, Araus V, Lu C, Parry G, Green PJ, Coruzzi GM, Gutierrez RA: **Nitrate-responsive miR393/AFB3 regulatory module controls root system architecture in Arabidopsis thaliana.** *Proc Natl Acad Sci U S A*

2010, **107**(9):4477-4482.

13. Okamoto M, Vidmar JJ, Glass AD: **Regulation of NRT1 and NRT2 gene families of Arabidopsis thaliana: responses to nitrate provision.** *Plant Cell Physiol* 2003, **44**(3):304-317.
14. Remans T, Nacry P, Pervent M, Girin T, Tillard P, Lepetit M, Gojon A: **A central role for the nitrate transporter NRT2.1 in the integrated morphological and physiological responses of the root system to nitrogen limitation in Arabidopsis.** *Plant Physiol* 2006, **140**(3):909-921.
15. Zhang HM, Forde BG: **An Arabidopsis MADS box gene that controls nutrient-induced changes in root architecture.** *Science* 1998, **279**(5349):407-409.
16. Engineer CB, Kranz RG: **Reciprocal leaf and root expression of AtAmt1.1 and root architectural changes in response to nitrogen starvation.** *Plant Physiol* 2007, **143**(1):236-250.
17. Himanen K, Boucheron E, Vanneste S, Engler JD, Inze D, Beeckman T: **Auxin-mediated cell cycle activation during early lateral root initiation.** *Plant Cell* 2002, **14**(10):2339-2351.
18. De Smet I, Signora L, Beeckman T, Inze D, Foyer CH, Zhang HM: **An abscisic acid-sensitive checkpoint in lateral root development of Arabidopsis.** *Plant J* 2003, **33**(3):543-555.
19. Lavenus J, Goh T, Roberts I, Guyomarc'h S, Lucas M, De Smet I, Fukaki H, Beeckman T, Bennett M, Laplace L: **Lateral root development in Arabidopsis: fifty shades of auxin.** *Trends Plant Sci* 2013, **18**(8):455-463.
20. Chandler JW, Werr W: **Cytokinin-auxin crosstalk in cell type specification.** *Trends Plant Sci* 2015, **20**(5):291-300.
21. Gou JQ, Strauss SH, Tsai CJ, Fang K, Chen YR, Jiang XN, Busov VB: **Gibberellins Regulate Lateral Root Formation in Populus through Interactions with Auxin and Other Hormones.** *Plant Cell* 2010, **22**(3):623-639.
22. Xia EH, Zhang HB, Sheng J, Li K, Zhang QJ, Kim C, Zhang Y, Liu Y, Zhu T, Li W *et al*: **The Tea Tree Genome Provides Insights into Tea Flavor and Independent Evolution of Caffeine Biosynthesis.** *Mol Plant* 2017, **10**(6):866-877.
23. Kim D, Landmead B, Salzberg SL: **HISAT: a fast spliced aligner with low memory requirements.** *Nat Methods* 2015, **12**(4):357-U121.
24. Ye J, Fang L, Zheng HK, Zhang Y, Chen J, Zhang ZJ, Wang J, Li ST, Li RQ, Bolund L *et al*: **WEGO: a web tool for plotting GO annotations.** *Nucleic Acids Res* 2006, **34**:W293-W297.
25. Schulze SK, Kanwar R, Golzenleuchter M, Therneau TM, Beutler AS: **SERE: Single-parameter quality control and sample comparison for RNA-Seq.** *Bmc Genomics* 2012, **13**:1-9.
26. Anders S, Huber W: **Differential expression analysis for sequence count data.** *Genome Biol* 2010, **11**(10).
27. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2(T)(-Delta Delta C) method.** *Methods* 2001, **25**(4):402-408.
28. Sun CH, Yu JQ, Hu DG: **Nitrate: A Crucial Signal during Lateral Roots Development.** *Front Plant Sci* 2017, **8**.

29. Lezhneva L, Kiba T, Feria-Bourrellier AB, Lafouge F, Boutet-Mercey S, Zoufan P, Sakakibara H, Daniel-Vedele F, Krapp A: **The Arabidopsis nitrate transporter NRT2.5 plays a role in nitrate acquisition and remobilization in nitrogen-starved plants.** *Plant J* 2014, **80**(2):230-241.
30. Krouk G, Lacombe B, Bielach A, Perrine-Walker F, Malinska K, Mounier E, Hoyerova K, Tillard P, Leon S, Ljung K *et al*: **Nitrate-Regulated Auxin Transport by NRT1.1 Defines a Mechanism for Nutrient Sensing in Plants.** *Dev Cell* 2010, **18**(6):927-937.
31. Stepanova AN, Robertson-Hoyt J, Yun J, Benavente LM, Xie DY, Dolezal K, Schlereth A, Jurgens G, Alonso JM: **TAA1-mediated auxin biosynthesis is essential for hormone crosstalk and plant development.** *Cell* 2008, **133**(1):177-191.
32. He WR, Brumos J, Li HJ, Ji YS, Ke M, Gong XQ, Zeng QL, Li WY, Zhang XY, An FY *et al*: **A Small-Molecule Screen Identifies L-Kynurenine as a Competitive Inhibitor of TAA1/TAR Activity in Ethylene-Directed Auxin Biosynthesis and Root Growth in Arabidopsis.** *Plant Cell* 2011, **23**(11):3944-3960.
33. Zhao YD: **Auxin Biosynthesis: A Simple Two-Step Pathway Converts Tryptophan to Indole-3-Acetic Acid in Plants.** *Mol Plant* 2012, **5**(2):334-338.
34. Tao Y, Ferrer JL, Ljung K, Pojer F, Hong FX, Long JA, Li L, Moreno JE, Bowman ME, Ivans LJ *et al*: **Rapid synthesis of auxin via a new tryptophan-dependent pathway is required for shade avoidance in plants.** *Cell* 2008, **133**(1):164-176.
35. Yuan LX, Graff L, Loque D, Kojima S, Tsuchiya YN, Takahashi H, von Wiren N: **AtAMT1;4, a Pollen-Specific High-Affinity Ammonium Transporter of the Plasma Membrane in Arabidopsis.** *Plant Cell Physiol* 2009, **50**(1):13-25.
36. Lima JE, Kojima S, Takahashi H, von Wiren N: **Ammonium Triggers Lateral Root Branching in Arabidopsis in an AMMONIUM TRANSPORTER1;3-Dependent Manner.** *Plant Cell* 2010, **22**(11):3621-3633.
37. Ruan L, Wei K, Wang LY, Cheng H, Zhang F, Wu LY, Bai PX, Zhang CC: **Characteristics of NH₄⁺ and NO₃⁻ fluxes in tea (*Camellia sinensis*) roots measured by scanning ion-selective electrode technique.** *Sci Rep-Uk* 2016, **6**.
38. Zhang SW, Li CH, Cao J, Zhang YC, Zhang SQ, Xia YF, Sun DY, Sun Y: **Altered Architecture and Enhanced Drought Tolerance in Rice via the Down-Regulation of Indole-3-Acetic Acid by TLD1/OsGH3.13 Activation.** *Plant Physiol* 2009, **151**(4):1889-1901.
39. Peat TS, Bottcher C, Newman J, Lucent D, Cowieson N, Davies C: **Crystal Structure of an Indole-3-Acetic Acid Amido Synthetase from Grapevine Involved in Auxin Homeostasis.** *Plant Cell* 2012, **24**(11):4525-4538.
40. Ulmasov T, Murfett J, Hagen G, Guilfoyle TJ: **Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements.** *Plant Cell* 1997, **9**(11):1963-1971.
41. Kumar R, Tyagi AK, Sharma AK: **Genome-wide analysis of auxin response factor (ARF) gene family from tomato and analysis of their role in flower and fruit development.** *Mol Genet Genomics* 2011, **285**(3):245-260.

42. Krouk G, Ruffel S, Gutierrez RA, Gojon A, Crawford NM, Coruzzi GM, Lacombe B: **A framework integrating plant growth with hormones and nutrients.** *Trends Plant Sci* 2011, **16**(4):178-182.
43. Brenner WG, Romanov GA, Kollmer I, Burkle L, Schmulling T: **Immediate-early and delayed cytokinin response genes of Arabidopsis thaliana identified by genome-wide expression profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through transcriptional cascades.** *Plant J* 2005, **44**(2):314-333.
44. Moubayidin L, Di Mambro R, Sabatini S: **Cytokinin-auxin crosstalk.** *Trends Plant Sci* 2009, **14**(10):557-562.
45. Schmelz EA, Alborn HT, Engelberth J, Tumlinson JH: **Nitrogen deficiency increases volicitin-induced volatile emission, jasmonic acid accumulation, and ethylene sensitivity in maize.** *Plant Physiol* 2003, **133**(1):295-306.
46. Tian QY, Sun P, Zhang WH: **Ethylene is involved in nitrate-dependent root growth and branching in Arabidopsis thaliana.** *New Phytol* 2009, **184**(4):918-931.
47. Zheng DC, Han X, An Y, Guo HW, Xia XL, Yin WL: **The nitrate transporter NRT2.1 functions in the ethylene response to nitrate deficiency in Arabidopsis.** *Plant Cell Environ* 2013, **36**(7):1328-1337.
48. Muday GK, Rahman A, Binder BM: **Auxin and ethylene: collaborators or competitors?** *Trends Plant Sci* 2012, **17**(4):181-195.
49. Ruzicka K, Ljung K, Vanneste S, Podhorska R, Beeckman T, Friml J, Benkova E: **Ethylene regulates root growth through effects on auxin biosynthesis and transport-dependent auxin distribution.** *Plant Cell* 2007, **19**(7):2197-2212.
50. Cai GH, Wang GD, Wang L, Pan JW, Liu Y, Li DQ: **ZmMKK1, a novel group A mitogen-activated protein kinase gene in maize, conferred chilling stress tolerance and was involved in pathogen defense in transgenic tobacco.** *Plant Sci* 2014, **214**:57-73.
51. Le Henanff G, Profizi C, Courteaux B, Rabenoelina F, Gerard C, Clement C, Baillieul F, Cordelier S, Dhondt-Cordelier S: **Grapevine NAC1 transcription factor as a convergent node in developmental processes, abiotic stresses, and necrotrophic/biotrophic pathogen tolerance.** *J Exp Bot* 2013, **64**(16):4877-4893.
52. Peng MS, Bi YM, Zhu T, Rothstein SJ: **Genome-wide analysis of Arabidopsis responsive transcriptome to nitrogen limitation and its regulation by the ubiquitin ligase gene NLA.** *Plant Mol Biol* 2007, **65**(6):775-797.
53. Quan XY, Zeng JB, Ye LZ, Chen G, Han ZG, Shah JM, Zhang GP: **Transcriptome profiling analysis for two Tibetan wild barley genotypes in responses to low nitrogen.** *Bmc Plant Biol* 2016, **16**.
54. Zhou J, Lee C, Zhong R, Ye ZH: **MYB58 and MYB63 are transcriptional activators of the lignin biosynthetic pathway during secondary cell wall formation in Arabidopsis.** *Plant Cell* 2009, **21**(1):248-266.
55. Piya S, Kihm C, Rice JH, Baum TJ, Hewezi T: **Cooperative Regulatory Functions of miR858 and MYB83 during Cyst Nematode Parasitism.** *Plant Physiol* 2017, **174**(3):1897-1912.

56. Zhong RQ, Lee C, McCarthy RL, Reeves CK, Jones EG, Ye ZH: **Transcriptional Activation of Secondary Wall Biosynthesis by Rice and Maize NAC and MYB Transcription Factors.** *Plant Cell Physiol* 2011, **52**(10):1856-1871.
57. Bollhoner B, Prestele J, Tuominen H: **Xylem cell death: emerging understanding of regulation and function.** *J Exp Bot* 2012, **63**(3):1081-1094.
58. Nakashima K, Takasaki H, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K: **NAC transcription factors in plant abiotic stress responses.** *Bba-Gene Regul Mech* 2012, **1819**(2):97-103.
59. Xie Q, Frugis G, Colgan D, Chua NH: **Arabidopsis NAC1 transduces auxin signal downstream of TIR1 to promote lateral root development.** *Gene Dev* 2000, **14**(23):3024-3036.
60. He XJ, Mu RL, Cao WH, Zhang ZG, Zhang JS, Chen SY: **AtNAC2, a transcription factor downstream of ethylene and auxin signaling pathways, is involved in salt stress response and lateral root development.** *Plant J* 2005, **44**(6):903-916.
61. Gu Q, Ferrandiz C, Yanofsky MF, Martienssen R: **The FRUITFULL MADS-box gene mediates cell differentiation during Arabidopsis fruit development.** *Development* 1998, **125**(8):1509-1517.
62. Ratcliffe OJ, Nadzan GC, Reuber TL, Riechmann JL: **Regulation of flowering in Arabidopsis by an FLC homologue.** *Plant Physiol* 2001, **126**(1):122-132.
63. Yu LH, Miao ZQ, Qi GF, Wu J, Cai XT, Mao JL, Xiang CB: **MADS-Box Transcription Factor AGL21 Regulates Lateral Root Development and Responds to Multiple External and Physiological Signals.** *Mol Plant* 2014, **7**(11):1653-1669.
64. Sun CH, Yu JQ, Wen LZ, Guo YH, Sun X, Hao YJ, Hu DG, Zheng CS: **Chrysanthemum MADS-box transcription factor CmANR1 modulates lateral root development via homo-/heterodimerization to influence auxin accumulation in Arabidopsis.** *Plant Sci* 2018, **266**:27-36.
65. Rushton PJ, Somssich IE, Ringler P, Shen QJ: **WRKY transcription factors.** *Trends Plant Sci* 2010, **15**(5):247-258.
66. Ali MA, Wieczorek K, Kreil DP, Bohlmann H: **The Beet Cyst Nematode *Heterodera schachtii* Modulates the Expression of WRKY Transcription Factors in Syncytia to Favour Its Development in Arabidopsis Roots.** *Plos One* 2014, **9**(7).
67. Yang WZ, Yoon J, Choi H, Fan YL, Chen RM, An G: **Transcriptome analysis of nitrogen-starvation-responsive genes in rice.** *Bmc Plant Biol* 2015, **15**.
68. Bashandy T, Guilleminot J, Vernoux T, Caparros-Ruiz D, Ljung K, Meyer Y, Reichheld JP: **Interplay between the NADP-Linked Thioredoxin and Glutathione Systems in Arabidopsis Auxin Signaling.** *Plant Cell* 2010, **22**(2):376-391.
69. Tyburski J, Tretyn A: **Glutathione and glutathione disulfide affect adventitious root formation and growth in tomato seedling cuttings.** *Acta Physiol Plant* 2010, **32**(2):411-417.
70. Wei K, Wang LY, Cheng H, Zhang CC, Ma CL, Zhang LQ, Gong WY, Wu LY: **Identification of genes involved in indole-3-butyric acid-induced adventitious root formation in nodal cuttings of *Camellia sinensis* (L.) by suppression subtractive hybridization.** *Gene* 2013, **514**(2):91-98.

71. Jiang HW, Liu MJ, Chen IC, Huang CH, Chao LY, Hsieh HL: **A Glutathione S-Transferase Regulated by Light and Hormones Participates in the Modulation of Arabidopsis Seedling Development.** *Plant Physiol* 2010, **154**(4):1646-1658.
72. Kepinski S, Leyser O: **The Arabidopsis F-box protein TIR1 is an auxin receptor.** *Nature* 2005, **435**(7041):446-451.
73. Ahkami AH, Lischewski S, Haensch KT, Porfirova S, Hofmann J, Rolletschek H, Melzer M, Franken P, Hause B, Druege U *et al.*: **Molecular physiology of adventitious root formation in *Petunia hybrida* cuttings: involvement of wound response and primary metabolism.** *New Phytol* 2009, **181**(3):613-625.
74. Rapaka VK, Faust JE, Dole JM, Runkle ES: **Endogenous carbohydrate status affects postharvest ethylene sensitivity in relation to leaf senescence and adventitious root formation in *Pelargonium* cuttings.** *Postharvest Biol Tec* 2008, **48**(2):272-282.

Tables

Table 1 | Quality assessment of clean data

Samples	Read number	Base number	GC content	%≥Q30
LN-1	22300039	6690011700	45.28	93.81
LN-2	22352271	6705681300	45.25	94.08
LN-3	71345574	21403672200	45.24	92.83
LN+NPA-1	22625999	6787799700	45.23	94.01
LN+NPA-2	22937652	6881295600	45.69	94.27
LN+NPA-3	20972742	6291822600	45.10	93.45
N-CK-1	50566454	15169936200	44.84	93.86
N-CK-2	26485890	7945767000	44.82	94.03
N-CK -3	51958794	15587638200	44.90	94.57
HN+IBA-1	24283790	7285137000	45.18	93.86
HN+IBA-2	25455012	7636503600	45.26	94.41
HN+IBA-3	24543852	7363155600	44.89	94.12
HN-1	22116811	6635043300	44.92	93.97
HN-2	23245627	6973688100	44.83	93.64
HN-3	22586710	6776013000	44.74	93.67

LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24h), HN (2.5 mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 24 h).

Table 2 | Corresponding annotated DEG number

DEG Set	Total	Swiss-Prot	GO	KEGG	COG	KOG	Pfam	NR
LN vs LN+NPA	4498	3248	2814	449	1304	2402	2728	4462
LN vs CK	4183	2977	2610	387	1108	2261	2427	4136
CK vs HN	2992	2107	1822	279	804	1558	1701	2956
HN vs HN+IBA	15274	11300	10235	1606	5037	8349	10328	15209

LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24 h), HN (2.5 mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 µM NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 µM IBA for 24 h).

Table 3 | DEGs related to nitrogen metabolism, plant hormone signal, transcription factors, Glutathione metabolism and F-box/kelch protein

Gene ID	Log2 (Fold change)				Description
	CK vs LN	CK vs HN	LN vs LN+NPA	HN vs HN+IBA	
CSA011051	1.84	1.53	-3.08	-3.24	High affinity nitrate transporter
MSTRG.51865	1.51	1.23	-2.73	-2.72	High affinity nitrate transporter
CSA015778	-	-	-	-2.56	Tryptophan aminotransferase-related protein 2
CSA018499	-2.32	-	2.61	-1.71	Ammonium transporter
CSA011327	3.86	-	-4.11	-	Auxin response factor
CSA012843	-	-	-	2.36	Auxin response factor
CSA006753	-1.33	-	-	-3.83	Adenylate isopentenyltransferase
CSA011288	1.16	-	-2.00	-	Cytokinin dehydrogenase
CSA017731	-2.62	-	-	-	Cytokinin hydroxylase
CSA028278	1.16	-	-	-	Ethylene biosynthetic
MSTRG.10434	-	-1.64	-	-	Ethylene biosynthetic
CSA00344	-	-	-1.68	-	Ethylene biosynthetic
CSA017586	-	-	-1.63	-	Ethylene biosynthetic
CSA017852	-	-	-	3.58	Ethylene biosynthetic
CSA003142	-	-	-	6.42	Ethylene biosynthetic
MSTRG.2855	-2.78	-	4.47	-4.25	P-type R2R3 Myb protein
MSTRG.31932	1.71	-	-1.31	-	F-box/kelch protein
CSA036587	-	-	-	1.17	F-box/kelch protein
CSA012447	-1.46	-1.26	2.74	9.59	Glutathione S-transferase

“ - ” represents no significant difference in gene expression. LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24 h), HN (2.5 mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 µM NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 µM IBA for 24 h).

Figures

Figure 1

Lateral roots growth performance of tea plant in hydroponics culture supplemented with 0, 0.25, 1 and 2.5 mM nitrogen for 10 weeks.

Figure 2

Lateral roots growth performance of tea plant in hydroponics culture supplemented with (a) 2.5 mM nitrogen for 10 weeks, then cultured with 2.5 mM nitrogen + 10 μ M IBA for 8 weeks (b) 1 mM nitrogen for 18 weeks and (c) 0.25 mM nitrogen for 10 weeks, then cultured with 0.25 mM nitrogen + 10 μ M NPA for 4 weeks.

Figure 3

(a) The length of lateral roots in hydroponics culture. (b) The number of lateral roots in hydroponics culture. NN (0 mM nitrogen for 10 weeks), LN (0.25 mM nitrogen for 10 weeks), CK-ten weeks (the control, 1 mM nitrogen for 10 weeks), HN (2.5mM nitrogen for 10 weeks), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 4 weeks), CK-eighteen weeks (the control, 1 mM nitrogen for 18 weeks), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 8 weeks).

Figure 4

Effect of nitrogen and auxin treatments on indole-3-acetic acid (IAA) and zeatin (ZT) concentration in lateral roots. LN (0.25 mM nitrogen for 18 weeks), CK-eighteen weeks (the control, 1 mM nitrogen for 18 weeks), HN (2.5mM nitrogen for 18 weeks), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 8 weeks).

Pearson Correlation of all samples

Figure 5

Pearson correlation analysis of CK, LN, HN, LN+NPA and HN+IBA treatments on lateral roots of *Camellia sinensis*. LN (T1, T2, T3); CK (T4, T5, T6); HN (T7, T8, T9); LN + NPA (T10, T11, T12); HN + IBA (T13, T14, T15). LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24 h), HN (2.5mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 24 h).

Figure 6

Functional annotation and classification of all unigenes identified in tea plant, as determined by Cluster of Orthologous Groups (COG), Non-redundant (NR) and gene ontology (GO) databases.

Figure 7

Number of DEGs between LN vs LN + NPA, LN vs CK, CK vs HN, HN vs HN + IBA treatments. LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24 h), HN (2.5mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 24 h).

Figure 8

Differentially expressed genes in tea plant lateral roots in response to LN vs CK, CK vs HN, LN vs LN + NPA and HN vs HN + IBA. (a) Venn diagram of common DEGs. (b) Functional distribution of common DEGs in tea plant transcriptome. LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24 h), HN (2.5mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 24 h).

Figure 9

Comparison of expression profiles of selected genes as determined by Real Time-PCR. LN (0.25 mM nitrogen for 10 weeks + 24 h), CK (the control, 1 mM nitrogen for 10 weeks + 24 h), HN (2.5 mM nitrogen for 10 weeks + 24 h), LN + NPA (0.25 mM nitrogen for 10 weeks, and then cultured with 0.25 mM nitrogen + 10 μ M NPA for 24 h), HN + IBA (2.5 mM nitrogen for 10 weeks, and then cultured with 2.5 mM nitrogen + 10 μ M IBA for 24 h).

Figure 10

Multiple signaling pathways regulating the formation of lateral roots to N deficiency. Nitrogen deficiency could induce the expression of NRT genes, inhibit cytokinin synthesis, induce synthesis and accumulation of auxin, promote lateral roots formation, and regulate growth and development of lateral roots by promoting metabolic synthesis of ethylene. The auxin accumulation and synthesis could control cytokinin production by promoting cytokinin hydroxylase and cytokinin dehydrogenase to further improve LR formation and development. Auxin and N deficiency would inhibit secondary cell wall biosynthesis by inhibiting MYB genes expression. Auxin and N deficiency also synergistically control the GSH/GSSG ratio by effecting Glutathione metabolism. TF families like NAC, WRKY and MADS-box genes were affected by nitrogen and auxin treatments, similar effect was observed in early auxin response genes including Aux/IAA, GH3.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Figuresupply.docx](#)
- [TableSupply.docx](#)