The Compensatory Increased BDNF and NGF in Patients With Multiple Sclerosis Following Home-based Aerobic Training and Vitamin D Supplementation During the COVID-19 Outbreak

Elahe Bahmani

Razi University

Rastegar Hoseini (✉ R.hoseini@razi.ac.ir)

Razi University

Ehsan Amiri

Razi University

Research Article

Keywords: Exercise, Vitamin D, Multiple sclerosis, BDNF, NGF

Posted Date: July 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-677336/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background: Lifestyle modifications (physical activity and diet) are among the most promising strategies in MS rehabilitation. This study aimed to investigate the effect of home-based aerobic training and vitamin D supplementation in patients with multiple sclerosis during the COVID-19 outbreak.

Methods: In this randomized, single-blinded, placebo-controlled trial, 38 females with Multiple sclerosis with EDSS: 3-5 (aged 20–40 years with body mass index [BMI] of 25–30 kg/m²) were randomly assigned into four groups: aerobic training + Vitamin D supplementation (AT+Vit D; n=10); aerobic training (AT; n=9); Vitamin D supplementation (Vit D; n=9), and Control (Placebo) (n=10). The AT program consisted of 50-70% of HR_{Max}, 25-40 min/day, three days/wk for eight weeks. Participants in the Vit D group consumed 50000 IU of Vitamin D supplement capsules per week for eight weeks. The data were analyzed through paired t-test and one-way analysis of variance, as well as Tukey's post hoc test at the signification level of P<0.05.

Results: BDNF and NGF levels improved significantly from pre-test to post-test within all experimental groups. BDNF and NGF decreased significantly in AT+Vit D, AT, and Vit D compared to C. Also, the results show that the AT+Vit D had significantly lower BDNF and NGF compared to AT (P=0.023 and P=0.011) and Vit D (P=0.001 and P=0.002).

Conclusions: These findings suggest that the combination of AT+Vit D improves BDNF and NGF status more effectively than AT or Vit D alone in female Multiple sclerosis patients.

Background

Multiple sclerosis (MS) is an autoimmune, inflammatory, chronic, and progressive disease in which the immune system attacks the nerve cells in the brain and spinal cord, causing damage to myelin and impairs the body system communication. Creates therapeutic and socio-economic for individuals [1]. MS is the most common neurological disease especially in women [2] occurring between the ages of 20 and 40 years [3]. In recent years, the MS cases are increasing daily and reaching over 35 per 100,000 individuals [3, 4]. According to the results of studies, the estimated prevalence of MS in Tehran is about 52 per 100,000 individuals, of which 72.2% are women and 27.3% are men [5]. The main cause of MS unknown, however a variety of factors, including the presence of genetic background, autoimmune mechanisms, and environmental factors (especially infections, latitude, vitamin D deficiency, and smoking) are effective in the development of MS [6]. In general, MS-related symptoms include fatigue, muscle weakness, spasm, ataxia, imbalance, and gait disturbance [7]. Aerobic training is among the approaches to improve MS symptoms [8]. Researchers have recently been focused on the effect of aerobic training on neuronal function, and especially on neurotrophins, neurogenic, and degenerative factors [9]. Neurotrophins are polypeptide growth factors that cooperate in the growth, survival, and function of the central and peripheral nervous system and play a significant role in nervous system disorders [10, 11]. The results of studies showed the beneficial effects of aerobic training on brain

function, partly due to enhanced function and increased levels of neurotrophins such as BDNF and NGF [12]. Cotman et al. (2002) observed increased neurotrophic factors expression including BDNF and NGF in the hippocampus of two animal models, Mice and Rats following aerobic training [12]. Although the role of BDNF in muscle has not yet been elucidated, it might increase the survival of damaged motor neurons. Serum BDNF levels have also been reported to increase immediately after a session of low intensity aerobic training in MS patients [13]. The NGF secretion, which is involved in biological functions such as neuroprotection, neuronal regeneration, and memory improvement, is affected by regular aerobic training; decreased neuronal apoptosis and demyelination have been also reported in spinal cord samples after NGF injection into the white matter [14]. Also, several studies have investigated the effects of dietary supplements in MS patients including vitamin D [15, 16]. As a non-genetic factor, the distribution of vitamin D receptors has been reported to play an important role in the susceptibility as well as the severity of MS [17]. The association between vitamin D receptor gene polymorphisms and MS was first demonstrated in the Japanese population [18]. Another study showed a direct link between mutation in the gene that controls the vitamin D level (causing vitamin D deficiency) and the development of MS in individuals with a family history; Individuals with low vitamin D were more likely to develop MS in the future [17, 19]. Several studies showed significantly lower serum vitamin D levels in MS patients compared with healthy individuals [20–22]; however, others have not reported [21]. Vitamin D plays a major role in the occurrence or severity of MS by regulating the immune system function, bone metabolism, cell regulation, proliferation, and differentiation, encephalomyelitis and T cell function [23], and the expression of many important genes in various pathways related to cancers and many autoimmune diseases [24]. Also, increased NGF and BDNF secretion induced by aerobic training might reduce the MS symptoms. However, to date, limited studies have been evaluated the simultaneous effect of aerobic training and vitamin D supplementation on nerve cell growth factors in patients with MS. On the one hand, due to the MS complications and limited studies on this individual, and on the other hand, the risk of exercising outside in sports facilities in the outbreak of COVID-19, this study aimed to investigate the effect of eight-week home-based aerobic training with vitamin D supplementation on serum BDNF and NGF in MS patients during COVID-19 outbreak.

Material And Methods

Participants and Study Design

This clinical trial study aimed to investigate the effects of 8-week home-based aerobic training and Vit D supplementation on BDNF and NGF in patients with multiple sclerosis. As detailed in Fig. 1, the study population consisted of 60 participants (aged 20–40 years) from the Kermanshah MS center with a disability scale of 3–5.

In this study, 40 women with multiple sclerosis (20–40 years) in Kermanshah were selected using purposive and available sampling method based on G. POWER 3.1 software with a statistical power of 99 %, the effect size of 95%, and significance level at 0.05.

Inclusion criteria included: at least 2 years MS history, disability scale of 3–5, lack of regular exercise, no history of other diseases, no smoking, and no immunosuppressive drug consumption. Exclusion criteria included a history of heart disease, hypertension, orthopedic disorders, and diabetes, consuming other drugs in addition to MS medications that might affect the individual's response to the intervention, lack of regular attendance in the exercise training and testing sessions muscle injuries, inability to perform the exercise, COVID-19 infection, and severe relapses during the study period.

The trial (IR.RAZI.REC.1400.002) was approved by the Ethics Committee of the Kermanshah Razi University and registered in the Iranian Clinical Trial Registration Center under the code IRCT IRCT20201129049525N; Written informed consent was obtained from all participants, including agreement of the patients to participate as volunteers and feasibility to leave the study. Forty subjects were randomly assigned into four equal ($n = 10$) groups; aerobic training plus vitamin D supplementation (AT + Vit D), aerobic training (AT), vitamin D supplementation (Vit D), and control (C) using the lottery method.

IRCT20201129049525N1

Intervention

The AT experimental groups exercised online three times per week for 2 months at home. All training sessions were supervised by an exercise physiologist. Vitamin D supplementation was performed based on the standard guidelines.

Aerobic Training

was executed online.

The subjects were instructed to do home-based walking aerobics three times per week, increasing the time and intensity as shown in Table 1 to 40 min and 70 % of HRmax [25] which was recommended by the Multiple Sclerosis Association of America (MSAA) [26] (Table 1). Every session included 10 min of warm-up and 10 min of cool-down.

The HRmax formula [$HR_{max} = 220 - \text{age}$] was used to determine the target heart rate [27]. The pulse palpation method was taught to the participants in an instructional session and they learned to monitor heart rate. Also, the 6–20 rating of perceived exertion (RPE) scale was considered to assure achieving and maintaining the desired exercise intensity during the walking aerobics phase, [28] (Table 1). The intervention continued for 8 weeks.

Table 1
Aerobic training protocol

Variables	Week							
	1	2	3	4	5	6	7	8
Intensity (HR _{max})	50– 55%	50– 55%	55– 60%	55– 60%	60– 65%	60– 65%	65–70%	65– 70%
Time (min)	20	25	25	30	30	35	35	40
Borg scale	10	10	10	11	11	11	12	12

Vitamin D supplementation and Diet

The participants filled the pre-participants physical activity readiness questionnaire (including demographic data, health status, and PA) before attending the training program, and a detailed semi-quantitative food frequency questionnaire to record and analyze 3-day food, recalls, adapted to the Iranian population (including common food items, serving sizes, and meals) before and at the end of the intervention. The Food Processor nutritionist 4 software (FPN4) was used to determine the amount of macronutrient consumption (protein, fat, and carbohydrates). Subjects were asked to consume the same food as the pre-test, the day before blood sampling in the post-test. In general, the subjects' diet consisted of 15% protein, 55% carbohydrates, and 30% fat.

In this study, 50000 IU of vitamin D (made by the Zahravi Pharmaceutical Company in Iran) were supplemented once per week for eight weeks in the Vit D groups [29, 30]. While, the C and AT received the placebo with the same shape, color, smell, and tastes as vitamin D supplement once per week (made by the Zahravi Pharmaceutical Company, Iran) for 8 weeks.

Body Composition and Anthropometric Measurements

Subjects were familiarized with the study procedure 3 days before the start of the intervention and primary measurements including anthropometric parameters and body composition were determined both at the start and end of the study. Waist circumference was measured to the nearest 0.5 cm with a non-elastic tape measure and Height to the nearest 0.5 cm using a stadiometer (DETECTO, Model 3PHTROD-WM, USA). Also, weight, fat mass, and BMI were determined by the INBODY test using bioelectric impedance analysis (Zeus 9.9 PLUS; Jawon Medical Co., Ltd., Kungsang Bukdo, South Korea). The body composition measurements were performed early in the morning (8–9 am) after at least 12 hours of fasting overnight. To minimize the effect of water consumption on the results, measurements were done after emptying the bladder, and subjects were asked to refrain from taking diuretic drugs and not to participate in intensive physical activity 48 hours before the test.

Outcome Variables and Biochemical Measurements

Subjects did not perform any exercise 48 hours before blood sampling and were on 12 hours of fasting before blood sampling. Then, 10 ml of blood samples were taken from the cubital vein, in the least possible time, at the beginning and the end of the study. For BDNF and NGF analyses Blood samples were collected in heparinized tubes, frozen, and stored at -80°C. Then, BDNF and NGF were measured using a human BDNF and NGF kit (made in China under the license of Boster Biological Technology, America Company with a sensitivity of 2 and 2.48 pg/ml, respectively) by the ELISA method. Serum 25-hydroxyvitamin D level was measured by quantitative luminescence method (DiaSorin kit made in the USA).

Statistical analysis

To evaluate the normality of distribution the Shapiro–Wilk’s test was used. ANOVA and t-test were used to compare the mean of variables between and within groups, respectively. If significant differences were found Tukey’s post hoc test was used. The SPSS statistical software (version 21; SPSS Inc., Chicago, IL, USA) was used to perform all statistical analyses at a significant level of $P < 0.05$.

Results

The findings on the anthropometric indices and their between-group comparison are presented in Table 2. Based on the results of the t-test, there were significant differences in the mean of BW, BMI, BFP, WHR, and Vit D in the post-test compared to the pre-test. After eight weeks, BW, BMI, BFP, and WHR significantly decreased in all intervention groups, while in the C, these variables increased significantly (Table 2). Also, after eight weeks Vit D increased significantly in all intervention groups, except C.

The results of one-way ANOVA showed no significant difference in BW, BMI, BFP, and WHR and Vit D between the groups in the pre-test, however, in the post-test, there was a significant difference in the above variables between the groups.

The results of Tukey’s post hoc test show that BW, BMI, BFP, and WHR, and Vit D were significant in AT + Vit D, AT, and Vit D compared to C. No significant difference in the BW, BFP, and WHR was observed between other groups. The results indicate no significant difference in Vitamin D levels in AT + Vit D compare to AT alone; Also, no significant difference was found in Vitamin D levels in AT compared to Vit D (Table 2).

Table 2
Mean $\pm$ SD of anthropometric indices before the intervention among the groups

Variables	AT + Vit D (n = 10)	AT (n = 9)	Vit D (n = 9)	C (n = 10)	P-Value ^a
Age (years)	27.70 $\pm$ 2.68	26.77 $\pm$ 2.27	25.44 $\pm$ 2.29	28.11 $\pm$ 3.62	0.563
Height	163.40 $\pm$ 2.63	166.03 $\pm$ 2.65	162.33 $\pm$ 1.50	164.20 $\pm$ 1.54	0.467
Body Weight (Kg)					
Before	71.90 $\pm$ 1.91	73.12 $\pm$ 1.93	71.11 $\pm$ 2.02	73.01 $\pm$ 1.33	0.086
After	68.11 $\pm$ 1.15 β	70.16 $\pm$ 1.22 β	69.33 $\pm$ 1.50 β	75.06 $\pm$ 1.15	0.001 $\yen$
P Value ^b	0.001*	0.001*	0.012*	0.004*	
BMI (kg/m ²)					
Before	26.94 $\pm$ 0.93	26.33 $\pm$ 1.07	26.98 $\pm$ 0.88	27.08 $\pm$ 0.66	0.279
After	25.48 $\pm$ 0.79 β	25.14 $\pm$ 0.99 β	26.32 $\pm$ 0.93 β	27.82 $\pm$ 0.71	0.001 $\yen$
P Value ^b	0.001*	0.001*	0.011*	0.004*	
Body Fat Percent (%)					
Before	28.37 $\pm$ 1.63	28.11 $\pm$ 1.26	27.66 $\pm$ 1.58	28.60 $\pm$ 1.50	0.213
After	24.90 $\pm$ 1.37 β	25.77 $\pm$ 0.83 β	26.55 $\pm$ 1.81 β	30.04 $\pm$ 1.33	0.001 $\yen$
P Value ^b	0.002*	0.002*	0.048*	0.003*	
WHR					
Before	0.84 $\pm$ 0.03	0.84 $\pm$ 0.01	0.83 $\pm$ 0.03	0.84 $\pm$ 0.02	0.150
After	0.81 $\pm$ 0.02 β	0.82 $\pm$ 0.03 β	0.82 $\pm$ 0.02 β	0.86 $\pm$ 0.01	0.001 $\yen$
P Value ^b	0.001*	0.001*	0.001*	0.002*	
Vitamin D (ng/mL)					
Before	25.80 $\pm$ 1.81	26.55 $\pm$ 1.50	26.44 $\pm$ 1.42	27.20 $\pm$ 3.45	0.588
After	35.60 $\pm$ 1.89 $\mu\beta$	29.66 $\pm$ 1.65 β	32.88 $\pm$ 1.90 β	25.10 $\pm$ 1.44	0.001 $\yen$

Variables	AT + Vit D (n = 10)	AT (n = 9)	Vit D (n = 9)	C (n = 10)	P-Value ^a
P Value ^b	0.001*	0.002*	0.001*	0.111	
AT + TS , Aerobic training + Turmeric supplement, AT , Aerobic training group; TS , Turmeric supplement group; C , the control group					
p values superscript with "a" is calculated using one-way analysis of variance test followed by post hoc Tukey's test; p values superscript with "b" is calculated using paired t-test					
*: Significantly different comparing pre and post-test within the groups					
¥: Significantly different comparing pre and post-test between groups					
μ: Significantly different between AT + TS compare to the AT group					
€: Significantly different between AT + TS compare to the TS group					
β: Significantly different between AT + TS, AT, and TS groups compare to the C group					

BDNF and NGF levels improved significantly from pre-test to post-test within all experimental groups (as detailed in Figs. 1 and 2). The results of one-way ANOVA showed no significant difference in BDNF and NGF between the groups in the pre-test, however in the post-test, there were significant differences in the above variables between the groups. BDNF and NGF decreased significantly in AT + Vit D, AT, and Vit D compared to C. Also, the results show that the AT + Vit D had significantly lower BDNF and NGF compared to AT and Vit D. Also, the results of this study showed significant differences between AT and Vit D groups in the BDNF and NGF levels.

Discussion

According to the results of the present study, eight weeks of aerobic training with vitamin D supplementation reduced anthropometric indices (body weight, body mass index, and fat percentage) and increased vitamin D levels significantly in women with MS. Although these changes were greater in the AT + Vit D compared with the AT or Vit D alone, only the vitamin D levels were significantly different between the AT + Vit D and AT. Interestingly, the anthropometric indices increased while vitamin D levels decreased significantly in the control group. Few studies have been done on the effects of aerobic training and vitamin D supplementation in MS patients, and the evidence for the preventive and therapeutic effects is largely dependent on epidemiological studies. However, our knowledge of the interactive effects of aerobic training and vitamin D supplementation on anthropometric indices is still limited.

Consistently, Babaei et al. (2014), [31] Hoseini et al. (2017), [32], and Hoseini et al. (2020) [33] reported that regular AT improves metabolic health parameters through increased daily energy intake, improved

and increased fat oxidation in skeletal muscles [34, 35] Also, based on the studies the prevalence of demyelinating diseases (e.g. MS) [36], metabolic syndrome, and chronic disease have been reported to be associated with Vit D deficiency [37, 38]. Ascherio et al. (2014) introduced Vitamin D as an early predictor of multiple sclerosis activity and progression [39]. Hosseini et al. (2017) reported improved metabolic indices following 8 weeks of high dose Vit D supplementation [32] The underlying mechanism might be increased parathyroid hormone (PTH) induced by hypovitaminosis D which leads to increased intracellular Ca^{++} . This, in turn, upregulates the insulin receptors and Glut-4 channel activity in the target tissues. Thus, vitamin D plays an important role in insulin function, glucose metabolism, and other metabolic processes [31, 40]. The findings of the present study showed the highest levels of BDNF, and NGF in the AT + Vit D after 8 weeks; Therefore, AT + Vit D had been more effective in improving BDNF, and NGF compared to Vit D or AT alone. The decrease in BDNF and NGF levels in the control group is probably due to the MS-induced Schwann cell demyelination. While BDNF and NGF levels increased significantly in other groups. According to the results of studies, deficiency of neurotrophic factor/receptor is involved in the progression of MS [41]. The increased gene and protein expression of proinflammatory factors, such as interferon-gamma ($\text{IFN-}\gamma$) and tumor necrosis factor-alpha ($\text{TNF-}\alpha$), have been observed with the progression of MS [42]. In turn, MAPK activates exon IV (inhibitor of BDNF and NGF protein synthesis) by CAMP receptor phosphorylation, which is a known factor in the regulation of BDNF and NGF expression [43, 44]. Various studies have shown that regular aerobic training increases the regeneration of sensory neurons after axon injury and stimulates gene expression of proteins required for axon growth and regeneration in MS patients [45]. Some of the beneficial effects of increased BDNF and NGF include direct stimulation, migration, and differentiation of oligodendrocytes (which are essential steps in myelinogenesis), upregulation of the activity and expression of cells necessary for myelin production (e.g., the main structural components of myelin, such as astrocytes and neurons), and regulatory proteins [46]. However, studies have reported that gene expression and secretion of neurodevelopmental factors, such as BDNF and NFG, function multiple molecular and non-molecular mechanisms [47, 48]; Aerobic training might affect BDNF and NFG possibly through increased activity of some signaling pathways (PERK, pAkt, and CaMKII), increased expression of growth proteins such as 43-GAP, altered concentrations of hormones such as norepinephrine and serotonin, as well as the secretion of heat shock proteins, HSP27 and HSP70 [49, 50]. In addition, vitamin D might affect the BDNF and NGF secretion and expression through nuclear signaling pathways secondary to the interaction of 1–25 (OH) 2D to vitamin D receptor (VDR) in the nuclear; which begins a genetic response after reacting with RXR and producing Vitamin D response element (VDRE) [51, 52]. VDRE binds to nuclear receptor coactivators (COACs) or corepressors (COREs) that increase BDNF and NGF gene expression [53]. Also, studies in MS patients shows increased free radical production, including nitric oxide that might exacerbate inflammatory reactions and brain lesions consequently due to the disrupted nicotinamide adenine dinucleotide phosphate (NADP^+) system in MS patients [54, 55]. Therefore, it is hypothesized that adequate vitamin D levels might help prevent the MS progression by inhibiting nitric oxide production [56].

Conclusions

In conclusion, despite the efficacy of aerobic training and vitamin D supplementation alone on the measured variables in MS patients, the combined aerobic training and vitamin D supplementation seems to induce greater improvements in BDNF and NGF levels; However, the mechanism of interactional effect is still not well understood. On one hand, due to the general recommendations for the application of non-pharmacologic therapies in the MS management, prevention of inactivity associated diseases, and on the other hand the anti-inflammatory and antioxidant effects of vitamin D supplementation during the outbreak COVID-19, combined aerobic training and vitamin D supplementation is recommended as an eligible approach in women with MS under the prescription and supervision of PCP or neurologists. However, further extensive studies are recommended.

Abbreviations

MS: Multiple sclerosis ; BDNF: Brain-derived neurotrophic factor; NGF: Nerve growth factor; AT; Aerobic training; Vit D: Vitamin D; RPE: Rating of perceived exertion; FPN4: Food Processor nutritionist 4 software; BW: Bodyweight, BMI: Body mass index, BFP: Body fat percentage; WHR: Waist–hip ratio; PTH: Parathyroid hormone; IFN- γ ; Interferon-gamma; TNF- α : Tumor necrosis factor-alpha; RXR: Retinoid X receptor; VDRE: vitamin D response element; nuclear receptor coactivators; NCOAs; Nuclear receptor coactivators; NCORs: Corepressors; NADP⁺: Nicotinamide adenine dinucleotide phosphate.

Declarations

Acknowledgments

The authors would like to thank the subjects for their willing participation in this study.

Authors' contributions

RH designed the study. EB experimented. RH analyzed the data and wrote the manuscript. EA involved in the interpretation of data, reviewed and edited the manuscript. All authors read and approved the final manuscript.

Funding

The author(s) received no specific funding for this work.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

This study was approved by the ethics committee of Razi University, Kermanshah, IRAN (IR.RAZI.REC.1400.002), conformed with the ethical standards of the Helsinki Declaration and registered in the Iranian Clinical Trial Registration Center under the code IRCT20201129049525N1. The written informed consent was obtained from all participants according to the protocols approved by the ethics committee of Razi University. Also, the fusibility to leave the study at any stage was guaranteed.

Consent for publication

Not applicable.

Competing interests

Authors do not have any competing interests

References

1. Bansi J, Bloch W, Gamper U, Kesselring J. Training in MS: influence of two different endurance training protocols (aquatic versus overland) on cytokine and neurotrophin concentrations during three week randomized controlled trial. *Multiple Sclerosis Journal*. 2013;19 5:613-21.
2. Mao P, Reddy PH. Is multiple sclerosis a mitochondrial disease? *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2010;1802 1:66-79.
3. Ransohoff RM, Hafler DA, Lucchinetti CF. Multiple sclerosis—a quiet revolution. *Nature Reviews Neurology*. 2015;11 3:134.
4. Ghadampour E, Radmehr P, Yousefvand L. The effectiveness of acceptance and commitment based treatment on social competence and life expectancy of patients with multiple sclerosis. *Armaghane danesh*. 2017;21 11:1100-14.
5. Sahraian MA, Khorramnia S, Ebrahim MM, Moinfar Z, Lotfi J, Pakdaman H. Multiple sclerosis in Iran: a demographic study of 8,000 patients and changes over time. *European neurology*. 2010;64 6:331-6.
6. Tofighi A, Saki Y, Razmjoo K. Effect of 12-week progressive resistance training on balance, fatigue and disability in women with MS. *Jundishapur Scientific Medical Journal*. 2013;12 2:159-67.
7. Jain RV, KP SG. BIOMARKERS, DRUG TREATMENT AND EXPERIMENTAL MODELS OF MULTIPLE SCLEROSIS. *Multiple sclerosis (MS)*. 2020;23:09.
8. Mackay CP, Kuys SS, Brauer SG. The effect of aerobic exercise on brain-derived neurotrophic factor in people with neurological disorders: a systematic review and meta-analysis. *Neural plasticity*. 2017;2017.
9. Rosenberg AJ, Wee SO, Shroeder EC, Grigoriadis G, Sandroff BM, Bunsawat K, et al. The effects of acute aerobic exercise on cerebral blood flow and cognition in persons with multiple sclerosis. *The FASEB Journal*. 2017;31:842.4-4.
10. White LJ, Castellano V. Exercise and brain health—implications for multiple sclerosis. *Sports medicine*. 2008;38 2:91-100.

11. Connor B, Dragunow M. The role of neuronal growth factors in neurodegenerative disorders of the human brain. *Brain Research Reviews*. 1998;27 1:1-39.
12. Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends in neurosciences*. 2002;25 6:295-301.
13. Gold SM, Schulz K-H, Hartmann S, Mladek M, Lang UE, Hellweg R, et al. Basal serum levels and reactivity of nerve growth factor and brain-derived neurotrophic factor to standardized acute exercise in multiple sclerosis and controls. *Journal of neuroimmunology*. 2003;138 1-2:99-105.
14. Romberg A, Virtanen A, Ruutiainen J, Aunola S, Karppi S-L, Vaara M, et al. Effects of a 6-month exercise program on patients with multiple sclerosis: a randomized study. *Neurology*. 2004;63 11:2034-8.
15. Beecham AH, Patsopoulos NA, Xifara DK, Davis MF, Kempainen A, Cotsapas C, et al. Analysis of immune-related loci identifies 48 new susceptibility variants for multiple sclerosis. *Nature genetics*. 2013;45 11:1353.
16. Munger KL, Levin LI, Hollis BW, Howard NS, Ascherio A. Serum 25-hydroxyvitamin D levels and risk of multiple sclerosis. *Jama*. 2006;296 23:2832-8.
17. Brown SJ. The role of vitamin D in multiple sclerosis. *Annals of Pharmacotherapy*. 2006;40 6:1158-61.
18. Hosseini S, Rakhshi N, Abdullah N, Jafari S, Nikraves A. Vitamin D, the healthy and inflamed Immune System in multiple sclerosis. *Journal of North Khorasan University of Medical Sciences*. 2014;5 5:935-41.
19. Deluca HF, Cantorna MT. Vitamin D: Its role and uses in immunology 1. *The FASEB journal*. 2001;15 14:2579-85.
20. Polachini CRN, Spanevello RM, Zanini D, Baldissarelli J, Pereira LB, Schetinger MRC, et al. Evaluation of delta-aminolevulinic dehydratase activity, oxidative stress biomarkers, and vitamin D levels in patients with multiple sclerosis. *Neurotoxicity research*. 2016;29 2:230-42.
21. Hejazi E, Amani R, SharafodinZadeh N, Cheraghian B. Comparison of antioxidant status and vitamin D levels between multiple sclerosis patients and healthy matched subjects. *Multiple sclerosis international*. 2014;2014.
22. MOSAYEBI G, Ghazavi A, Payani M. The effect of vitamin D3 on the inhibition of experimental autoimmune encephalomyelitis in C57BL/6 mice. 2006.
23. Pierrot-Deseilligny C, Souberbielle J-C. Contribution of vitamin D insufficiency to the pathogenesis of multiple sclerosis. *Therapeutic advances in neurological disorders*. 2013;6 2:81-116.
24. Nabavi SM, Sabet Z, Morsalee D, Aminzadeh M. Considering the Safety Issues Vitamin D3 Supplementation in Relapsing-Remitting Multiple Sclerosis. *Clinical Toxicology*. 2012;3 2.
25. Monazamnezhad A, Habibi A, Majdinasab N, Ghalvand A. The effects of aerobic exercise on lipid profile and body composition in women with multiple sclerosis. *Jundishapur Journal of Chronic Disease Care*. 2015;4 1.

26. Rae-Grant A, Day GS, Marrie RA, Rabinstein A, Cree BA, Gronseth GS, et al. Practice guideline recommendations summary: Disease-modifying therapies for adults with multiple sclerosis: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;90 17:777-88.
27. Branco BHM, de Oliveira Mendes F, Ladeia GF, Bertolini SMMG, Badilla PV, Andreato LV. Maximum heart rate predicted by formulas versus values obtained in graded exercise tests in Brazilian jiu-jitsu athletes. *Sport Sciences for Health*. 2020;16 1:39-45.
28. Kalb R, Brown TR, Coote S, Costello K, Dalgas U, Garmon E, et al. Exercise and lifestyle physical activity recommendations for people with multiple sclerosis throughout the disease course. *Multiple Sclerosis Journal*. 2020;26 12:1459-69.
29. VanAmerongen B, Dijkstra C, Lips P, Polman C. Multiple sclerosis and vitamin D: an update. *European journal of clinical nutrition*. 2004;58 8:1095-109.
30. Tuyserkani H, Rezaei M. Role of Vitamins D and A in Immune System of Multiple Sclerosis Patients: An Updated Mini-Review. *Journal of Reviews in Medical Sciences*. 2020;1 1:45-8.
31. Babaei P, Damirchi A, Hoseini R. The interaction effects of aerobic exercise training and vitamin D supplementation on plasma lipid profiles and insulin resistance in ovariectomized rats. *Journal of exercise nutrition & biochemistry*. 2015;19 3:173.
32. Hoseini R, Damirchi A, Babaei P. Vitamin D increases PPAR γ expression and promotes beneficial effects of physical activity in metabolic syndrome. *Nutrition*. 2017;36:54-9.
33. Hoseini Z, Behpour N, Hoseini R. Co-treatment with Vitamin D Supplementation and Aerobic Training in Elderly Women with Vit D Deficiency and NAFLD: A Single-blind Controlled Trial. *Hepatitis Monthly*. 2020;20 2.
34. Smart N, King N, McFarlane JR, Graham P, Dieberg G. Effect of exercise training on liver function in adults who are overweight or exhibit fatty liver disease: a systematic review and meta-analysis. *British journal of sports medicine*. 2018;52 13:834-43.
35. Van Gemert WA, May AM, Schuit AJ, Oosterhof BY, Peeters PH, Monninkhof EM. Effect of weight loss with or without exercise on inflammatory markers and adipokines in postmenopausal women: the SHAPE-2 trial, a randomized controlled trial. *Cancer Epidemiology and Prevention Biomarkers*. 2016;25 5:799-806.
36. Burton JM, Costello FE. Vitamin D in multiple sclerosis and central nervous system demyelinating disease—a review. *Journal of Neuro-ophthalmology*. 2015;35 2:194-200.
37. Nakhl S, Sleilaty G, El Samad S, Saliba Y, Chahine R, Farès N. Association between vitamin D deficiency and lipid and non-lipid markers of cardiovascular diseases in the middle east region. *European journal of clinical nutrition*. 2019;73 6:850-8.
38. Mogili KD, Karuppusami R, Thomas S, Chandy A, Kamath MS, Aleyamma T. Prevalence of vitamin D deficiency in infertile women with polycystic ovarian syndrome and its association with metabolic syndrome—A prospective observational study. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2018;229:15-9.

39. Ascherio A, Munger KL, White R, Köchert K, Simon KC, Polman CH, et al. Vitamin D as an early predictor of multiple sclerosis activity and progression. *JAMA neurology*. 2014;71 3:306-14.
40. Barker T, Martins TB, Hill HR, Kjeldsberg CR, Dixon BM, Schneider ED, et al. Circulating pro-inflammatory cytokines are elevated and peak power output correlates with 25-hydroxyvitamin D in vitamin D insufficient adults. *European journal of applied physiology*. 2013;113 6:1523-34.
41. Eyileten C, Sharif L, Wicik Z, Jakubik D, Jarosz-Popek J, Soplinska A, et al. The relation of the brain-derived neurotrophic factor with microRNAs in neurodegenerative diseases and ischemic stroke. *Molecular neurobiology*. 2021;58 1:329-47.
42. Farina G, Magliozzi R, Pitteri M, Reynolds R, Rossi S, Gajofatto A, et al. Increased cortical lesion load and intrathecal inflammation is associated with oligoclonal bands in multiple sclerosis patients: a combined CSF and MRI study. *Journal of neuroinflammation*. 2017;14 1:1-11.
43. Jeon SJ, Rhee SY, Seo JE, Bak HR, Lee SH, Ryu JH, et al. Oroxylin A increases BDNF production by activation of MAPK–CREB pathway in rat primary cortical neuronal culture. *Neuroscience research*. 2011;69 3:214-22.
44. Cao C, Xiao J, Liu M, Ge Z, Huang R, Qi M, et al. Active components, derived from Kai-xin-san, a herbal formula, increase the expressions of neurotrophic factor NGF and BDNF on mouse astrocyte primary cultures via cAMP-dependent signaling pathway. *Journal of ethnopharmacology*. 2018;224:554-62.
45. Molteni R, Zheng J-Q, Ying Z, Gómez-Pinilla F, Twiss JL. Voluntary exercise increases axonal regeneration from sensory neurons. *Proceedings of the National Academy of Sciences*. 2004;101 22:8473-8.
46. Linker RA, Lee D-H, Demir S, Wiese S, Kruse N, Siglienti I, et al. Functional role of brain-derived neurotrophic factor in neuroprotective autoimmunity: therapeutic implications in a model of multiple sclerosis. *Brain*. 2010;133 8:2248-63.
47. Khademosharie M, Tadibi V, Behpoor N, Hamedinia MR. The effect of 12-weeks resistance and endurance training on the serum levels NGF, BDNF, and VDBP in women with multiple sclerosis. *International Journal of Applied Exercise Physiology*. 2018;7 1:76-86.
48. Mondal AC, Fatima M. Direct and indirect evidences of BDNF and NGF as key modulators in depression: role of antidepressants treatment. *International Journal of Neuroscience*. 2019;129 3:283-96.
49. Currie J, Ramsbottom R, Ludlow H, Nevill A, Gilder M. Cardio-respiratory fitness, habitual physical activity and serum brain derived neurotrophic factor (BDNF) in men and women. *Neuroscience letters*. 2009;451 2:152-5.
50. Santoso D, Yolanda S, Redjeki S, Andraini T, Ivanali K. Continuous environmental enrichment and aerobic exercise improves spatial memory: focus on rat hippocampal BDNF and NGF. *Comparative Exercise Physiology*. 2020;16 2:121-8.
51. Geng C, Shaikh AS, Han W, Chen D, Guo Y, Jiang P. Vitamin D and depression: mechanisms, determination and application. *Asia Pacific journal of clinical nutrition*. 2019;28 4:689-94.

52. Eyles DW, Burne TH, McGrath JJ. Vitamin D, effects on brain development, adult brain function and the links between low levels of vitamin D and neuropsychiatric disease. *Frontiers in neuroendocrinology*. 2013;34 1:47-64.
53. Medhat E, Rashed L, Abdelgwad M, Aboulhoda BE, Khalifa MM, El-Din SS. Exercise enhances the effectiveness of vitamin D therapy in rats with Alzheimer's disease: emphasis on oxidative stress and inflammation. *Metabolic brain disease*. 2020;35 1:111-20.
54. García-Ortiz A, Serrador JM. Nitric oxide signaling in T cell-mediated immunity. *Trends in molecular medicine*. 2018;24 4:412-27.
55. Miller ED, Dziedzic A, Saluk-Bijak J, Bijak M. A review of various antioxidant compounds and their potential utility as complementary therapy in multiple sclerosis. *Nutrients*. 2019;11 7:1528.
56. Sintzel MB, Rametta M, Reder AT. Vitamin D and multiple sclerosis: a comprehensive review. *Neurology and therapy*. 2018;7 1:59-85.

Figures

Figure 1

Flow chart of the study population AT+Vit D, Aerobic training + vitamin D supplement, AT, Aerobic training group; Vit D, vitamin D supplement; C, the control group

Figure 2

Comparison between mean $\pm$ SD of BDNF between groups AT+Vit D, Aerobic training + vitamin D supplement, AT, Aerobic training group; Vit D, vitamin D supplement; C, the control group p values superscript with "a" is calculated using one-way analysis of variance test followed by post hoc Tukey's test; p values superscript with "b" is calculated using paired t-test *: Significantly different comparing pre and post-test within the groups μ : Significantly different between AT+Vit D compare to the AT group ϵ : Significantly different between AT+Vit D compare to the Vit D group α : Significantly different between AT compare to the Vit D group. β : Significantly different between AT+Vit D, AT, and Vit D groups compare to the C group

Figure 3

Comparison between mean $\pm$ SD of NGF between groups AT+Vit D, Aerobic training + vitamin D supplement, AT, Aerobic training group; Vit D, vitamin D supplement; C, the control group p values superscript with "a" is calculated using one-way analysis of variance test followed by post hoc Tukey's test; p values superscript with "b" is calculated using paired t-test *: Significantly different comparing pre and post-test within the groups μ : Significantly different between AT+Vit D compare to the AT group ϵ : Significantly different between AT+Vit D compare to the Vit D group α : Significantly different between AT compare to the Vit D group. β : Significantly different between AT+Vit D, AT, and Vit D groups compare to the C group