Variation of Actual Corn (Zea Mays L.) Evapotranspiration, Single and Dual Crop Coefficient Under Different Point Source Irrigation Systems in a Semiarid Region

Elahe Kanani

Bu-Ali Sina University

Hossein Dehghanisanij (✉ dehghanisanij@yahoo.com)

Agricultural Engineering Research Institute <https://orcid.org/0000-0002-5617-9375>

Samira Akhavan

Bu-Ali Sina University

Research Article

Keywords: Basal crop coefficient, Soil evaporation, Lysimeter, Point source irrigation, Porous capsule

Posted Date: July 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-682889/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Theoretical and Applied Climatology on January 26th, 2022. See the published version at <https://doi.org/10.1007/s00704-022-03932-w>.

Variation of actual corn (*Zea mays* L.) evapotranspiration, single and dual crop coefficient under different point source irrigation systems in a semiarid region

Elahe Kanani ¹, Hossein Dehghanisanij ^{2*}, Samira Akhavan¹

¹ Bu-Ali Sina University, faculty of agriculture, Dept. of Water Engineering, Hamedan, Iran. P.O. Box 65178-33131

² Agricultural Engineering Research Institute, Agricultural Research, Education and Extension Organization, Karaj, Alborz, Iran. P.O. Box 31585-845

*Corresponding author: Hossein Dehghanisanij, Agricultural Engineering Research Institute, Agricultural Research, Education and Extension Organization, Karaj, Alborz, Iran. P.O. Box 31585-845, Email: h.dehghanisanij@areeo.ac.ir

Abstract

Paying attention to irrigation of corn could have significant attention on water resources. The experiment was carried out to consider the influence of porous capsule irrigation (*PCI*), surface drip irrigation (*DI*) and subsurface drip irrigation (*SDI*) systems under mulch condition on actual evapotranspiration ($ET_{c\ act}$), crop coefficients ($K_{c\ single}$ and $K_{c\ dual}$), yield and water use efficiency (*WUE*) of corn in a semi-arid region of Iran. The experiment was arranged in a split-plot design with three irrigation systems as the main-factor and two soil coverage with mulch (M_1) and non-mulch (M_2) as sub- factor. The results showed that corn $ET_{c\ act}$ varied significantly with different irrigation systems ($P<0.05$), and was reported 389.8, 377.0 and 372.8 mm for *PCI*, *DI* and *SDI* systems, respectively. The highest value of $K_{c\ average}$ and K_{cb} (0.82 and 1.22, respectively) and the minimum value of K_e (0.12) were seen in *PCI* system. The dry and wet biological yield were the highest in (*PCI* + M_1) treatment (29.98 and 107 ton/ ha, respectively), and the lowest dry and wet biological yield were recorded in (*DI*+ M_2) treatment (23.19 and 58.54 ton/ha, respectively). The highest *WUE* (7.89 kg/m³) were recorded in (*PCI* + M_1) treatment. Using *PCI* system leads to higher biological yield, *WUE* and *IWUE* in comparison to *DI* and *SDI* systems. Accordingly, *PCI* system as an alternative to drip irrigation system can be a viable option for water scarce area particularly for smallholder farmers.

Keywords: Basal crop coefficient, Soil evaporation, Lysimeter, Point source irrigation, Porous capsule

1. Introduction

Agriculture section is the main consumer in arid and semiarid countries. This is the first source of exploitation to compete for water (Postel 1992; Ayars et al. 2015). Evapotranspiration (ET_c) loss is one of the major challenges of irrigated agriculture. Evaporation occurs predominantly from the exposed soil fraction in the field and free water surface (Allen et al., 1998). Recently, mulching is a recommended for water management practice to reduce water loss due to evaporation from the soil surface and to create more favourable condition for plant transpiration (Tarara 2000; Tiwari et al. 2014, Reddy et al. 2018). Improving irrigation management methods under mulching conditions is a new agricultural water-saving irrigation technology, which can be effective to prevent water loss due to evaporation from the soil surface. In recent years, drip irrigation under mulch condition has been widely used in arid and semiarid regions (Qin et al. 2016). The effect of drip irrigation method under mulch makes soil temperature changeable, reduces the amount of water loss due to decrease soil evaporation, maintains soil moisture and improves soil fertility, prevents the growth of weeds and also helps to evenly distribute moisture to reduce plant stress (Singh 2005; Sharma et al. 2010; Paul et al. 2013; Panigrahi et al. 2016). A large number of researchers recorded that an increase of crop yield and water use efficiency are improved by application drip irrigation with mulch in different agro-climatic region and soil condition (Biswas et al. 2015; Thentu et al. 2016; Liu et al. 2017; Zhang et al. 2017). But there is gap between the research sites and the actual farm conditions. Moreover our knowledge about some traditional irrigation system and its relation with crop development is limited. Drip irrigation systems are well known, and researchers showed that intensive irrigation management of drip irrigation increased water use efficiency significantly. There is a traditional irrigation system in arid and semiarid regions with same specification while are not well studied and promoted (Bainbridge et al. 2008). Porous capsule irrigation is an effective promoted adaptation of buried clay pot irrigation (Silva et al. 1981a, b; Bainbridge 2002; Bainbridge et al. 2008). Supply of water with small porous capsules is one of the methods that can be used in small scale farms in arid and semi-arid regions. Porous capsule irrigation method introduced for planting of a wide range of annual and perennial plants including: horticultural crops (Setiawan et al. 1998), corn (Bainbridge 2001), watermelons (Soomro 2002), tomatoes (Tesfaye et al. 2012), turnip, okra and eggplant (Siyal et al. 2011). Crop coefficient (K_c) is slightly affected by climate, but significantly affected by crop and Irrigation scheduling or environmental conditions, that can lead to water stress conditions for crop (Zhang et al. 2013). The K_c can be applied as single crop coefficient ($K_{c\ single}$) which is influenced by effect of evaporation (E) and plant transpiration (T_p) together and dual crop coefficient ($K_{c\ dual}$) that is expressed by soil evaporation coefficient (K_e) and

basal crop coefficient (K_{cb}), separately (Allen et al. 1998). The $K_{c\ dual}$ can provide a more accurate assessment of the effects of soil moisture by rainfall or irrigation and also the effects of using mulches to control soil evaporation. Hence, to some extent, the $K_{c\ dual}$ can improve the accuracy of ET_c estimation when used with appropriate calibrated base data (Allen et al. 2011; Rosa et al. 2012; Zhang et al. 2013). Corn is one of the main cereal crop in the world and it ranks third in cereal production in cultivated area and production (FAO 2011). Accordingly paying attention to irrigation of corn could have significant attention on water resources. To improve our ability on corn (*Zea mays L.*) irrigation management under different localized irrigation systems in arid and semiarid condition, this study aimed to investigate variation of evaporation and transpiration during growth stages, soil evaporation coefficient, and basal crop coefficient in different irrigation systems and climate condition.

2. Materials and methods

2.1. Experimental site

The experiment was conducted at the farm of Agricultural Engineering Research Institute, Karaj, Iran. The site was located at 35°46' N latitude, 50°55' E longitude and at an elevation of 1260 m above sea level. The local climate is semiarid, with mean annual precipitation of 279.3 mm, mean annual temperature of 16.6 °C, cold and dry winters, and hot and humid summers. The daily climate data for growing season (August to November 2014) were collected from a synoptic meteorology station located in 5 km far from the farm. Table 1 summarizes the monthly average climatic data for the corn growing seasons.

(Table. 1 here)

The seasonal average air temperature was 24°C during corn growing season, however, air temperature were hotter in August, and cooler in November. On average, wind speed and relative humidity were higher in August and November, respectively. The average solar radiation, however was higher in August.

The soil samples were taken from 0-10, 10-20 and 20-30 cm soil depths. The soil was loam with a mean bulk density of 1.42 gr/cm³ and slightly alkaline with a pH of 7.9. Soil characteristics and irrigation water for experimental site are given in Tables 2 and 3.

(Table. 2 here)

(Table. 3 here)

In order to monitor water consumption by corn under different irrigation systems, 21 volumetric lysimeters were applied in the soil of a corn farm with an area of 18 ha, that was filled with excavated soil. The lysimeters were in

cylindrical shape with diameter of 40 cm and depth of 70 cm and is considered as a mini-lysimeter (Dugas and Bland 1989; Kong et al. 2012). Inside each mini lysimeter, three forage corn seeds (Single Cross 704) were planted with 13 cm seeds spacing on 6th August 2014 (Fig. 1).

(Fig. 1 here)

2.2. Irrigation management

The corn was irrigated with porous capsule irrigation (*PCI*), surface drip irrigation (*DI*) and subsurface drip irrigation (*SDI*) system. The *DI* and *SDI* system were equipped with 40 cm emitter spacing and discharge of 4 L/h, and also *SDI* system was buried 30 cm below the soil surface. In this design, porous capsule buried at 30 cm soil depth. Each porous capsule has a length of 30 cm and with a diameter of 6.5 cm as shown in Fig. 2. Irrigation interval was twice a week for *DI* and *SDI* system and daily for *PCI* system.

(Fig. 2 here)

The daily irrigation requirement was estimated using FAO-Penman-Monteith model (Allen et al. 1998; Eqs. (1) and (2)) confirmed for Karaj region by Dehghanisanij et al. (2004) and Akhavan et al. (2018).

$$ET_0 = \frac{0.408 \Delta (R_n - G) + \gamma [890 (T + 273)] U_2 (e_s - e_a)}{\Delta + \gamma (1 + 0.34 u_2)} \quad (1)$$

$$ET_c = ET_0 \times K_c \quad (2)$$

Where ET_c is crop evapotranspiration; ET_0 is reference evapotranspiration (mm/day); R_n is net radiation (MJ/m²/day); G is soil heat flux density (MJ/m²/day); T is mean temperature (°C); U_2 is wind speed at 2 m height (m/s); γ is psychrometric constant (k Pa/°C); Δ is slope vapor pressure curve (k Pa/°C); e_a is actual vapor pressure (k Pa); e_s is saturation vapor pressure (k Pa). In this study, corn K_c is crop coefficient recommended for Karaj by Farshi et al. (1997).

Fertilizer injected to the irrigation water, and was started from the stage of 3 and 4 leafed of corn growth to 45 days before harvesting. The amount of 250 kg/ha ammonium phosphate fertilizer and 200 kg/ha nitrogen fertilizer was delivered to the crop.

2.3. Experimental design

The experiment was laid out in a split plot design keeping three irrigations methods (i) *PCI*, (ii) *DI* and (iii) *SDI* system as main- factor and two soil coverage (i) with mulch (M_1) and (ii) non- mulch (M_2) as sub-factor. Each treatment was replicated three times in a randomized complete block design.

2.4. Measurements

2.4.1. Actual corn evapotranspiration ($ET_{c\ act}$)

Daily actual corn evapotranspiration ($ET_{c\ act}$) of each mini-lysimeter was calculated by applying the water balance method using the Eqs. (3) and (4) (Allen et al., 1998):

$$ET_{c\ act} = P + I - D - R - \Delta S \quad (3)$$

$$\Delta S = S_t - S_{t-1} \quad (4)$$

Where P is the rain (mm); I is the irrigation depth (mm); D is the water loss through drainage from the mini-lysimeter (mm); R is the runoff (mm); and ΔS is the change of soil water storage in the mini- lysimeter (mm), S_t and S_{t-1} are the amount of water in the root zone at the beginning and end of the period (mm), respectively.

2.4.2. Soil surface evaporation (E) and plant transpiration (T_p)

In order to measure evaporation (E) from the soil surface, another three mini-lysimeters were placed on the farm which filled-out with same soil but without cultivating corn. The E was measured from the difference between the amount of intake and drainage water in mini-lysimeter at the irrigation interval. Finally, plant transpiration (T_p) was estimated from the difference between $ET_{c\ act}$ and E , by the following general equation (Moran et al. 2009; Dehghanisanij et al. 2020):

$$T_p = ET_{c\ act} - E \quad (5)$$

2.4.3. Crop coefficient (K_c)

Two forms of K_c are presented for calculate of crop coefficient including single and dual crop coefficient forms. In this study, single crop coefficient ($K_{c\ single}$) and dual crop coefficient ($K_{c\ dual}$) by FAO-56 method (Allen et al. 1998) were modified based on the climatic conditions of the study area.

2.4.4. Water use efficiencies

Water use efficiency (WUE , kg/m³) and irrigation water use efficiency ($IWUE$, kg/m³) were calculated by using Eqs. (6) and (7) (Sakthivadivel et al. 1999; Ati et al. 2012):

$$WUE = \frac{\text{Yield (kg)}}{ET_c \text{ (m}^3\text{)}} \quad (6)$$

$$IWUE = \frac{\text{Yield (kg)}}{\text{Total applied water (m}^3\text{)}} \quad (7)$$

2.4.5. Plant and yield sampling

leaf area index (LAI), plant height (H) and stem diameter (D) dry and wet biological yield was measured during growth stages. Leaf area index (LAI) was measured with the electronic leaf area-meter CI – 202, seven times during the growing season.

2.4.6. Statistical analyses

Analysis of variance was carried out to evaluate the effect of the three irrigations systems and mulch treatments on the evapotranspiration, yield and water use efficiency of corn by using the SAS package. Compare and rank the treatment means was done by Least Significant difference (LSD) tests. Differences were declared significant at $P \leq 0.05$.

3. Results and discussion

3.1. Actual corn evapotranspiration ($ET_{c\ act}$) and crop coefficient (K_c)

Table 4 shows the total corn $ET_{c\ act}$ during the growing season of corn. The total corn $ET_{c\ act}$ was 379.8, 377.0 and 371.9 mm for PCI , DI and SDI systems in the growing seasons, respectively.

(Table 4. here)

Lowest values of corn $ET_{c\ act}$ occurred at the initial stage with a value of 3.31, 2.34 and 2.34 mm/day for PCI , DI and SDI systems, during the growing season, respectively. Daily corn $ET_{c\ act}$ increased rapidly and reached its maximum value at the mid-season stage. Highest values of corn $ET_{c\ act}$ occurred 44 days after planting at the mid-season stage with a value of 11.61, 9.33 and 9.24 mm/day for PCI , DI and SDI systems during the growing season, respectively. Previous studies showed that seasonal variation of corn $ET_{c\ act}$ ranged from 200 to 663 mm for different climatic and regional conditions (Dehghanisani et al. 2006; Chuanyan and Zhongren 2007; Liu et al. 2017; Zhang et al. 2017; Akhavan et al. 2018; Dehghanisani et al 2020). Variation of corn $ET_{c\ act}$ during the growing season was shown in Fig. 3.

(Fig. 3 here)

Daily corn $ET_{c\ act}$ varied from 3.31 to 11.61 mm/ day, 2.34 to 9.33 mm/ day and 2.34 to 9.24 mm/ day in PCI , DI and SDI systems, respectively. Highest corn $ET_{c\ act}$ was observed for PCI system. It should be noted that the total actual corn evapotranspiration under the PCI system was as crop transpiration (T_p). Because of the type of system, the wet soil surface was not exposed to air and the total amount of irrigation water applied to the plant was the T . There was no E , drainage and deep percolation in PCI system, and results in zero percent of E loss from the soil surface (Fig. 4).

(Fig. 4 here)

When porous capsule is used, the soil surface usually remains dry than *DI* and *SDI* systems, this leads to the reduction of E and an increase in transpiration, WUE and $IWUE$ (Romero et al. 2004; Badr et al. 2010). The lowest $ET_{c\ act}$ values were obtained in *SDI* system in the growing season of corn, because less soil moisture faced with the free air surface, and accordingly the rate of E is lower in comparison with *DI* system.

Statistical analysis of $ET_{c\ act}$ and K_c indicated that the effects of irrigation systems were significant on corn $ET_{c\ act}$ and K_c ($P \leq 0.01$). However, there were no significant differences between the mulch and interaction of irrigation systems and mulch on corn $ET_{c\ act}$ and K_c (Table 5).

(Table 5. here)

The results showed that *PCI* system produced higher corn K_c values in comparison with *DI* and *SDI* system over the whole crop season ($P < 0.01$). Highest mean K_c value (0.82) was obtained in *PCI* system. Also, the lowest K_c values were obtained under with mulch condition, because using mulch reduced E compared with non-mulch treatments (Table 5). These results are in agreement with the findings of Vickers (2001), Mata et al. (2002) and Yaghi et al. (2013). Compare the results with the findings of other researchers show the crop coefficient changes in different regions or in specific region can be different with different planting dates, so that K_c extraction of regional and the calculated corn $ET_{c\ act}$ for different climatic conditions and different planting times is essential. The changing of K_c and LAI for the corn crop is plotted as a function of days after planting (*DAP*) (Fig. 5).

(Fig. 5 here)

Analysis of changes in K_c values was done by using the relationship between LAI and K_c observed on the same days when LAI was measured. On the basis of these data, a new K_c curve was plotted for seven days of LAI sampling as illustrated (Fig. 5). This was done to divest days with wet plants and soil because it is assumed that there were not irrigation and precipitation when LAI was sampled in middle stage of growing season. Fig. 5 showed that with the increase of LAI , the K_c increases markedly and LAI generally peaked, as did K_c .

3.2. Variation in single and dual crop coefficient ($K_{c\ single}$ and $K_{c\ dual}$)

Single crop coefficient ($K_{c\ single}$) and dual crop coefficient ($K_{c\ dual}$) values for corn growth stage given in Table 6, and the variation of $K_{c\ single}$ and $K_{c\ dual}$ for the corn are plotted as a function of days after planting (*DAP*) and are presented in Fig. 6.

(Table 6. and Fig. 6. here)

The corn $K_{c\ single}$ values suggested by FAO-56 were 0.3, 0.3-0.9 and 1.2 for the initial stage, crop development and mid-season stages, respectively. In this study, the recommended $K_{c\ single}$ values were modified based on the climatic conditions of the study area and changed to 0.3, 0.88 and 1.35 for PCI, 0.3, 0.86 and 1.34 for DI and 0.3, 0.87 and 1.35 for SDI systems for the initial, development-season and mid -season stages, respectively (Table 6).

The $K_{c\ dual}$ included the basal crop coefficient (K_{cb}) and evaporation coefficient (K_e). The amounts of K_{cb} suggested by FAO-56 for the initial stage, crop development and mid-season stages ($K_{cb\ ini}$, $K_{cb\ dev}$ and $K_{cb\ mid}$) were 0.15, 0.15- 1.15 and 1.15, respectively. The amount of $K_{cb\ dev}$ and $K_{cb\ mid}$ modified on the basis of plant height, wind speed and relative humidity in different regions. The recommended K_{cb} values were adjusted to 0.15, 0.77, and 1.22 for PCI, 0.15, 0.70, and 1.20 for DI and 0.15, 0.71, and 1.21 for SDI system for the initial, development-season and mid -season stages, respectively (Table 6).

Maximum value of K_{cb} was obtained equal to 1.22 for PCI system at the mid-season stage. The K_e varied temporally during the corn growing season. Average K_e value was higher at initial stage, and decreased gradually in all treatments, and reaching a minimum value as low as 0.12 for PCI system at the mid-season stage (Table 6). These results indicated that soil evaporation coefficient decreased during corn growth season, because of increase in more shade of the plant. The $K_{c\ dual}$ was higher than $K_{c\ single}$ in initial, development and mid-season stages. Also, Fig. 6 showed that the E was higher than the transpiration from crop in initial -season stage, and with increase in plant shading E was lowest than the transpiration from crop in development and mid-season stages.

3.3. Biological characteristics of corn

For each treatment, leaf area index (LAI), plant height (H) and stem diameter (D) during growth season are listed in Table 5. The result showed that irrigation systems had effect ($P < 0.05$) on LAI , H and D , and mulching treatments had significant effect ($P < 0.05$) on LAI and D for the growth season, but had no effect on H . The highest values of LAI , H and D were belonged to PCI system and recorded 4, 182 and 2.9 cm, respectively. When PCI system is used, the percentage of E loss was zero and due reduce water loss from the soil surface created more favorable condition for plant transpiration and also increase in biological characteristics of corn (Siyal et al. 2011; Tiwari et al. 2014). The LAI , H and D increased with mulching and recorded 4.1, 18 and 2.9 cm, respectively (Table 5). The interaction effects of irrigation methods and mulching treatments had no effect on LAI , H and D . PCI system in combination with mulch (PCI M₁) treatment had the highest values of LAI , H and D with 4.23, 191.6 and 3.0 cm. The lowest values obtained in DI system and non –mulch (M₂) and recorded 3.4, 169.2 and 2.9 cm. In all irrigation and mulch treatments, the

lowest values of *LAI*, *H* and *D* during the growing season, occurred at the initial stage, then increased rapidly and reached its maximum value at mid-season stage (Fig 7).

(Fig. 7 here)

Larger value of *LAI*, *H* and *D* of corn was found at during mid-season stage (70 days after sowing) of growing season. Positive effect of mulching on *LAI* in wheat was previously reported by Xie et al. (2005). Interaction of irrigation methods and mulching showed no significant effect on the *LAI*, *H* and *D* in during growing season. The temporal variations of the *H* and *D* of corn for all treatments are shown in Fig. 7.

3.4. Yield (biomass)

PCI, *DI* and *SDI* systems had significant effect ($p < 0.05$) on total dry and wet biological yield of corn (Table 5). Overall average dry and wet biological yield resulted from *PCI* system was significantly higher (28.41 and 97.70 ton/ ha) over *DI* (25.13 and 63.21 ton/ ha) and *SDI* (26.67 and 81.99 ton/ha). The better performance under *PCI* system explained maintenance of favorable moisture status in the root zone and help the plants to utilize moisture and efficient nutrients from the wetted area (Zotarelli et al. 2008; Badr et al. 2010). The M_1 treatment had also significant effect ($p < 0.05$) on dry and wet biological yield and recorded the highest biological yield which was 27.86 (ton/ ha). There were significant reductions on total yield under M_2 treatment condition. Previous studies have also reported significant yield with using mulch (Nijamudeen and Dharmasena 2002; Seyfi and Rashidi 2007). Mulching is known for reducing the weed dry matter due to physical hindrance and by reducing the amount of solar radiation reaching the weeds on soil surface and thus maintaining the optimal yield have shown the multiplier effect of mulching on biological yield crop in corn (Khurshid 2006). The *PCI* system with mulch (M_1) prevent the E from the soil surface. With the reduction in E is a general increase in T from corn. Increase of corn yield with mulch (M_1) in comparison with non- mulch (M_2) treatment was observed and is in agreement with the findings of Stein et al. (1998), Chakraborty et al. (2010), Balwinder et al. (2011) and Ram et al. (2013). *PCI* system with mulch (*PCI* M_1) produced highest total dry and wet biological yield in comparison with *DI* and *SDI* system with non- mulch cover. The lowest value of total dry and wet biological yield was obtained in *DI* in combination to non-mulch (*DI* M_2) treatment (23.19 and 58.54 ton /ha), respectively. *E* increased during the first week after transplanting when plants did not have enough canopies to shade the soil. Mulch directly affected the microclimate around the plant (absorptivity vs. reflectivity) of the surface and decreasing the soil water loss. These results are in agreement with the findings of El-Nemr (2006), Korir et al. (2006), Ertek et al. (2007), Dagdelen et al. (2009) and Tesfaye et al. (2016).

3.5. Water use efficiency (WUE)

Differences in irrigation methods caused significant effect ($p < 0.05$) in the water use efficiency (WUE) and irrigation water use efficiency ($IWUE$) (Table 5). *PCI* system produced higher WUE and $IWUE$ in comparison to *DI* and *SDI* systems. With using porous capsule the soil surface usually remains dry than *DI* and *SDI* systems, and this leads to an increase in T , WUE and $IWUE$ (Romero et al., 2004). The M_1 treatment had also significant effect ($p < 0.05$) on WUE and $IWUE$ which was 7.46 and 7.33 (kg/m^3). Interaction of irrigation methods and mulching showed no significant effect on the $IWUE$ and WUE in during growing season (Table 5). The $IWUE$ of all the treatment ranges was from 6.10 to 7.89 kg/m^3 , while the WUE of all the treatment ranges from 6.14 to 7.89 kg/m^3 , which this finding can indicate the difference in irrigation $IWUE$ and WUE compared between irrigation methods. *PCI* system with mulch (*PCI M₁*) treatment produced higher values of $IWUE$ and WUE , because *PCI* system had E and deep percolation losses than other irrigation methods. The lowest $IWUE$ and WUE values were obtained in *DI* system in combination with non-mulch (*DI M₂*) treatment which was 6.10 and 6.14 kg/m^3 in the growing season of corn. Uniform mulching on the soil surface reduces E , increases soil moisture content, and decreases water consumption, saving water without reducing grain yield and leading to a higher WUE (Jin et al. 2009; Wang et al. 2015; Shen et al. 2012).

4. Conclusions

This study evaluated the influence of porous capsule irrigation (*PCI*), surface drip irrigation (*DI*) and subsurface drip irrigation systems (*SDI*) under mulch condition on actual evapotranspiration ($ET_{c\ act}$), single and dual crop coefficient ($K_{c\ single}$ and $K_{c\ dual}$), yield and water use efficiency (WUE) of corn (*Zea mays L.*) in the semiarid climate of Iran. The resulted indicated that the effects of irrigation systems were significant on evapotranspiration. Maximum value of K_{cb} was obtained for *PCI* treatment at the mid-season stage. The K_e varied temporally during the corn growing season. The mean K_c was estimated to be 0.83, 0.74 and 0.73 for *PCI*, *DI* and *SDI* systems in the growing seasons, respectively. Highest total dry and wet biological yield was recorded in the *PCI* system in comparison to *DI* and *SDI* systems. Application of mulch was effective on the increase dry and wet biological yield of corn. *PCI* system under mulch conditions (*PCI M₁*) produced highest total dry and wet biological yield, WUE and $IWUE$ in compared to non-mulch. The better performance under *PCI* system can be explained to the maintenance of favorable moisture status in the root zone, that helped the plants to utilize moisture and efficient nutrients from the wetted area According to the results of this study, it can be concluded that using *PCI* system leads to higher biological yield, WUE and $IWUE$ in comparison to *DI* and *SDI* system.

Acknowledgements

We would like to thank the Agricultural Engineering Research Institute, Agricultural Research, Education and Extension Organization, Karaj, Alborz, Iran, for providing assistance to conduct this study.

Funding

The authors received support to conduct the study but the support did not include support for the submitted work.

Availability of data and material

The datasets generated during and/or analyzing for the current study are available from the corresponding author on reasonable request.

Authors contributions: Elahe Kanani: Performed the experiments and drafted the manuscript. Hossein Dehghanisanij: Conceived and designed the experiments, supervised the work, interpreted the data and co-wrote the paper. Samira Akhavan: Substantial contributions to the conception or design of the work; analysis, or interpretation of data for the work. All authors read and approved the final manuscript.

Code availability

For this publication we did not use any code.

Ethics approval

This approval is attached.

Compliance with ethical standards

Work is approved by technical committee in Agricultural Engineering Research Institute.

Conflict of interest The authors declare they have no conflict of interest.

Consent to participate The authors declare they have consent to participate.

Consent for publication The authors declare they have consent to publication.

References

- Akhavan S, Kanani E, Dehghanisanij H (2018) Assessment of different reference evapotranspiration models to estimate the actual evapotranspiration of corn (*Zea mays* L.) in a semiarid region (case study, Karaj, Iran). *Theor Appl Climatol* 137: 1403-1419. <https://doi.org/10.1007/s00704-018-2634-y>
- Allen RG, Pereira LS, Howell TA, Jensen ME (2011) Evapotranspiration information reporting: I. Factors governing measurement accuracy. *Agric Water Manag* 98: 899–920. <https://doi.org/10.1016/j.agwat.2010.12.015>
- Allen R G, Pereira L S, Raes D, Smith M (1998) Crop evapotranspiration: guidelines for computing crop water requirements. *FAO Irrig. Drain. Pap. No, 56*. FAO, Rome.
- American Soil Conservation Service (ASCS) (1992) National engineering handbook. Section 7, Part 15.

- Ati A S, Iyada A D, Najim S M (2012) Water use efficiency of potato (*Solanum tuberosum* L.) under different irrigation methods and potassium fertilizer rates. *AOAS* 57(2): 99-103. <https://doi.org/10.1016/j.aos.2012.08.002>
- Ayars J E, Fulton A, Taylor B (2015) Subsurface drip irrigation in California-Here to stay?. *Agric Water Manage* 157: 39-47. <https://doi.org/10.1016/j.agwat.2015.01.001>
- Badr M A., Hussein S A., El-Tohamy W A, Gruda N (2010) Efficiency of subsurface drip irrigation for potato production under different dry stress conditions. *Gesunde Pelandz* 62(2): 63-70. DOI:10.1007/s10343-010-0222-x
- Bainbridge D A (2001) Buried clay pot irrigation: A little known but very efficient traditional method of irrigation. *Agric Water Manage* 48 (2): 79-88. DOI: 10.1016/S0378-3774(00)00119-0
- Bainbridge D A (2002) Alternative irrigation systems for arid land restoration. *Ecol Restor* 20(1): 23-30.
- Bainbridge D A, Ramirez A, Jose J (2008) More efficient irrigation systems for desert and dryland restoration. *Expo zara goza, Land Use Planning, Forest Cover and Afforestation*.
- Bai J, Wang J, Chen X, Luo G, Shi H, Li, L, Li J (2015) Seasonal and inter-annual variations in carbon fluxes and evapotranspiration over cotton field under drip irrigation with plastic mulch in an arid region of Northwest China. *J. Arid Land* 7: 272–284. DOI: 10.1007/s40333-014-0012-x
- Balwinder-Singh Humphreys E, Eberbach P L, Katupitiya A, Yadvinder-Singh Kukkal S S (2011) Growth yield and water productivity of zero till wheat as affected by rice straw mulch and irrigation schedule. *Field Crop Res* 121: 209–225. <https://doi.org/10.1016/j.fcr.2010.12.005>
- Biswas S K, Akanda A R, Rahman M S, Hossain M A (2015) Effect of drip irrigation and mulching on yield, water-use efficiency and economics of tomato. *Plant Soil Environ* 61: 97–102. <https://doi.org/10.17221/804/2014-PSE>
- Chakraborty D, Garg R N, Tomar R K, Singh R, Sharma S K, Singh R K, Trivedi S M, Mittal R B, Sharma P K., Kamble K H (2010) Synthetic and organic mulching and nitrogen effect on winter wheat (*Triticum aestivum*) in a semi-arid environment. *Agric Water Manage* 97: 738–748. <https://doi.org/10.1016/j.agwat.2010.01.006>
- Chuanyan Z, Zhongren N (2007) Estimating water needs of maize (*Zea mays* L.) using the dual crop coefficient method in the arid region of northwestern China. *Afric. J. Agric. Res* 2(7): 325-333.
- Dagdelen N, Basal H, Yilmaz E, Gurbuz T, Akcay S (2009) Different drip irrigation regimes affect Cotton yield, water use efficiency and fiber quality in western Turkey. *Agric. Water Manage* 69: 111–120. <https://doi.org/10.1016/j.agwat.2008.07.003>
- Dehghanisanij H, Kanani E, Akhavan S (2020) Evapotranspiration and components of corn (*Zea mays* L.) under micro irrigation systems in a semi-arid environment. *SJAR* 18 (2): e1202. <http://dx.doi.org/10.5424/sjar/2020182-15647>
- Dehghanisanij H, Yamamoto T, Rasiah V (2004) Assessment of evapotranspiration estimation models for use in semi-arid environments. *Agric Water Manage* 64 (2): 91–106. [https://doi.org/10.1016/S0378-3774\(03\)00200-2](https://doi.org/10.1016/S0378-3774(03)00200-2)
- Dehghanisanij H, Agassi M, Anyoji H, Yamamoto T, Inoue M, Eneji, A.E (2006) Improvement of saline water use under drip irrigation system. *Agric Water Manage* 85(3): 233-242. <https://doi.org/10.1016/j.agwat.2006.05.005>
- Dugas W A, Bland W L (1989) The accuracy of evaporation measurements from small lysimeters. *Agric. For. Meteorol* 46(1-2): 119-129. [https://doi.org/10.1016/0168-1923\(89\)90116-0](https://doi.org/10.1016/0168-1923(89)90116-0)
- El-Nemr A M (2006) Effect of mulch types on soil environmental conditions and their effect on the growth and yield of cucumber plants. *J. Appl Sci* 2 (2): 67–73.

- Ertek A, Sensoy S, Gedik I, Kuciikyumuk C (2007) Irrigation scheduling for green capsicum (*Capsicum annum* L.) grown by field condition by using class A pan evaporation value. *American- Eurasian J. Agril. Environ. Sci* 2: 349-358.
- FAO (2011) Subset production crops database. Accessed 18 Feb 2015. FAO, Rome. <http://faostat3.fao.org/home/index.html>.
- Farshi A A, Shariati M H, Jarollahi R, Ghaemi M H, Shabifar M, tolaei M M (1997) Estimated water requirement major plants agricultural and horticultural of country. Soil and Water Research Institute, Publication of Agriculture Education in Karaj. 394p. (In Persian).
- Jin H, Qingjie W, Hongwen L, Lijin L, Huanwen G (2009) Effect of alternative tillage and residue cover on yield and water use efficiency in annual double cropping system in North China Plain. *Soil Till Res* 104(1): 198-205. <https://doi.org/10.1016/j.still.2008.08.015>
- Khurshid K, Iqbal M, Arif . S, Nawaz A (2006) Effect of tillage and mulch on soil physical properties and growth of maize. *IJAB* (5): 593–596.
- Kong Q, Li G, Wang Y, Huo H (2012) Bell pepper response to surface and subsurface drip irrigation under different fertigation levels. *Irrig. Sci* 30 (3): 233–245. DOI: 10.1007/s00271-011-0278-0
- Korir N, Aguyoh K, Gaqiong JN (2006) Enhanced growth and yield of greenhouse produced cucumber under high altitude areas of Kenya. *ATS* 39 (4): 249–254. <http://ir-library.ku.ac.ke/handle/123456789/7384>
- Liu H, Wang X, Zhang X, Zhang L, Li Y, Huang G (2017) Evaluation on the responses of maize (*Zea mays* L.) growth, yield and water use efficiency to drip irrigation water under mulch condition in the Hetao irrigation District of China. *Agric Water Manage* 179: 144-157. <https://doi.org/10.1016/j.agwat.2016.05.031>
- Mata V H, Nunez E R, Sanchez G P (2002) Soil temperature and soil moisture in Serrano pepper (*Capsicum annum* L.) with fertigation and mulching. In: Proceeding of the 16th International Pepper Conference, Tamaulipas, Mexico, Available at: [http://www.pepperconference.org/proceedings/soil temperature and soil moisture.pdf](http://www.pepperconference.org/proceedings/soil%20temperature%20and%20soil%20moisture.pdf).
- Moran M S, Scott R L, Keefer T O, Emmerich W E., Hernandez M, Nearing G S, Paige G B, Cosh M H, O'Neill P E (2009) Partitioning evapotranspiration in semiarid grassland and shrubland ecosystems using time series of soil surface temperature. *AGR FOREST METEOROL* 149(1): 59-72. <https://doi.org/10.1016/j.agrformet.2008.07.004>
- Nijamudeen M S, Dharmasena P B (2002) Performance of chilli under drip irrigation with mulch. *ASDA* 4: 89-94.
- Paul J C, Mishra J N, Pradhan P L, Panigrahi B (2013) Effect of drip and surface irrigation on yield, water-use-efficiency and economics of capsicum (*capsicum annum* L.) Grown under mulch and no mulch conditions in eastern coastal India. *IJSD* 2(1): 99-108. DOI: <https://doi.org/10.14207/ejsd.2013.v2n1p99>
- Panigrahi H K, Agrawal N, Agrawal R, Dubey S, Tiwari S P (2016) Effect of drip irrigation and polythene mulch on the fruit yield and quality parameters of mango (*Mangifera indica* L.). *J. Hortic Sci Biotech* 5 (2): 140-143.
- Postel S (1992) The last oasis: facing water scarcity. Worldwatch Institute. Washington D.C, USA.
- Qin S, Li S, Kang S, Du T, Tong L, Ding R (2016) Can the drip irrigation under film mulch reduce crop evapotranspiration and save water under the sufficient irrigation condition? *Agric Water Manage* 177: 128-137. DOI: 10.1016/j.agwat.2016.06.022
- Ram H, Dadhwal V, Vashist K K, Kaur H (2013) Grain yield and water use efficiency of wheat (*Triticum aestivum* L.) in relation to irrigation levels and rice straw mulching in North West India. *Agric Water Manage* 128: 92-101. <https://doi.org/10.1016/j.agwat.2013.06.011>
- Reddy M, Ayyanagowdar M S, Patil M G., Polisgowdar B S, Nemichandrappa M, Patil J R (2018) Performance of Water Melon under Mulching, Subsurface and Surface Drip Irrigation Systems in Semi-Arid Region. *Int. J. Pure App. Biosci* 6(1): 488-496. DOI: <http://dx.doi.org/10.18782/2320-7051.6089>

- Romero P, Botia P, Garcia F (2004) Effects regulated deficit irrigation under subsurface drip irrigation conditions on water relations of mature almond trees. *Plant Soil* 260: 155–168. <https://doi.org/10.1023/B:PLSO.0000030193.23588.99>
- Rosa R D., Paredes P, Rodrigues G C, Fernando R M, Alves I, Pereira L S, Allen R G (2012) Implementing the dual crop coefficient approach in interactive software. 2. Model testing. *Agric Water Manag* 103: 62–77. <https://doi.org/10.1016/j.agwat.2011.10.018>
- Sakthivadivel R, de Fraiture C, Molden D J., Perry C, Kloezen W (1999) Indicators of land and water productivity in irrigated agriculture. *Int. J. Water Resour D* 15: 161–179. <https://doi.org/10.1080/07900629948998>
- Setiawan B I., Saleh E, Nurhidayat Y (1998) Pitcher irrigation system for horticulture in dry lands. *Proceedings water and land resources development and management for sustainable use*, 2.
- Seyfi K, Rashidi M (2007) Effect of drip irrigation and plastic mulch on crop yield and yield components of cantaloupe. *Int. J. Agric. Biol* 9(2): 247–249.
- Sharma A R, Singh R, Dhyani SK, Dube R K (2010) Moisture conservation and nitrogen recycling through legume mulching in rainfed maize (*Zea mays*)- wheat (*Triticum aestivum*) cropping system. *Nutr Cycl Agroecosys* 87 (2): 187–197. <https://doi.org/10.1007/s10705-009-9327-y>
- Shen J Y, Zhao D D, Han H F, Zhou X B, Li Q Q (2012) Effects of straw mulching on water consumption characteristics and yield of different types of summer maize plants. *Plant Soil Environ* 58(4): 161–166. <https://doi.org/10.17221/404/2011-PSE>
- Silva D A da, Silva S A de, Gheyi H R (1981a) Irrigacao por capsulas porosas III: Avaliacao tecnica do metodo por pressao hidrostatica. *Boletim de pesquisa* 3: 20–42.
- Silva D A da, Gheyi H R, Silva S A de, Magalhaes A A (1981b) Irrigacao por capsulas porosas IV: Efeitos das diferentes pressoes hidrostaticas e populacoes de plantas sobre a producao do milho. *Boletim de pesquisa* 3: 43–59.
- Singh R (2005) Influence of mulching on growth and yield of tomato (*Lycopersicon esculentum*) in north India plains. *Vegetable Science* 32 (1): 55–58.
- Siyal A A, Siyal A G, Hasini M Y (2011) Crop Production and Water Use Efficiency under Subsurface Porous Clay Pipe Irrigation. *Pakistan. J. Agric. Eng'g., Vet. Sc* 27 (1): 39–50.
- Soomro A A (2002) Viability of pitcher irrigation. *Daily Dawn Karachi*. 6th May issue. <http://www.dawn.com/2002/05/06/eb12.htm>. Accessed on May 11, 2009.
- Stein T (1998) Development and evaluation of design criteria for pitcher irrigation systems. *Beiheft no. 66, selbstverlag des verbandes der tropenlandwirte*. Witzenhausen ev, witzenhausen.
- Tarara J M (2000) Microclimate modification with plastic Mulch. *Hortscience* 3 (5): 169–180.
- Tesfaye T, Tigabu E, Germadu Y, Lemma H (2016) Effect of colored polyethylene mulch on soil temperature, growth, fruit quality and yield of tomato (*Lycopersicon esculentum* Mill.). *World J. Agric. Sci* 12: 161–166. DOI: 10.5829/idosi.wjas.2016.12.3.1910
- Tesfaye T, Tesfaye K, Woldetsadik K (2012) Clay pot irrigation for tomato (*lycopersicon esculentum* mill) production in the north east semiarid region of ethiopia. *JARTS* 112 (1): 11–18. <http://nbn-resolving.de/urn:nbn:de:hebis:34-2011101139325>
- Thentu T L, Dutta D, Mudi D D, Saha A (2016) Performance of broccoli (*Brassica oleracea* var. *italica*) under drip irrigation and mulch. *J. ANSF* 8(3): 1410–1415. DOI: 10.31018/jans.v8i3.974
- Tiwari K N, Kumar M, Santosh D T, Singh V K, Maji M K, Karan A K (2014) Influence of drip irrigation and plastic mulch on yield of sapota (*achras zapota*) and soil nutrients. *Irrig Drain Syst Eng* 3: 116. DOI: 10.4172/2168-9768.1000116

- Vickers A (2001) Water use Conservation. Water Plow Press, Amherst, MA, US. 215–380.
- Wang H, Wang C, Zhao X, Wang F (2015) Mulching increases water-use efficiency of peach production on the rainfed semiarid Loess Plateau of China. *Agric Water Manage* 154: 20-28. <https://doi.org/10.1016/j.agwat.2015.02.010>
- Xie Z K, Wang Y J, Li F M (2005) Effect of plastic mulching on soil water use and spring wheat yield in arid region of northwest China. *Agric Water Manage* 75: 71–83. <https://doi.org/10.1016/j.agwat.2004.12.014>
- Yaghi T, Arslan A, Naoum F (2013) Cucumber (*Cucumis sativus*, L.) water use efficiency (WUE) under plastic mulch and drip irrigation. *Agric Water Manage* 128: 149-157. <https://doi.org/10.1016/j.agwat.2013.06.002>
- Zhang B, Liu Y, Xu D, Zhao N, Lei B, Rosa R D, Paredes P, Paço T A, Pereira L S (2013) The dual crop coefficient approach to estimate and partitioning evapotranspiration of the winter wheat–summer maize crop sequence in North China Plain. *Irrig Sci*, 31(6), 1303-1316. <https://doi.org/10.1007/s00271-013-0405-1>
- Zhang H, Xiong Y, Huang G, Xu X, Huang Q (2017) Effects of water stress on processing tomatoes yield, quality and water use efficiency with plastic mulched drip irrigation in sandy soil of the Hetao Irrigation District. *Agric Water Manage* 179: 205-214. <https://doi.org/10.1016/j.agwat.2016.07.022>
- Zotarelli L, Dukes M D, Scholberg J.M, Hanselman T, Le Femminella K, Munoz-Carpena R (2008) Nitrogen and water use efficiency of zucchini squash for a plastic mulch bed system on a sandy soil. *Sci. Hortic* 116: 1–8. <https://doi.org/10.1016/j.scienta.2007.10.029>

Figure Captions

Fig. 1. The layout of experimental setup (mulch, plants, and the mini-lysimeter).

Fig. 2. Porous capsule used for installation in the farm.

Fig. 3. Variation of actual corn evapotranspiration in irrigation methods: a – Porous capsule irrigation (*PCI*), b- Surface drip irrigation (*DI*), c- Subsurface drip irrigation (*SDI*) during the corn growing season.

Fig. 4. Variation of soil evapotranspiration and corn transpiration in irrigation methods: a – Porous capsule irrigation (*PCI*), b- Surface drip irrigation (*DI*), c- Subsurface drip irrigation (*SDI*) during the corn growing season.

Fig. 5. Variation of crop coefficient and leaf area index (*LAI*) of corn in irrigation system: a – Porous capsule irrigation (*PCI*), b- Surface drip irrigation (*DI*), c- Subsurface drip irrigation (*SDI*) during the corn growing season.

Fig. 6. Daily single and dual crop coefficient during the corn growing season.

Fig. 7. Variation of crop height (*H*) and stem diameter (*D*) of corn in irrigation system: a – Porous capsule irrigation (*PCI*), b- Surface drip irrigation (*DI*), c- Subsurface drip irrigation (*SDI*) during the corn growing season.

Fig. 1

Fig. 2

Fig. 3

a

b

c

Fig. 4

a

b

c

Fig. 5

Fig. 6

a

b

c

Fig. 7

Table 1. Average climatic parameters during the 2014 corn growing season at the experimental site.

Month	T _{min} (°C)	T _{max} (°C)	T _{avg} (°C)	RH _{avg} (%)	U ₂ (m/s)	RS (Mj/m ² day)
August	20.4	36.5	28.4	32	5.0	24.9
September	18.2	33.5	25.8	38	4.1	23.7
November	11.5	24.7	17.8	49	4.8	22.7
Total	16.7	31.5	24	39.6	4.7	23.8

T: air temperature. *RH*: relative humidity. *U*₂: wind speed. *RS*: solar radiation

Table 2. Physicochemical properties of the experimental site soil

Soil depth (cm)	BD (g/cm ³)	FC (%)	WP (%w)	PH	EC (dS/m)	Soil textures
0-20	1.42	22.5	9.8	7.8	1.41	loam
20-40	1.42	22.4	9.6	7.9	1.21	loam
40-60	1.42	22.1	9.5	8.1	2.46	loam

BD: bulk density. *FC*: field capacity. *WP*: permanent wilting. *EC*: electrical conductivity

Table 3. Chemical characteristics of irrigation water

Water source	pH	EC (dS/m)	Cations and anions (meq/L)					SAR (%)
			Cations			Anions		
			Na+	Mg ²⁺	Ca ²⁺	SO ₄ ²⁻	HCO ₃ ⁻	
Well	7.8	0.8	3	2.4	2.4	1.8	2.4	1.82

Table 4. Amounts of actual corn evapotranspiration ($ET_{c\ act}$) during growing season

stage after planting	$ET_{c\ act-(PCI)}$ (mm)	$ET_{c\ act-(DI)}$ (mm)	$ET_{c\ act-(SDI)}$ (mm)	Irrigation amount (mm)
Initial	79.7	79	77.8	79.7
Development	143.4	142.5	140.1	143.4
Middle	156.7	155.5	153.9	156.7
sum	379.8	377.0	371.9	379.8

PCI: Porous capsule irrigation, *DI*: Surface drip irrigation, *SDI*: Subsurface drip irrigation

Table 5. Influence of irrigation systems and mulching on actual crop evapotranspiration ($ET_{c \text{ act}}$), crop coefficient (K_c), leaf area index (LAI), plant height (H) and stem diameter (D) dry and wet biological yield, water use efficiency (WUE) and irrigation water use efficiency (IWUE)

Treatmen	$ET_{c \text{ act}}$ (mm)	$K_c \text{ average}$	LAI (m ² /m ²)	H (cm)	D (cm)	Dry biological yield (ton/ha)	Wet biological yield (ton/ha)	IWUE (kg/m ³)	WUE (kg/m ³)
irrigation									
Systems									
PCI	379.9 ^a	0.82 ^a	4.0	182	2.9	28.41 ^a	97.70 ^a	7.47 ^a	7.47 ^a
DI	374.1 ^b	0.74 ^b	3.7	179	2.8	25.13 ^b	63.21 ^b	6.56 ^b	6.67 ^b
SDI	370.2 ^b	0.73 ^b	3.9	182.5	2.9	26.67 ^{ab}	81.99 ^{ab}	7.07 ^{ab}	7.26 ^{ab}
Prob > F	<0.004	<0.0001	Ns	Ns	Ns	0.040	0.054	0.034	0.042
Mulching									
M ₁	373.1	0.77	4.1 ^a	189	2.9 ^a	27.86 ^a	84.13 ^a	7.33 ^a	7.46 ^a
M ₂	376.1	0.76	3.6 ^b	172.5	2.8 ^b	25.61 ^b	73.78 ^b	6.74 ^b	6.80 ^b
Prob > F	NS	NS	0.014	NS	0.01	0.023	0.22	0.031	0.017
Interaction									
PCI M ₁	379.9	0.82	4.2	191.6	3.0	29.98	107.8	7.89	7.89
PCI M ₂	379.9	0.82	3.7	183.5	2.9	26.83	75.63	7.05	7.05
DI M ₁	370.8	0.74	3.5	173.1	2.8	26.80	67.88	7.02	7.19
DI M ₂	377.2	0.75	3.4	169.2	2.7	23.19	58.54	6.10	6.14
SDI M ₁	368.5	0.73	3.7	189.2	2.9	27.07	87.15	7.09	7.30
SDI M ₂	371.9	0.74	3.7	179.3	2.8	26.53	76.64	7.06	7.21
Prob > F	NS	NS	NS	NS	NS	NS	NS	NS	NS

M₁: with mulch, M₂: non- mulch and NS: not significant

Table 6. Average values of soil evaporation coefficient (k_e), basal crop coefficient (k_{cb}), single crop coefficient ($k_{c\ single}$) and dual crop coefficient ($k_{c\ dual}$) at different growing stages of corn

Irrigation Methods	Factor	Initial	Development	Mid-season
Porous capsule irrigation (<i>PCI</i>)	K_{cb}	0.15	0.77	1.22
	K_e	0.33	0.26	0.12
	$K_{c\ single}$	0.30	0.88	1.35
	$K_{c\ dual}$	0.48	1.03	1.34
Surface drip irrigation (<i>DI</i>)	K_{cb}	0.15	0.70	1.20
	K_e	0.45	0.36	0.16
	$K_{c\ single}$	0.30	0.86	1.34
	$K_{c\ dual}$	0.60	1.06	1.36
Subsurface drip irrigation (<i>SDI</i>)	K_{cb}	0.15	0.71	1.21
	K_e	0.40	0.36	0.14
	$K_{c\ single}$	0.30	0.87	1.35
	$K_{c\ dual}$	0.55	1.07	1.35