Comparative Analysis of Postoperative Sexual Dysfunction and Quality of Life in Type A Aortic Dissection Patients of Different Ages

Zeng-Rong Luo

Xiehe Affiliated Hospital of Fujian Medical University

Dong-Shan Liao (✉ liaodongshan3031@163.com)

Xiehe Affiliated Hospital of Fujian Medical University

Liang-Wan Chen

Xiehe Affiliated Hospital of Fujian Medical University

Research article

Keywords: sexual dysfunction, quality of life, type A aortic dissection, surgery, different age

Posted Date: September 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-68396/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Cardiothoracic Surgery on May 1st, 2021. See the published version at <https://doi.org/10.1186/s13019-021-01468-0>.

Abstract

Background: To compare postoperative sexual dysfunction (SD) and quality of life (QOL) in Type A Aortic Dissection (AAD) Patients of Different Ages.

Methods: From January 2018 to December 2019, 204 AAD postoperative survivors in Union Hospital of Fujian Medical University were selected and were divided into youth group (less than 50 years old) and elderly group (more than 50 years old). We evaluated SD according to the male International Erectile Dysfunction Index (IIEF-5) and female sexual function index (FSFI) . The Short Form 12 Health Survey Questionnaire (SF-12) and Quality of Life Enjoyment and Satisfaction Questionnaire (Q-LES-Q) were used to investigate the QOL, Quick Inventory Depressive Symptomatology-Self Report (QIDS-SR) and the Beck Depression Inventory-II (BDI-II) to investigate depressive symptoms.

Results: 175 patients completed all the questionnaire (85.8%). The total SD prevalence rate was 38.9% (68 cases), with 27.4% of the youth (20 cases) and 47.1% of the elderly (48 cases). The age of non-SD and SD patients was 49.0 ± 11.5 and 56.9 ± 10.8 years, respectively ($P=0.03$). Compared with non-SD patients, the total physical health of SD patients was significantly worse ($P=0.04$), however, the mental health was not significantly worse ($P=0.77$); the depressive symptoms did not expressed a significant difference between the SD and non-SD groups (QIDS-SR $P=0.15$, BDI-II $P=0.06$) . Total physical health in the youth SD group did not show significant better ($P = 0.24$), however, total mental health showed significantly worse ($P=0.04$), depressive symptoms scores were significantly higher (QIDS-SR $P=0.03$, BDI-II $P=0.04$).

Conclusion: The postoperative AAD SD prevalence of elderly is higher than that of youth, and the total physical health of SD patients is poorer than those without SD patients. The youth SD patients did not show a significant higher scores than the elderly SD patients, instead, the youth SD patients were more psychologically affected than the elderly SD patients, whose mental health was worse, and depression symptoms were more obvious, suggesting that the factors affecting the QOL of postoperative SD patients are related to physical factors, but the youth postoperative SD patients mainly affected by psychological factors.

Background

Acute aortic dissection (AAD) is an age-dependent, life-threatening cardiovascular disease associated with high mortality due to various fatal complications¹⁻⁷. Studies have shown that long-term survival after initial treatment is good⁸. With many advances in surgical techniques in the past decade, efforts have been made to improve the survival rate after AAD surgery, and a lack of attention has been paid to improving quality of life (QOL). QOL covers physical, emotional, mental, professional and sexual aspects⁹. However, rare research has been conducted to evaluate the impact of changes in sexual function and emotion on the QOL of postoperative AAD survivors. Resuming sexual activity is one of the important factors for psychosocial recovery after cardiac surgery¹⁰. The aim of this study is to better understand the impact of sexual function status of postoperative AAD survivors at different ages on the QOL, and to provide possible suggestions for improving physical health and psychological health.

Methods

1. Research object

1.1 We totally distributed related questionnaires by mail or by outpatient appointments to 204 type A aortic dissection (AAD) survivors who successfully underwent “modified triple-branched stent graft implantation technique” surgery¹¹ in the Union Hospital of Fujian Medical University at the end of the 6th month after discharge from January 2018 to December 2019. This study was approved by the ethics committee of Fujian Medical University, China.

1.2 Inclusion criteria: (1) Patients who survived more than 6 months after being discharged from the hospital after successful AAD surgery; (2) 18 to 70 years old; (3) Fixed sex partners; (4) No mental illness, no cognitive dysfunction in reading, writing and communicating, able to read and comprehend the contents of the questionnaire; (5) Informed consent to this study, volunteered to participate in the study; (6) People who had regular sex before surgery .

1.3 Exclusion criteria: (1) Those with mental illness, language communication barriers, dyslexia, and non-cooperation; (2) Reproductive system diseases and organic diseases of other vital organs; (3) Exclude those with severe preoperative sexual dysfunction; (4) Exclude those who take drugs that seriously affect sexual function; (5) Exclude those who fail to fill in the questionnaire.

2. Demographics

We collected the following demographic and clinical variables from all effective subjects: age, gender, chronic disease, marital status and the sexual partners situation.

3. Research tools

3.1 Sexual Dysfunction(SD)

We defined SD for men by using the International Index of Erectile Function-5 (IIEF-5): The International Index of Erectile Function (IIEF) was designed in 1997 by Rosen et al.¹², a doctor of American psychology. It originally contained 15 items to form IIEF-15. Rosen et al.¹³ deleted 10 items after a series of analyses and the remaining 5 scoring items consist of IIEF-5 (including the confidence to maintain erection, the hardness of erection, the frequency of maintaining erection, the ability of maintaining erection, satisfaction with sexual intercourse). Because IIEF-5 and IIEF-15 have the same good evaluation effect and are highly recommended, it has become the most widely used and most authoritative questionnaire on male sexual function in the world. For the diagnosis of erectile dysfunction (ED) and the severity of ED, IIEF-5 can be used as an accurate and effective screening tool. A number of studies have shown that IIEF-5 also showed good reliability and validity in the Chinese population for the diagnosis of ED divided by 21^{13,14}. Patients were scored according to IIEF-5 based on their erectile function in the past 6 months. IIEF-5 score ≥ 22 is classified as normal erectile function, ≤ 21 is classified as erectile dysfunction (ED), of which, scores 12-21 are classified as mild ED, 8-11 are classified as moderate ED, and ≤ 7 are classified as severe ED. The Female Sexual Function Index (FSFI)

was used to define female SD which is a multi-dimensional scale used to assess female sexual function with score ranging from 2-36¹⁵. SD for women was defined as obtaining a total FSFI score of < 26.5¹⁶.

3.2 Quality of Life (QOL)

The Quality of Life Enjoyment and Satisfaction Questionnaire (Q-LES-Q)¹⁷ were used to assess the QOL. The total score is calculated as a percentage of the total possible scores and reported as a score between 0 and 100. The lower the score, the worse the QOL. We also measured QOL using the Short Form 12 Health Survey Questionnaire (SF-12)¹⁸, with scoring based on the RAN-12 score, which calculates the average score of physical and mental health among the general population in the United States, with an average score of 50 and a standard deviation of 10¹⁹. Scoring higher or lower than 50 indicate better or worse function, respectively, as compared with the general population.

3.3 Depression

We used the Quick Inventory Depressive Symptomatology-Self Report (QIDS-SR)²⁰ to assess the severity of depressive symptoms, with a score ranging from 0 to 27. The severity of depression was also measured using the Beck Depression Inventory II (BDI-II)²¹, with scores ranging from 0 to 63. For both assessments, higher scores represent more severe depressive symptoms.

4. Statistical analysis

In this study, SPSS 20.0 statistical software was used to perform data statistical analysis. We performed normality test on measurement data firstly, measurement data conform to normal distribution were expressed as mean $\pm$ standard deviation ($\bar{x} \pm s$), analyzed with independent sample t test; corrected P value is used if the variance between groups was uneven; the enumeration data was described by percentage, the comparison of rates between groups was by χ^2 test. $P < 0.05$ indicated that the difference was statistically significant.

Results

1. General Information of The Research Object

The response rate to the questionnaire was high, 85.8% of the participants completed the SD measurement, 92% completed the SF-12, and 97% completed the Q-LES-Q, QIDS-SR and BDI-II. Effective data analysis was performed on 175 patients who had completely answered all questionnaires, aged 23 to 70 years old, with an average age of 53.4 ± 8.5 years, 73 cases in the youth group (mean age 43.4 ± 8.4 years) and 102 cases in the elderly group (mean age 59.4 ± 9.6 years); 144 cases were married or had a fixed sexual partner (82.29%), 31 cases were unmarried or divorced but had a fixed sexual partner; 119 cases of smoking (all men), 43 cases in the youth group and 76 cases in the elderly group; of the total number of chronic diseases, 29 cases of diabetes, 143 cases of hypertension, 33 cases of dyslipidemia, 10 cases of coronary heart disease, 40 cases with renal dysfunction, 10 cases with COPD; 143 cases with medication (hypertensive, hypoglycemic and

lipid-lowering drugs). The demographic characteristics are shown in Table 1. All patients underwent AAD three-branch stent implantation surgery about 6 months before starting the study.

2. SD Prevalence

Participants are mainly elderly patients (102, 58.3%) and youth patients (73, 41.7%). The total SD prevalence was 38.9% (68 cases), with the youth prevalence of 27.4% (20 cases, 29% of the SD patients) versus the elderly prevalence of 47.1% (48 cases, 71% of the SD patients) (Figure 1).

3. SD and QOL

The age of non-SD and SD patients is 49.0 ± 11.5 and 56.9 ± 10.8 years, respectively. The total physical health of SD patients is significantly worse ($P=0.04$), including general health ($P=0.02$), physical function ($P=0.04$); However, the QOL reported on mental health measures was not significantly worse ($P=0.77$), including vitality ($P=0.16$), social function ($P=0.59$), emotional role limitation ($P=0.59$) and mental health ($P=0.75$); the average score of depressive symptoms did not expressed a significant difference between the SD group and the non-SD group (QIDS-SR $P=0.15$, BDI- Φ $P=0.06$) (Table 2). Total physical health ($P=0.24$) in the youth SD group did not show a significant higher scores than the elderly SD group, including physical function ($P=0.43$), role limitation physical ($P=0.50$), pain ($P=0.71$) and general health ($P=0.64$), however, the scores of total mental health in the youth group was significantly lower than that of the elderly group ($P=0.04$), including emotional role limitation ($P=0.04$) and mental health ($P=0.03$), the average score of depressive symptoms was significantly higher than that of the elderly group (QIDS-SR $P=0.03$, BDI- Φ $P=0.04$) (Table 3).

Discussion

Very few postoperative survivors will consult a doctor "when can they resume sex?" when they are discharged from the hospital, and few doctors will explain the precautions related to the patient's postoperative sexual behavior. After all, this is an embarrassing question. Normal sexual function is a biological psychological process, usually has been found to be related to age, depressive symptoms, disease and hormone²². Some survey results show that sexual dysfunction (SD) after cardiac surgery is widespread²³. SD can be multi-factorial in etiology with hormonal, anatomical, physiological, and psychological effects⁹. Previous studies have shown that sexual function changes after organ transplantation: For example, patients with end-stage renal failure undergoing kidney transplantation have been found to get improved sexual function²⁴⁻²⁶; however, despite the improvement in total quality of life (QOL) and physical function, SD seems to persist after heart transplantation. Tabler and colleagues found that the causes of SD after transplantation included drug side effects, fear of death during intercourse, depression, body-image concerns, uncertainty about the gender of the donor, and changes in roles and responsibilities in the family²⁷. There is little information about the quality of sexual life and mood disorders of survivors after AAD surgery.

1. SD Prevalence

Our research shows that there is a certain prevalence of SD after AAD surgery (38.9%), and the prevalence of 47.1% in the elderly is significantly higher than 27.4% in the youth. This may be because the majority of our research subjects are elderly people, most of whom are associated with chronic diseases. With age, the level

of androgens in the body decreases. The decline of testosterone can reduce erectile response, sexual satisfaction and sexual frequency and also has a certain effect on ejaculation function ²⁸. In addition, the incidence of chronic diseases such as hypertension, diabetes, and dyslipidemia gradually increases with age, and the total number of combined chronic diseases is related to low interest in sexual activities or decreased satisfaction ²⁹. In view of this, in the cardiac rehabilitation program for patients after AAD surgery, instruct patients to actively treat various chronic diseases and strictly control blood pressure, blood sugar and blood lipid levels may be an effective way of improving the quality of sexual life.

2. SD and QOL

(1) SD is closely related to physical health

Our results show that the AAD postoperative SD patients showed obvious impairment of QOL, and we found that the impaired QOL was mainly manifested in physical health items, associated without depressive symptoms or clinically measurable mental health, which is consistent with the trend of QOL changes after heart transplantation ⁹. The results of SF-12 show that the total physical health of AAD postoperative SD patients, including general health, physical function, and physical role limitations are lower than those of non-SD patients, but the total mental health items, including social function, emotional limitations and mental health, are not found significant difference between AAD postoperative SD patients and non-SD patients. Although this does not imply a causal relationship, this study suggests that AAD postoperative SD is more strongly related to physical health than mental health and may indicate that the occurrence of SD in this population is more likely to be a physical cause rather than a psychological cause. This is consistent with the results of previous studies: the study of kidney transplant recipients found a link between sexual function and physical health ³⁰. Wolpowitz and Barnard ³¹ found that after a heart transplant, a quarter of male subjects were unable to obtain or maintain an erection, despite their desire to have sex with their partners. Heart transplant patients found increased libido but impaired erectile function, resulting in a mismatch between libido and performance ³².

(2) SD in the youth group is closely related to mental health

In addition to aging and other confounding factors such as drugs and lifestyle factors, studies have shown that poor mental health is also related to SD. For example, there was a strong relationship between depression and SD ³³. Such patients mainly manifested as decreased libido, erectile function and decreased sexual activity ^{34,35}. The relationship between mental health with AAD postoperative SD prevalence needs to be further clarified. Although self-reported depression and anxiety after AAD surgery were common ²³, in our study, SD patients were not found a significantly closer relationship with mental health or depressive symptoms than non-SD patients. But interestingly, it was found that the impaired QOL of AAD postoperative SD patients in the youth group is mainly reflected in the mental health items rather than physical health items. The mental health score of the youth group was lower than that of the elderly group, and the depression symptoms scores were higher than that of the elderly group, although the physical health was not impaired as severe as that of the elderly group. These results indicate that although our previous study found that SD was generally more strongly related to psychological than mental factors after AAD surgery, the

opposite is true for young SD patients, with more psychological factors than physical factors. Mild aerobic activity has been shown to promote physical and mental health and lower resting blood pressure after AAD surgery³⁶. Clinicians will encourage postoperative AAD survivors to perform mild to moderate aerobic exercise when discharged from the hospital. Previous investigations have shown that postoperative AAD survivors have significantly reduced physical activity after surgery. Analyzing the reasons of increase in physical inactivity, it was unlikely due to impaired physical function after AAD, because most of our patients showed adequate physical function status, and only 1% of patients needed home care²³. This most likely results from psychological depression or anxiety or fear. Studies have shown that serious health conditions such as ATAAD are often associated with fear and can trigger a post-traumatic stress disorder (PTSD)³⁷, which is diagnosed more frequently in the case of female and younger patients^{38,39}. Our research also shows that the physical health of AAD postoperative SD patients in the youth group did not show better than that of the elderly group, instead, the total mental health score is significantly lower than that of the elderly group, and the depression score is significantly higher than that of the elderly group. The questionnaire suggests that young patients think sexual activity is strenuous exercise and are fear of corresponding complications resulting from sexual activity. For example, due to the implantation of the stent, some young patients may worry about the displacement and shedding of the stent during sexual activity, and these psychological burdens will affect the patient's sexual function such as erection and ejaculation to a certain extent and sexual dysfunction aggravate the impairment of mental health in turn. Therefore, when postoperative AAD survivors recovers the physical health, it is unnecessary to deliberately avoid sexual activities⁴⁰. Health care providers should evaluate sexual health and encourage patients eliminate this unfounded psychological fear after AAD surgery. In recent years, many previous studies⁴¹⁻⁴⁴ have shown that providing sex education and counseling to patients and their spouses can improve the quality of sexual life of patients with cardiovascular disease and patients after cardiac surgery. It can be seen that it is necessary for postoperative AAD survivors to receive sexual health education provided by medical staff during hospitalization or discharge.

Limitations

The sample size included is relatively small, especially in the youth group. It is difficult to obtain accurate preoperative data about sexual function and quality of life (QOL). Selection bias may have occurred as patients were not randomized and the majority of patients were enrolled from outpatient clinic follow-up appointments or investigated by mailing questionnaires. Recall bias can also affect results as the data was acquired retrospectively. Some patients might have been experiencing SD, although we only include patients who deny preoperative SD. It was also a limitation that questionnaires were subjective assessment. Objective measures, such as activity, strength and ejaculation levels, in association with SD have not been well studied in the postoperative AAD population. Also, sex in the general population has appeared to be correlated with SD in heart transplant recipients⁹. This study could not rule out sex as a cause for SD, as it was not powered to address the effects of different sex groups on SD due to the small sample size.

Conclusions

There is a considerable proportion of SD patients after AAD surgery, with higher ratio in elderly than youth. SD seemed to be linked to physical health rather than mental health or QOL. Interestingly, in contrast, youth SD patients might experience worse mental health rather than physical health or QOL.

Declarations

Ethical Approval and Consent to participate

The present study was approved by the ethics committee of Fujian Medical University, China and adhered to the tenets of the Declaration of Helsinki.

Consent to publish

Not applicable

Availability of Data and Materials

Data sharing not applicable to this article as no data sets were generated or analyzed during the current study.

Competing interests

All authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author Contributions

LZR and LDS designed the study and submitted the manuscript. LZR collected, analyzed data and drafted the article. CLW supervised this study. All authors read the final version of this article and approved for publication.

Acknowledgement

We highly acknowledge the contribution by the participating doctors: Dao-Zhong Chen, Xiao-Fu Dai, Feng Lin, Qi-Min Wang, Han-Fan Qiu, Xue-Shan Huang, Zhong-Yao Huang.

Abbreviations

AAD: acute aortic dissection; SD: sexual dysfunction; S.Dev: standard deviation; Q-LES-Q: Quality of Life Enjoyment and Satisfaction Questionnaire-Short Form; QIDS-SR: Quick Inventory Depressive Symptomatology-Self Report; BDI-II: Beck Depression Inventory-II; SF-12: Short Form 12 Health Survey

References

1. P G Hagan, C A Nienaber, E M Isselbacher, D Bruckman, D J Karavite, P L Russman¹, et al. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA*, 2000;283:897-903.
2. De Leon Ayala IA, Chen YF. Acute aortic dissection: an update. *Kaohsiung J Med Sci*. 2012;28:299-305.
3. Golledge J, Eagle KA. Acute aortic dissection. *Lancet*. 2008;372:55-66.
4. Moon MR. Approach to the treatment of aortic dissection. *Surg Clin North Am*. 2009;89:869-93.
5. Suzuki T. Cardiovascular diagnostic biomarkers: the past, present and future. *Circ J*. 2009;73:806-9.
6. Tsai TT, Trimarchi S, Nienaber CA. Acute aortic dissection: perspectives from the International Registry of Acute Aortic Dissection (IRAD). *Eur J Vasc Endovasc Surg*. 2009;37:149-59.
7. Parthenakis F, Koutalas E, Patrianakos A, Koukouvas M, Nyktari E, Vardas P. Diagnosing acute aortic syndromes: the role of specific biochemical markers. *Int J Cardiol*. 2010;145:3-8.
8. Tsai TT, Fattori R, Trimarchi S, Isselbacher E, Myrmel T, Evangelista A, Hutchison S, Sechtem U, Cooper JV, Smith DE, Pape L, Froehlich J, Raghupathy A, Januzzi JL, Eagle KA, Nienaber CA; International Registry of Acute Aortic Dissection. Long-term survival in patients presenting with type B acute aortic dissection: insights from the International Registry of Acute Aortic Dissection. *Circulation*. 2006;114:2226-31.
9. Anita Phan, Waguih William Ishak, Bing-Jiun Shen, Justin Fuess, Kiran Philip, Catherine Bresee, Lawrence Czer, Ernst R Schwarz. Persistent Sexual Dysfunction Impairs Quality of Life after Cardiac Transplantation. *J Sex Med*, 2010;7:2765-73.
10. Seyed Khalil Foruzan-Nia¹, Mohammad Hassan Abdollahi, Seyed Hossein Hekmatimoghaddam, Seyedeh Mahdijeh Namayandeh, Mohammad Hadi Mortazavi. Incidence of sexual dysfunction in men after cardiac surgery in Afshar hospital, Yazd. *Iran J Reprod Med*, 2011;9:89-94.
11. Chen LW, Dai XF, Wu XJ, et al. Ascending Aorta and Hemiarch Replacement Combined With Modified Triple-Branched Stent Graft Implantation for Repair of Acute DeBakey Type I Aortic Dissection. *Ann Thorac Surg*. 2017;103:595-601.
12. Rosen R C, Riley A, Wagner G, et al. The international index of erectile function(IIEF): a multidimensional scale for assessment of erectile dysfunction. *Urology*, 1997,49:822-830.
13. Rosen R, Cappelleri JM, Lipsky J, et al. Development and evaluation of an abridged, 5-item version of the International Index of Erectile Function (IIEF-5) as a diagnostic tool for erectile dysfunction. *International Journal of Impotence Research*, 1999,11:319-326.
14. Ying Zhang, Qin Zhang, Ziyang Shi, et al. Sexual dysfunction in patients with neuromyelitis optica spectrum disorder. *Journal of Neuroimmunology*, 338 (2020) 577093.
15. Rosen R, Brown C, Heiman J, et al. The Female Sexual Function Index (FSFI): A multidimensional self-report instrument for the assessment of female sexual function. *J Sex Marital Ther*, 2000;26:191–208.
16. Wiegel M, Meston C, Rosen R. The female sexual function index (FSFI): Cross-validation and development of clinical cutoff scores. *J Sex Marital Ther*, 2005;31:1–20.

17. Endicott J, Nee J, Harrison W, Blumenthal R. Quality of Life Enjoyment and Satisfaction Questionnaire: A new measure. *Psychopharmacol Bull*, 1993;29:321-6.
18. Ware J Jr, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: Construction of scales and preliminary tests of reliability and validity. *Med Care*, 1996;34:220–33.
19. Windsor TD, Rodgers B, Butterworth P, Anstey KJ, Jorm AF. Measuring physical and mental health using the SF-12: Implications for community surveys of mental health. *AustNZJ Psychiatry*, 2006;40:797–803.
20. Rush AJ, Trivedi MH, Ibrahim HM, et al. The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): A psychometric evaluation in patients with chronic major depression. *Biol Psychiatry*, 2003;54:573–83.
21. Beck AT, Steer RA, Ball R, Ranieri W. Comparison of Beck Depression Inventories -IA and -II in psychiatric outpatients. *J Pers Assess*, 1996;67:588–97.
22. Seidman SN. Exploring the relationship between depression and erectile dysfunction in aging men. *J Clin Psychiatry*, 2002;63:5–12.
23. Ashish Chaddha, Eva Kline-Rogers, Alan C Braverman, Steven R Erickson, Elizabeth A Jackson, Barry A Franklin, Elise M Woznicki, Justin T Jabara, Daniel G Montgomery, Kim A Eagle. Survivors of Aortic Dissection: Activity, Mental Health, and Sexual Function. *Clin Cardiol*, 2015;38:652-9.
24. Barroso LV, Miranda EP, Cruz NI, et al. Analysis of sexual fu21. Filocamo MT, Zanazzi M, Li Marzi V, et al. Sexual dysfunction in women during dialysis and after renal transplantation. *J Sex Med*, 2009;6:3125–31.
25. Filocamo MT, Zanazzi M, Li Marzi V, et al. Sexual dysfunction in women during dialysis and after renal transplantation. *J Sex Med*, 2009;6:3125–31.
26. Nassir A. Sexual function in male patients undergoing treatment for renal failure: A prospective view. *J Sex Med*, 2009;6:3407–14.
27. J B Tabler, R L Frierson. Sexual concerns after heart transplantation. *J Heart Transplant*, 1990;9:397-403.
28. Nascimento E R, Maia A C O, Pereira V, et al. Sexual dysfunction and cardiovascular diseases: a systematic review of prevalence. *Clinics*, 2013;68:1462-1468.
29. Appa AA, Creasman J, Brown JS, et al. The Impact of Multimorbidity on Sexual Function in Middle-Aged and Older Women: Beyond the Single Disease Perspective. *Journal of Sexual Medicine*, 2015,11: 2744-2755.
30. TaValaii SA, Fathi Ashtiani A, Nasiri M, AsARS S, Maleki P, Einollahi B. The correlation between sexual function and quality of life after kidney transplantation: Does gender matter? *J Sex MD*, 2007; 4:1610-8.
31. Wolpowitz A, Barnard CN. Impotence after heart transplantation. *S. AFR MED J*, 1978; 53:693.
32. David Mulligan T, Sheehan H, Hanrahan J. Sexual function after heart transplantation. 1991 cases of heart and lung transplantation, 10:125
33. Araujo AB, Durante R, Feldman HA, Goldstein I, McKinlay JB. The relationship between depressive symptoms and male erectile dysfunction: Cross-sectional results from the Massachusetts Male Aging Study. *Psychosom Med*, 1998;60:458–65.

34. Reynolds CF 3rd, Frank E, Thase ME, et al. Assessment of sexual function in depressed, impotent, and healthy men: Factor analysis of a Brief Sexual Function Questionnaire for men. *Psychiatry Res*, 1988;24:231–50.
35. Williams K, Reynolds MF. Sexual dysfunction in major depression. *CNS*, 2006;11:19–23.
36. Brooke Road, Appel LJ, Ruben Fei M, et al. American Heart and Hypertension Research Council Professional Education Committee, Cardiovascular and Stroke Care Council, Epidemiology and Prevention Council, and Nutrition and Sports Activities Council. In addition to medication and diet: alternative ways to lower blood pressure: a scientific statement from the American Heart Association. *Hypertension*, 2013; 61: 1360–1383.
37. Jackson JC, Hart RP, Gordon SM, Hopkins RO, Girard TD, Ely EW. Post traumatic stress disorder and post-traumatic stress symptoms following critical illness in medical intensive care unit patients: assessing the magnitude of the problem. *Crit Care*, 2007 ; 11: R27.
38. Stein MB, Walker JR, Hazen AL, Forde DR. Full and partial post traumatic stress disorder: findings from a community survey. *Am J Psychiatry*, 1997;154:1114–9.
39. Ditlevsen DN, Elklit A. The combined effect of gender and age on post traumatic stress disorder: do men and women show differences in the lifespan distribution of the disorder? *Ann Gen Psychiatry*, 2010;9:32.
40. Corone S, Iliou MC, Pierre B, et al. The Heart Rehabilitation Working Group of the French Society of Cardiology. French registered type I acute aortic dissection entered the cardiac rehabilitation center after surgery. *Yul George Prehobill*, 2009; 16: 91–95.
41. Byrne M, Doherty S, Fridlund B G, et al. Sexual counselling for sexual problems in patients with cardiovascular disease. *Cochrane Database of Systematic Reviews*, 2016,2:D10988.
42. Steinke E E, Jaarsma T. Sexual counseling and cardiovascular disease: practical approaches. *Asian Journal of Andrology*, 2015,17:32-39.
43. Murphy P J, Mc S J, Casey D, et al. Sexual counselling for patients with cardiovascular disease: protocol for a pilot study of the CHARMS sexual counselling intervention. *Bmj Open*, 2016,6:e11219.
44. Steinke EE, Tiny J, Barnason SA, et al. Sexual counseling for individuals with cardiovascular disease and their partners: a consensus document from the American Heart Association and the ESC Council on Cardiovascular Nursing and Allied Professions (CCNAP). *Circulation*, 2013,34:3217-3235.

Tables

Table 1. Preoperative characteristics							
	Total	Youth	Elderly	P1	SD	Non-SD	P2
Patients (n)	175	73(41.7%)	102(58.3%)	-	68(38.9%)	107(61.1%)	-
Age (y)	53.4 ± 8.5	43.4±8.4	59.4±9.6	0.000	56.9±10.8	49.0±11.5	0.033
Male (n)	148(84.6%)	56	92	0.015	57	91	0.827
Chronic diseases							
Diabetes (n)	29(16.6%)	5	24	0.003	19	10	0.001
Hypertension (n)	143(81.7%)	49	94	0.000	62	81	0.010
Hyperlipidemia (n)	33(18.9%)	6	27	0.002	17	16	0.098
Coronary heart disease (n)	10(5.7%)	2	8	0.152	6	4	0.158
Renal dysfunction* (n)	40(22.9%)	12	28	0.087	21	19	0.044
COPD (n)	10(5.7%)	2	8	0.152	5	5	0.457
Drugs (n)	143(81.7%)	55	88	0.065	62	81	0.010
Smoking(n)	119(68.0%)	43	76	0.029	59	60	0.000
* Defined as preoperative creatinine greater than 1.5 mg/dl.							
P1: P value of the youth group versus the elderly group P2: P value of the SD group versus the non-SD group.							

Table 2. SD is closely related to physical health					
	Non-SD		SD		
	Mean	S.Dev.	Mean	S.Dev.	P value
Age (y)	49.0	11.5	56.9	10.8	0.03
QIDS-SR score	6.8	5.4	8.5	4.6	0.15
BDI-II score	6.5	6.8	9.7	6.6	0.06
Q-LES-Q score	78.2	10.6	70.4	12.3	0.08
SF-12					
Physical health total	60.0	10.8	49.6	11.8	0.04
Mental health total	68.0	12.2	60.3	10.9	0.77
Physical Functioning	60.0	11.0	50.2	11.1	0.04
Role limitation physical	62.3	11.5	59.5	9.6	0.36
Pain	65.5	9.5	54.3	13.3	0.49
General health	55.6	11.0	45.9	12.4	0.02
Vitality	53.5	14.8	48.3	9.9	0.16
Role limitation emotional	70.2	8.9	68.8	10.6	0.59
Social functioning	69.2	10.8	63.9	11.4	0.59
Mental health	62.3	10.8	60.1	12.1	0.75
* SD = sexual dysfunction; S.Dev = standard deviation; Q-LES-Q = Quality of Life Enjoyment and Satisfaction Questionnaire-Short Form; QIDS-SR = Quick Inventory Depressive Symptomatology-Self Report; BDI-II = Beck Depression Inventory-II; SF-12 = Short Form 12 Health Survey.					

Table 3. SD in the youth group is closely related to mental health					
	Youth SD		Elderly SD		
	Mean	S.Dev.	Mean	S.Dev.	P value
Age (y)	45.4	11.9	60.7	11.9	0.02
IIED score	16.6	5.8	16.8	8.8	0.49
FSFI score	19.5	10.6	21.7	11.0	0.68
QIDS-SR score	11.2	4.8	8.3	5.6	0.03
BDI-II score	12.0	7.5	8.1	5.8	0.04
Q-LES-Q score	69.6	10.3	74.9	15.0	0.05
SF-12 score					
Physical health total	52.4	10.8	48.1	9.2	0.24
Mental health total	55.5	10.8	63.7	10.2	0.04
Physical Functioning	54.9	12.8	49.6	9.9	0.43
Role limitation physical	60.0	14.2	60.2	13.8	0.50
Pain	56.8	12.5	52.6	14.6	0.71
General health	49.6	10.4	44.4	10.0	0.64
Vitality	43.1	8.5	50.2	10.6	0.10
Role limitation emotional	61.8	10.5	69.8	12.8	0.04
Social functioning	60.8	11.8	64.2	9.9	0.36
Mental health	55.2	10.2	61.1	11.8	0.03
* S.Dev = standard deviation; IIED = International Index of Erectile Dysfunction; FSFI = Female Sexual Function Index; Q-LES-Q = Quality of Life Enjoyment and Satisfaction Questionnaire; QIDS-SR = Quick Inventory Depressive Symptomatology-Self Report; BDI-II = Beck Depression Inventory-II; SF-12 = Short Form 12 Health Survey.					

Figures

Figure 1

The percentage of postoperative AAD sexual dysfunction patients by different age groups.