Metastatic Gastroesophageal Cancer in Older Patients – Is this Patient Cohort Represented in Clinical Trials?

Maeve A. Hennessy

University Hospital Waterford

Munzir Hamid

University Hospital Waterford

Niamh Keegan

University Hospital Waterford

Lynda Corrigan

University Hospital Waterford

Caitriona Goggin

University Hospital Waterford

Nay MyoOo

University Hospital Waterford

Marie Carrigan

St Lukes Radiation Oncology Oncology Network

David Mockler

John Stearne Library, Trinity Centre for Health Sciences, Dublin

Anita O'Donovan

Applied Radiation Therapy Trinity (ARTT), Trinity St James's Cancer Institute, Trinity College

Anne M Horgan (✉ annem.horgan@hse.ie)

University Hospital Waterford

Research Article

Keywords: Gastroesophageal, metastatic, phase III clinical trials

Posted Date: August 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-684646/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Cancer on January 3rd, 2022. See the published version at <https://doi.org/10.1186/s12885-021-09103-w>.

Abstract

Background: Older patients are underrepresented in the clinical trials that determine the standards of care for oncological treatment. We conducted a review to identify whether there have been age-restrictive inclusion criteria in clinical trials over the last twenty five years, focusing on patients with metastatic gastroesophageal cancer.

Methods: A search strategy was developed encompassing Embase, PubMed and The Cochrane Library databases. Completed phase III randomised controlled trials evaluating systemic anti-cancer therapies in metastatic gastroesophageal malignancies from 1st January 1995 to 18th November 2020 were identified. These were screened for eligibility using reference management software (Covidence; Veritas Health Innovation Ltd). Data including age inclusion/exclusion criteria and median age of participants were recorded. The percentage of patients ≥ 65 enrolled was collected where available. The change over time in the proportion of studies using an upper age exclusion was estimated using a linear probability model.

Results: 363 phase III studies were identified and screened, with 66 trials remaining for final analysis. The majority of trials were Asian (48%; $n = 32$) and predominantly evaluated gastric malignancies, (86%; $n = 56$).

The median age of participants was 62 (range 18-94). Thirty-two percent ($n = 21$) of studies specified an upper age limit for inclusion and over half of these were Asian studies. The median age of exclusion was 75 (range 65-80). All studies prior to 2003 used an upper age exclusion ($n = 12$); whereas only 9 that started in 2003 or later did (17%). Among later studies, there was a very modest downward yearly-trend in the proportion of studies using an upper age exclusion (-0.02 per year; 95%CI -0.05 to 0.01 ; $p = 0.31$). Fifty-two percent ($n = 34$) of studies specified the proportion of their study population who were ≥ 65 years. Older patients represented only 36% of the trial populations in these studies (range 7-60%).

Conclusions: Recent years have seen improvements in clinical trial protocols, with many no longer specifying restrictive age criteria. Reasons for poor representation of older patients are complex and ongoing efforts are needed to broaden eligibility criteria and prioritise the inclusion of older adults in clinical trials.

Background

Cancer is a disease of older age, however, historically the older population has not been well-represented in the clinical trials that determine the standards of care for oncological treatment. The relative lack of representation of older patients in clinical trials is well documented [1, 2] with the proportion of participation reported to be as low as 25% [3]. While older patients are more likely to have a poorer performance status or comorbidity that renders them ineligible for trials on the basis of strict enrollment criteria, there is increasing evidence to demonstrate that enrollment of older patients in trials is safe [4]. Clinician bias may also impact the low accrual rate, however older patients appear to have similar

perspectives to younger patients regarding clinical trials [5]. The lack of clinical trial evidence for older patients has generated a significant challenge in translating trial results into clinical practice for a substantial proportion of patients.

Older patients are a heterogeneous cohort, with varying levels of functional status and comorbidities. Historically patients were excluded who were above a certain age or individuals who had a high morbidity burden. Therefore the subjects enrolled in clinical trials, even in the oldest cohort, often don't represent older patients in the general population [2, 4].

Gastroesophageal cancers remain one of the most lethal malignancies, with 5 year survival rates of approximately 20–30% [6]. Typically these cancers affect older patients; 59% and 61% of patients with esophageal and gastric cancer respectively are 65 years or older at diagnosis [7]. Cancers of the upper gastrointestinal tract have the propensity for early dissemination, and 35–38% of cases are metastatic at presentation [7]. The treatment of advanced gastric and oesophageal cancers is similar. In general, combination chemotherapy regimens provide higher response rates than single agents, however, this translates into only modestly longer durations of disease control and survival, typically measured in weeks to short months [8]. For those with HER2 positive disease, the anti-HER2 directed monoclonal antibody trastuzumab has proven benefit [9]. In the second-line setting, a survival benefit of chemotherapy over best supportive care has been demonstrated, although there is no consensus on the optimal regimen [8]. The role of immunotherapy in the treatment of metastatic gastric and oesophageal cancers continues to evolve, with new data showing significant survival benefits, challenging standard chemotherapy and targeted therapies in this setting [10].

Against this background, this study reviews all phase III trials of systemic therapy for advanced gastroesophageal cancer over the last 25 years. We aim to identify whether age restrictive criteria was specified in the trial protocols and thus to determine whether or not the current evidence base is applicable to the majority of patients diagnosed with this disease.

Materials And Methods

Search strategy and study identification:

A search strategy was developed by a health information specialist (M.C.) encompassing the following databases: Embase, PubMed and The Cochrane Library. The population of interest was patients with metastatic gastroesophageal malignancies and the intervention was systemic anti-cancer therapies. Medical Subject Headings (MeSH) terms and text words were identified for these components and linked using the AND operator. The search was filtered for phase III randomised clinical trials published between January 1st 1995 and November 18th 2020. Search terms were reviewed by A.H. and M.A.H. to ensure that the search strategy was comprehensive (for full details of all search terms, see appendix). Additionally, we conducted a search of currently enrolling trials on clinicaltrials.gov to see if there has been any recent improvement in enrolment criteria.

Selection Criteria:

The retrieved articles were imported into EndNote (version X9; Clarivate Analytics) and subsequently exported into a reference management software (Covidence; Veritas Health Innovation Ltd) for study selection. Two reviewers, (M.A.H., M.H.) independently screened study titles and abstracts for eligibility. Studies that were deemed eligible by title and abstract screening then underwent a full-text review by M.A.H. and M.H. using the same criteria. Any conflicts arising from this process were settled by discussion and with the help of a third reviewer (A.H.). Inclusion criteria were as follows: 1) English language; 2) full text available; 3) phase III randomised controlled trials; 4) trials evaluating outcomes for systemic therapies in advanced gastric, oesophageal or gastroesophageal cancer. Studies were excluded for the following reasons: 1) trials conducted in the neoadjuvant or adjuvant setting; 2) trials involving surgery or radiation; 3) no results published.

Data Extraction:

The relevant information from eligible studies was extracted using a standardised template and this process was carried out by two reviewers (M.A.H. and M.H.). The following details were recorded: country of study, date of study onset, date of publication, patient number, age inclusion and exclusion criteria. In addition to the median age of the participants, the percentage of older patients, defined as age ≥ 65 , enrolled in each study was collected where available. Where this information was not reported, it was sought from the corresponding authors via e-mail request. In cases where eligibility criteria were not directly available from the primary publication, the clinicaltrials.gov website was searched for this information.

Statistical Analysis:

The change over time in the proportion of studies using an upper age exclusion was estimated using a linear probability model (i.e. linear regression with a binary outcome coded as [0,1]). The resulting model coefficients were reported with 95% confidence intervals and p-values, with p-values < 0.05 significant. All analyses were conducted using R (version 4.0.3). Figures were produced using ggplot2.

Results

A total of 363 phase III studies of systemic chemotherapy in advanced gastric, oesophageal or gastroesophageal cancer were identified and screened for eligibility. One hundred and fifty two studies were eligible for full text review, 86 were excluded for reasons including duplication, no full text available, study design and non-English language, leaving 66 trials for final analysis (Figure I). Eighty six percent ($n = 56$) included gastric, 9% ($n = 6$) oesophageal and 5% ($n = 3$) oesophagogastric malignancies. Asian studies represented 48% ($n = 32$) of trials included, 29% ($n = 19$) were worldwide and 23% ($n = 15$) were European (Table I).

The median age of trial participants was 62, (range 18–94) (Figure II). Thirty two percent (n = 21) of studies specified an upper age limit for inclusion and 57% of these were Asian studies. In the studies with an upper age limit, the median age of exclusion was 75 years (range 65–80). All studies starting before 2003 used an upper age exclusion (n = 12); whereas only 9 of the 52 that started in 2003 or later did (17%). Among these later studies, there was a very modest downward yearly-trend in the proportion of studies using an upper age exclusion (-0.02 per year; 95%CI -0.05 to 0.01; p = 0.31) (Figure III).

Fifty-two percent (n = 34) of studies specified the proportion of their study population who were over 65 years. Older patients represented only 36% of the trial populations in these studies (range 7–60%). Recruitment of older patients did not appear to change over time. Additionally, we conducted a search of currently enrolling trials on clinicaltrials.gov which yielded 11 active phase III studies in metastatic gastroesophageal cancer, none of which specified an upper age limit.

Discussion

With an ageing population, where 60% of cancer diagnoses are made in patients over the age of 65, the inclusion of older patients in clinical trials is a priority [7, 11]. In our evaluation of phase III trials assessing systemic therapy for advanced gastroesophageal malignancies, 32% of the studies excluded patients based on older age alone. Furthermore, the median age of patients included was just 62, and those aged over 65 made up only 36% of the total study population. Consequently, it is difficult to apply this evidence to our everyday clinical practice, where we frequently encounter older, frailer and more complex patients than the individuals included in these trials. This may lead to both suboptimal treatment of some 'fitter' patients and over-treatment of those who may be more frail than their biological age, resulting in detrimental patient outcomes.

Reasons for poor representation of older adults in clinical trials are complex and multifactorial. They relate to a mix of patient, physician and system factors. There have been a few studies examining patient perspectives and attitudes towards clinical trial participation. Townsley et al. conducted a study focusing on understanding the attitudes of elderly patients with cancer towards clinical trial enrolment [5]. Over 80% of respondents were between the ages of 70 and 79 and the majority of patients stated they would participate in clinical trials to prevent or screen for cancer, to compare a new drug to a 'standard' drug, and 70% would participate in clinical trials to test a new drug in situations where there is no 'standard' drug [5]. However, while most were willing to consider participation when offered, few elderly patients actively sought clinical trials and overall were less well informed regarding the availability of relevant clinical trials [5]. A study by Yellen et al. looked at age and clinical decision-making in oncology patients using clinical vignettes in an interview situation. They found that older patients were as likely as the younger cohort to agree to chemotherapy for both curative and disease control purposes [12]. Ayodele et al. carried out a similar study to compare the attitudes of younger and older patients to clinical trials and found that older patients were as willing as younger patients to participate in clinical trials, yet significantly less were enrolled [13]. In reality, the inclusion of elderly patients is not always straightforward and may not be feasible, due to comorbidities, cognitive issues or social circumstances.

Multiple clinic visits, paperwork and travel to medical appointments are well-documented barriers to trial accrual, and reducing trial participation burden is an area where further progress is needed [11].

Misconceptions among physicians can also act as a barrier to trial enrollment. In a survey of American oncologists, 50% indicated that they declare patients unsuitable for clinical trials based on age alone [14]. Another study examining barriers to clinical trial participation in older women with breast cancer, found that the physicians' perceptions about age and tolerance of toxicity were the greatest obstacle to enrolling older women onto trials [15]. Sedrek et al. carried out semi-structured interviews with 44 medical oncologists (24 academic-based and 20 community-based) in an attempt to explore oncologists' perceptions of barriers to clinical trial enrollment of older adults with cancer. The most common barriers identified by oncologists were stringent eligibility criteria and concerns for treatment toxicities [16]. A better awareness of clinical trials must be promoted amongst healthcare providers in order to increase older adult participation.

Efforts to increase the representation of older adults has been recognised as a priority by a number of international oncological organisations. The International Society of Geriatric Oncology (SIOG) published updated guidance in January 2021 relating to 'Priorities for global advancement of care for older adults with cancer' [17]. This policy document discusses priorities relating to education, clinical practice, research, and collaborations in an attempt to improve healthcare for this rapidly growing patient cohort.

In February 2021, the American Society of Clinical Oncology (ASCO) and the Friends of Cancer Research (Friends) group issued new recommendations to further broaden eligibility criteria for clinical trials, with the aim of expanding patient access [18]. Although the underlying rationale for eligibility criteria is to protect the safety of trial participants and to exclude patients who may have an unacceptably high risk of toxicity, this must be balanced with need to include a representative group of individuals [18, 19]. Interestingly, an analysis conducted from CancerLinQ Discovery® (CancerLinQ's deidentified real-world data product for researchers) found that the number of lung cancer patients potentially suitable for clinical trials almost doubled, when three common eligibility criteria (renal function, presence of brain metastases, history of prior malignancy) were relaxed [20].

In addition to the broadening of eligibility criteria, other suggestions to promote inclusion of older patients in trials focus on addressing the study design, statistical analysis and reporting of trial results. These include assessing more 'age-relevant' endpoints such as functional status and quality of life, as well as the traditional outcomes of progression free survival (PFS) and overall survival (OS). Similarly, expanding treatment-related toxicities to include adverse effects relevant to elderly patients, such as incontinence and falls has been recommended [21, 22, 11]. In terms of reporting of results, it is advised that age-specific subgroup analyses should be powered to detect any age related differences, and in situations where sub-group analysis is not pre-specified, any conclusions should be described as exploratory [23]. Tackling the above issues will help to strengthen and develop our evidence base and allow better decision making for complex, older patients.

Clinical trials can also specifically focus on older adults with cancer and indeed there have been a number of 'elderly-specific' trials in the last few years, highlighting that it is possible to conduct phase III trials in this patient population [24–26]. In the context of gastroesophageal cancer, the GO2 trial was a large phase III study which included 514 older patients with advanced gastroesophageal cancer who were unfit for full dose chemotherapy and aimed to find the optimal dosing strategy. Patients were randomised (1:1:1) to oxaliplatin and capecitabine (xelox) on 3 different dose schedules. The lowest dose demonstrated decreased rates of toxicity and improved quality of life, without shortening survival [26]. These studies illustrate that large randomized studies on older patients are feasible and contribute to creating a body of evidence that guides clinical decision making in this setting.

Overall, there is a slow shift towards including older and multimorbid patients in clinical trials, and certainly the creation of 'elderly-specific' trials as mentioned above is important and encouraging. Interestingly, in our study, when we analysed the patterns of enrollment over the time, we found that age limits were much more common pre-2003. Perhaps this was influenced by the publication by Hutchins et al. in the New England Journal of Medicine in 1999, 'Underrepresentation of patients ≥ 65 in cancer treatment trials.' In this study, data on 16,396 patients enrolled in clinical trials between 1993 and 1996 particularly focusing on sex, race and age, were analysed and compared with rates of cancer in the general population, according to the US Census and the National Cancer Institute [3]. Efforts had been made previously to address the under-representation of women and minority ethnic groups, and the overall proportions of these cohorts were found to be similar. In contrast, patients 65 years or older were dramatically under represented (25% versus 63%, $p < 0.001$) [3]. Since then a number of policies have been developed in an attempt to remedy this. As part of our study, we accessed clinicaltrials.gov and we found that none of the currently active and enrolling phase III clinical trials in advanced gastroesophageal cancer specified an upper age limit for inclusion. It is encouraging that trials have reduced the explicit upper age limits and now we must move to focusing on the other barriers that disproportionately exclude older individuals.

In conclusion, recent years have seen modest improvements in clinical trial protocols, with many no longer specifying restrictive age criteria. With an ageing population there is a growing need to include older, frailer patients who are more reflective of the 'real world' oncology patient in clinical trials. We must advocate for a more inclusive culture and strive to generate the best data and evidence that will allow us to make informed treatment decisions for our patients.

Abbreviations

Medical Subject Headings (MeSH), International Society of Geriatric Oncology (SIOG), American Society of Clinical Oncology (ASCO), Progression Free Survival (PFS), Overall Survival (OS)

Declarations

Ethics approval and consent to participate:

Not applicable

Consent for publication:

Not applicable

Availability of data and material:

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests:

The authors declare that they have no competing interests

Funding:

Not applicable

Authors' contributions:

MAH, AH, LC, NK, MH, AOD contributed to study conception and design, data collection and manuscript preparation. DM, MC contributed to search strategy and acquisition of data. NM and CG contributed to data collection and manuscript preparation. All authors reviewed and approved the final manuscript.

Acknowledgements:

Not applicable

References

1. Talarico L, Chen G, Pazdur R (2004) Enrollment of Elderly Patients in Clinical Trials for Cancer Drug Registration: A 7-Year Experience by the US Food and Drug Administration. *Journal of Clinical Oncology* 22 (22):4626-4631. doi:10.1200/JCO.2004.02.175
2. Scher KS, Hurria A (2012) Under-Representation of Older Adults in Cancer Registration Trials: Known Problem, Little Progress. *Journal of Clinical Oncology* 30 (17):2036-2038. doi:10.1200/JCO.2012.41.6727
3. Hutchins LF, Unger JM, Crowley JJ, Coltman CA, Albain KS (1999) Underrepresentation of Patients 65 Years of Age or Older in Cancer-Treatment Trials. *New England Journal of Medicine* 341 (27):2061-2067. doi:10.1056/NEJM199912303412706
4. Lewis JH, Kilgore ML, Goldman DP, Trimble EL, Kaplan R, Montello MJ, Housman MG, Escarce JJ (2003) Participation of Patients 65 Years of Age or Older in Cancer Clinical Trials. *Journal of Clinical*

5. Townsley CA, Chan KK, Pond GR, Marquez C, Siu LL, Straus SE (2006) Understanding the attitudes of the elderly towards enrolment into cancer clinical trials. *BMC Cancer* 6:34. doi:10.1186/1471-2407-6-34
6. Domper Arnal MJ, Ferrández Arenas Á, Lanas Arbeloa Á (2015) Esophageal cancer: Risk factors, screening and endoscopic treatment in Western and Eastern countries. *World J Gastroenterol* 21 (26):7933-7943. doi:10.3748/wjg.v21.i26.7933
7. Ries LAG EM, Kosary CL, Hankey BF, Miller BA, Clegg L, Mariotto A, Fay MP, Feuer EJ, Edwards BK (eds) SEER Cancer Statistics Review, 1975-2000. https://seer.cancer.gov/csr/1975_2000/, 2003.
8. Wagner AD, Syn NL, Moehler M, Grothe W, Yong WP, Tai B-C, Ho J, Unverzagt S (2017) Chemotherapy for advanced gastric cancer. *Cochrane Database Syst Rev* 8 (8):CD004064-CD004064. doi:10.1002/14651858.CD004064.pub4
9. Bang Y-J, Van Cutsem E, Feyereislova A, Chung HC, Shen L, Sawaki A, Lordick F, Ohtsu A, Omuro Y, Satoh T, Aprile G, Kulikov E, Hill J, Lehle M, Rüschoff J, Kang Y-K, To GATI (2010) Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet (London, England)* 376 (9742):687-697. doi:10.1016/S0140-6736(10)61121-X
10. Zeng Z, Yang B, Liao Z (2021) Progress and prospects of immune checkpoint inhibitors in advanced gastric cancer. *Future Oncol* 17 (12):1553-1569. doi:10.2217/fon-2020-0829
11. Sedrak MS, Freedman RA, Cohen HJ, Muss HB, Jatoi A, Klepin HD, Wildes TM, Le-Rademacher JG, Kimmick GG, Tew WP, George K, Padam S, Liu J, Wong AR, Lynch A, Djulbegovic B, Mohile SG, Dale W, the C, Aging Research G (2021) Older adult participation in cancer clinical trials: A systematic review of barriers and interventions. *CA: A Cancer Journal for Clinicians* 71 (1):78-92. doi:<https://doi.org/10.3322/caac.21638>
12. Yellen SB, Cella DF, Leslie WT (1994) Age and Clinical Decision Making in Oncology Patients. *JNCI: Journal of the National Cancer Institute* 86 (23):1766-1770. doi:10.1093/jnci/86.23.1766
13. Ayodele O, Akhtar M, Konenko A, Keegan N, Calacsan F, Duggan L, O'Connor M, Calvert P, Townsley CA, Horgan AM (2016) Comparing attitudes of younger and older patients towards cancer clinical trials. *Journal of Geriatric Oncology* 7 (3):162-168. doi:10.1016/j.jgo.2016.03.005
14. Benson AB, Pregler JP, Bean JA, Rademaker AW, Eshler B, Anderson K (1991) Oncologists' reluctance to accrue patients onto clinical trials: an Illinois Cancer Center study. *Journal of Clinical Oncology* 9 (11):2067-2075. doi:10.1200/JCO.1991.9.11.2067
15. Kemeny MM, Peterson BL, Kornblith AB, Muss HB, Wheeler J, Levine E, Bartlett N, Fleming G, Cohen HJ (2003) Barriers to Clinical Trial Participation by Older Women With Breast Cancer. *Journal of Clinical Oncology* 21 (12):2268-2275. doi:10.1200/JCO.2003.09.124
16. Sedrak MS, Mohile SG, Sun V, Sun C-L, Chen BT, Li D, Wong AR, George K, Padam S, Liu J, Katheria V, Dale W (2020) Barriers to clinical trial enrollment of older adults with cancer: A qualitative study of

- the perceptions of community and academic oncologists. *Journal of geriatric oncology* 11 (2):327-334. doi:10.1016/j.jgo.2019.07.017
17. Extermann M, Brain E, Canin B, Cherian MN, Cheung K-L, de Glas N, Devi B, Hamaker M, Kanesvaran R, Karnakis T, Kenis C, Musolino N, O'Donovan A, Soto-Perez-de-Celis E, Steer C, Wildiers H (2021) Priorities for the global advancement of care for older adults with cancer: an update of the International Society of Geriatric Oncology Priorities Initiative. *The Lancet Oncology* 22 (1):e29-e36. doi:10.1016/S1470-2045(20)30473-3
 18. Magnuson A, Bruinooge SS, Singh H, Wilner KD, Jalal S, Lichtman SM, Kluetz PG, Lyman GH, Klepin HD, Fleury ME, Hirsch B, Melemed A, Arnaldez FI, Basu Roy U, Schenkel C, Sherwood S, Garrett-Mayer E (2021) Modernizing Clinical Trial Eligibility Criteria: Recommendations of the ASCO-Friends of Cancer Research Performance Status Work Group. *Clinical Cancer Research* 27 (9):2424. doi:10.1158/1078-0432.CCR-20-3868
 19. Kim ES, Bruinooge SS, Roberts S, Ison G, Lin NU, Gore L, Uldrick TS, Lichtman SM, Roach N, Beaver JA, Sridhara R, Hesketh PJ, Denicoff AM, Garrett-Mayer E, Rubin E, Multani P, Prowell TM, Schenkel C, Kozak M, Allen J, Sigal E, Schilsky RL (2017) Broadening Eligibility Criteria to Make Clinical Trials More Representative: American Society of Clinical Oncology and Friends of Cancer Research Joint Research Statement. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 35 (33):3737-3744. doi:10.1200/JCO.2017.73.7916
 20. Harvey RD, Bruinooge SS, Chen L, Garrett-Mayer E, Rhodes W, Stepanski E, Uldrick TS, Ison G, Khozin S, Rubinstein WS, Schenkel C, Miller RS, Komatsoulis GA, Schilsky RL, Kim ES (2021) Impact of Broadening Trial Eligibility Criteria for Patients with Advanced Non–Small Cell Lung Cancer: Real-World Analysis of Select ASCO–&Friends& Recommendations. *Clinical Cancer Research* 27 (9):2430. doi:10.1158/1078-0432.CCR-20-3857
 21. Loh KP, Soto-Perez-de-Celis E, Hsu T, de Glas NA, Battisti NML, Baldini C, Rodrigues M, Lichtman SM, Wildiers H (2018) What Every Oncologist Should Know About Geriatric Assessment for Older Patients With Cancer: Young International Society of Geriatric Oncology Position Paper. *Journal of oncology practice* 14 (2):85-94. doi:10.1200/JOP.2017.026435
 22. Guida JL, Ahles TA, Belsky D, Campisi J, Cohen HJ, DeGregori J, Fuldner R, Ferrucci L, Gallicchio L, Gavrillo L, Gavrillova N, Green PA, Jhappan C, Kohanski R, Krull K, Mandelblatt J, Ness KK, O'Mara A, Price N, Schrack J, Studenski S, Theou O, Tracy RP, Hurria A (2019) Measuring Aging and Identifying Aging Phenotypes in Cancer Survivors. *Journal of the National Cancer Institute* 111 (12):1245-1254. doi:10.1093/jnci/djz136
 23. Zulman DM, Sussman JB, Chen X, Cigolle CT, Blaum CS, Hayward RA (2011) Examining the evidence: a systematic review of the inclusion and analysis of older adults in randomized controlled trials. *J Gen Intern Med* 26 (7):783-790. doi:10.1007/s11606-010-1629-x
 24. Perry JR, Laperriere N, O'Callaghan CJ, Brandes AA, Menten J, Phillips C, Fay M, Nishikawa R, Cairncross JG, Roa W, Osoba D, Rossiter JP, Sahgal A, Hirte H, Laigle-Donadey F, Franceschi E, Chinot O, Golfopoulos V, Fariselli L, Wick A, Feuvret L, Back M, Tills M, Winch C, Baumert BG, Wick W, Ding

- K, Mason WP (2017) Short-Course Radiation plus Temozolomide in Elderly Patients with Glioblastoma. *New England Journal of Medicine* 376 (11):1027-1037. doi:10.1056/NEJMoa1611977
25. Battaglin F, Schirripa M, Buggin F, Pietrantonio F, Morano F, Boscolo G, Tonini G, Lutrino ES, Lucchetti J, Salvatore L, Passardi A, Cremolini C, Arnoldi E, Scartozzi M, Pella N, Boni L, Bergamo F, Zagonel V, Loupakis F, Lonardi S (2018) The PANDA study: a randomized phase II study of first-line FOLFOX plus panitumumab versus 5FU plus panitumumab in RAS and BRAF wild-type elderly metastatic colorectal cancer patients. *BMC Cancer* 18 (1):98-98. doi:10.1186/s12885-018-4001-x
26. Hall PS, Swinson D, Waters JS, Wadsley J, Falk S, Roy R, Tillett T, Nicoll J, Cummings S, Grumett SA, Kamposioras K, Garcia A, Allmark C, Marshall H, Ruddock S, Katona E, Velikova G, Petty RD, Grabsch HI, Seymour MT (2019) Optimizing chemotherapy for frail and elderly patients (pts) with advanced gastroesophageal cancer (aGOAC): The GO2 phase III trial. *Journal of Clinical Oncology* 37 (15_suppl):4006-4006. doi:10.1200/JCO.2019.37.15_suppl.4006

Tables

Table I: Trial characteristics for included phase III studies of systemic anti-cancer therapy in advanced gastric, oesophageal or gastroesophageal cancer

	n (Total n = 66)	%
Geographic Location		
Asian	32	48
European	15	23
Worldwide	19	29
Disease Site		
Gastric	56	86
Gastroesophageal Junction	3	5
Oesophageal	6	9
Line of Treatment		
≥1	42	64
≥2	21	32
≥3	3	4
Year of Publication		
1995-2004	7	12
2005-2014	25	38
2015- to date	34	51
Median Age of Participants	62	(Range 18-94)
Upper Age Restriction		
Yes	21	32
No	43	65
Not Specified	2	3
Median Cut Off Age	75	(Range 65-80)

Figures

Figure 1

Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Flow Diagram

Figure 2

Observed ages of study participants (median and range) by year of study start (n = 66 studies; 1986 to 2017)

Figure 3

Numbers of studies with and without an upper age exclusion

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterialBMCCancer.docx](#)