Microbacterium Sulfonyleureivorans Sp. Nov., Isolated From A Sulfonyleurea Herbicides Degrading Consortium

Qingyun Ma

Institute of Agricultural Resources and regional planning

Delong Kong

Institute of Agricultural Resources and Regional Planning

Qi Zhang

Xinjiang Urumqi Maternal and Child Care Hospital

Miaomiao Li

China Agricultural University College of Geosciences Resources and Environmental Sciences

Xiaoyan Han

Institute of Agricultural Resources and Regional Planning

Juan Che

College of Life Science, Xinjiang Normal University

Yiqing Zhou

Regional Centre for Mapping of Resources for Development

Wei Zhang

College of Life Science, Xinjiang Normal University

Xu Jiang

Institute of Agricultural Resources and Regional Planning

Zhiyong Ruan (✉ ruanzhiyong@caas.cn)

Institute of Agricultural Resources and Regional Planning <https://orcid.org/0000-0002-1302-4832>

Research Article

Keywords: *Microbacterium sulfonyleureivorans*, polyphasic taxonomy, herbicide degradation, plant-growth promoting

Posted Date: July 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-685177/v1>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Archives of Microbiology on January 13th, 2022. See the published version at <https://doi.org/10.1007/s00203-021-02750-4>.

Abstract

A novel Gram-stain positive, aerobic, motile, rod-shaped bacterium, designated strain LAM7116^T, was isolated from a sulfonylurea herbicides degrading consortium enriched with the birch forest soil from Xinjiang. The optimal temperature and pH for the growth of strain LAM7116^T were 35 °C and 7.0, respectively. Strain LAM7116^T could grow in the presence of NaCl up to 4% (w/v). Phylogenetic analysis based on the 16S rRNA gene sequences indicated that strain LAM7116^T was closely related to the members of the genus *Microbacterium*, with the highest similarity to *Microbacterium flavescens* DSM 20643^T (98.1%) and *Microbacterium kitamiense* Kitami C2^T (98.1 %). Strain LAM7116^T formed a distinct subclade with *M. flavescens* DSM 20643^T within the genus *Microbacterium* in the 16S rRNA gene phylogenetic trees. The genomic DNA G+C content of LAM7116^T was 69.9 mol%. The DNA-DNA hybridization (DDH) value between strain LAM7116^T and *M. flavescens* DSM 20643^T was 27.20 %. The average nucleotide identity (ANI) value was 83.96 % by comparing the draft genome sequences of strain LAM7116^T and *M. flavescens* DSM 20643^T. The major fatty acids were anteiso-C_{15:0}, anteiso-C_{17:0}, iso-C_{17:0} and iso-C_{16:0}. The respiratory quinones of strain LAM7116^T were MK-13 and MK-14. The main polar lipids were diphosphatidylglycerol, phosphatidylglycerol, an unidentified lipid and an unidentified glycolipid. Based on the phenotypic characteristics and genotypic analyses, we propose that strain LAM7116^T represents a novel species, for which the name *Microbacterium sulfonylureivorans* sp. nov. was proposed. The type strain is LAM7116^T (=CGMCC 1.16620^T =JCM 32823^T). Strain LAM7116^T secreted auxin IAA and grew well in Ashby nitrogen-free culture medium. Genomic results showed that strain LAM7116^T carried the nitrogenase iron protein (*nifU* and *nifR3*) gene, which indicated that strain LAM7116^T has the potential to fix nitrogen and promote plant growth. At the same time, the strain LAM7116^T can degrade nicosulfuron (a kind of sulfonylurea herbicides) by using glucose as carbon source, microbial degradation of nicosulfuron is the primary process of removing nicosulfuron from environments, and biodegradation is the most effective and environmentally friendly. *Microbacterium sulfonylureivorans* sp. nov. LAM7116^T is a potential candidate for the biofertilizers of organic agriculture areas, having broad prospects for using, and may possess potential to be used in bioremediation of nicosulfuron-contaminated environments.

Introduction

The genus *Microbacterium* includes a very large number of species (over 100) that have been isolated from a wide diversity of environments including plant-associated habitats. At the time of writing, the genus *Microbacterium* consists of 206 species with validly published names in the genus *Microbacterium* recorded on LPSN (<http://www.bacterio.net/microbacterium.html>). For example, this genus have been isolated from various sources, such as foods, freshwater, seawater, soil, sediment, activated sludge, milk products, animals, plants, insects and humans (Collins 1992; Shivaji et al. 2007). Which indicated that they can live in a wide range of habitats. In addition, species in the genus *Microbacterium* have a wide range of physiological and biochemical characteristics from psychrophilic to thermophilic, members of

this genus are Gram-stain-positive, asporogenous and rod-shaped. The colonies of most type species of the genus are yellow, yellow-brown, light yellow, pale yellow or yellow-white with optimum growth temperatures of 20-30 °C. The cell-wall peptidoglycan often contains lysine or ornithine as the diagnostic amino acid (Chen et al. 2016). The genus *Microbacterium* belong to family *Microbacteriaceae*, order *Actinomycetales*, class *Actinobacteria*, the genus *Microbacterium* was first described by Orla-Jensen (Orla-Jensen 1919), based on studies of lactic-acid-producing bacteria, and the description was later emended by Collins et al. (1983) , and the genus redefinition of Takeuchi and Hatano (1998), with the unification of the genus *Microbacterium* and *Aureobacterium*, to include 40 species. It belonging to the class of high GC content *Actinobacteria*, In the latter emendment, the genus *Aureobacterium* was united with the genus *Microbacterium*. To determine the phylogenetic significance of the peptidoglycan type in members of the genus *Microbacterium*. Their cells are generally strictly aerobic except for some species, the cell-wall peptidoglycan and sugars principally contain lysine or ornithine and galactose and rhamnose, respectively, the major menaquinones are unsaturated MK11, MK12, MK13 and/or MK14, polar lipids usually consist of diphosphatidylglycerol, phosphatidylglycerol and one or more glycolipids, cellular fatty acids are predominantly anteiso-C_{15:0}, anteiso-C_{17:0} and iso-C_{16:0} (Minnikin et al. 1979), DNA G+C content range is 61-75 mol% (Artur et al. 2014). Richert et al. (2006) performed phylogenetic analysis of 27 species using four housekeeping genes in addition to 16S rRNA genes. The authors concluded that those species with lysine form a coherent cluster in the tree based on the *ppk* gene and in the concatenated tree and that the genus *Aureobacterium* seems to be embedded within the genus *Microbacterium*.

In this context, we found a novel bacterial strain LAM7116^T, which was isolated from the birch forest soil in Xinjiang. The isolated bacterium is classified as a novel species in the genus *Microbacterium* based on its phylogenetic, chemotaxonomic, and biochemical analysis results.

Materials And Methods

Bacterial Strains, Growth Conditions, and Cultivation

The strain LAM7116^T was isolated from a soil sediment sample taken from Haba river birch forest, located in Kulebai township, Haba river county, Altay region, Xinjiang (GPS coordinates 48° 04' N, 86° 20' E), is the largest naturally growing birch forest belt in northwest China. Five grams of soil sediment was added to 250 mL Erlenmeyer flasks containing 100 mL of modified mineral salts medium (MSM) supplemented with 100 mg/L nicosulfuron as a sole carbon source, and enrichment cultures were incubated on a rotary shaker (160 rpm) at 30 °C in the dark. After incubation for two months, the cultures were called “the degrading consortium”. 2 mL the degrading consortium was suspended in 18 mL sterile water and mixed, the liquid phase was diluted 10⁻⁵ and 100 mL samples were spread onto the surface of each cultivation plate. Aliquots (0.2 mL) of the suspension were plated on Luria-Bertani (LB; BD) agar adjusted to pH 7.0, and the isolation plates were incubated at 35 °C for 5 days. Following incubation, pure cultures of strains LAM7116^T was purified for further characterization based on their low 16S rRNA gene

sequence similarities with known members of the genus *Microbacterium*. The strain was maintained on LB agar and preserved as glycerol suspensions (20%, v/v) at -80°C . The basal growth condition of the strains for all experiments was maintained at pH 7.0 and 35°C , unless otherwise stated.

Phenotypic Characterization tests of LAM7116^T

The strain LAM7116^T morphological characteristics of the isolates were examined after cells were grown on LB medium at 35°C for the initial stationary phase. To standardize the physiological age, growth curve for each strain was tested respectively. The optimum temperature for growth was determined by growing the isolate on LB medium at eight different temperatures (10, 15, 20, 25, 30, 35, 37, 40°C). Growth at various NaCl concentrations was tested over the range 0–10% (w/v) NaCl (at 1 % intervals) by incubating at 35°C . The optimal pH for growth was tested and pH values 4–13 (at 0.5 intervals pH unit) using LB as the basal medium. In the case of pH tests, using the buffer system described by Xu et al. (2005). The pH values of <6, 6–9 and >9 were adjusted by using sodium acetate/acetic acid, Tris/HCl and Na_2CO_3 buffers, respectively (Moraine et al. 2010). Anaerobic growth was examined in serum bottles with cysteine (0.001 g) added to nutrient broth (0.5 % tryptone, 0.1 % yeast extract and 3.4 % sodium chloride) and the upper air layer was replaced with nitrogen. Gram staining was performed using the Gram-stain kit (Meierkolthoff et al. 2013). Cell motility was examined on motility agar (Chen et al. 2007). Catalase activity was determined by observing bubble production in 3% (v/v) hydrogen peroxide solution and oxidase activity was determined using 1% (v/v) tetramethyl-p-phenylenediamine. Cell morphology was observed under a Nikon phase-contrast microscope at $\times 1000$ magnification using cells grown for 48 h at 35°C on LB plates. Endospores were examined according to the Schaeffer–Fulton staining method (Murray et al. 1994). Biochemical characteristics, sugar fermentation tests, enzyme activities, oxidase activity and utilization of carbon and nitrogen sources were performed using the API 20NE, ZYM and 50CH, according to the manufacturer's instructions (bioMérieux). Casein and starch hydrolysis was tested as described by Smibert & Krieg (1994). Hydrolysis of cellulose, gelatin, starch, casein, tyrosine and Tweens 20, 40, 60 and 80, milk coagulation and peptonization, utilization of urea and nitrate reduction were performed as described by Uchida et al. (1994). The type strains of *M. flavescens* DSM 20643^T (= JCM 3877^T) were used as reference strains.

Chemotaxonomic Analysis

The cellular fatty acid is a useful and functional tool to identify species in the genus *Microbacterium* and related genera. For fatty acid analysis, all strains were incubated in LB agar plates and cells were harvested when their optical density reached 70 % of the values to stationary phase in each growth curve. *Microbacterium flavescens* JCM 3877^T was used as a reference strain and was processed under the same conditions. Fatty acid methyl esters were prepared and analysed by using the standard method of the Microbial Identification System and were analysed using a Hewlett Pckard HP 6890 and Microbial Identification software (Sasser et al. 1990). The strain was cultured in LB at 35°C for 48 h, and isoprenoid quinones were extracted and analysis through HPLC (Collins et al. 1977; Dworzanski et al. 1990). The cell walls were prepared by the method of Kawamoto et al. (1981), cell-wall sugars were

determined following procedures described by Stanek & Roberts (1974) and cell-wall characteristic diamino acids were examined using high-speed amino acid analyzer L-8900 (Hitachi). The peptidoglycan was isolated after disruption of the cells by shaking with glass beads and subsequent trypsin digestion, according to the method of Schleifer et al. (1985). Polar lipids were extracted, separated by two-dimensional TLC and identified according to Minnikin et al. (1977). TLC plates were sprayed with various specific reagents for the detection of sugars (α-naphthol/sulfuric acid), phosphates (Zinzadze reagent, Schiff reagent), amino groups (ninhydrin/n-butyl alcohol, Dragendoff's reagent) and total lipids (molybdatophosphoric acid), respectively.

Phylogenetic 16S rRNA Gene Analysis

Genomic DNA was extracted from a single colony of the strain grown on LB plates at 35°C for 48 h. The 16S rRNA gene PCR amplification and sequencing were performed as described by Cai et al. (2011). The 16S rRNA gene was amplified by PCR by using the universal primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTTACCTGTTACGACTT-3') (Lane et al. 1991). The amplification reaction and purification of the product were performed according to Pandey et al. (2002). The identification of phylogenetic neighbors was done by searching against the EzTaxon database (<http://www.ezbiocloud.net/eztaxon>) (Kim et al. 2012). The CLUSTAL_W algorithm was used for sequence alignments using the neighbor-joining (Saitou et al. 1987; Rzhetsky et al. 1992) and maximum-likelihood (Felsenstein et al. 1981) and maximum-parsimony (Miya et al. 2000) methods that were implemented with MEGA 7.0 software for analysis evolutionary status of strains. Evolutionary distance was computed by using the Kimura two-parameter model. The robustness of the tree branches was estimated by bootstrap analysis with 1000 replications (Efron, 1987). The GenBank accession number of 16S rRNA sequence is MG009458.

Genome Sequencing and Analysis

The draft genome sequence of strain LAM7116^T was used to detect genes sequence similarity analysis. The genome was sequenced by Magigene Bioinformatics Technology Co., Ltd. (Guang dong, PR China). Genomes of the most closely related species chosen above were retrieved from the GenBank database in NCBI. Reads of each data set were filtered by using AdapterRemoval SOAPdenovo v1.05 (Mikkel et al. 2016; Li et al. 2010) with default parameters was used to assemble the trimmed reads. The size of the genome was estimated by k-mer analysis (Liu et al. 2013) and genome completeness was determined with CheckM.v1.0.3 (Parks et al. 2015). The genomic DNA for G+C content analysis and DNA–DNA hybridization was prepared according to the protocol of Meier-Kolthoff et al. (2014). The partial genome files were uploaded to the GGDC 2.0 web interface (<http://ggdc.dsmz.de/ggdc.php#>), and Formula 2 was used as recommended for the calculation of DDH values. Average nucleotide identity (ANI) values (<https://ani.jgi.doe.gov/html/calc.php?>) of the total genomic sequences shared between the genomic sequences of strain LAM7116^T with closely related genomic sequences from GenBank were determined according to Goris et al. (2007). As a further extension of genome-based phylogeny, the GGDC website

was used to establish the phylogenomic tree of strain LAM7116^T and other closely related *Microbacterium* species.

Degradation ability of Nicosulfuron by strain LAM7116^T

Several reports have shown that herbicide residues influence the activity and diversity of soil microbes and increase the resistance of weeds, which may endanger food production security due to the phytotoxicity to sensitive crops (Thayer et al. 2012; Regitano et al. 2008). The ability of individual isolates to degrade nicosulfuron in liquid GSM medium (MSM supplemented with 5 g/L glucose) containing 50 mg/L nicosulfuron was analyzed by HPLC, as described by Zhou et al. (2017).

Plant Growth-Promoting Function of LAM7116^T

The functional of secreting plant growth hormone indoleacetic acid (IAA) of strain LAM7116^T was measured by the method described by Glickmann and Dessaux (1995). The qualitative and quantitative analyses of siderophore production were conducted by the method described by Machuca and Milagres (2003). Siderophores are small molecular weight iron chelates secreted by microorganisms under iron stress, which have high affinity for Fe³⁺. Which assist microorganisms to obtain essential iron from the environment. Microorganisms with the ability to synthesize iron carriers can not only improve their own nutritional status by competing for iron, but also supply iron nutrition to plants or achieve biological control by competing with plant pathogens (Barns et al. 2007). By using CAS blue detection plate, qualitative and quantitative methods indentification the ability of strain LAM7116^T to produce iron carrier (Schwyn et al. 1987). The utilization of phosphorus by the strain was determined by organic phosphorus and inorganic phosphorus plates (Robert et al. 1999). All experimental analyses were performed in triplicate to ensure reproducibility. The results were expressed as the mean value of these determinations.

Results And Discussion

Phenotypic Characterization

Strain LAM7116^T showed visible colonies (2 mm in diameter) on LB after 48 h of incubation at 35 °C. Colonies were creamish yellow, circular and opaque with entire edges (Figure S1b). The size of the cells was observed by light microscopy (Figure S1a). Cells were Gram-positive, non-motile, non-spore-forming rods. Interestingly, when the strain was cultured in LB liquid medium, the medium was yellow in the first 48 hours pink in the second 72 hours, but the cell color was still yellow. Strain LAM7116^T differed from *M. flavescens* JCM 3877^T in terms of colony colour, acid production from carbohydrates and a few other biochemical characteristics (Table 1). Strain LAM7116^T catalase activity was positive, but oxidase was negative. The reactions of salicin, d-raffinose, starch and glycogen were positive but the reference strains were negative. And the reactions of d-melezitose and d-turanose were negative but the reference strain was positive. With API 20NE test, strain LAM7116^T was negative for maltose and malic acid but the reference strain was positive. d-Glucose fermentation and β-glucosidase were positive but the reference

was negative. The phenotypic properties differentiating between strain LAM7116^T and its closest phylogenetic neighbors are shown in Table 1.

Isoprenoid Quinones, Polar Lipids and Cellular Fatty Acid

The main isoprenoid quinone of strain LAM7116^T was identified as MK-13 (98.7%) and MK-14 (1.3%). The major polar lipids detected were diphosphatidylglycerol, phosphatidylglycerol and one unidentified glycolipid and one unidentified lipid (Fig. S2). The major cellular fatty acids of strain LAM7116^T were anteiso-C_{15:0} (47.8%), anteiso-C_{17:0} (13.4 %) and iso-C_{17:0} (12.1%), which were consistent with the related species in genus *Microbacterium*, but the presence of anteiso-C_{17:1} (1.0 %) in the two isolates distinguished them from others (Table 2). Cell wall sugars contained galactose and ribose, which indicated that strain LAM7116^T is similar to genus *Microbacterium*, but there are also some differences. Therefore, the strain LAM7116^T is a new member of the genus *Microbacterium*.

Phylogenetic analysis

The 16S rRNA gene sequence (1473bp) was discovered of LAM7116^T. Phylogenetic analysis based on 16S rRNA gene sequences showed that strain LAM7116^T was related most closely to *Microbacterium flavescens* DSM 20643^T (98.1% similarity). As it is generally accepted that 16S rRNA gene sequence similarity less than 98.6% of the new species. Phylogenetic trees were reconstructed using the maximum-likelihood, neighbour-joining, and maximum-parsimony methods. All three treeing methods yielded a similar phylogeny. The phylogenetic relationships of strain LAM7116^T with other closely related species belonging to the genus *Microbacterium* are shown in Figure 1.

Genome Sequencing Analyses

The draft genome of strain LAM7116^T was 379,171,8 bp and N₅₀ value was 214,759,8 bp. The genome was predicted to contain a total of 3591 genes in the plasmid, and gene average length (973 bp). The chromosome encodes 5 rRNA genes. The genomic DNA G+C content of strain LAM7116^T was 69.9 mol%, a value that within the range described for the genus *Microbacterium* [50]. Functional annotation of the genome was performed with a RAST sever (Krishnamurthi et al. 2012; Aziz et al. 2008). Analysis process based on RAST default settings (Overbeek et al. 2014), there were 235 of the genes were associated with carbohydrate metabolism, 190 genes, with metabolism of amino acids, and 122 genes, with metabolism of cofactors and vitamins, 91 genes, with metabolism energy. The genome of strain LAM7116^T was compared with that of *Microbacterium flavescens* DSM 20643^T. The genome of strain DSM 20643^T has been upload to NCBI (accession number JAFIWA000000000).

Base on phylogenomic tree, strain LAM7116^T formed the branch with *M. flavescens* DSM 20643^T. The ANI and DDH value of strain 83.96% and 27.20%, to the genome of the closest species *M. flavescens* DSM 20643^T was below the species cutoff (95% for ANI and 70% for DDH). Therefore, draft genome analysis combined with 16S rRNA phylogenetic, physiological, and biochemical properties all

supported the conclusion that strain LAM7116^T should be considered a novel species in the genus *Microbacterium*.

Biodegradation of Nicosulfuron by strain LAM7116^T

The results suggested that strain LAM7116^T with the excellent nicosulfuron degradation ability, when nicosulfuron was provided at an initial concentration of 50 mg/L, strain LAM7116^T could degrade 69.56 % of the nicosulfuron within 7 days (Figure S10). At the same time, the pH value of the system decreased from 7.0 to 4.24. It was speculated that the degradation process was caused by the acid substance produced by the glucose metabolism of strain LAM7116^T.

Growth Promoting Characteristics

Through the qualitative determination show that strain LAM7116^T secreted auxin IAA by tryptophan. And the color reaction was pink on the white porcelain plate, and at a concentration of 4.97 mg/L (Figure S5). Strain LAM7116^T grow well in nitrogen medium (Ashby) indicated that strain has the ability of nitrogen fixation (Figure S6). Strain LAM7116^T did generate a color ring on the CAS flat plate (Figure S9), but did not a clear zone was observed around each of the colonies of strain LAM7116^T on inorganic or organic phosphate media (Figure S7, S8), which indicated that it could produce siderophores but don't dissolve phosphate.

The results of genome annotation showed that strain LAM7116^T carried the nitrogenase iron protein (*nifU* and *nifR3*) gene, which is a key enzyme for fixing nitrogen in bacteria. The *nifU* gene of LAM7116^T had a relatively low similarity with published sequences based on Blast in Genbank (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). It was mostly related to *Microbacterium* sp. NY27 with a similarity of 82.17%, and the *nifR3* gene of LAM7116^T had a low similarity with published sequences. It was mostly related to *Microbacterium* sp. 4R-513 with a similarity of 88.97%, which indicated that strain LAM7116^T has the potential to fix nitrogen.

Conclusions

Based on the present polyphasic analysis, strain LAM7116^T is considered to represent a novel species within the genus *Microbacterium*, for which we propose the name *Microbacterium sulfonylureivorans* sp. nov. It was found that strain LAM7116^T had multiple ecological functions. It mainly includes the secreted auxin IAA, produce iron carrier and nitrogen fixation. These functions can promote plant growth and resist pathogens. At the same time, the strain LAM7116^T has a certain degradation ability of nicosulfuron, which can be used for remediation of nicosulfuron pollution of environment.

Description of *Microbacterium sulfonylureivorans* sp. nov

Microbacterium sulfonylureivorans (sul.fo.nyl.u.re.i.vo'rans. N.L. fem. n. *sulfonylurea* sulfonylurea; L. v. voro to eat; N.L. part. adj. *sulfonylureivorans* eating sulfonylurea herbicides)

Strain LAM7116^T is Gram-stain-positive, aerobic short rods. non-spore-forming and rod-shaped colonies on LB are circular, smooth and yellowish after 3 days incubation at 35°C. Colonies on LB are circular to slightly irregular and light brown after incubation at 35°C for 3 days. The temperature range for growth is from 15 to 40°C (optimum: 35°C). The pH range for growth is from 5.5 to 11.5 (optimum: 7.0). The NaCl concentration range for growth is from 0 to 4.0 % (optimum: 2.0 %). Oxidase-negative and catalase-positive, methyl red-positive. Tween 20 and gelatin are hydrolysed. Negative for hydrolysis of Tweens 40, 60 and 80, starch and casein. In the API ZYM system, strain LAM7116^T is found to be negative for alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase, α -glucosidase and *N*-acetyl- β -glucosaminidase, weak for lipase (C14), cystine arylamidase and naphthol-AS-BI-phosphohydrolase, negative for trypsin, chymotrypsin, α -galactosidase, β -galactosidase, β -glucosidase, β -glucuronidase, α -mannosidase and α -fucosidase. In the API 20NE system, strain LAM7116^T is found to be positive for d-glucose fermentation, β -glucosidase, β -galactosidase, glucose, mannitol, *N*-acetyl-glucosamine, gluconate, negative for nitrate reduction, maltose, indole production, arabinose, arginine dihydrolase, urease, arabinose, capric acid, adipic acid, phenylacetic acid, malic acid and citrate acid, weak for gelatinase. In the API 50CH system, strain LAM7116^T is found to be positive for d-glucose, d-fructose, d-mannitol, aesculin, salicin, d-cellobiose, d-maltose, d-sucrose, d-raffinose, starch, glycogen. l-arabinose, d-xylose, d-galactose, l-rhamnose, d-mannitol, amygdalin, arbutin, d-melibiose, d-trehalose (weakly) and glycogen, negative for others. The predominant respiratory quinone are MK-13 and MK-14. The main amino acids are glycine, lysine, alanine and glutamic acid are detected in cell-wall preparations of strain LAM7116^T. The acyl type of the peptidoglycan is glycolyl, the whole-cell-wall sugars are galactose and ribose. The major fatty acids are anteiso-C_{15:0}, anteiso-C_{17:0}, iso-C_{17:0} and iso-C_{16:0}. The major polar lipids are diphosphatidylglycerol, phosphatidylglycerol, an unidentified glycolipid and an unidentified lipid.

The type strain is LAM7116^T (= CGMCC 1.16620^T = JCM 32823^T), which is isolated from a sulfonylurea herbicides degrading consortium enriched with the birch forest soil, and the soil collected from Xinjiang County, Xinjiang Province, China. The DNA G + C content is 69.9 mol%. Based on phylogenetic, biochemical, chemotaxonomic, and genome analyses, the strain LAM7116^T is considered a novel species within the genus *Microbacterium*. The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain LAM7116^T is MG009458. The GenBank accession number for the draft genome sequence of strain LAM7116^T is RJAD00000000.

Declarations

Author Contributions: Q.Y.M., D.L.K., study design; Q.Z, and M.M.L.; conceived and supervised the study; Q.Y.M.; analyzed the data; X.Y.H., J.C., prepared the figures; Y.Q.Z., J.K., wrote the manuscript; Z.Y.R., and

W.Z.; edited the manuscript and reviewed the literature; X.J.; manuscript revision. All authors have read and agreed to the published version of the manuscript.

Funding information: This research was supported by the National Natural Science Foundation of China (NSFC 32070004, 31670006). This work was also supported by Central Public interest Scientific Institution Basal Research Fund (No. Y2021GH18).

Conflicts of interest: The authors declare that there are no conflicts of interest.

References

1. Artur A, António C, José MI, Martha ET (2014) *Microbacterium endophyticum* sp. nov. and *Microbacterium halimionae* sp. nov., endophytes isolated from the salt-marsh plant *Halimione portulacoides* and emended description of the genus *Microbacterium*. *Systematic and Applied Microbiology* 37:474-479
2. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA et al (2008) The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9:75-75
3. Barns SM, Cain EC, Sommerville L, Kuske CR, Barns SM (2007) *Acidobacteria* phylum sequences in uranium-contaminated subsurface sediments greatly expand the known diversity within the phylum. *Applied and Environmental Microbiology* 73:3113-3119
4. Cai M, Wang L, Cai H, Li Y, Wang Y, Tang Y, et al (2011) *Salinarimonas ramus* sp nov and *Tessaracoccus oleiagri* sp nov., isolated from a crude oil-contaminated saline soil. *Int J Syst Evol Microbiol* 61:1767-1774
5. Chen X, Li G, Li Q, Xu F, Jiang C, Han L, et al (2016) *Microbacterium faecale* sp. nov., isolated from the faeces of *Columba livia*. *Int J Syst Evol Microbiol* 66:11
6. Chen Y, Cui X, Pukall R, Li H, Yang Y, Xu L, et al (2007) *Salinicoccus kunmingensis* sp. nov., a moderately halophilic bacterium isolated from a salt mine in Yunnan, south-west China. *Int J Syst Evol Microbiol* 57:2327-2332
7. Collins MD, Jones D, Kroppenstedt RM (1983) Reclassification of *Brevibacterium imperiale* (Steinhaus) and "*Corynebacterium laevaniformans*" (Dias and Bhat) in a Redefined Genus *Microbacterium* (Orla-Jensen), as *Microbacterium imperiale* comb. nov. and *Microbacterium laevaniformans* nom. rev.; comb. nov. *Systematic and Applied Microbiology* 4:65-78
8. Collins M, Pirouz T, Goodfellow M, Minnikin D (1977) Distribution of Menaquinones in *Actinomycetes* and *Corynebacteria*. *Journal of General Microbiology* 100:221
9. Collins MD (1992) The genera *Agromyces*, *Aureobacterium*, *Clavibacter*, *Curtobacterium*, and *Microbacterium*. *The Prokaryotes*
10. Dworzanski JP, Berwald L, Meuzelaar HL, Dworzanski JP (1990) Pyrolytic methylation-gas chromatography of whole bacterial cells for rapid profiling of cellular Fatty acids. *Applied and Environmental Microbiolog* 56:1717

11. Efron B (1987) Better bootstrap confidence intervals. J Am Stat Assoc 82:171–185
12. Felsenstein J (1981) Evolutionary trees from DNA sequences: A maximum likelihood approach. J Mol Evol 17:368–376
13. Glickmann E, Dessaux Y (1995) A critical examination of the specificity of the salkowski reagent for indolic compounds produced by phytopathogenic bacteria. Applied and Environmental Microbiology 61:793-799
14. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T (2007) Dna-dna hybridization values and their relationship to whole-genome sequence similarities. Int J Syst Evol Microbiol 57
15. Kawamoto I, Oka T, Nara T (1981) Cell wall composition of *Micromonospora olivoasterospora*, *Micromonospora sagamiensis*, and related organisms. J Bacteriol 146:527–534
16. Kim OS, Cho YJ, Lee K, Yoon SH, Kim M, Na H, Park SC, Jeon YS, Lee JH, Yi H, et al (2012) Introducing EzTaxon-e: A prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. Int J Syst Evol Microbiol 62:716–721
17. Krishnamurthi S, Bhattacharya A, Schumann P, Dastager SG, Tang SK, Li WJ, Chakrabarti T (2012) *Microbacterium immunditiarum* sp. nov., an actinobacterium isolated from landfill surface soil, and emended description of the genus *Microbacterium*. Int J Syst Evol Microbiol 62:2187-2193
18. Lane DJ (1991) 16S/23S rRNA Sequencing. In Nucleic Acid Techniques in Bacterial Systematics; Stackebrandt, E., Goodfellow, M., Eds.; John Wiley and Sons: New York, NY, USA, pp. 115–175
19. Li R, Zhu H, Ruan J, Qian W, Fang X, Shi Z, et al (2010) De novo assembly of human genomes with massively parallel short read sequencing. Genome Res 20:265
20. Liu B, Shi Y, Yuan J, Hu X, Zhang H, Li N, et al (2013) Estimation of genomic characteristics by analyzing k-mer frequency in de novo genome projects
21. Machuca A, Milagres A (2003) Use of CAS-agar plate modified to study the effect of different variables on the siderophore production by *Aspergillus*. Journal of Applied Microbiology 3:177-181
22. Meier-Kolthoff JP, Klenk HP, Göker M (2014) Taxonomic use of DNA G+C content and DNA-DNA hybridization in the genomicage. Int J Syst Evol Microbiol 64:352–356
23. Meierkolthoff JP, Auch AF, Klenk HP, Markus G (2013) Genome sequence-based species delimitation with confidence intervals and improved distance functions. BMC Bioinform 14:60
24. Mikkelsen SB, Stinus L, Ludovic O (2016) AdapterRemoval v2: rapid adapter trimming, identification, and read merging. BMC Research Notes 9:88
25. Minnikin DE, Collins MD (1979) Goodfellow, M. Fatty acid and polar lipid composition in the classification of *Cellulomonas*, *Oerskovia* and related taxa. J Appl Bacteriol 47:87–95
26. Minnikin DE, Patel PV, Alshamaony L, Goodfellow M (1977) Polar Lipid Composition in the Classification of *Nocardia* and Related Bacteria. Int J Syst Bacteriology 27(2):104
27. Miya M, Nishida M (2000) Use of mitogenomic information in teleostean molecular phylogenetics: a tree-based exploration under the maximum-parsimony optimality criterion. Molecular Phylogenetics Evolution 17:437-455

28. Moraine RA, Rogovin P (2010) Kinetics of Polysaccharide B-1459 Fermentation. *Biotechnology Bioengineering* 8:511-524
29. Murray R, Doetsch R, Robinow C (1994) Determinative and cytological light microscopy
30. Orla-Jensen S (1919) *The Lactic Acid Bacteria*. Copenhagen: Høst & Sons
31. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, et al (2014) The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:206
32. Pandey KK, Mayilraj S, Chakrabarti T (2002) *Pseudomonas indica* sp nov., a novel butane-utilizing species. *Int J Syst Evol Microbiol* 52:1559
33. Parks D, Imelfort M, Skennerton C, Hugenholtz P, Tyson G (2015) CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Research* 25:1043-1055
34. Regitano JB, Koskinen WC (2008) Characterization of nicosulfuron availability in aged soils. *J Agric Food Chem* 56:5801– 5805
35. Richert KE, Brambilla E, Stack E (2006) The phylogenetic significance of peptidoglycan types: Molecular analysis of the genera *Microbacterium* and *Aureobacterium* based upon sequence comparison of gyrB, rpoB, recA and ppk and 16S rRNA genes. *Systematic and Applied Microbiology* 30:102-108
36. Robert EC (1999) Chapter 1 Introduction: Nature, occurrence and functioning of plant hormones. *New Comprehensive Biochemistry* 33
37. Rzhetsky A, Nei M (1992) A simple method for estimating and testing minimum evolution trees. *Mol Biol Evol* 9:945–967
38. Saitou N, Nei M (1987) The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
39. Sasser M (1990) Identification of bacteria by gas chromatography of cellular fatty acids. *USFCC News* 20:16
40. Schleifer KH (1985) Analysis of the Chemical Composition and Primary Structure of Murein. In: Bergan T, editor. *Methods in Microbiology: Academic Press* 123
41. Schwyn B, Neilands JB (1987) Universal chemical assay for the detection and determination of siderophores. *Anal Biochem* 160:47
42. Shivaji S, Bhadra B, Rao RS, Chaturvedi P, Pindi PK, Raghukumar C, et al (2007) *Microbacterium indicum* sp. nov., isolated from a deep-sea sediment sample from the Chagos Trench, Indian Ocean. *Int J Syst Evol Microbiol* 57:1819-1822
43. Smibert RM, Krieg NR (1994) Phenotypic characterization. In *methods for general and molecular bacteriology* 607-654
44. Stanek JL, Roberts GD (1974) Simplified approach to identification of aerobic actinomycetes by thin-layer chromatography. *Appl Microbiol* 28:226-232

45. Takeuchi M, Hatano K (1998) Union of the genera *Microbacterium* Orla-Jensen and *Aureobacterium* Collins et al. in a redefined genus *Microbacterium*. *Int J Syst Bacteriol* 48:739-47
46. Thayer KA, Heindel JJ, Bucher JR, Gallo MA (2012) Role of environmental chemicals in diabetes and obesity: a national toxicology program workshop review. *Environ. Health Perspect* 120:779–789
47. Uchida K, Kudo T, Suzuki K, Nakase T (1999) A new rapid method of glycolate test by diethyl ether extraction, which is applicable to a small amount of bacterial cells of less than one milligram. *Journal of General and Applied Microbiology* 45:49-55
48. Xu P, Li WJ, Tang SK, Zhang YQ, Chen GZ, Chen HH, Xu LH, Jiang CL (2005) *Naxibacter alkalitolerans* gen. nov., sp. nov., a novel member of the family “Oxalobacteraceae” isolated from China. *Int J Syst Evol Microbiol* 55:1149
49. Zhou S, Song J, Dong W, Mu Y, Zhang Q, Fan Z, et al (2017) Nicosulfuron Biodegradation by a Novel Cold-Adapted Strain *Oceanisphaera psychrotolerans* LAM-WHM-ZC. *J Agric Food Chem* 65:10243-10251

Tables

Table 1. Differential characteristics of strain LAM7116^T and its relative reference strains in the genus *Microbacterium*. Strains: 1, LAM7116^T; 2, *Microbacterium flavescens* JCM 3877^T. All strains were cultivated with the same medium and growth conditions. In the Biolog system both strains were symbols: +, positive; -, negative; w, weakly reaction. All are positive for catalase and negative for oxidase, alkaline phosphatase, esterase (C4), esterase lipase (C8), lipase (C14), valine arylamidase, cystine arylamidase, trypsin, chymotrypsin, acid phosphatase and α-galactosidase, β-galactosidase, β-glucuronidase β-glucosidase, N-acetyl-β-glucosaminidase, α-mannosidase, α-fucosidase, inulin, xylitol and d-gentiobiose are negative. Nitrate reduction, indole production, urease, arginine dihydrolase, sorbitol, methyl-α-d-mannopyranoside, N-acetyl-glucosamine, d-ribose, d-lactose, d-lyxose, d-tagatose, d-fucose, l-fucose, d-arabitol, l-arabitol, potassium gluconate, 2-ketogluconate, 5-ketogluconate are all negative. β-galactosidase, glucose and gluconate are positive. Capric acid, adipic acid phenylacetic acid, and glycerol, erythritol, d-arabinose l-xylose, d-adonitol, methyl-β-d-xylopyranoside, dulcitol and inositol are negative. d-fructose, d-mannose, l-sorbose, mannitol, d-cellobiose and d-maltose are all positive. d-galactose, d-mannitol and arbutin are all weakly reaction.

Characteristic	1	2
Temperature range for growth (°C)	15-40	20-35
Methyl red	+	-
Enzyme activities:(API ZYM)		
Naphthol-AS-BI-phosphohydrolase	-	+
α -Glucosidase	-	w
API 20NE		
d-Glucose fermentation	+	-
β -Glucosidase	+	-
Gelatinase	w	+
Mannose	w	+
N-Acetyl-glucosamine	+	w
Maltose	-	+
Malic acid	-	+
Citrate	-	w
API 50CH		
l-Arabinose	w	-
d-Xylose	w	-
d-Glucose	+	w
l-Rhamnose	w	-
Methyl- α -d-glucopyranoside	-	w
Amygdalin	w	-
Aesculin	+	w
Salicin	+	-
d-Melibiose	w	-
d-Sucrose	+	w
d-Trehalose	w	+
d-Melezitose	-	+
d-Raffinose	+	-

Starch	+	-
Glycogen	+	-
d-Turanose	-	+
DNA G+C content (mol%)	69.9	66.9*

*Data from Takeuchi et al. (1998).

Table 2. Cellular fatty acid contents of strain LAM7116^T and its relative reference strains in the genus *Microbacterium*.

Strains: 1, LAM7116^T; 2, *Microbacterium flavescens* JCM 3877^T. All the data were obtained from this study. All strains were cultivated with the same medium and growth conditions. Values were percentages of total fatty acids. tr, Trace amount (<0.5 %); nd, not detected. The data marked in bold represented the major fatty acids (>10 %) of all strains.

Fatty acids	1	2
C _{16:0}	0.9	0.8
iso-C _{14:0}	1.0	0.7
iso-C _{15:0}	9.0	14.8
iso-C_{16:0}	10.8	14.7
iso-C _{17:0}	12.1	5.9
iso-C _{18:0}	1.9	tr
anteiso-C_{15:0}	47.8	37.4
anteiso-C_{17:0}	13.4	22.6
anteiso-C _{17:1} ω 9c	nd	0.8
anteiso-C _{17:1} A	1.0	nd

*Summed features represent groups of two or more fatty acids that could not be separated using the MIDI system. Summed feature 1 contained iso-C_{15:1} H/ C_{13:0} 3-OH, summed feature 2 contained C_{14:0} 3-OH and/or iso-C_{16:1} I, summed feature 3 contained C_{16:1} ω 7c and/or C_{16:1} ω 6c and summed feature 8 contained C_{18:1} ω 7c and/or C_{18:1} ω 6c.

Figures

Figure 1

Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the relationships between strain LAM7116T and the type strains of all recognized *Microbacterium* species. Bootstrap values as a percentage of 1000 replications are shown at nodes; only values >70% are shown. The sequence from *Actinoplanes abujensis* was used as an outgroup. Filled circles indicate nodes also recovered by using the maximum-parsimony method. Bar, 0.02 substitutions per site.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementarymaterials.pdf](#)