Trends Over Two Decades in Life Expectancy With Complex Health Problems Among Older Swedes: Implications for The Provision of Integrated Medical and Social Care

Bettina Meinow (✉ bettina.meinow@aldrecentrum.se)

Aging Research Center, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet and Stockholm University, Solna

Peng Li

Max Planck Institute for Demographic Research, Rostock

Domantas Jasilionis

Max Planck Institute for Demographic Research, Rostock

Anna Oksuzyan

Max Planck Institute for Demographic Research, Rostock

Louise Sundberg

Aging Research Center, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet and Stockholm University, Solna

Susanne Kelfve

Division of Ageing and Social Change, Department of Social and Welfare Studies, Linköping University, Linköping

Jonas W Wastesson

Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Solna

Research Article

Keywords: Sullivan method, health expectancies, complex health problems, multimorbidity, multi-domain health measures, integrated care

Posted Date: September 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-689257/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Due to population aging, it is essential to examine to what extent rises in life expectancy (LE) consist of healthy or unhealthy years. Most health expectancy studies have been based on single health measures and have shown divergent trends. We used a multi-domain indicator, complex health problems (CHP), indicative of the need for integrated medical and social care, to investigate how LE with and without CHP developed in Sweden between 1992 and 2011. We also addressed whether individuals with CHP more commonly lived in the community in 2011 compared to earlier years.

Methods

CHP were defined as having severe problems in at least two of three health domains related to the need for medical and/or social care: symptoms/diseases, cognition/communication, and mobility. The Swedish Panel Study of Living Conditions of the Oldest Old (SWEOLD), a nationally representative survey of the Swedish population aged ≥ 77 years with waves in 1992, 2002 and 2011 ($n \approx 2000$), was used to estimate the prevalence of CHP. Age- and gender-specific death rates were obtained from the Human Mortality Database. The Sullivan method was deployed to calculate the remaining life expectancy with and without CHP. The estimates were decomposed to calculate the contribution of changes from morbidity and mortality to the overall trends in LE without CHP.

Results

Between 1992 and 2011, both total LE (+ 1.69 years [95% CI 1.56;1.83] and LE without CHP (+ 0.84 years [-0.87;2.55]) at age 77 increased for men, whereas LE at age 77 increased for women (+ 1.33 [1.21;1.47]) but not LE without CHP (-0.06 years [-1.39;1.26]). When decomposing the trend, we found that the increase in LE with CHP was mainly driven by an increase in the prevalence of CHP. Among individuals with CHP the proportion residing in residential care was lower in 2011 (37%) compared to 2002 (58%) and 1992 (53%).

Background

Life expectancy is increasing internationally and most deaths in low-mortality countries now occur around the ages of 85 to 90 years [1] and are seldom sudden [2, 3]. Deaths after age 80 are often preceded by a period with chronic conditions, functional loss, and dependency [2, 4–6]. For example, many age-related conditions are not curable but can be managed for years [5, 7, 8]. The extent to which longer lives are accompanied by a compressed, postponed, or expanded period with health problems and care needs has broad ramifications for individuals and families, as well as for financing and organizing appropriate medical and social care.

In international comparison, Sweden has a comprehensive, universal, and largely tax-financed system for social care for older people. Social care is available for all inhabitants aged 65 years or older whose need of support with personal care and/or household chores is approved by municipal need assessors. For decades, *“aging in place”*, i.e., supporting older people in their own homes, instead of in residential care, has been a key and guiding strategy in addressing older adults’ care needs [9]. Most people enter the social care system through home care, as residential care is only approved if more intense home care would still be insufficient. Yet, during the past two decades, due to economic constraints, the *“aging in place”* policy has been further strengthened, involving a drastic reduction of beds in residential care, as well as hospital beds [9]. The downsizing of residential care entailed a shorter length of stay in recent years, with a rapid increase in the proportion of people who moved in and died shortly afterwards [10, 11].

If older people survive longer with extensive and complex care needs, and if they also to a greater extent remain in their own homes, despite such comprehensive care needs, this development will reinforce challenges for the provision of well-coordinated and coherent medical and social care in the community.

To address the question to what extent increased life expectancy entails healthy and unhealthy years, the concept of health expectancies has been developed and estimates the interaction of mortality and morbidity [12]. Divergent time trends in health expectancies have been reported across countries, subgroups of populations (e.g., regarding gender, age, and ethnic groups), time periods and the health indicator used [13]. The Global Burden of Disease Study 2016 provided healthy life expectancy estimates for 195 countries and territories between 1990 and 2016 [13]. At a global level, healthy life expectancy continued to show improvements, although many populations were facing an increase in time with functional limitations [13]. Most health expectancy studies that have suggested a trend towards shorter time spent with poor health at the end of life – a compression of morbidity – have used health indicators based on disability (e.g., activities of daily living, ADLs) [14–18] or cognitive impairment [19–22]. This trend has been attributed to decreasing rates of disability among more recent cohorts of older people, especially those under the age of 85 [14, 23, 24].

In contrast, health expectancy studies finding support for an expansion of morbidity during the last two decades have frequently focused on chronic diseases/conditions [14, 22, 25–27]. That more people are getting diagnosed at an earlier stage in later cohorts has probably contributed to this development.

However, while trends in health expectancies for specific single health indicators have developed differently, health problems often exist simultaneously and are interrelated in old age [28, 29]. Hence, in this study we use a composite health measure including both chronic diseases/conditions and functional limitations to address the remaining lifetime with and without combined medical conditions and functional limitations. Composite health measures, including both medical conditions and functional limitations, may to a larger extent reflect the broader impact of health problems experienced by individuals and their families [29]. From a policy and planning perspective, such indicators can be useful to reflect the need for services that are organized to meet multiple complex care needs, including both medical care and social care [30].

Yet, only a few studies have analyzed years lived with multiple morbid conditions and most of them focus on multimorbidity (the presence of two or more diseases) or frailty (a state that is characterized by cumulative decline in many physiological systems, vulnerability to physiologic stressors and risk of adverse outcomes) [31, 32]. Both concepts rely primarily on clinical data, such as diagnosed diseases and physiological markers. Therefore, they may indicate a need for medical care or a risk for *subsequent* disability and associated social care needs. Yet, neither multimorbidity nor frailty capture *actual* needs for combined medical care and social care [29]. One exception is a study based on the British Cognitive Function and Ageing Studies that integrated geriatric conditions and dependency in activities of daily living (ADL) into a measure of dependency [33]. Results showed that years lived with dependency from age 65 increased between 1991 and 2011 [33].

Based on representative samples of the Swedish population aged 77 or older, we have previously developed a composite measure of complex health problems (CHP) that reflects the simultaneous presence of medical conditions, and physical and cognitive functional limitations, implying a need for both medical care and social care [29, 34]. The presence of CHP has also been shown to be highly predictive of 4-year mortality [35]. The objective of this study was to investigate how life expectancy with and without CHP has developed between 1992 and 2011 in Swedish men and women aged 77 or older. We also addressed whether it has become more common among individuals with CHP to live in the community instead of in residential care.

Methods

Data

Prevalence of CHP was based on three waves (1992, 2002 and 2011) of the Swedish Panel Study of Living Conditions of the Oldest Old (SWEOLD), which covers representative samples of the Swedish population aged 77 or older (n=2016) [36]. The SWEOLD samples are nationally representative with respect to the distribution of age, gender, geography and proportion of people living in residential care. Age- and sex- specific death rates for the survey years were obtained from the Human Mortality Database [37].

In 2011, 85- to 99-year-olds were oversampled to increase the precision of measurements in the oldest age groups. Sample weights have been applied to adjust for this oversampling [36]. Response rates across the survey waves were high (between 86% and 90%) due to intensive fieldwork that was designed to include very frail and cognitively impaired older persons. Whenever possible, direct face-to-face interviews were conducted in the respondent's dwelling. Proxy interviews were carried out with a relative or caregiver if the respondent was too frail and/or cognitively impaired to participate.

Health dimensions and indicators

The previously developed measure of CHP was based on three health domains that are related to the need for different types of care services: symptoms/diseases, cognition/communication and mobility [34,

38]. Typically, symptoms and diseases imply need for medical care. Functional limitations, such as mobility problems are mainly handled by social services and/or informal caregivers. Serious limitations in cognitive/communication skills often require help from medical and social care providers. Persons with serious problems in at least two of the three health domains were considered to have CHP [34].

To provide comparable and nationally representative estimates, we used measures that were available for the complete sample, including proxy interviews. This excluded subjective evaluations of health and tests of function (except for the cognition test, see below). Severe problems in each of the health domains were identified as briefly described below, while details can be found elsewhere [34, 38].

Symptoms/diseases: Respondents were asked whether they have had any of 14 common symptoms and diseases during the previous 12 months. For each item, mild problems were coded as 1 and severe problems were coded as 3. This resulted in a summed index that ranged from 0 to 42. A cut-off for severe problems for this domain was set at the highest quintile for the 1992 sample (corresponding to 9 on the summed scale) [34]. Accordingly, persons belonging to the highest quintile had, e.g., at least three severe symptoms/diseases or two severe and three mild. The same cut-off was used for the survey waves in 2002 and 2011.

Mobility: This domain was an index of four items: the reported ability to walk 100 meters fairly briskly without difficulties, walk up and down stairs, rise from a chair without difficulty, and stand without support. Limitations in at least three of the four activities were coded as serious mobility problems.

Cognition/communication: Cognition was assessed using an abridged version of the Mini-Mental-State Examination [39] that has been validated against the full MMSE scale and clinical dementia diagnoses [40]. Interviewer notes showed that almost all proxy-interviews were due to cognitive problems or, in a few cases, aphasia. Respondents who could not be interviewed directly, scored below the cutoff (< 7 out of 11) in the test or did not do the test were coded as having serious cognitive/communication problems.

Use of social care: The interviewer noted whether an individual lived in residential care or not. Respondents were asked whether they received home care or not.

Statistical analyses

The Sullivan method was used to estimate life expectancy at age 77 with and without CHP for women and men correspondingly [41]. Age-, gender- and period-specific prevalence of CHP was combined with standard period life tables. Age-specific prevalence of CHP was smoothed using a logistic regression model with linear and quadratic terms of age [42]. Using a decomposition method by Andreev et al. [43], the changes in life expectancy with and without CHP between the survey years were decomposed into the contributions a) attributable to mortality change and b) the contributions attributable to changes in the prevalence of CHP. Microsoft Excel and R (version 3.5) were used for life table analysis, smoothing, and decomposition analysis. STATA SE15 was used to obtain prevalence of health problems, tests for

statistical significance and 95% confidence intervals (95% CI). All statistical tests were two-sided; the differences in prevalence between the selected time points were statistically significant if $p < 0.05$.

Results

The proportion of people with severe problems increased in each of the three health domains between 1992 and 2011 among Swedes aged 77 or older (Table 1). The most distinct increase, from 20.5% percent to 34.2% ($p < 0.001$), occurred for the severe symptoms/diseases domain. Severe mobility limitations increased from 19.5–24.4% ($p < 0.05$). The slight increase of cognition/communication problems from 25.7–29.8% was not statistically significant ($p = 0.116$). The proportion of the population with CHP, i.e., severe problems in two or three health domains, increased by 41%, from 18.2% in 1992 to 25.6% in 2011 ($p < 0.01$). As for the three health domains separately, the prevalence of CHP increased mainly during the first 10-year period and remained stable thereafter. Patterns were similar for men and women, although women generally had more health problems (not shown).

Table 1

Sample characteristics and prevalence of severe problems in three health domains 1992, 2002 and 2011

	1992 n = 537	2002 n = 621	2011^a n = 831	p-value^b 1992– 2002	p-value^b 1992– 2011
Proxy interview	11.7	13.2	17.8	0.414	0.003
Females	60.8	59.3	62.6	0.581	0.528
Institutional care	18.6	22.5	11.7	0.088	0.003
Age range	77–98	77–99	77–101		
Mean/median age	82.4/82.0	83.3/83.0	83.4/83.0	0.001 ^c	< 0.001 ^c
Multiple severe symptoms/diseases	20.5	32.3	34.2	< 0.001	< 0.001
Severe mobility limitations	19.5	26.0	24.4	0.008	0.048
Poor cognition/ Communication	25.7	33.3	29.8	0.003	0.116
No serious problems in any domain	56.4	41.7	45.1	< 0.001	< 0.001
Serious problems in one domain	25.5	32.1	29.3	0.011	0.146
Serious problems in 2–3 domains	18.1	26.1	25.6	0.001	0.002
Notes: Between 2002 and 2011, the prevalence of severe problems did not change significantly in any of the health domains. ^a Percentages were weighted for an oversampling (n = 257) of persons aged 85 years or older in the 2011 survey. ^b based on chi2 test. ^c p-value refers to t-test of mean differences. Data: The Swedish Panel Study of Living Conditions of the Oldest Old 1992, 2002, 2011. Own calculations.					

Use of social care

Between 1992 and 2002, the proportion of individuals with CHP who lived in residential care increased slightly from 53–58%, whereas the proportion who used home care slightly decreased. During the following decade, among individuals with CHP, the proportion living in residential care dropped to 37%, while the proportion with home care in the same group remained constant. The proportion of individuals with CHP who did not use any social care services more than doubled, from 16–38%. Few individuals without CHP lived in residential care.

Life expectancy with and without complex health problems

Between 1992 and 2011, remaining life expectancy at age 77 increased by 1.69 years for men (from 8.08 to 9.77 years) and by 1.33 years for women (from 10.38 years to 11.71 years). For both genders and at all

time points, most of the remaining life expectancy consisted of years without CHP (Table 2).

For men, half of the additional expected life years (0.85 years) consisted of years with CHP. While the increase of years with CHP mainly occurred during the first half of the period (1992–2002), years without CHP (0.84) increased essentially during the second half (2002–2011).

Among women, the entire increase of expected life years (1.33) consisted of years with CHP. Life expectancy without CHP decreased by 0.64 years between 1992 and 2002 and then increased again by 0.58 years between 2002 and 2011, resulting in almost no change over the 19-year period. This resulted in an increase of years with CHP for men and women, and more so for women, who could expect to live with CHP about twice as long as men at all three time points.

Table 2

Remaining life expectancy at age 77 and estimated remaining life expectancy with and without complex health problems, by gender and survey year

	1992	2002	2011	Diff 2011 – 1992
Men				
LE	8.08 (8.01–8.14)	8.78 (8.72–8.84)	9.77 (9.70–9.84)	1.69
LE without CHP	6.78 (5.83–7.73)	6.76 (5.94–7.57)	7.62 (6.86–8.38)	0.84
Years (95% CI)				
LE with CHP	1.30 (1.10–1.50)	2.02 (1.75–2.29)	2.15 (1.96–2.35)	0.85
Years (95% CI)				
Women				
LE	10.38 (10.31–10.44)	10.78 (10.72–10.84)	11.71 (11.65–11.78)	1.33
LE without CHP	7.77 (7.03–8.51)	7.13 (6.58–7.68)	7.71 (7.12–8.29)	-0.06
Years (95% CI)				
LE with CHP	2.61 (2.29–2.93)	3.65 (3.31–3.99)	4.01 (3.69–4.33)	1.40
Years (95% CI)				

Data: The Swedish Panel Study of Living Conditions of the Oldest Old 1992, 2002, 2011. Human Mortality Database. Own calculations. Abbreviations: LE life expectancy, CI confidence interval

Figure 2. (Title) Estimated proportions of remaining life expectancy at age 77 to be lived with and without complex health problems 1992–2011, by gender. (Legend) The bars show estimated proportions of remaining life years to be lived with and without complex health problems for the years 1992, 2002 and

2011, for men and women. Data: The Swedish Panel Study of Living Conditions of the Oldest Old 1992, 2002, 2011, Human Mortality Database. Own calculations.

Decomposition of changes in life expectancy with and without complex health problems into mortality and morbidity effects

Among men, between 1992 and 2011, the increase in life expectancy without CHP (0.84 years) was mainly driven by mortality effect (1.27 years), which was partially offset by the effect of increased prevalence of CPH (-0.43 years). For women, years without CHP decreased somewhat during the same period (0.06 years) because the positive mortality effect (0.79 years) was completely offset by a slightly greater negative morbidity effect (-0.85 years), i.e., an increase in the prevalence of CHP.

During the first part of the period (1992 and 2002), the opposite contributions of mortality effect (0.51 years) and effect of CPH prevalence (-0.53 years) to increasing LE with CPH were similar among men. Among women, the contribution of effects of CPH prevalence (-0.89 years) was much larger than that of mortality effects (0.26 years). Between 2002 and 2011, on the other hand, for both sexes the increase of life expectancy without CHP was predominantly driven by mortality effects.

Table 3

Contribution of the effects of mortality and prevalence of complex health problems to changes in life expectancy without complex health problems, 1992–2011

	Men			Women		
	1992– 2002	2002– 2011	1992– 2011	1992– 2002	2002– 2011	1992– 2011
Change in LE	0,70	0,99	1,69	0,40	0,93	1,33
Change in LE without CHP	-0.02	0.86	0.84	-0.64	0.58	-0.06
Mortality effect	0.51	0.66	1.27	0.26	0.47	0.79
CHP-effect	-0.53	0.20	-0.43	-0.89	0.11	-0.85

Data: The Swedish Panel Study of Living Conditions of the Oldest Old 1992, 2002, 2011, Human Mortality Database. Own calculations. Abbreviations: LE life expectancy, CHP complex health problems.

Discussion

In this nationally representative study of older adults in Sweden, we found that life expectancy at age 77 continuously increased between 1992 and 2011. Overall, the number of years lived with CHP and the share of life with CHP increased for both men and women. In 2011, from age 77, men spent on average 2.15 years and women 4.01 years with CHP. For women, the added life years consisted only of years with CHP, while the added years for men were equally distributed between years with and without CHP. Fewer individuals with CHP resided in residential care in 2011 compared to one and two decades before.

The finding that there was no sign of a decrease in time spent with CHP has important implications for resource needs and funding of care services. Moreover, the composite measure of CHP is highly indicative of the need of both social care and medical care, usually involving several providers. Our results, that an increasing proportion and number of older adults with CHP live in the community, and not in residential care, underlines the importance of the development and improvement of integrated medical and social services.

The trend of an increase of years lived with CHP can mainly be attributed to an increasing prevalence of CHP in the first part of the period, which all three health dimensions contributed to. Most previous studies on time trends in health expectancies have used single health indicators and divergent trends have been shown for different indicators, countries and time-points compared [21]. Given limits in comparability, our results of an increase in years lived with complex health problems point towards a similar trend as a few previous studies that have used composite health measures. For example, expected life years spent with cognitive impairment and disability combined increased slightly in Hong Kong between 2001/02 and 2011/12 [19]. A study based on the British Cognitive Function and Ageing Studies integrated dependency in activities of daily living and geriatric conditions into a measure of dependency. Results showed a significant increase of years lived from age 65 with both low dependency (requiring help less often than daily) and high dependency (24-h care) between 1991 and 2011 [33]. As in the present study, the development was more pronounced for women than for men [33]. On the other hand, results from the Longitudinal Aging Study Amsterdam reported an increase in years lived with combined multimorbidity and disability between 1993 and 2016 for men, but no clear trend for women [22]. Studies focusing on diseases and/or physiological indicators found an increase in time spent with six or more chronic conditions and polypharmacy between 2005 and 2014 in Germany, both in absolute years and as an increasing proportion of life [32].

A study among community-dwelling people conducted in 2011 revealed substantial differences in life expectancy with frailty at age 70 across 15 European countries, with estimates ranging between 0.1 to 1.8 years for men and from 0.4 to 5.5 years for women [31]. However, to our knowledge, there is a lack of studies on time trends in life expectancy with frailty.

That women in our study at all three time points could expect to live twice as long as men with CHP reflects the male-female health-survival paradox [44], that has been observed for most other single as well as composite health measures, such as multimorbidity [32, 45], frailty [31, 46], disability [33] as well as poor physical and cognitive health combined [19, 22].

Our decomposition analyses indicated that the increase in expected life years with complex health problems among women in the period of 1992–2002 was mainly driven by change in the prevalence of CHP, rather than mortality, while the contributions of these forces were similar among men. The increase of years without CHP, on the other hand, was predominantly driven by change in mortality, rather than the prevalence complex health problems, especially during the latter part of the period (2002–2011), and among men.

Increased prevalence of diseases and chronic conditions, as well as some functional limitations, have also been widely reported in Sweden [47] and internationally [14, 25, 33], although evidence on trends in the prevalence of disability seems more favorable [47]. Earlier diagnoses, successful life-saving interventions and better disease-control probably contribute to an increase of the prevalence of chronic conditions and functional limitations. Improved survival with disease may also entail an increased risk for further diseases [48].

Some strengths and limitations of this study must be acknowledged when interpreting the results. Although the sample sizes of the three SWEOLD waves are relatively small, a main strength of the study is that prevalence estimates can be assumed to be highly representative. All survey waves comprise nationally representative samples of the population aged 77 or older with high response rates (> 85%), including people with poor cognition and those living in residential care. Identical study design and methods were used over the decades. In order to avoid non-response, and thereby an underestimation of health problems on population level, telephone interviews or proxy interviews with a close family were conducted for those individuals who were too frail and/or cognitively impaired to participate by themselves or who did not want to participate in a face-to-face interview [36].

The health indicators are mostly self-reported (except the cognitive test), which could introduce bias in the health assessments. In national population surveys, for practical reasons, clinical precision cannot be accomplished. However, crude health measures that are not clinically relevant, but allow the inclusion of proxy interviews, can be highly suitable for national surveys aimed at informing decision makers for policy and planning.

To cover the most vulnerable subset of the older population, we made our definition of complex health problems restrictive, setting high thresholds for severe problems within each domain. However, people with severe problems in only one domain could also need care from a variety of providers. Thus, rates of CHP as measured in this study, could to some extent underestimate actual needs for integrated medical and social care in the older population.

Moreover, the Sullivan method, which we used to calculate health expectancies, does not take transitions between health states into account, which could potentially introduce bias in the health estimates [41]. However, recovery from complex health problems, involving severe problems in several health domains, is likely to be limited in high age, and the estimates based on the Sullivan method have proven reliable as long as prevalence is regular and smooth over time [49, 50].

Conclusion And Implications For Social Policy

Assuming unchanged prevalence of CHP since 2011, based on population statistics and projections from Statistics Sweden [37], the number of individuals aged 77 or older with CHP in Sweden can be expected to have risen from around 172,000 in 2011 to 196,000 in 2019. An additional 43% increase to 280,000 individuals with CHP could be expected by 2030, when the large cohorts born in the 1940s will reach 80 to 90 years of age, still assuming unchanged prevalence of CHP [37].

In parallel with population aging and an increase in the number and proportion of older adults with CHP, there have been substantial cutbacks in publicly financed and universally available eldercare in Sweden during the past few decades [9]. Following the principle of “aging in place”, as well as due to economic constraints, since 2000, 30 percent (37,000) of the 121,000 beds in residential care have disappeared. Also, half the hospital beds have been cut since the 1990s [9]. Shorter hospital stays and fewer beds in residential care have entailed that very frail old people increasingly are cared for at home, typically living alone or with a partner [9]. Our results confirmed that a substantially higher proportion of those with CHP lived in the community 2011 compared to previous decades. Also, among older adults *without* CHP, very few individuals lived in residential care, indicating that this most comprehensive form of social care is concentrated to those with the greatest care needs. This has also been reflected in a decrease in the length of stay in residential care and an increase in the proportion of people who died shortly after they moved in [10]. Since another 10,000 beds in residential care have disappeared since 2011, the trend of “aging in place” among individuals with substantial and complex care needs has probably continued during the past years [9].

While our results show that more people remain in their own homes despite CHP, the ability to navigate the health- and social care system, and to manage care contacts with different providers, generally decreases with poor health [51]. This is particularly challenging in a fragmented system, where medical care is financed and organized at the regional level while municipalities are responsible for social care [52]. The highly decentralized Swedish care system, as well as the large number of independent care providers, who for example may have different journal systems, has been reported to result in suboptimal care for individuals who need a variety of both medical and social care [52]. Thus, our results, that more people live longer with CHP, and increasingly remain in their own homes, reinforce the importance of coordination and communication between providers to provide coherent medical and social care for the most vulnerable individuals living in the community.

Abbreviations

SWEOLD – The Swedish Panel Study of Living Conditions of the Oldest Old

LE – life expectancy

CHP – complex health problems

CI – confidence interval

Declarations

Ethics approval and consent to participate

The project is covered by ethical approval from the Uppsala University Hospital Ethical Committee (1992; Dnr 247/91), the Karolinska Institutet Ethical Research Committee (2002; Dnr 03-413), and the Regional

Ethical Review Board in Stockholm (2011; Dnr 2010/403-31/4). A relative (normally a spouse or an adult child) signed the consent form for participants who were physically or cognitively too impaired to sign the form at the time of the interview.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request. Public access to the database may be given if applicants have a scientific affiliation, sign a statement that the data will only be used for scientific purposes, and that the scientific project have ethical approval. Applicable sections of The Swedish Research Council (VR) principles for conducting research in humanities and the social sciences must be adhered to (<http://www.codex.vr.se/en/forskninghumsam.shtml>). Data will be available after the above-mentioned documents have been received, by email at the SWEOLD Research Data Center dataaccess@sweold.se. More information can be found at www.sweold.se.

Competing interest

The authors declare no conflicts of interest.

Funding

This work was supported by the Swedish Research Council for Health, Working Life and Welfare (Grant 2016-00197) and the Swedish Research Council (Grant 2016-01072).

Authors' contributions

BM managed and prepared the data, drafted and finalized the manuscript. PL and DJ performed the statistical analyses. BM, PL, DJ, AO, LS, SK, and JWW participated in the conception and design of the study and interpretation of the data and revised the manuscript. BM, PL, DJ, AO, LS, SK, and JWW have read and approved the manuscript.

Acknowledgements

Not applicable

Authors' information

BM: Aging Research Center, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet and Stockholm University, Solna, Sweden

Stockholm Gerontology Research Center, Stockholm, Sweden

PL, DJ and AO: Max Planck Institute for Demographic Research, Rostock, Germany

SK: Division of Ageing and Social Change, Department of Social and Welfare studies, Linköping University, Linköping, Sweden

Aging Research Center, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet and Stockholm University, Solna, Sweden

JWW: Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Solna, Sweden

Aging Research Center, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet and Stockholm University, Solna, Sweden

References

1. Robine JM. Age patterns in adult mortality. In: Rogers RG, Crimmins EM, editors. International handbook of adult mortality. New York: Springer; 2011. p. 207–226.
2. Lunney JR, Lynn J, Foley DJ, Lipson S, Guralnik JM. Patterns of functional decline at the end of life. *JAMA*. 2003; 289(18):2387–2392.
3. Baltes MM. The psychology of the oldest-old: the fourth age. *Curr. Opin. Psychiatry*. 1998; 11(4):411–415.
4. Smith AK, Walter LC, Miao Y, Boscardin WJ, Covinsky KE. Disability during the last two years of life. *JAMA Intern Med*. 2013; 173(16):1506–1513.
5. Romoren TI, Blekesaune M. Trajectories of disability among the oldest old. *J Aging Health*. 2003; 15(3):548–566.
6. Meinow B, Wastesson JW, Kåreholt I, Kelfve S. Long-Term Care Use During the Last 2 Years of Life in Sweden: Implications for Policy to Address Increased Population Aging. *J Am Med Dir Assoc*. 2020; 21(6):799–805.
7. Laslett P. *A Fresh Map of Life*. London: Palgrave; 1996.
8. Violan C, Foguet-Boreu Q, Flores-Mateo G, Salisbury C, Blom J, Freitag M, Glynn L, Muth C, Valderas JM. Prevalence, determinants and patterns of multimorbidity in primary care: a systematic review of observational studies. *PloS one*. 2014; 9(7):e102149.
9. Schön P, Heap J. European Social Policy Network thematic report on challenges in long-term care: Sweden. Brussels: European Commission; 2018.
10. Schön P, Lagergren M, Kåreholt I. Rapid decrease in length of stay in institutional care for older people in Sweden between 2006 and 2012: results from a population-based study. *Health Soc Care Community* 2016; 24(5):631–638.
11. National Board of Health and Welfare. Statistik om särskilt boende [Statistics on municipal institutional care]. Stockholm: National Board of Health and Welfare; 2016.

12. Jagger C, Crimmins EM, Saito Y, Yokota RTDC, Van Oyen H, Robine J-M. International handbook of health expectancies. Springer Nature; 2020.
13. Hay SI, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, Abdulkader RS, Abdulle AM, Abebo TA, Abera SF. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017; 390(10100):1260–1344.
14. Crimmins EM, Beltrán-Sánchez H. Mortality and morbidity trends: is there compression of morbidity? *J Gerontol B Psychol Sci Soc Sci*. 2011; 66(1):75–86.
15. Christensen K, Thinggaard M, Oksuzyan A, Steenstrup T, Andersen-Ranberg K, Jeune B, McGue M, Vaupel JW. Physical and cognitive functioning of people older than 90 years: a comparison of two Danish cohorts born 10 years apart. *Lancet*. 2013; 382(9903):1507–1513.
16. Lagergren M, Johnell K, Schön P, Danielsson M. Towards a postponement of activities of daily living dependence and mobility limitations: Trends in healthy life years in old age in Sweden. *Scand J Public Health*. 2017; 45(5):520–527.
17. Jeune B, Eriksen ML, Andersen-Ranberg K, Bronnum-Hansen H. Improvement in health expectancy at ages 50 and 65 in Denmark during the period 2004–2011. *Scand J Public Health*. 2015; 43(3):254–259.
18. Freedman VA, Wolf DA, Spillman BC. Disability-free life expectancy over 30 years: a growing female disadvantage in the US population. *Am J Public health*. 2016; 106(6):1079–1085.
19. Yu R, Leung J, Lum C, Auyeung T, Lee JS, Lee R, Woo J. A comparison of health expectancies over 10 years: implications for elderly service needs in Hong Kong. *Int J Public health*. 2019; 64(5):731–742.
20. Crimmins EM, Saito Y, Kim JK. Change in cognitively healthy and cognitively impaired life expectancy in the United States: 2000–2010. *SSM Popul Health*. 2016, 2:793–797.
21. Jagger C, Matthews FE, Wohland P, Fouweather T, Stephan BC, Robinson L, Arthur A, Brayne C, Medical Research Council Cognitive Function and Ageing Collaboration. A comparison of health expectancies over two decades in England: results of the Cognitive Function and Ageing Study I and II. *Lancet*. 2016; 387(10020):779–786.
22. Deeg DJ, Comijs HC, Hoogendijk EO, van der Noordt M, Huisman M. 23-year trends in life expectancy in good and poor physical and cognitive health at age 65 years in the Netherlands, 1993–2016. *Am J Public Health*. 2018; 108(12):1652–1658.
23. Falk H, Johansson L, Ostling S, Thogersen Agerholm K, Staun M, Host Dorfinger L, Skoog I. Functional disability and ability 75-year-olds: a comparison of two Swedish cohorts born 30 years apart. *Age Ageing*. 2014; 43(5):636–641.
24. Lagergren M, Johnell K, Schon P, Danielsson M. Healthy life years in old age: Swedish development 1980–2010 according to different health indicators. *Scand J Public Health*. 2016; 44(1):55–61.
25. Chatterji S, Byles J, Cutler D, Seeman T, Verdes E. Health, functioning, and disability in older adults-present status and future implications. *Lancet*. 2014; 385(9967):563–575.

26. Walter S, Beltrán-Sánchez H, Regidor E, Gomez-Martin C, del-Barrio JL, Gil-de-Miguel A, Subramanian SV, Gil-Prieto R. No evidence of morbidity compression in Spain: a time series study based on national hospitalization records. *Int J Public Health* 2016; 61(7):729–738.
27. van Oostrom SH, Gijsen R, Stirbu I, Korevaar JC, Schellevis FG, Picavet HSJ, Hoeymans N. Time trends in prevalence of chronic diseases and multimorbidity not only due to aging: data from general practices and health surveys. *PloS one*. 2016; 11(8):e0160264.
28. Marengoni A, Angleman S, Meinow B, Santoni G, Mangialasche F, Rizzuto D, Fastbom J, Melis R, Parker M, Johnell K. Coexisting chronic conditions in the older population: Variation by health indicators. *Eur J Intern Med*. 2016; 31:29–34.
29. Meinow B, Kåreholt I, Thorslund M, Parker MG. Complex health problems among the oldest old in Sweden: increased prevalence rates between 1992 and 2002 and stable rates thereafter. *Eur J Ageing*: 2015; 12(4):285–297.
30. Murray CJ, Salomon JA, Mathers CD, Lopez AD, Organization WH. Summary measures of population health: concepts, ethics, measurement and applications. World Health Organization; 2002.
31. Romero-Ortuno R, Fouweather T, Jagger C: Cross-national disparities in sex differences in life expectancy with and without frailty. *Age ageing*. 2014; 43(2):222–228.
32. Tetzlaff J, Muschik D, Epping J, Eberhard S, Geyer S. Expansion or compression of multimorbidity? 10-year development of life years spent in multimorbidity based on health insurance claims data of Lower Saxony, Germany. *Int J Public Health*. 2017; 62(6):679–686.
33. Kingston A, Wohland P, Wittenberg R, Robinson L, Brayne C, Matthews FE, Jagger C, Green E, Gao L, Barnes R. Is late-life dependency increasing or not? A comparison of the Cognitive Function and Ageing Studies (CFAS). *Lancet*. 2017; 390(10103):1676–1684.
34. Meinow B, Parker MG, Kåreholt I, Thorslund M. Complex health problems in the oldest old in Sweden 1992–2002. *Eur J Ageing*. 2006; 3(2):98–106.
35. Meinow B, Parker MG, Thorslund M. Complex health problems and mortality among the oldest old in Sweden: decreased risk for men between 1992 and 2002. *Eur J Ageing*. 2010; 7(2):81–90.
36. Lennartsson C, Agahi N, Hols-Salen L, Kelfve S, Kareholt I, Lundberg O, Parker MG, Thorslund M. Data resource profile: The Swedish Panel Study of Living Conditions of the Oldest Old (SWEOLD). *Int J Epidemiol*: 2014; 43(3):731–738.
37. Statistical database. Statistics Sweden, Stockholm/Örebro. 2021.
<https://www.statistikdatabasen.scb.se/pxweb/en/ssd/> Accessed 23 April 2021.
38. Meinow B, Kareholt I, Thorslund M, Parker MG. Complex health problems among the oldest old in Sweden: Increased prevalence rates between 1992 and 2002 and stable rates thereafter. *Eur J Ageing*: 2015; 12(4):285–297.
39. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12(3):189–198.
40. Ericsson MC, Gatz M, Kåreholt I, Parker MG, Fors S. Validation of abridged mini-mental state examination scales using population-based data from Sweden and USA. *Eur J Ageing*. 2017;

14(2):199.

41. Jagger C, van Oyen H, Robine JM. Health Expectancy Calculation by the Sullivan Method: A Practical Guide. 4th ed. Newcastle: Newcastle University, Institute for Ageing; 2014.
42. Yang Y. Long and happy living: Trends and patterns of happy life expectancy in the US, 1970–2000. *Soc Sci Res.* 2008; 37(4):1235–1252.
43. Andreev EM, Shkolnikov VM, Begun AZ. Algorithm for decomposition of differences between aggregate demographic measures and its application to life expectancies, healthy life expectancies, parity-progression ratios and total fertility rates. *Demog Res.* 2002; 7:499–522.
44. Oksuzyan A, Juel K, Vaupel JW, Christensen K. Men: good health and high mortality. Sex differences in health and aging. *Aging Clin Exp Res.* 2008; 20(2):91–102.
45. Hoogendijk EO, van der Noordt M, Onwuteaka-Philipsen BD, Deeg DJ, Huisman M, Enroth L, Jylhä M. Sex differences in healthy life expectancy among nonagenarians: A multistate survival model using data from the Vitality 90 + study. *Exp Gerontol.* 2019; 116:80–85.
46. Herr M, Arvieu J-J, Ankri J, Robine J-M. What is the duration of life expectancy in the state of frailty? Estimates in the SIPAF study. *Eur J Ageing.* 2018; 15(2):165–173.
47. Lindgren B. The rise in life expectancy, health trends among the elderly, and the demand for health and social care. Working Paper No 142. Stockholm: National Institute of Economic Research; 2016.
48. Ebeling M, Meyer AC, Modig K. The rise in the number of long-term survivors from different diseases can slow the increase in life expectancy of the total population. *BMC Public Health.* 2020; 20(1):1–8.
49. Sullivan DF. A single index of mortality and morbidity. *HSMHA Health Rep.* 1971; 86:347–354.
50. Mathers CD, Robine J-M. How good is Sullivan's method for monitoring changes in population health expectancies? *J Epidemiol Community Health.* 1997; 51(1):80–86.
51. Fried LP, Ferrucci L, Darer J, Williamson JD, Anderson G. Untangling the concepts of disability, frailty, and comorbidity: Implications for improved targeting and care. *J Gerontol A Biol Sci Med Sci.* 2004; 59(3):255–263.
52. Liljas AE, Brattström F, Burström B, Schön P, Agerholm J: Impact of integrated care on patient-related outcomes among older people—a systematic review. *Int J Integr Care.* 2019; 19(3):6,1–16.

Figures

Figure 1

(Title) Long-term care use among individuals with and without complex health problems, 1992-2011.
 (Legend) The bars show the proportions of individuals living in residential care, using home care, or no social care, stratified according to the presence of complex health problems or not. Data: The Swedish Panel Study of Living Conditions of the Oldest Old 1992, 2002, 2011. Own calculations.

Figure 2

(Title) Estimated proportions of remaining life expectancy at age 77 to be lived with and without complex health problems 1992-2011, by gender. (Legend) The bars show estimated proportions of remaining life years to be lived with and without complex health problems for the years 1992, 2002 and 2011, for men and women. Data: The Swedish Panel Study of Living Conditions of the Oldest Old 1992, 2002, 2011, Human Mortality Database. Own calculations.