Platelet generation from circulating megakaryocytes is triggered in the lung vasculature

Xiaojuan Zhao

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Dominic Alibhai

University of Bristol <https://orcid.org/0000-0001-5421-4424>

Tony Walsh

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Nathalie Tarassova

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Semra Birol

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Christopher Williams

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Chris Neil

Wolfson Bioimaging Facility, University of Bristol

Elizabeth Aitken

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Amie Waller

University of Cambridge / NHS Blood and Transplant, Wellcome-MRC Cambridge Stem Cell Institute, Jeffrey Cheah Biomedical Centre, University of Cambridge

Jose Ballester-Beltran

Department of Haematology, University of Cambridge

Peter Gunning

University of New South Wales <https://orcid.org/0000-0003-0833-3128>

Edna Hardeman

University of New South Wales Sydney <https://orcid.org/0000-0003-1649-7712>

Ejaife Agbani

Cumming School of Medicine, University of Calgary <https://orcid.org/0000-0002-0043-9115>

Ingeborg Hers

School of Physiology, Pharmacology and Neuroscience, University of Bristol

Cedric Ghevaert

Stem Cell Institute and Department of Haematology, University of Cambridge, and National Health Service Blood and Transplant

Alastair Poole (✉ a.poole@bris.ac.uk)

Article

Keywords:

Posted Date: December 2nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-690639/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Platelet generation from circulating megakaryocytes is triggered in the lung vasculature

Xiaojuan Zhao^{1*}, Dominic Alibhai², Tony G. Walsh¹, Nathalie Tarassova¹, Semra Z. Birol¹,
Christopher M. Williams¹, Chris R. Neal², Elizabeth W. Aitken¹, Amie K. Waller³, Jose Ballester-
Beltran³, Peter W. Gunning⁴, Edna C. Hardeman⁴, Ejaife O. Agbani⁵, Ingeborg Hers¹, Cedric
Ghevaert³ & Alastair W. Poole^{1*}

¹School of Physiology, Pharmacology and Neuroscience, Biomedical Sciences Building,
University of Bristol, Bristol, BS8 1TD, UK.

²Wolfson Bioimaging Facility, Biomedical Sciences Building, University of Bristol, Bristol, BS8
1TD, UK.

³University of Cambridge / NHS Blood and Transplant, Wellcome-MRC Cambridge Stem Cell
Institute, Jeffrey Cheah Biomedical Centre, Cambridge Biomedical Campus, University of
Cambridge, Cambridge, CB2 0AW, UK.

⁴School of Medical Sciences, University of New South Wales, Sydney, NSW 2052, Australia.

⁵Cumming School of Medicine, University of Calgary, Calgary, AB, T2N 1N4, Canada.

*Corresponding Authors. Email: xz14926@bristol.ac.uk or a.poole@bristol.ac.uk

SUMMARY

Platelets are small anucleate blood cells^{1,2} with critical roles in haemostasis, thrombosis, inflammation, vascularization, innate immunity and tissue regeneration^{3,4}. In vitro-derived platelets, as an alternative to native platelets, are attractive for fundamental research because of their rapid genetic tractability, as vectors for drug and genetic component delivery⁵ and in clinical

platelet transfusion. At present, however, their very low production rate, and poor agonist responsiveness, are major obstacles. Platelets are formed by fragmentation from mature polyploid megakaryocytes (MKs), their precursor cells⁶, although the process of their generation remains incompletely understood^{7,8}. Bone marrow is proposed to be the main site of platelet production, however indirect evidence since the 1930s^{9,10} and recent direct observation¹¹ has shown that the lung can also be a primary site of platelet biogenesis. Here we established an *ex vivo* mouse heart-lung model (Fig. 1a) through which we were able to perfuse murine MKs. Remarkably, we could show for the first time that MKs, despite their large size, can pass multiple times through the lung vasculature, and that this leads to the generation of very large numbers of fully functional platelets (up to 3,000 per megakaryocyte^{7,12}). Using this system and a novel *in vitro* microfluidic chamber we show roles for ventilation, oxygenation and healthy pulmonary endothelial cells in platelet generation. We show that MKs undergo enucleation upon repeated passage through pulmonary vasculature before fragmentation to generate platelets, with this final process dependent on the actin regulator TPM4. This advances our understanding of platelet formation in the body and establishes a novel approach to generate large numbers of them outside the body.

The *ex vivo* mouse heart-lung model (Fig. 1a) is based upon isolating the heart and lungs as a single unit, ligating the venae cavae and the aortic arch and perfusing CD41-PE or FITC-labelled MKs through the pulmonary circulation from the right ventricle, collecting the perfusate from the left ventricle. This allows quantitation and imaging of cells passaged through the pulmonary vasculature. In the first instance, lungs were artificially ventilated with air. We were expecting that the vast majority of MKs would be trapped within the lung vasculature due to their large size (around 50-100 μm diameter¹³), but unexpectedly more than 50% of the MKs emerged in the perfusate after the first passage (Fig. 1b). The perfusate could be re-injected through the lungs and

although upon multiple passages the number of intact MKs (showing a circular shape and central nucleus) continued to decrease, it was evident that MKs were able to pass through the lung vasculature multiple times. It was also apparent that CD41-labelled particles also appeared in the perfusate that were similar in size and granularity to mouse platelets. We term these “generated” platelets (CD41-positive events in gate P1 in Fig. 1c). The numbers of generated platelets per MK gradually increased with increasing passages, up to around 1134.7 ± 597.2 platelets/MK after 18 passages (Fig. 1d and f). Two-photon microscopy of fixed lung sections after 18 passages showed that some generated platelets were retained in the lung microvasculature (Fig. 1e and Extended Data Video 1-3). These were quantified in a defined lung volume and the total lung volume measured using a fluid displacement method (Extended Data Fig.1b) to estimate numbers of retained platelets. We calculated this to be 1950 ± 682 per MK injected (Fig. 1f). Adding this to the number contained in the perfusate (Fig. 1d), we estimate that after 18 passages through the lung approximately 3084.7 ± 1279.2 platelets were generated per MK (Fig. 1f), in keeping with previous estimates that each MK *in vivo* produces approximately 1000-4000 platelets^{7,12}. Platelets were not generated simply as a consequence of passage through small-bore needles, since we showed that repeated passage of MKs through 21G needles 18 times is not sufficient to generate platelets (Fig. 1c and Fig. 1f). Altogether, we could generate physiological levels of platelets after multiple recirculation of MKs through the lung microvasculature. We also infer that MKs substantially, rapidly and reversibly, deform their shape to passage through the capillary network of the lung.

The *ex vivo* mouse heart-lung model (Fig. 1a) can be a useful tool to allow artificial ventilation with either ambient air or with pure nitrogen (N₂), or no ventilation, to assess the roles of physical ventilation and gaseous oxygen in regulating MK biology and thrombogenesis. We first explored whether air ventilation is essential for platelet generation in our model. Strikingly, when lungs are

ventilated with pure N₂ to completely de-oxygenate the heart-lung preparation, the number of generated platelets in the perfusate was almost ablated, reduced to just 43 ± 20 platelets/MK (Fig. 1d). Two-photon imaging of nitrogen-ventilated lungs showed mature MKs were trapped in the lung vasculature (Fig. 1e and Extended Data Video 4), a feature not observed under air ventilation.

5 Pulmonary endothelial cells (ECs), which play key roles in gas exchange in the lung¹⁴, interact closely with MKs as they passage through the vasculature, and we therefore compared their viabilities (determined by the Calcein Deep Red retention assay) and mitochondrial membrane potential (an indicator of healthy cells, determined by accumulation of tetramethyl rhodamine methyl ester (TMRM) in active mitochondria) after 18 passages under air- or N₂- ventilation for
10 approximately 2 hours. Surprisingly, ECs from preparations ventilated under N₂ were fully viable, and comparable with those ventilated under air (Fig. 1g). However, the mean intensity of TMRM of ECs from N₂-ventilated lungs was approximately half that of air-ventilated lung or fresh lung (Fig. 1h). These data suggest that air ventilation and fully healthy ECs are essential for efficient platelet generation in the heart-lung preparation, and the capability of MKs to pass through the
15 lung vasculature is lost in heart-lung preparations that were N₂-ventilated.

We then investigated whether healthy pulmonary ECs are required for the passage of MKs through the heart-lung preparation and the importance of ventilation for platelet generation. After flushing the lung under air ventilation, we then stopped ventilation while perfusing MKs through the preparation 18 times. In the absence of ventilation, the number of platelets generated per MK in
20 the perfusate still gradually increased with increasing passage number, but the numbers generated were substantially lower than in the air-ventilated condition (498.4 ± 235.7 platelets/MK, Fig. 1d). Importantly, the mean intensity of endothelial cell TMRM was approximately halved by lack of ventilation relative to air-ventilated controls, equivalent to the N₂-ventilated lung (Fig. 1h). In contrast to the N₂-ventilated lung however, two-photon imaging of unventilated lungs showed no

MKs trapped in the lung vasculature (Fig. 1e). The data suggest therefore that passage of MKs through the pulmonary vasculature may not require a fully healthy endothelium, but clearly oxygenation plays a critical role at some level, since MKs cannot passage through the vasculature in N₂-ventilated lungs. This effect may reflect a direct requirement of MKs for oxygenation, but the data also suggest that the passage of MKs through structural arrangement of the lung microcirculation is also a major contributor to platelet generation.

To verify whether this structural arrangement could mediate platelet generation, we designed a polydimethylsiloxane (PDMS)-based (gas permeable) microfluidic chamber with channel arrangement mimicking tissue microcirculation. The channels were of uniform depth of 10 μ m throughout, where the entry and exit channels had a width of 100 μ m and where branches emerged halving the channel width each time, to a minimal width of 12.5 μ m, as per the diagram shown in Fig. 1i and Extended Data Fig. 1c. This channel arrangement allowed us to flow through cells and determine platelet generation in the perfusate after repeated passage. Fig. 1d shows that the numbers of generated platelets per MK gradually increased with increasing passages, similar to the numbers generated in the unventilated lung, with 492.3 ± 95.1 platelets/MK after 18 passages (Fig. 1d).

Altogether, these data suggested that (1) air-ventilation and healthy ECs are required for MKs to generate physiological levels of platelets in the heart-lung preparation; (2) the structural arrangement of the pulmonary microcirculation plays a role in platelet generation; (3) lack of ventilation or N₂-ventilation for 2 hours caused partial loss of the mitochondrial membrane potential of pulmonary ECs; (4) the exposure of pure N₂ for 2 hours impaired the mobility of MKs.

We determined that generated platelets were *bona fide* live platelets rather than cellular fragments, using the vital dye calcein-AM (Fig. 2a) and showing mitochondrial function comparable to

normal mouse platelets using TMRM (Fig. 2b-c). Generated platelets were anuclear, showing no staining with the DNA dye Draq5 (Fig. 2a). The proportion of cells expressing CD61 and CD42b, and the mean fluorescence intensity of those markers, were comparable between generated and control platelets (Fig. 2d). However, the mean fluorescence intensity of three collagen receptors GPVI, CD49b¹⁵ and CD42d¹⁶ was higher in generated platelets than controls (Fig. 2d), while the proportion of cells expressing these receptors was lower (Extended Data Fig. 1d). Platelets display an almost uniquely characteristic sub-plasma membrane microtubular ring, running circumferentially in resting platelets^{17,18}. Our generated platelets, immunolabelled for α -tubulin, display this characteristic ring structure (Fig. 2e), and the cells are larger than controls (3.5 ± 1.2 μm vs 1.9 ± 0.3 μm , Fig. 2f). Younger platelets have been shown to be larger than the mean platelet diameter^{19,20}, and our generated platelets are therefore consistent with this observation. Finally, we visualized the ultrastructures of generated platelets by transmission electron microscopy (TEM, Fig. 2g), after depletion of host platelets using anti-GPIb α antibodies. Generated platelets displayed a discoid shape with classical characteristics including α -granules, σ -granules or dense bodies, mitochondria, open canalicular system, and microtubule coils.

We then demonstrated that the functionality of generated platelets is also similar to that of mouse platelets (controls). Both generated and control platelets showed equivalent responses to agonists (CRP and thrombin) in terms of integrin $\alpha_{\text{IIb}}\beta_3$ activation and degranulation (P-selectin expression, Fig. 3 a-b). Thrombus formation *in vitro* was also assessed, determining how generated platelets mixed into whole blood interact with a collagen-coated surface under flow. Generated platelets (stained with both DiOC6 and CellTracker™ Red CMTPX dye, white) occupied all levels of the thrombus whilst control platelets (stained with CellTracker™ Red CMTPX dye alone, magenta) were mainly situated on top of the thrombus, suggesting generated platelets showed a higher responsiveness to collagen, or were primary reactors to it (Fig. 3c-e and Extended Data Video 5).

This may suggest that generated platelets are early interactors with collagen, displaying the greater adhesive functionality of younger platelets^{19,20}, possibly due to higher levels of collagen adhesive receptors (GPVI and CD49b¹⁵, and CD42d¹⁶) (Fig. 2d).

5 We wanted to explore the details of the release of platelets from MKs upon repeated passage through the pulmonary vasculature. The cells in the perfusate were imaged after collected over a defined numbers of passages (3, 6, 9, 12 and 18), and strikingly, upon repeated passages, MKs gradually move their nuclei to the periphery and subsequently enucleate, generating both naked nuclei and enucleated MKs. Although small numbers of enucleated round MKs were found, we
10 saw the gradual accumulation of larger anuclear objects. Figure 1b, 4 and Extended Data Video 6-7 show the steps involved in the process, with images shown in Fig. 4a, quantified in Fig. 1b and 4b and summarized in Fig. 4j. As shown in Fig. 1b, the percentage of intact MKs decreased from 54% after 1 passage (P1) to 6.7% after 9 passages (P9), while in Fig. 4b the percentage of large naked nuclei (>20 μm diameter) increased from 5% after P1 to 22% after P9. At the same time,
15 large anuclear objects (>10 μm) also increased, as a proportion of total MKs and derivatives, from 13% after P1 to 45% after P9. These data suggested that the large polyploid nucleus moves from a central position to the periphery of the cell, in a process of polarization. The nucleus is then extruded from the cell upon further passages through the lung vasculature, until by approximately 9 passages there are very few nucleated MKs left. After 12 passages, large naked nuclei (>20 μm)
20 also became rare, being replaced by irregular small sub-nuclei. These sub-nuclei appeared connected to each other, probably by membranous structures or thin DNA bridges²¹ that have been described previously between segregated chromosomes which were too fine for visualization by light microscopy (Fig. 4c and Extended Data Video 8). Sub-nuclei were characterized by depth (Fig. 4d) and aspect ratio (width: height, Fig. 4e), all of which decreased substantially with

increasing passage number (P3 to P18 parameters: depth 8.9 μm to 5.3 μm , aspect ratio 2 to nearly 1, major axis 14 μm to 6.3 μm , minor axis 8 μm to 5.7 μm). The extruded naked nuclei therefore undergo a process of division into multiple component sub-nuclei, which proceed to condense into compact sub-nuclear units with a greater circularity. The anucleate MK proceeds to fragment into platelets after multiple passages, to reach plateau numbers by 15-18 passages (see Fig. 1d). This is the first time these enucleating behaviors have been observed for MKs when releasing platelets.

The release of platelets is understood to require cytoskeletal re-organisation involving the actin cytoskeleton. Tropomyosins form co-polymers with actin filaments and regulate filament function in an isoform-specific manner²². Tropomyosin TPM4 has been shown to have a role in platelet formation¹⁴. In our *ex vivo* lung system, *Tpm4*^{-/-} MKs generate no platelets (Fig. 4f-g), in contrast to the *in vivo* setting where platelet counts in *Tpm4*^{-/-} mice drop by approximately 50% compared to wild-type²³. This suggests that the remaining 50% platelet generation may take place in the bone marrow and may not fully require TPM4. This is consistent with the findings of LeFrancais *et al.*¹¹, who have shown that approximately 50% of circulating platelets are generated in the mouse lung. We then wanted to explore further the steps in platelet generation requiring TPM4. Compared to wild type MKs, during the first 3 passages, fewer *Tpm4*^{-/-} MKs underwent transformation to large anuclear objects. However, this represented only a delay in these events, since by 6-9 passages the numbers of enucleated cells were comparable with wild-type (Fig. 4h). Two-photon microscopy of fixed lung sections after 18 passages showed abundant anuclear fluorescent objects, sized 10 μm (Fig. 4i and Extended Data Video 9). These data suggest a small and non-essential role for TPM4 in regulating enucleation, but a critical role in regulating both fragmentation of MKs into platelets with normal sizes (summarized in Fig. 4j).

This study is the first to demonstrate platelet generation from cultured MKs outside the body with an output similar to *in vivo* estimated MK capabilities. Several unexpected findings were observed during the study. First, approximately 50% of MKs pass through the lung microvasculature. This suggests a substantial ability to reversibly deform as they passage through the lung microvasculature. Pulmonary capillaries have average internal diameters of 5-8 μm ²⁴ which decrease with lung inflation²⁵. This challenges an assumption that MKs may be trapped in the lung microvasculature, despite the mismatch in sizes of cell and capillary diameter²⁶. However, lung ventilation with pure N₂ effectively ablated platelet formation, and caused MKs to lodge in the lung vasculature, indicating a requirement for oxygenation for MK motility. MK numbers are increased in the lungs of patients with acute respiratory distress syndrome caused by Covid-19, for example, and pulmonary MK levels may represent a histologic variable of interest in this disease²⁷. Although some of these MKs may be resident cells in the stroma of the lung²⁸, some may be MKs trapped in lung vasculature.

Second, repeated passage of MKs through the pulmonary vasculature induces a reproducible sequence of events leading to platelet generation. Our data suggest that O₂ tension and maintaining healthy pulmonary ECs, as well as repeated physical contraction of the lung are important to support platelet generation. This might suggest an essential and dynamic conversation between the pulmonary microvasculature environment and MKs, we term ‘education’, to induce reversible deformation of MKs, stimulate motility through the microvasculature and promote platelet release.

This education makes the lung a primary and unique site for platelet biogenesis. However, since MKs pass through the pulmonary vasculature, it is possible that an important educational signal is induced simply by passage through the small channels of the vasculature, and therefore that some platelet generation may take place as they passage through other organs of the body. This is

supported by the observation that we can generate substantial numbers of platelets by passage of MKs through the microfluidic chamber that mimics microvasculature.

These pieces of evidence suggest that there is synergy between all the unique features of the lung vasculature (oxygenation, ventilation, healthy pulmonary ECs and the microvascular structure) that makes this organ likely to be the most important site of vascular production of platelets. This may partially explain differences in platelet count reported in patients with lung disease^{29,30}, or people living at the different altitudes^{31,32}.

Third, MKs enucleate intravascularly, as part of the process leading to platelet generation. Few cells are known to enucleate, but importantly these include erythrocytes. This may be important because MKs and erythrocytes are developmentally closely related^{33,34}. It is possible therefore that there may be shared machinery important for enucleation in both related cell lineages. Our observation could be consistent with the phenomena that naked nuclei of denuded megakaryocytes were found in the bone marrow and the lung of patients with severe Covid-19³⁵.

Fourth, another striking observation was that in the *ex vivo* lung preparation, *Tpm4*^{-/-} murine MKs generated no platelets. Mechanistically we showed this to be by abolishing the fragmentation of MKs into platelets. This demonstrated not only a critical role for TPM4 in MK education in the lung, but also suggested that platelet generation in the lung may be mechanistically distinct from platelet generation in the bone marrow. These observations are consistent with Lefrancais *et al*¹¹ who had proposed that approximately 50% of platelet generation takes place in the lung vasculature. The findings suggest that the two systems of platelet generation, in lung and bone marrow, may each operate, and that at least in the mouse each system contributes approximately half of the circulating platelets. Studies in genetically modified mice and correlates with genetic-based causes of thrombocytopenias in humans show that both microtubules⁶ and actin^{8,36,37} play important roles in platelet generation from MKs. TPM4 has a prominent role in actin regulation,

therefore our data suggest that the lung vasculature educates MKs to generate platelets by an actin-based mechanism.

This work identifies a mechanism of platelet generation, by repeated passage of megakaryocytes through lung vasculature under air ventilation, involving enucleation and TPM4-dependent fragmentation of cells. Importantly, this may help to set up new approaches to large scale in vitro generation of human platelets, valuable for establishing novel ways to understand the molecular control of platelet generation and function, and for developing a variety of cell-mediated therapies and supply of donor cells for transfusion.

METHODS

Materials

Cell culture reagents including IMDM-Glutamax and penicillin-streptomycin were from Invitrogen (Paisley, UK). Recombinant murine stem cell factor (rSCF) and recombinant murine thrombopoietin (rTPO) were purchased from PeproTech EC (London, UK). PE or FITC - conjugated anti-CD41 and isotype-nonspecific IgG were from BD Biosciences (Berkshire, UK). FITC-conjugated anti-mouse CD31, CD102 antibodies and isotype-nonspecific IgG were from Biolegend (San Diego, USA). SYLGARD™ 184 Silicone Elastomer Kit was from Dow Corning (Michigan, USA). Anti-mouse GPIb α antibody R300, FITC-conjugated anti-mouse CD61, CD42b, CD42d, CD49b, Glycoprotein VI (GPVI) antibodies and isotype-nonspecific IgG, PE-conjugated anti-CD62P antibodies, JON/A anti-integrin $\alpha_{IIb}\beta_3$ and isotype-nonspecific IgG were from Emfret Analytics (Würzburg, Germany). Hoechst 33324, DRAQ5, Calcein AM, Calcein Deep Red, tetramethylrhodamine (TMRM) and CellTracker™ Red CMTPX dye were from Molecular Probes™ (Loughborough, UK). DiOC₆ iodide was supplied by Enzo Life Sciences

(Exeter, UK). Ibidi slides were from Thistle Scientific (Glasgow, UK). Horm fibrillary collagen was from Takeda Pharma (Linz, Austria). Cross-linked collagen-related peptide (CRP-XL) was provided by Richard Farndale (University of Cambridge, UK). Unless stated, all other reagents were from Sigma-Aldrich (Dorset, UK).

5

Animals

C57BL/6 mice were purchased from Harlan UK. All animal studies were approved by local research ethics committee (AWERB) and licensed under UK Home Office project license PPL 30/3445.

10

Culture and Differentiation of Murine Megakaryocytes

Briefly, bone marrow from C57BL/6 or *TPM4^{-/-}* mice was isolated and dispersed prior to centrifugation at 200 x g for 10 minutes. Bone marrow was re-suspended in IMDM- Glutamax containing 1% penicillin/streptomycin and 2% serum replacement 1. Cells were cultured for 3 days in the presence of 20 ng/mL rSCF and a further 13 days in the presence of 10 ng/mL rTPO at 37 °C and 5% CO₂. From day 8 to day 16, cells were transmitted to fresh cell culture dishes every day to reduce megakaryocytes (MKs) to contact with fibroblast cells.

15

On day 16, MKs were harvested and enriched with a 1.5%-3% albumin gradient for one hour in cell incubator, following stained with either IgG-PE/ or FITC, or CD41-PE or -FITC for 3 hours and further with DNA dye Hoechst 33342 for 20 mins. Then MKs were washed and resuspended in 2 mL medium containing 10 U/mL heparin prior to use.

20

Mouse *ex vivo* heart-lung preparation

Mice aged at 8-12 weeks were sacrificed by a Schedule 1 process, by exposure to rising CO₂. A tracheostomy was then performed, and lungs ventilated with room air or pure nitrogen via a rodent ventilator (Minivent type 845, Harvard Apparatus, USA). Respiratory rate was maintained at 200
5 breaths/min and tidal volume was 10 mL/kg (~250 μ L). The inferior vena cava was then exposed by laparoscopy, and the chest opened by median sternotomy and fat tissue carefully removed.

A 21-gauge catheter was passed into the right ventricle and approx. 100 mL warm (36-38 °C) perfusion buffer (Krebs-Hepes buffer (KHB): 140 mM NaCl, 3.6 mM KCl, 0.5 mM NaH₂PO₄, 0.2 mM MgSO₄, 1.5 mM CaCl₂, 10 mM Hepes (pH 7.4), 2 mM NaHCO₃) containing 10 U/mL heparin
10 was perfused at 3-4 mL/min into the lung vasculature. Once the colour of mouse lung turned pale, suggesting most of blood in the lung circulation was flushed, the catheter was disconnected.

The left and right superior venae cavae, ascending aorta, inferior vena cava and descending aorta were ligated using 6-0 silk, and a small incision was made in the left ventricle for collection of the perfusate. A suspension of mouse MKs (usually labelled with a fluorescently tagged anti-CD41
15 Ab) was pumped into the right ventricle, whilst the lungs were either ventilated with air or N₂ or without ventilation, and the flow rate was maintained at 0.35 mL/min by a SyringeOne Programmable Syringe Pump (Product SKU: NE-1000-ES). The perfusate was collected and re-pumped into the right ventricle after samples were collected for imaging and FACS analysis. The perfusate was recirculated through the pulmonary vasculature in this manner for a total of 18
20 passages.

After the final passage, 1 mL perfusion buffer was injected into the system to remove some of the remaining cells in the lung vasculature, and all perfusate collected for evaluation of platelet

function. The mouse lung was immediately removed and fixed in 4% PFA/PBS. The volume of the lung was measured by fluid displacement.

Flow cytometry

5 Platelets derived from IgG-PE stained MKs were set up as negative controls. Washed mouse platelets were used to set gates for generated platelets (P1) at the FSC/SSC density plot by size and granularity. 25 μ L of CD41-PE-stained MK suspension, or perfusates from lung or suspensions from microfluidic chambers after 1, 2, 3, 6, 9, 12, 15 or 18 passages, were fixed in 2% PFA/PBS before FACS. DNA content, viability and mitochondrial membrane potential of
10 generated platelets were determined by Draq5, Calcein AM and TMRM staining, respectively.

To detect the viability and mitochondrial membrane potential of pulmonary endothelial cells (ECs), assays for Calcein Deep Red retention in ECs and TMRM accumulation in active mitochondria were conducted. Pulmonary ECs were isolated from perfused lungs under air- or
15 pure-N₂-ventilation or unventilated for approximately 2 hours and marked with either anti-CD31/PECAM-1 or anti-CD102/ICAM-2 antibodies, followed by loading with Calcein Deep Red and TMRM dyes, respectively, for 25 mins at room temperature. Pulmonary ECs from fresh lungs served as control.

Platelet surface glycoproteins were measured by incubating with FITC-conjugated anti-mouse
20 CD61, CD42b, CD42d, CD49b, Glycoprotein VI (GPVI) antibodies and isotype-nonspecific IgG for 20 mins at room temperature before fixation.

To investigate the function of generated platelets (platelets derived from CD41 stained MKs), assays for $\alpha_{IIb}\beta_3$ integrin activation (JON/A-binding) and P-selectin exposure were performed. Washed platelets were stimulated with 2 U/ml thrombin or 5 μ g/mL CRP for 10 mins followed by

incubation with PE-JON/A or PE-P selectin antibodies for 20 mins at room temperature before fixation. Tirofiban and PE-IgG were used as negative controls for the measurement of PE-conjugated JON/A or PE-conjugated P-selectin exposure, respectively.

Samples were analysed on a BD Accuri™ C6 Plus flow cytometer (BD) with 50,000 gated events/sample.

Two Photon Imaging

After overnight fixation at 4 °C, the lung was staged on a microslicer device (Type: DTK-1000N, UK) and cut into small slices with 800 µm thickness and flat surface. Lung slices from different lobes were fixed into a 100 mm cell culture dish and immersed in PBS for two photon imaging.

Imaging was performed using a DeepSee multiphoton laser (Spectra Physics) attached to an upright SP8 confocal microscope (Leica Microsystems). All images were collected using a 25x/0.95NA water dipping lens. For CD41-FITC or CD41-PE, excitation was provided by tuning the multiphoton laser to 927nm, for Hoechst33342, excitation was provided by tuning the multiphoton laser to 750nm. The resultant fluorescence for both scans passing off a SP500 dichroic beam splitter and through both a SP680 filter and either a 525/50 nm bandpass filter to selected only CD41-FITC signal, a 630/75 nm bandpass filter to select only CD41-PE signal or a 460/50 nm bandpass to select only Hoechst33342 signal. CD41-FITC or CD41-PE fluorescence was detected using non-descanned Hybrid detectors (Leica Microsystems) and Hoechst33342 fluorescence was detected using a non-descanned PMT.

Images were acquired with an additional zoom of 2.5x with 1704x1704 pixels (XY) resulting in an effective pixel size of 100 nm. Z stacks were captured with a z-step spacing of 2 µm. All images were capture using a scan speed of 400 Hz with a bidirectional scan.

Confocal Microscope Imaging

Confocal images were obtained on an inverted SP8 confocal microscope (Leica Microsystems) attached to a DMI6000 microscope frame (Leica Microsystems). All images were acquired using a 100x/1.44 NA oil immersion objective. Excitation was provided by either a 405 nm laser (Hoechst33324) or 488 nm laser (for CD41-FITC) or 561 nm laser (for CD41-PE) and the resultant fluorescence was detected using a Hybrid detector in the range 410-470 nm for Hoechst 33342, 495-570 nm for CD41-FITC or 571-650 nm for CD41-PE. A total of 50 randomly chosen fields of view were imaged and where z-stacks were acquired, a 2 μ m z-step spacing was used for Extended Data Video 6-7, or a 0.5 μ m z-step spacing for Extended Data Video 8.

To obtain an estimate of the dimensions of each nucleus in XY, z-stacks were loaded into FIJI and subjected to a maximum intensity projection and a region of interest (ROI) was manually drawn around an individual nucleus. The inbuilt measure function within imageJ was used. To obtain the dimensions of the major and minor axis, the 'fit ellipse' option was enabled. To estimate the nuclear depth, analysis was performed in FIJI. Image stacks were loaded, and a cell nucleus was selected using the ROI tool. The average intensity profile of this nucleus was plotted in z and then fitted with a gaussian profile using the built-in plot and fitting tools of FIJI. From the fitted parameters the depth of the nucleus was estimated using the full width at half maximum (FWHM) of the gaussian fit. The FWHM was calculated as $2\sqrt{(2\ln 2)} \sigma$. This process was manually repeated for multiple cell nuclei.

Microfluidic chamber design, fabrication, and experimental protocol

A set of channels mimicking a physiological vascular system was constructed using standard PDMS approach. The design shows a branching structure such that as branching progressing, the channel diameter decreases by half. All channels are rectangular in cross-section and 10 μ m deep,

with the largest channels being 100 μm across, decreasing to the smallest channels which are 12.5 μm across. From each larger channel, 16 smaller channels branch off, allowing for maintenance of flow resistance due to the r^4 power relationship between resistance and channel diameter. Fluid then flows from larger diameter channels to smaller diameter channels, and in reverse on the way out of the system. Cells are passed through the system, repeatedly. The system is scaled up, through multiplexing in parallel, to allow greater cell volumes to be used, as shown in Fig. 1i and Extended Data Fig. 1c.

The mask and SU-8 master mold were fabricated by NuNano (Bristol, UK). The microfluidic channels were fabricated by soft lithography. The mixture of PDMS in a 10:1 ratio was poured over the SU-8 master mold after being degassed in a vacuum desiccator. The PDMS mixture was cured at 80 °C for 2 h and incubated in the oven overnight. The PDMS mold was removed from the SU-8 master. Input (Cell-in) and output (Cell-out) holes were punched using a 0.5 mm OD biopsy puncher (Elveflow, Paris, France). Finally, the PDMS microchannels were irreversibly bonded to glass slides using oxygen plasma for 3 min in a plasma device (Diener Plasma Systems, Ebhausen, Germany).

A suspension of mouse MKs pre-labelled with CD41-PE was pumped into the microfluidic chamber at 0.30 mL/min flow rate. Suspensions were collected from the Cell Out-hole and re-pumped into the microfluidic chamber, after 25 μL samples were taken for FACS analysis. Suspensions were then recirculated through the microfluidic chamber in this manner for a total of 18 passages.

Transmission Electron Microscope Imaging

To visualize and compare the ultrastructures of generated and control platelets by transmission electron microscopy (TEM), host platelets were first depleted by intraperitoneal administration of

anti-GPIIb α antibody R300 (2 μ g/g bodyweight) prior to MK infusion through the heart-lung preparation. After 18 passages, the generated platelets were pelleted by centrifuging the collected perfusate. The platelet pellet was then resuspended in a cacodylate-buffered glutaraldehyde fixative and fixed at 4-8°C overnight and then post-fixed in osmium ferrocyanide. Fixed cells were then embedded in a solidifying BSA/glutaraldehyde gel. Gel-embedded platelets were stained with uranyl acetate and lead aspartate followed by dehydration with ethanol and infiltrated with Epon resin in a Tissue Processor (Leica EMTP). 70nm sections were cut from blocks with a Reichert Ultracut E and imaged with a Tecnai 12 electron microscope (ThermoFisher UK).

***In Vitro* Thrombus Formation**

In brief, ibidi μ -Slide VI^{0.1} chips were coated with 50 μ g/mL collagen overnight at 4 °C before being flushed and blocked with 2% fatty acid-free BSA prepared in HEPES-Tyrode's buffer. Freshly drawn mouse blood was collected from the inferior vena cava using 10 U/mL heparin and 20 μ M PPACK (D-phenylalanyl-prolyl-arginyl chloromethyl ketone) as anticoagulant, following euthanasia by rising CO₂. Mouse blood was mixed with 2 mL NaCl (150 mM) and centrifuged to remove PRP. Megakaryocytes stained with 2 μ M DiOC₆ were passaged through the mouse-lung preparation 18 times, and approx. 0.7 mL perfusate was mixed with 1.3 mL mouse blood lacking PRP and incubated with CellTracker™ Red CMTPX dye (1:1000 dilution) for 10 mins.

The mixed sample was then transferred to a 5 mL syringe and perfused using an Aladdin AL-1000 syringe pump (World precision instruments, United Kingdom) through the ibidi slide, at a shear rate of 1000/s for 20 minutes. Platelets were fixed by perfusion of 4% paraformaldehyde through channels for 4 minutes before nonadherent cells were flushed away with HEPES Tyrode buffer.

Thrombus formation was determined by generating confocal z-stacks (1024×1024 pixels, 0.787 μ m z stack distance) from 5 randomly chosen fields of view using a Leica SP8 confocal microscope. Images were acquired using a 20x/0.7 NA air immersion objective. Excitation was

provided by either a 488 nm (DiOC₆) or 561 nm (CellTracker™ Red CMTPX) laser with the resultant fluorescence being detected by Hybrid detectors in the range 498-551 nm (DiOC₆) or 571-623 nm (CellTracker Red CMTPX).

5 **Analysis and statistics**

All data were analyzed using GraphPad Prism 7 software (GraphPad Software Inc., San Diego, CA, USA). All image analysis was performed using ImageJ-win 64 software. Quantified data are presented as mean ± SEM from at least 3 independent experiments. A value of *P < 0.05 was considered statistically significant and was determined using either a paired Student's t-test or two-
10 way ANOVA with Dunnett's or Bonferroni post hoc test for multiple comparison.

Data and materials availability: The datasets generated during and/or analyzed during the current study are available from the corresponding authors on reasonable request.

Fig. 1: Mouse platelets are generated in physiological quantities from megakaryocytes passed multiple times through mouse pulmonary vasculature *ex vivo*. **(a)** Diagram illustrating the approach to generating mouse platelets from megakaryocytes (MKs) by infusion through the heart-lung preparation. **(b)** Intact MKs (inset: showing a circular shape and central nucleus, scale bar: 10 μ m) were imaged from samples of perfusates after passing the indicated number of times through lung vasculature. Quantification was from at least 250 fields of view, counting at least 230 cells in total and displayed as a percentage of total number of cells. Numbers above each column are passage number against which the column show a significant difference. Data are from at least 5 independent experiments. **(c)** Gating strategy for quantification of generated platelets. The number of generated platelets in the perfusate collected after the 18th passage was determined by the number of CD41(+) events in gate P1. **(d)** The number of generated

platelets per MK present in the perfusates from different passage numbers in lungs either ventilated with air, 100% nitrogen, or without ventilation (as indicated), or from microfluidic chamber (shown diagrammatically in **i**) were measured by FACS. Data are mean and S.E.M. (n as indicated). **(e)** Stained MKs were passaged through pulmonary vasculature *ex vivo* 18 times, and lung tissue visualized at 20 stacked focal planes by two-photon microscopy. During the cell passage lungs were either ventilated with air or 100% nitrogen, or were not ventilated, as indicated. Mouse lung without MKs passed through served as control. Images shown are representative of at least 4 independent experiments. scale bar: 20 μ m. **(f)** Numbers of platelets generated per MK in perfusate and remaining in mouse lung under air ventilation, were calculated and displayed as means $\pm$ S.E.M (n as indicated). As a control, MKs passed through 21G needles 18 times generated no platelets, as indicated. **(g-h)** The viability **(g)** and mitochondrial membrane potential **(h)** of pulmonary endothelial cells (ECs) were measured by FACS. Data are from 5 independent experiments. **(g)** The viabilities of pulmonary ECs were determined by Calcein Deep Red retention and displayed as means of % $\pm$ S.E.M. **(h)** The mitochondrial membrane potential was determined by accumulation of tetramethyl rhodamine methyl ester (TMRM) in active mitochondria and displayed as mean intensity $\pm$ S.E.M. **(i)** Design of microfluidic chamber.

Fig. 2: Generated platelets demonstrate typical physical features comparable to control platelets. (a) Events in P1 (red) are defined as generated platelets, with higher mean fluorescence compared to those derived from control IgG-PE- treated MKs. The viability of the generated platelets, and whether they contain DNA, were checked by Calcein AM and DRAQ5 dyes, respectively. (b) Mitochondrial membrane potential was monitored by FACS in generated and control platelets. (c) TMRM signals from (b) were quantified and displayed as mean $\pm$ S.E.M

(n=5). **(d)** Platelets were defined by staining with anti-CD41 PE conjugated antibody. Surface glycoproteins of platelets were measured by FACS on generated platelets and control platelets after staining with different FITC-conjugated antibodies as indicated. Data are presented as mean $\pm$ S.E.M (n=6) of mean FITC fluorescence intensities. **(e)** Perfusates, containing both generated platelets (stained green with CD41-FITC, white arrow) and host platelets, were stained for α -tubulin (stained magenta, blue arrow), and confocal images shown as a mixed population in the top panels. More detailed images of α -tubulin rings are shown in the magnified images in the middle panel (generated platelets) and bottom panel (control platelets). Images are representative of 3 independent experiments. Scale bars: 2 μ m. **(f)** The diameter of platelets (40 platelets from 3 independent experiments) from (E) was measured using Fiji (ImageJ-Win64). **(g)** Ultrastructures of generated platelets and control platelets visualized by transmission electron microscopy (TEM). Low magnification images are shown for generated platelets in I-III, and control platelets in IV. Detailed images, taken from the low magnification images, are shown in (A)-(D), as indicated. Subcellular structures are shown and annotated as abbreviations: α -G, α -granules; σ -G, σ -granules or dense bodies; Mit, mitochondria; OCS, open canalicular system; MTC, microtubule coils; RBC, red blood cells. Scale bars: 2 μ m in I-IV, 500 nm in A-D. Images shown are representative of 5 independent experiments.

Fig. 3: Generated platelets are functionality comparable to control platelets. Mouse MKs, labelled with FITC-conjugated anti-CD41 antibody, were passaged repeatedly through the pulmonary vasculature *ex vivo*. Lungs were ventilated with air throughout. **(a)** Representative FACS histograms of integrin α IIb β 3 activation (JON/A binding) and P-selectin expression induced by 2 U/ml thrombin (Magenta) or 5 μ g/ml CRP (green) in washed generated platelets under air ventilation in comparison to washed control platelets. Tirofiban (80 ng/ml) was used as control for non-specific binding in measurements of integrin activation. **(b)** Data from (A) were expressed as mean fluorescence intensity (a.u.). Data are mean and S.E.M. (n as indicated). **(c)** Images of 3 planes of a representative platelet-rich thrombus. Cross-sectional image is shown on the far right. Generated platelets (stained with both DiOC6 and CellTracker™ Red CMTPIX dye, white) occupied all levels of the thrombus while host platelets (stained with CellTracker™ Red CMTPIX dye alone, magenta) were mainly situated on top of thrombus. Images are representative of 5 independent experiments. Scale bars as indicated. **(d-e)** MFI (mean fluorescence intensity) profiles

of R1 and R2 (Region 1 and Region2 from Fig. 3C) along the z-axis. **(d)** MFI profile of R1 along the z-axis. In R1, generated platelets occupied the lower part of the thrombus up to $\sim 12\ \mu\text{m}$ (both green and magenta signals increased simultaneously), whilst beyond this point host platelets were predominant (magenta signals were stronger than green beyond $12\ \mu\text{m}$). **(e)** MFI profile of R2 along the z-axis. In R2, this part of thrombus was composed only of generated platelets as both green and magenta signals changed simultaneously along z-axis.

Fig. 4: Megakaryocytes show nuclear marginalization and enucleation prior to TPM4-dependent fragmentation. (a-e) MKs from C57BL/6 mice, labelled with PE-conjugated anti-CD41 antibody, (f-i) MKs from *Tpm4*^{-/-} mice, labelled with FITC-conjugated anti-CD41 antibody. MKs were passaged repeatedly through the pulmonary vasculature of a C57BL/6 mouse *ex vivo*. Lungs were ventilated with air throughout. (a) Representative images of MK derivatives during the process of platelet generation: (I) nuclear polarization and enucleation, where the nucleus is marginalized, of irregular shape or in the process of ejection from the cell; (II) naked nuclei, where the ejected nucleus is larger than 20 μm in diameter and free from the parent cell and may be partially encased in a thin cytoplasmic layer; (III) MKs without nuclei, where the MKs have an approximately circular shape; and (IV) large anuclear objects, where ghost cells are of irregular shape and larger than 10 μm in their longer axis. (b) & (h) Cells were imaged from samples of perfusates after passage numbers 1, 2, 3, 6 & 9 through murine lung vasculature *ex vivo*. Cells

were morphologically classified as 5 subgroups: intact Mks (Fig.1b) and (I)-(IV) above and quantified as a percentage of total number of cells. Quantification was from at least 250 **(b)** or 150 **(h)** fields of view, counting at least 230 **(b)** or 170 **(h)** cells in total for each of the subgroups. Numbers above each column are passage numbers against which the column shows a significant difference (p<0.05). Asterisk above columns in (h) represents significant difference to corresponding wild-type column in **(b)**. Data are from at least 5 **(b)** or 3 **(h)** independent experiments. **(c-e)** Nuclear lobes of MKs fragment into small condensed sub-nuclei. **(c)** Representative images, from n=3, of naked nuclei generated after multiple passages (as indicated) of mouse MKs through pulmonary vasculature. Scale bars: 5 μ m. **(d-e)** The depth **(d)** and aspect ratio **(e)** of sub-nuclei decreased substantially with increasing passage number. For depth **(d)**, the decrease was from ~8.9 μ m after P3 to 5.3 μ m after 18 passages (P18). For aspect ratio **(E)**, the decrease was from ~2 after P3 to nearly 1 after P18. Each symbol represents one sub-nucleus. Numbers above each column are passage numbers against which the column shows a significant difference. Data are from 3 independent experiments. **(f-g)** Numbers of generated platelets per *Tpm4*^{-/-} MKs, or control WT MKs, in perfusates after different passage numbers through WT lung, were quantified by FACS. **(f)** Representative FACS dot plot images for generated platelets (CD41-FITC positive events within the red square). **(g)** *Tpm4*^{-/-} platelets were consistently undetectable after up to 18 passages, in the perfusate. Data shown are platelets generated per MK from either *Tpm4*^{-/-} MKs or control WT MKs and displayed as mean +/- S.E.M. (n as indicated). **(h)** see above, with **(b)**. **(i)** Abundant fluorescent objects, 10-12 μ m diameter, were visible in sections of mouse lung after 18 passages of stained *Tpm4*^{-/-} MKs, as shown in extended focus stacks of 20 continuous two-photon planes of lung. Images shown are representative of 3 independent experiments. Scale bar: 20 μ m. **(j)** Schematic diagram of the steps

in platelet generation from mature MK. These include nuclear polarization, enucleation, cytoplasmic fragmentation into platelets and nuclear condensation and fragmentation.

5

10

15

20

References

- 1 Denis, M. M. *et al.* Escaping the nuclear confines: signal-dependent pre-mRNA splicing in anucleate platelets. *Cell* **122**, 379-391, doi:10.1016/j.cell.2005.06.015 (2005).
- 2 van der Meijden, P. E. J. & Heemskerk, J. W. M. Platelet biology and functions: new
5 concepts and clinical perspectives. *Nature reviews. Cardiology* **16**, 166-179,
doi:10.1038/s41569-018-0110-0 (2019).
- 3 Gaertner, F. *et al.* Migrating Platelets Are Mechano-scavengers that Collect and Bundle Bacteria. *Cell* **171**, 1368-1382.e1323, doi:10.1016/j.cell.2017.11.001 (2017).
- 4 Heimark, R. L., Twardzik, D. R. & Schwartz, S. M. Inhibition of endothelial regeneration
10 by type-beta transforming growth factor from platelets. *Science (New York, N.Y.)* **233**,
1078-1080, doi:10.1126/science.3461562 (1986).
- 5 Li, Y. J. *et al.* From blood to brain: blood cell-based biomimetic drug delivery systems. *Drug delivery* **28**, 1214-1225, doi:10.1080/10717544.2021.1937384 (2021).
- 6 Machlus, K. R. & Italiano, J. E., Jr. The incredible journey: From megakaryocyte
15 development to platelet formation. *The Journal of cell biology* **201**, 785-796,
doi:10.1083/jcb.201304054 (2013).
- 7 Brown, E., Carlin, L. M., Nerlov, C., Lo Celso, C. & Poole, A. W. Multiple membrane
extrusion sites drive megakaryocyte migration into bone marrow blood vessels. *Life*
science alliance **1**, doi:10.26508/lsa.201800061 (2018).
- 20 8 Potts, K. S. *et al.* Membrane budding is a major mechanism of in vivo platelet biogenesis. *The Journal of experimental medicine* **217**, doi:10.1084/jem.20191206 (2020).
- 9 Howell, W. H. & Donahue, D. D. THE PRODUCTION OF BLOOD PLATELETS IN
THE LUNGS. *The Journal of experimental medicine* **65**, 177-203,
doi:10.1084/jem.65.2.177 (1937).
- 25 10 Zucker-Franklin, D. & Philipp, C. S. Platelet production in the pulmonary capillary bed:
new ultrastructural evidence for an old concept. *The American journal of pathology* **157**,
69-74, doi:10.1016/s0002-9440(10)64518-x (2000).
- 11 Lefrançois, E. *et al.* The lung is a site of platelet biogenesis and a reservoir for
haematopoietic progenitors. *Nature* **544**, 105-109, doi:10.1038/nature21706 (2017).
- 30 12 R M Kaufman, R. A., S Pollack, W H Crosby. Circulating megakaryocytes and platelet
release in the lung. *Blood Rev* **26**, 12 (1965).
- 13 Kaushansky, K. *et al.* Promotion of megakaryocyte progenitor expansion and
differentiation by the c-Mpl ligand thrombopoietin. *Nature* **369**, 568-571,
doi:10.1038/369568a0 (1994).
- 35 14 Huertas, A. *et al.* Pulmonary vascular endothelium: the orchestra conductor in respiratory
diseases: Highlights from basic research to therapy. *The European respiratory journal*
51, doi:10.1183/13993003.00745-2017 (2018).
- 15 Nurden, A. T. Clinical significance of altered collagen-receptor functioning in platelets
with emphasis on glycoprotein VI. *Blood Rev* **38**, 100592,
doi:10.1016/j.blre.2019.100592 (2019).
- 40 16 Moog, S. *et al.* Platelet glycoprotein V binds to collagen and participates in platelet
adhesion and aggregation. *Blood* **98**, 1038-1046, doi:10.1182/blood.v98.4.1038 (2001).
- 17 Sadoul, K. Tubulin acetylation a valuable accessory of the platelet cytoskeleton. Focus on
"Histone deacetylase 6-mediated deacetylation of α -tubulin coordinates cytoskeletal and
45 signaling events during platelet activation". *American journal of physiology. Cell*
physiology **305**, C1211-1213, doi:10.1152/ajpcell.00309.2013 (2013).

- 18 Dmitrieff, S., Alsina, A., Mathur, A. & Nédélec, F. J. Balance of microtubule stiffness and cortical tension determines the size of blood cells with marginal band across species. *Proceedings of the National Academy of Sciences of the United States of America* **114**, 4418-4423, doi:10.1073/pnas.1618041114 (2017).
- 5 19 Hirsh, J. Platelet age: its relationship to platelet size, function and metabolism. *British journal of haematology* **23**, Suppl:209-214, doi:10.1111/j.1365-2141.1972.tb03520.x (1972).
- 20 21 Thompson, C. B., Jakubowski, J. A., Quinn, P. G., Deykin, D. & Valeri, C. R. Platelet size and age determine platelet function independently. *Blood* **63**, 1372-1375 (1984).
- 10 22 Vainchenker, W. & Raslova, H. Megakaryocyte polyploidization: role in platelet production. *Platelets* **31**, 707-716, doi:10.1080/09537104.2019.1667497 (2020).
- 22 23 Gunning, P. W., Hardeman, E. C., Lappalainen, P. & Mulvihill, D. P. Tropomyosin - master regulator of actin filament function in the cytoskeleton. *Journal of cell science* **128**, 2965-2974, doi:10.1242/jcs.172502 (2015).
- 15 24 Pleines, I. *et al.* Mutations in tropomyosin 4 underlie a rare form of human macrothrombocytopenia. *The Journal of clinical investigation* **127**, 814-829, doi:10.1172/jci86154 (2017).
- 24 25 Townsley, M. I. Structure and composition of pulmonary arteries, capillaries, and veins. *Comprehensive Physiology* **2**, 675-709, doi:10.1002/cphy.c100081 (2012).
- 20 26 Mazzone, R. W. Influence of vascular and transpulmonary pressures on the functional morphology of the pulmonary microcirculation. *Microvascular research* **20**, 295-306, doi:10.1016/0026-2862(80)90030-8 (1980).
- 26 27 Dickinson, C. J. & Martin, J. F. Megakaryocytes and platelet clumps as the cause of finger clubbing. *Lancet (London, England)* **2**, 1434-1435, doi:10.1016/s0140-6736(87)91132-9 (1987).
- 25 28 Valdivia-Mazeyra, M. F. *et al.* Increased number of pulmonary megakaryocytes in COVID-19 patients with diffuse alveolar damage: an autopsy study with clinical correlation and review of the literature. *Virchows Archiv : an international journal of pathology*, 1-10, doi:10.1007/s00428-020-02926-1 (2020).
- 30 29 Pariser, D. N. *et al.* Lung megakaryocytes are immune modulatory cells. *The Journal of clinical investigation* **131**, doi:10.1172/jci137377 (2021).
- 29 30 Zinellu, A. *et al.* Platelet Count and Platelet Indices in Patients with Stable and Acute Exacerbation of Chronic Obstructive Pulmonary Disease: A Systematic Review and Meta-Analysis. *Copd* **18**, 231-245, doi:10.1080/15412555.2021.1898578 (2021).
- 35 31 Zhang, L. *et al.* The common risk factors for progression and mortality in COVID-19 patients: a meta-analysis. *Archives of virology* **166**, 2071-2087, doi:10.1007/s00705-021-05012-2 (2021).
- 31 32 Lehmann, T. *et al.* Platelet count and function at high altitude and in high-altitude pulmonary edema. *Journal of applied physiology (Bethesda, Md. : 1985)* **100**, 690-694, doi:10.1152/japplphysiol.00991.2005 (2006).
- 40 33 Prabhakar, A. *et al.* Venous thrombosis at altitude presents with distinct biochemical profiles: a comparative study from the Himalayas to the plains. *Blood advances* **3**, 3713-3723, doi:10.1182/bloodadvances.2018024554 (2019).
- 33 34 McDonald, T. P. & Sullivan, P. S. Megakaryocytic and erythrocytic cell lines share a common precursor cell. *Experimental hematology* **21**, 1316-1320 (1993).
- 45 35 Ji, P., Murata-Hori, M. & Lodish, H. F. Formation of mammalian erythrocytes: chromatin condensation and enucleation. *Trends in cell biology* **21**, 409-415, doi:10.1016/j.tcb.2011.04.003 (2011).

- 35 Roncati, L. *et al.* A proof of evidence supporting abnormal immunothrombosis in severe COVID-19: naked megakaryocyte nuclei increase in the bone marrow and lungs of critically ill patients. *Platelets* **31**, 1085-1089, doi:10.1080/09537104.2020.1810224 (2020).
- 5 36 Bender, M. *et al.* ADF/n-cofilin-dependent actin turnover determines platelet formation and sizing. *Blood* **116**, 1767-1775, doi:10.1182/blood-2010-03-274340 (2010).
- 37 Eckly, A. *et al.* Abnormal megakaryocyte morphology and proplatelet formation in mice with megakaryocyte-restricted MYH9 inactivation. *Blood* **113**, 3182-3189, doi:10.1182/blood-2008-06-164061 (2009).

Acknowledgments: We gratefully acknowledge the Wolfson Bioimaging Facility for their support and assistance in this work. We are also grateful to Professor Jack Mellor, University of Bristol, for the use of the tissue slicer machine. This work was supported by a Wellcome Trust Investigator Award to A.W.P and C.G (219472/Z/19/Z) and grants from the British Heart Foundation (RG/15/16/31758 to A.W.P, PG/16/102/32647 to A.W.P and E.O.A).

Author contributions:

XZ, CG and AWP contributed to conceptualization.

XZ, DA, TGW, NT, SZB, CRN, EWA, AKW and J B-B contributed to methodology.

XZ and AWP contributed to investigation and supervision.

XZ, DA and CRN contributed to cell imaging.

AWP, CG and EOA contributed to funding acquisition.

AWP was responsible for project administration.

XZ, AWP, DA and CRN contributed to writing the original draft.

XZ, AWP, DA, CMW, PWG, ECH, EOA, IH and CG reviewed and edited the manuscript.

Competing interests: The authors declare no competing interests.

Extended Data Fig. 1

10 Approach to generating mouse platelets from MKs by infusion through the pulmonary vasculature or a novel microfluidic chamber. (a) Experimental protocol for generating platelets from repeated infusion of mouse heart-lung preparation. (b) The lung volume after 18 passages was estimated by a fluid displacement technique. This was used to allow accurate estimation of

numbers of platelets generated that were still retained within the circulation of the lung. **(c)** Photograph of the microfluidic chamber, made in PDMS by soft lithography, with the arrangement of microchannels visible. Description of the channel architecture is provided in Fig. 1I. **(d)** Platelets generated in the mouse lung vasculature were defined by staining with anti-CD41 PE conjugated antibody. Surface glycoproteins of platelets were measured by FACS on generated platelets and control platelets after staining with different FITC-conjugated antibodies as indicated. Data are presented as % of FITC- positive events.

Extended Data Video 1.

3D reconstructed image of *ex vivo* mouse lung – air ventilation. The demarcation membrane system (DMS) and plasma membrane of mouse MKs were stained with anti-CD41-PE (red) and nuclei stained with Hoechst 33342 (cyan) and passaged through *ex vivo* mouse pulmonary circulation 18 times, whilst lungs were artificially ventilated with air. Images were taken by 2-photon microscopy, and reconstructions made from 20 z-stacks (in 2 μm steps) and reconstructed into 3D space shown in the video. Red events with $\sim 2\text{-}4\ \mu\text{m}$ in dimension were generated platelets. Images are representative of 4 independent experiments. Scale bar in this video: 20 μm .

Extended Data Video 2.

3D reconstructed image of *ex vivo* mouse control lung – air ventilation. To demonstrate background autofluorescence, lungs were artificially ventilated with air and imaged by 2-photon microscopy, and reconstructions made from 20 z-stacks (in 2 μm steps) and reconstructed into 3D space shown in the video. Images are representative of 4 independent experiments. Scale bar in this video: 20 μm .

Extended Data Video 3.

3D reconstructed image of *ex vivo* mouse lung – air ventilation. Mouse MK DMS and plasma membranes were stained with anti-CD41-FITC (green) and passaged through *ex vivo* mouse pulmonary circulation 18 times, whilst lungs were artificially ventilated with air. Images were taken by 2-photon microscopy, and reconstructions made from 20 z-stacks (in 2 μm steps) and reconstructed into 3D space shown in the video. Images are representative of 6 independent

experiments. Green events with ~ 2-4 μm in dimension were generated platelets. Scale bar in this video: 20 μm .

Extended Data Video 4.

3D reconstructed image of ex vivo mouse lung – N₂ ventilation. Mouse MK DMS and plasma membranes were stained with anti-CD41-FITC (green) and passaged through ex vivo mouse pulmonary circulation 18 times, whilst lungs were artificially ventilated with 100% N₂. Images were taken by 2-photon microscopy, and reconstructions made from 20 z-stacks (in 2 μm steps) and reconstructed into 3D space shown in the video. Images are representative of 4 independent experiments. Several MKs with very bright green fluorescence are trapped in the pulmonary circulation. Scale bar in this video: 20 μm .

Extended Data Video 5.

3D reconstructed image of in vitro thrombus formation under flow. Generated platelets (stained with DiOC6, plus CellTracker™ Red CMTPX, blue) and control platelets (stained with CellTracker™Red CMTPX alone, magenta) were mixed and perfused through the ibidi slide pre-coated with collagen, at a shear rate of 1000/s for 20 minutes. Images were taken by confocal fluorescence microscopy, and reconstructions made from 30 z-stacks (in 0.787 μm steps) and reconstructed into 3D space shown in the video. Images are representative of 5 independent experiments. Scale bar in this video: 20 μm .

Extended Data Video 6.

3D reconstructed confocal image of MK with an extruding giant lobulated nucleus adhering to a large cytoplasmic fragment. The DMS and plasma membrane of a MK was stained with

anti-CD41-PE (red) and nuclei stained with Hoechst 33342 (blue). Images were obtained on an inverted SP8 confocal microscope and 3D reconstruction image was made from 20 z-stacks at 2 μm z-step spacing. Images are representative of 8 independent experiments, Scale bar in the video: 10 μm .

5

Extended Data Video 7.

3D reconstructed confocal image of a naked lobulated nucleus with a completely separated large cytoplasmic fragment. The DMS and plasma membrane of a MK was stained with anti-CD41-PE (red) and nuclei stained with Hoechst 33342 (blue). Images were obtained on an inverted SP8 confocal microscope and 3D reconstruction image was made from 20 z-stacks at 2 μm z-step spacing. Images are representative of 8 independent experiments, Scale bar in the video: 10 μm .

10

Extended Data Video 8.

3D reconstructed confocal image of sub-nuclei. The DMS and plasma membrane of MKs were stained with anti-CD41-PE (red) and nuclei stained with Hoechst 33342 (blue) and passaged through *ex vivo* mouse pulmonary circulation 3 times, whilst lungs were artificially ventilated with air. Images were obtained on an inverted SP8 confocal microscope and 3D reconstructed image was made from 20 z-stacks at 0.5 μm z-step spacing. Nuclear lobes were partially fragmenting from a naked MK nucleus. Images are representative of 8 independent experiments. Scale bar in the video: 10 μm .

15

20

Extended Data Video 9.

3D reconstructed image of *ex vivo* mouse lung – air ventilation. The DMS and plasma membranes of *Tpm4^{-/-}* MKs were stained with anti-CD41-FITC (green) and passaged through ex

vivo mouse pulmonary circulation 18 times, whilst lungs were artificially ventilated with air. Images were taken by 2-photon microscopy, and reconstructions made from 20 z-stacks at 2 μm step spacing and reconstructed into 3D space shown in the video. Abundant fluorescent objects sized 10-12 μm , were visible in focus stacking of 20 continuous 2-photon planes. Images are representative of 3 independent experiments. Scale bar in this video: 20 μm .

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [ExtendedDataFig.1.jpg](#)
- [ExtendedDataVideo1.avi](#)
- [ExtendedDataVideo2.avi](#)
- [ExtendedDataVideo3.avi](#)
- [ExtendedDataVideo4.avi](#)
- [ExtendedDataVideo5.avi](#)
- [ExtendedDataVideo6.avi](#)
- [ExtendedDataVideo7.avi](#)
- [ExtendedDataVideo8.avi](#)
- [ExtendedDataVideo9.avi](#)