

# Psychological resilience and health-related quality of life in Swedish women with newly diagnosed breast cancer

Åsa Mohlin (✉ [asa.mohlin@med.lu.se](mailto:asa.mohlin@med.lu.se))

Lunds Universitet <https://orcid.org/0000-0002-2969-9981>

**Ulrika Axelsson**

Lunds Universitet

**Pär-Ola Bendahl**

Lunds Universitet

**Carl A. K. Borrebaeck**

Lunds Universitet

**Cecilia Hegardt**

Lunds Universitet

**Per Johnsson**

Lunds Universitet

**Ingalill Rahm Hallberg**

Lunds Universitet

**Lisa Rydén**

Lunds Universitet

---

## Research article

**Keywords:** Breast cancer, Psychological resilience, Health-related quality of life, Connor-Davidson Resilience Scale 25 (CD-RISC25), Short Form Health Survey (SF-36)

**Posted Date:** October 21st, 2019

**DOI:** <https://doi.org/10.21203/rs.2.16283/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Cancer Management and Research on November 1st, 2020. See the published version at <https://doi.org/10.2147/CMAR.S268774>.

# Abstract

**Purpose** Psychological resilience appears to be an important influencing factor in various aspects of health-related quality of life (HRQoL) in a context of adversity, e.g., being informed of a cancer diagnosis. The purpose was to investigate psychological resilience and health-related quality of life in Swedish women with newly diagnosed breast cancer in relation to demographic and clinicopathological characteristics.

**Methods** A population-based cross-sectional study was conducted including 517 women with breast cancer in the South Swedish Health Care Region. Participants were enrolled at the time of consultation for the diagnosis. Psychological resilience was assessed with the Connor-Davidson Resilience Scale 25 (CD-RISC25), and HRQoL was assessed with the Short Form Health Survey. The participants responded to questions regarding demographic variables. Clinicopathological data were collected from the Swedish National Quality Register for Breast Cancer.

**Results** The mean score for psychological resilience was 70.6, identifying 15% of included patients with a score lower than 58 (-1 standard deviation). The study cohort had significantly lower mean scores for several aspects of HRQoL compared with Swedish normative data. Multiple regression analyses demonstrated that psychological resilience was significantly associated with all domains of HRQoL after adjustment for demographic and clinicopathological factors.

**Conclusions** Higher levels of psychological resilience were significantly related to higher levels of HRQoL in Swedish women with newly diagnosed breast cancer and no modifying factor was identified. The assessment of psychological resilience at the time of breast cancer diagnosis might allow for early identification of women in need of more intense psychosocial support. Future studies are needed to identify a clinically relevant threshold of the CD-RISC25.

## Background

Breast cancer (BC) is the most prevalent cancer in women in the Western world [1]. Although the prognosis has dramatically improved, BC still causes considerable physical and psychological consequences for many women [2–6]. Receiving a BC diagnosis means emotional turmoil for the patient, although the reactions may differ due to, e.g., the personal stability of the individual [2]. Previous studies have shown that BC has a major impact on women's health-related quality of life (HRQoL) when measured by valid instruments, e.g., the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC-QLQ), the Functional Assessment of Cancer Therapy Breast Cancer (FACT-B) and the Short Form Health Survey (SF-36) [3]. As the incidence of BC increases and survivors live longer, HRQoL has become an increasingly important outcome measure for BC patients [2–6].

There has been a growing interest in determining the psychosocial factors that may enhance HRQoL in BC patients [4–7], and the positive impact of a supportive social network has been reported [8]. Additionally, psychological resilience, defined as the capacity to cope successfully with external stress,

has been described as a protective factor associated with HRQoL in BC patients [4, 6, 8–10]. Psychological resilience is presumed to be a dynamic process and may fluctuate during treatment [11–13]. Higher levels of psychological resilience do not make the patient resistant to distress, but they do help the patient adapt and recover when facing adversities [11, 14]. In studies including BC patients, higher levels of psychological resilience have been found to be associated with higher levels of HRQoL [4, 8, 10]. The present study used the commonly used Connor-Davidson Resilience Scale 25 (CD-RISC25) [15–16]. Prior BC studies using the CD-RISC25 are limited, and the reported mean scores on the CD-RISC25 (range 0–100) vary between 54.7 and 74.7 [8, 11, 12, 18–20]. These studies are characterized by small sample sizes or different time points of assessment, making it difficult to draw any firm conclusions.

To enhance the life situation of patients in severe distress, such as women with newly diagnosed BC, a deeper understanding of the factors that contribute to HRQoL is needed. The Swedish version of the SF-36 was used in the present study to assess HRQoL [21].

The purpose of this study was (a) to investigate psychological resilience, as measured by the CD-RISC25, at the time of BC diagnosis, (b) to investigate HRQoL, as measured by the SF-36, at the time of BC diagnosis and (c) to investigate the association between psychological resilience and HRQoL in relation to demographic and clinicopathological characteristics in Swedish women with newly diagnosed BC.

## Methods

### Study design and study cohort

This cross-sectional study was conducted within the greater prospective study *SCAN-B Resilience* [7] (NCT03430492) as part of the *Sweden Cancerome Analysis Network–Breast (SCAN-B) initiative* [22].

The SCAN-B is a large population-based study that includes almost 90% of all new BC patients from southern Sweden [22, 23] (NCT02306096). The SCAN-B cohort has good representativity in comparison to all women with BC reported in the Swedish National Quality Registry for Breast Cancer (NKBC) during the same time period [24]. Patients enrolled in SCAN-B at Blekinge County Hospital, Central Hospital Växjö, Hallands Hospital Halmstad and Helsingborgs Hospital were also invited to be enrolled in SCAN-B Resilience. Karlskrona (Blekinge), Växjö, Halmstad and Helsingborg are urban cities; however, Karlskrona, Växjö and Halmstad have more rural areas than Helsingborg. Helsingborg is a larger, multi-cultural city that was included in SCAN-B Resilience at a later stage.

In all, 607 Swedish women with primary BC diagnosed between February 2016 and September 2018 were included in SCAN-B Resilience. A total of 517 patients with complete assessments (CD-RISC25 and SF-36) and NKBC data (clinicopathological variables) were selected for this cross-sectional study (Figure 1). The participants were enrolled at the time they were informed of the BC diagnosis at Blekinge County Hospital (N = 149), Central Hospital Växjö (N = 175), Hallands Hospital Halmstad (N = 163) and

Helsingborgs Hospital (N = 30). The diagnostic work-up was performed approximately two to three weeks before the visit to the Breast Unit and included mammography, ultrasound and biopsy.

The inclusion rate of the present study was estimated to be approximately 70%. Approximately one-third of those not included gave no specific reasons for not wanting to participate. Another one-third declined to participate because of limitations due to physical or mental problems; e.g., some reported experiencing too much stress or shock after the diagnosis. For the remaining third, the reasons were more varied, e.g., language problems or a lack of time.

The patients were given oral and written information about SCAN-B Resilience by BC nurses after consenting to participate in SCAN-B. Written consent was provided before the participants completed the assessments. The participants completed the assessments before leaving the Breast Units after the cancer consultation at which they were informed about the diagnosis and the treatment plan. The participants completed the assessments electronically or on paper.

SCAN-B and SCAN-B Resilience were approved by the Ethics Committee at Lund University (Dnr 2009/658, 2010/383, 2012/58, 2013/459, 2015/277, 2015/522, 2016/944, 2017/875).

## Instruments

Social network, educational level and financial situation data were collected by questions added to the standardized instruments. The questions about educational level and financial situation are commonly used in Swedish population studies [25]. The question about social network was designed for the SCAN-B Resilience study and aimed to capture the presence of adults or children living in the same or separate households with whom the participant has an ongoing relationship.

Information on age, menstrual status, mode of detection, stage of BC, type of BC and primary therapy was collected from the NKBC register, which holds information on almost 100% of Swedish women diagnosed with BC since 2008 [24].

Psychological resilience was measured with the CD-RISC25 [15–16]. The Swedish version of the instrument was used; it was obtained from its author, who gave permission for its use. The instrument consists of 25 items that are ranked on a Likert scale from 0 to 4 points, from “Not true at all” to “True nearly all the time”. Examples of items are “Able to adapt to change”, “Think of self as strong person” and “Tend to bounce back after illness or hardship”. The full range of the scale is 0–100 points. Higher scores reflect greater psychological resilience. The CD-RISC25 has shown good validity and reliability in previous studies in which psychological resilience was measured in relation to different health problems, including cancer [15–17]. To our knowledge, no population-based studies have been conducted in Sweden; however, Connor and Davidson, in their original study, reported a mean score of 80.4 for psychological resilience in a US general population (N = 577) [15]. The Cronbach’s alpha of the CD-RISC25 was 0.91 in this study.

The Swedish version of the SF-36 was used to assess HRQoL [21]. The instrument consists of 36 items that are ranked on scales of varying types. Examples of items are “Compared to one year ago, how would you rate your health in general now?” and “During the past 4 weeks, to what extent has your physical health or emotional problems interfered with your normal social activities with family, friends, neighbours, or groups?”. The 36 items are grouped into eight domains: physical functioning (PF), role limitations due to physical problems (RP), bodily pain (BP), general health (GH), vitality (VT), social functioning (SF), role limitations due to emotional problems (RE) and mental health (MH). The items are recoded in the domains to scores between 0 and 100, where 0 represents the worst HRQoL and 100 the best HRQoL. The Swedish version of the SF-36 has been proven to be a valid and reliable instrument [21, 26–27]. Swedish norm data are available [21]. Permission to use the SF-36 was obtained from Optum (Optum Circle, Eden Prairie, MN USA). The Cronbach’s alpha of the SF-36 was 0.86 in this study.

## Statistics

Percentage, mean and standard deviation (SD), were calculated for the demographic variables, clinicopathological variables, and CD-RISC25 and SF-36 scores. The median was also calculated for age. Independent-samples t-tests were used to compare two groups, and one-way analysis of variance was used to compare three or more groups for the comparisons of the CD-RISC25 scores to different demographic and clinicopathological variables. Independent-samples t-test was also used to compare the mean SF-36 scores across the study cohort with Swedish normative data. Pearson correlation coefficient analyses were used to investigate the association between the CD-RISC25 and the SF-36. Multiple linear regression analyses were used to test the association between the CD-RISC25 and the SF-36 scores, both unadjusted and adjusted for demographic and clinicopathological variables. P-values < 0.05 were considered statistically significant, but because no adjustment for multiple testing was performed, some caution is warranted when interpreting the level of evidence for a specific test. The statistical analyses were conducted using the *Statistical Package for the Social Sciences* (SPSS) version 25.0 (IBM, Armonk, NY, USA).

## Results

### Study cohort

The median age of the women was 64 years (range 31–89), 81% were postmenopausal, 52% were diagnosed with stage I BC and 91% had invasive BC (Table 1). In 65%, BC was detected within a screening programme. Most patients (95%) planned to undergo surgery as primary therapy, whereas 5% planned to neoadjuvant therapy. Regarding demographic variables, 63% of the women were living with an adult/adults, 22% were living alone, 13% were living with an adult/adults and a child/children under 18 years, and 2% were living with a child/children only. In addition, 39% had more than two years of post-secondary education, and 90% answered yes to the question about their ability to pay an unexpected bill of SEK 11 000/EUR 1100.

## Psychological resilience

The mean score for psychological resilience was 70.6 (SD±12.7) (Figure 2). As depicted in Figure 2, 70% of the women in the cohort had a level of psychological resilience within 1 SD of the mean score. Approximately 15% of the women had a level of psychological resilience lower than 58 (1 SD below the mean score). Scores for psychological resilience were significantly higher in premenopausal women than in postmenopausal women ( $P = 0.005$ ) (Table 1). Women who lived with a child/children but with or without an adult/adults had significantly higher psychological resilience than women who lived alone or with an adult/adults only ( $P = 0.001$ ). None of the other demographic and clinicopathological variables included were significantly related to the CD-RISC25 score.

## Health-related quality of life

The study cohort had significantly lower mean scores for several aspects of HRQoL compared with Swedish normative data (Table 2). Except for VT ( $P = 0.780$ ), the psychological aspects of the SF-36; SF, RE and MH ( $P < 0.001$ ), were lower in the study cohort than in the normative sample. Among the physical domains of the SF-36; GH ( $P < 0.001$ ) scores were significantly lower in the study cohort. The mean score for BP ( $P < 0.001$ ) was significantly higher, indicating lower levels of pain in the study cohort compared to the normative data. For PF ( $P = 0.056$ ) and RP ( $P = 0.846$ ), there was no significant differences across the cohorts.

## Correlation between psychological resilience and health-related quality of life

Pearson's correlation coefficient analyses presented positive correlations between the CD-RISC25 and all eight domains of the SF-36 (Table 3). The correlations ranged between 0.20 and 0.37, indicating a low to moderate correlation between psychological resilience and all aspects of HRQoL ( $P < 0.001$ ).

## Multiple regression analyses between psychological resilience and health-related quality of life

Psychological resilience was found to be significantly associated with each of the eight domains of HRQoL ( $P < 0.001$ ). Unadjusted models and models adjusted for clinically relevant factors and covariables associated with psychological resilience are presented in Table 4. The coefficients for the CD-RISC25 with each of the eight domains of the SF-36 did not change substantially in the different models, although social network and menstrual status tended to have more confounding effects than the other factors. Psychological resilience was shown to have the strongest impact on GH and on the psychological aspects of the SF-36 (VT, SF, RE and MH).

## Discussion

To the best of our knowledge, this is the first time data from a population-based consecutive cross-sectional Swedish study cohort has been used to show that psychological resilience is associated with HRQoL in women with newly diagnosed BC. The study indicates that at the time of receiving the diagnosis, both psychological resilience and HRQoL scores are lower than those from population-based data.

The results are consistent with the findings of prior studies [8, 11, 12, 18–20] demonstrating the possible need for psychosocial support of BC patients as most of the women presented lower levels of psychological resilience compared to normative data reported by Connor and Davidson [15]. The present study included a cohort of 517 Swedish women with a mean psychological resilience score of 70.6 at the time of diagnosis. The mean CD-RISC25 score varied between 54.7 and 74.7 in earlier studies [8, 11, 12, 18, 19, 20]. Markovitz et al. (2015) reported higher levels of psychological resilience, 93.8, compared to other BC studies [14]. However, they used a 1- to 5-point Likert scale instead of the recommended 0- to 4-point Likert scale [15–16]. On the website of the CD-RISC25 authors, the mean score for the BC cohort in the study by Markovitz et al. [16] was recalculated to 68.8, which is close to the mean score in this study.

As psychological resilience is assumed to be a dynamic process, minor variations across BC studies can be explained by differences in the timing of the CD-RISC25 assessment [11, 13]. The time span from diagnosis to CD-RISC25 assessment differs widely in the reported studies [8, 11, 12, 18]. This indicates that some women were affected by events linked to the treatment, while others more or less had recovered their levels of psychological resilience at the time they completed the CD-RISC25. Likewise, the women included in the present study may already have been affected by the diagnostic work-up for suspected BC. Although the information about diagnosis and treatment was given right before they responded to the instruments, it is reasonable to assume that they were aware at that time that something was wrong. The diagnostic work-up was conducted two to three weeks before the consultation, and the women were advised to bring someone close to them. Similarly, BC patients in previous studies scored even lower on psychological resilience when it was measured close to therapy [11, 19, 20]. Other than the present study, Markovitz et al. (2015) is the only BC study to report a distinct time of inclusion [14]. In this Belgic BC cohort, psychological resilience was measured at the time of primary surgery, one to two weeks after the women had been informed of the diagnosis, and the mean score is in line with the results of this study. Our results indicate that as soon as at the time of the BC diagnosis, some women may need psychosocial support because the levels of psychological resilience at this early time point were lower than those based on population data [15].

HRQoL (GH, SF, RE and MH) was lower in the investigated women compared with Swedish normative data [21]. It is fair to assume, as with psychological resilience, that the fear of BC during the diagnostic work-up had already had an impact on the patients' HRQoL at the time of the diagnosis. Only a few of the previous BC studies have explored HRQoL in relation to psychological resilience, although they did not use the SF-36 [8, 10]. Despite differences in the instruments chosen, the similarities between the results

of our study and those of Ristevska-Dimitrovska et al. (2015) and Zhang et al. (2017) are a striking demonstration that higher levels of psychological resilience correspond to higher levels of HRQoL [8, 10]. In this study, significant correlations between the CD-RISC25 and all eight domains of the SF-36 were found. This agrees with findings reported by Ristevska-Dimitrovska et al. that psychological resilience is correlated to most aspects of HRQoL in Macedonian BC patients [10]. Additionally, Zhang et al. showed that psychological resilience was correlated with HRQoL in a Chinese cohort [8]. Using another measure for psychological resilience, Harms et al. (2018) found an association between the Protective Factors for Resilience Scale (PFRS) and the SF-36 in a cancer cohort that included BC patients [4]. Harms et al. demonstrated that the correlations between the psychological aspects of the SF-36 and the PFRS were stronger than the correlations between most of the physical aspects of the SF-36 [4]. The results of this study are consistent with those of Harms et al. as the strongest correlations in Swedish women were noted between the CD-RISC25 and GH, followed by the psychological aspects of the SF-36 (MH, VT, SF and RE).

The findings also indicate a significant relationship between psychological resilience and social network and menstrual status. Consequently, variables assumed to be related to psychological resilience, e.g., social network, were included in the multiple regression analyses. Wu et al. (2017) [11] investigated predictors of psychological resilience among Chinese women and found that age was negatively correlated with psychological resilience, which is in line with our finding that postmenopausal women had lower scores than premenopausal women. Similarly, Zhang et al. (2017), Huang et al. (2019) and Alizadeh et al. (2018) presented significant relationships between psychological resilience and social support/network in Chinese and Iranian women [8, 19, 20]. The regression analyses in this study demonstrated that the CD-RISC was significantly associated with each of the eight domains of the SF-36 in terms of both unadjusted models and models adjusted for demographic and clinicopathological factors. Social network and menstrual status tended to have more confounding effects than the other variables, including the stage of BC. After adjustment for social network and menstrual status, most of the coefficients for the CD-RISC25 did not distinctly change. Although psychological resilience was significantly regressed on all domains of the SF-36, the strongest impact of psychological resilience was again demonstrated on GH and the psychological aspects of the SF-36. Higher levels of psychological resilience can be considered to be a potential protective factor among Swedish BC patients for preserving their HRQoL in the context of adversity. Lower levels are a risk factor for emotional distress and impaired HRQoL [10, 14, 18].

The mean score of 70.6 with a SD of 12.7 in the study cohort indicates that a fair share of the women included had a level of psychological resilience lower than 58 (approximately 15% of the population). The application of a threshold of 58, which corresponds to less than 1 SD below the mean score, may identify patients in need of psychosocial support at this early stage. The understanding that psychological resilience is associated with HRQoL in Swedish BC patients could be clinically important in the future development of rehabilitation interventions. Improved knowledge of this association would present an opportunity to develop evidence-based interventions for those presenting lower levels of psychological resilience at the time of BC diagnosis.

More knowledge on psychological resilience in Swedish women with BC is needed, such as Swedish normative data for the CD-RISC25. Longitudinal studies are required in which changes in psychological resilience can be observed over time. Qualitative studies are necessary to describe psychological resilience more extensively. Because of the cross-sectional design of the present study, causal relationships between psychological resilience and other variables of BC patients cannot be established. Further limitations are the non-inclusion rate of approximately 30%. One reason women declined to take part in this study was experiencing too much stress after the BC diagnosis. It is possible that the most stressed patients might be the women with the lowest levels of psychological resilience, which could bias the results of this study towards patients with higher levels of psychological resilience.

Although this study had limitations, it is one of the largest studies published to date regarding psychological resilience and HRQoL in BC patients. Another strength is the adjustment for other variables that may influence HRQoL in these women. All patients were included at the time of diagnosis in the population-based SCAN-B study [22, 23]. Such early and coherent assessment of psychological resilience in relation to BC diagnosis has not been previously presented.

## Conclusions

This population-based study is the first to investigate the association between psychological resilience and HRQoL in Swedish women with newly diagnosed BC. At the time of diagnosis, psychological resilience and HRQoL appear to be impaired in these women. Higher levels of psychological resilience were related to higher levels of HRQoL independent of demographic and clinicopathological characteristics. The assessment of psychological resilience at the time of BC diagnosis might allow for the early identification of women in need of extended psychosocial support. The results of this study show that 15% of the patients had a level of psychological resilience lower than 58, which might be a putative level for stratifying patients for individualized rehabilitation to prevent impairment of HRQoL related to BC diagnosis.

## List Of Abbreviations

BC: Breast cancer

HRQoL: Health-related quality of life

SF-36: Short Form Health Survey

CD-RISC25: Connor-Davidson Resilience Scale 25

SCAN-B: Sweden Cancerome Analysis Network—Breast initiative

NKBC: Swedish National Quality Registry for Breast Cancer

PF: Physical functioning

RP: Role limitations due to physical problems

BP: Bodily pain

GH: General health

VT: vitality

SF: Social functioning

RE: Role limitations due to emotional problems

MH: Mental health

SD: Standard deviation

## Declarations

*Ethics approval and consent to participate:* All procedures performed were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Ethical approvals was obtained from the Regional Ethics Committee at Lund University, Sweden (2009/658, 2010/383, 2012/58, 2013/459, 2015/277, 2015/522, 2016/944, 2017/875). Written informed consent was obtained from all individual participants included in the study.

*Consent for publications:* Not applicable.

*Availability of data and materials:* The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

*Competing interest:* The authors declare that they have no competing interests.

*Funding:* The study was supported by grants from the Birgit Rausing Donation, the Faculty of Medicine, Lund University, the Mats Paulsson Foundation, the Mrs Berta Kamprad Foundation, The Biltema Foundation and the CREATE Health Cancer Center.

*Authors' contributions:* Study concept and design (UA, CAKB, PJ, IRH, LR); acquisition of data (UA, CH, PJ, LR); analysis and interpretation of data (ÅM, POB, IRH, LR); drafting of manuscript (ÅM); critical revision of manuscript for important intellectual content (UA, POB, CAKB, CH, PJ, LR); supervision (IRM, LR). All authors read and approved the final manuscript.

*Acknowledgements:* We acknowledge the SCAN-B steering group, the SCAN-B Resilience study group, the Regional Cancer Center South and the South Swedish Breast Cancer Group for all their support. We thank all patients and personnel at the participating hospitals.

# References

1. National guidelines Breast cancer. Regionalt cancercentrum Stockholm-Gotland; 2019. <https://www.cancercentrum.se/samverkan/cancerdiagnoser/brost/vardprogram/>. Accessed May 2019.
2. Izci F, Sarsanov D, Erdogan ZI, Ilgun AS, Celebi E, Alco G, et al. Impact of Personality Traits, Anxiety, Depression and Hopelessness Levels on Quality of Life in the Patients with Breast Cancer. *Eur J Breast Health*. 2018;14(2):105–11.
3. Montazeri A. Health-related quality of life in breast cancer patients: a bibliographic review of the literature from 1974 to 2007. *J Exp Clin Cancer Res*. 2008;27:32.
4. Harms CA, Cohen L, Pooley JA, Chambers SK, Galvao DA, Newton RU. Quality of life and psychological distress in cancer survivors: The role of psycho-social resources for resilience. *Psychooncology*. 2018.
5. Karlsen RV, Frederiksen K, Larsen MB, von Heymann-Horan AB, Appel CW, Christensen J, et al. The impact of a breast cancer diagnosis on health-related quality of life. A prospective comparison among middle-aged to elderly women with and without breast cancer. *Acta Oncol*. 2016;55(6):720–7.
6. Zhang T, Li H, Liu A, Wang H, Mei Y, Dou W. Factors promoting resilience among breast cancer patients: a qualitative study. *Contemp Nurse*. 2018;54(3):293–303.
7. Axelsson U, Rydén L, Johnsson P, Eden P, Mansson J, Hallberg IR, et al. A multicenter study investigating the molecular fingerprint of psychological resilience in breast cancer patients: study protocol of the SCAN-B resilience study. *BMC Cancer*. 2018;18(1):789.
8. Zhang H, Zhao Q, Cao P, Ren G. Resilience and Quality of Life: Exploring the Mediator Role of Social Support in Patients with Breast Cancer. *Med Sci Monit*. 2017;23:5969–79.
9. Haglund ME, Nestadt PS, Cooper NS, Southwick SM, Charney DS. Psychobiological mechanisms of resilience: relevance to prevention and treatment of stress-related psychopathology. *Dev Psychopathol*. 2007;19(3):889–920.
10. Ristevska-Dimitrovska G, Filov I, Rajchanovska D, Stefanovski P, Dejanova B. Resilience and Quality of Life in Breast Cancer Patients. *Open Access Maced J Med Sci*. 2015;3(4):727–31.
11. Wu Z, Liu Y, Li X, Li X. Resilience and Associated Factors among Mainland Chinese Women Newly Diagnosed with Breast Cancer. *PLoS One*. 2016;11(12):e0167976.
12. Fradelos EC, Papathanasiou IV, Veneti A, Daglas A, Christodoulou E, Zyga S, et al. Psychological Distress and Resilience in Women Diagnosed with Breast Cancer in Greece. *Asian Pac J Cancer Prev*. 2017;18(9):2545–50.
13. Dooley LN, Slavich GM, Moreno PI, Bower JE. Strength through adversity: Moderate lifetime stress exposure is associated with psychological resilience in breast cancer survivors. *Stress Health*. 2017;33(5):549–57.
14. Markovitz SE, Schrooten W, Arntz A, Peters ML. Resilience as a predictor for emotional response to the diagnosis and surgery in breast cancer patients. *Psychooncology*. 2015;24(12):1639–45.

15. Connor KM, Davidson JR. Development of a new resilience scale: the Connor-Davidson Resilience Scale (CD-RISC). *Depress Anxiety*. 2003;18(2):76–82.
16. The Connor-Davidson Resilience Scale. <http://www.connordavidson-resiliencescale.com>. Accessed May 2019.
17. Windle G, Bennett KM, Noyes J. A methodological review of resilience measurement scales. *Health Qual Life Outcomes*. 2011;9:8.
18. Ristevska-Dimitrovska G, Stefanovski P, Smichkoska S, Raleva M, Dejanova B. Depression and Resilience in Breast Cancer Patients. *Open Access Maced J Med Sci*. 2015;3(4):661–5.
19. Huang Y, Huang Y, Bao M, Zheng S, Du T, Wu K. Psychological resilience of women after breast cancer surgery: a cross-sectional study of associated influencing factors. *Psychol Health Med*. 2019:1–13.
20. Alizadeh S, Khanahmadi S, Vedadhir A, Barjasteh S. The Relationship between Resilience with Self-Compassion, Social Support and Sense of Belonging in Women with Breast Cancer. *Asian Pac J Cancer Prev*. 2018;19(9):2469–74.
21. Sullivan M, Karlsson J, Taft C, Ware JE. SF–36 Health Survey: Swedish manual and interpretation guide. Göteborg: Sahlgrenska sjukhuset, Sektionen för vårdforskning; 2002.
22. Saal LH, Vallon-Christersson J, Hakkinen J, Hegardt C, Grabau D, Winter C, et al. The Sweden Cancerome Analysis Network - Breast (SCAN-B) Initiative: a large-scale multicenter infrastructure towards implementation of breast cancer genomic analyses in the clinical routine. *Genome Med*. 2015;7(1):20.
23. Rydén L, Loman N, Larsson C, Hegardt C, Vallon-Christersson J, Malmberg M, et al. Minimizing inequality in access to precision medicine in breast cancer by real-time population-based molecular analysis in the SCAN-B initiative. *Br J Surg*. 2018;105(2):e158-e68.
24. Swedish National Quality Registry for Breast Cancer (NKBC) <https://www.cancercentrum.se/samverkan/cancerdiagnoser/brost/kvalitetsregister/>. Accessed May 2019.
25. Statistics Sweden. Surveys about living conditions (ULF/SILC). <https://www.scb.se/>. Accessed May 2019.
26. Sullivan M, Karlsson J, Ware JE, Jr. The Swedish SF–36 Health Survey I. Evaluation of data quality, scaling assumptions, reliability and construct validity across general populations in Sweden. *Soc Sci Med*. 1995;41(10):1349–58.
27. Sackey H, Sandelin K, Frisell J, Wickman M, Brandberg Y. Ductal carcinoma in situ of the breast. Long-term follow-up of health-related quality of life, emotional reactions and body image. *Eur J Surg Oncol*. 2010;36(8):756–62.

## Tables

**Table 1. Mean scores of psychological resilience (CD-RISC25) according to clinicopathological and demographic characteristics (N = 517)**

Variables		N	%	CD-RISC25	
				Mean (SD)	P-value
<b>All participants</b>				70.6 (12.7)	
<b>Age (years)</b>	Mean (SD): 62±11				
	≤ 64 (median)	269	52	71.3 (12.4)	0.239
	> 64 (median)	248	48	70.0 (13.0)	
<b>Menstrual status</b>	Premenopausal	93	19	74.0 (11.7)	0.005
	Postmenopausal	397	81	69.8 (12.7)	
	Unknown	27			
<b>Stage of breast cancer</b>	0	43	8	69.5 (12.8)	0.196
	I	269	52	69.8 (13.1)	
	II	164	32	71.4 (11.5)	
	III	39	8	73.9 (13.6)	
	Unknown	2			
<b>Type of cancer</b>	Carcinoma <i>in situ</i>	46	9	69.6 (13.0)	0.620
	Invasive cancer	450	91	70.6 (12.7)	
	Unknown	21			
<b>Mode of detection</b>	Screening	334	65	69.9 (12.4)	0.084
	Symptomatic	182	35	72.0 (13.0)	
	Unknown	1			
<b>Primary therapy</b>	Surgery	491	95	70.5 (12.8)	0.201
	Systemic therapy	26	5	73.7 (9.7)	
<b>Study site</b>	Halmstad	163	31	70.1 (13.1)	0.464
	Helsingborg	30	6	73.8 (14.5)	
	Karlskorna	149	29	71.1 (12.6)	
	Växjö	175	34	70.2 (12.0)	
<b>Social network</b>	Living alone	114	22	72.0 (12.1)	0.001
	Living with child/children < 18 years old only	11	2	74.4 (11.7)	

	Living with adult/adults and child/children < 18 years old	65	13	75.4 (11.4)	
	Living with adult/adults only	327	63	69.1(12.9)	
<b>Educational level</b>	Primary school < 9 years	70	14	70.9 (14.6)	0.734
	Primary school completed	74	14	69.3 (13.6)	
	Upper secondary education	92	18	70.8 (11.7)	
	Post-secondary education < 2 years	66	13	70.0 (12.3)	
	Post-secondary education ≥ 2 years	204	39	71.0 (12.1)	
	PhD (doctoral education)	11	2	75.5 (15.2)	
<b>Financial situation</b>	Able to pay an unexpected bill of SEK 11 000/EUR 1100	466	90	70.8 (12.5)	0.300
	Unable to pay an unexpected bill of SEK 11 000/EUR 1100	51	10	68.9 (14.3)	

Table 2. Health-related quality of life (SF-36) (N = 517)

Variables	Study cohort: Mean (SD)	Norm: Mean (SD)	P-values for differences between study cohort and norm data for SF-36
<b>SF-36</b>			
Physical functioning	84.4 (19.4)	86.2 (20.4)	0.056
Role-physical	81.3 (34.7)	81.6 (33.1)	0.846
Bodily pain	80.8 (20.9)	72.7 (26.5)	<b>&lt;0.001</b>
General health	70.2 (19.4)	75.1 (22.7)	<b>&lt;0.001</b>
Vitality	67.0 (23.1)	66.7 (23.2)	0.780
Social functioning	83.9 (22.4)	87.5 (20.8)	<b>&lt;0.001</b>
Role-emotional	76.6 (36.7)	84.0 (30.9)	<b>&lt;0.001</b>
Mental health	70.1 (21.5)	79.6 (19.4)	<b>&lt;0.001</b>

**Table 3. Correlations between psychological resilience (CD-RISC25) and health-related quality of life (SF-36) (N = 517)**

Variables	
	<b>CD-RISC25</b>
<b>SF-36</b>	<b>Pearson correlation coefficient***</b>
Physical functioning	0.21
Role-physical	0.20
Bodily pain	0.21
General health	0.37
Vitality	0.33
Social functioning	0.27
Role-emotional	0.26
Mental health	0.34


\*\*\* *P*-value < 0.001 for all correlation coefficients

**Table 4. Multiple linear regression analyses for psychological resilience (CD-RISC25) and health-related quality of life (SF-36) (N=517)**

	Model 1 - unadjusted	Model 2 - adjusted for social network	Model 3 - adjusted for social network and menstrual status	Model 4 - adjusted for social network, menstrual status and age	Model 5 - adjusted for social network, menstrual status, age and mode of detection	Model 6 - adjusted for social network, menstrual status, age, mode of detection and stage of breast cancer
Variables	CD-RISC25	CD-RISC25	CD-RISC25	CD-RISC25	CD-RISC25	CD-RISC25
	Coefficient***	Coefficient***	Coefficient***	Coefficient***	Coefficient***	Coefficient***
Physical functioning	0.320	0.277	0.259	0.265	0.270	0.272
Role-physical	0.548	0.503	0.479	0.485	0.494	0.490
Bodily pain	0.349	0.319	0.308	0.310	0.315	0.315
General health	0.559	0.554	0.596	0.595	0.600	0.599
Vitality	0.603	0.611	0.606	0.604	0.610	0.609
Social functioning	0.474	0.487	0.511	0.509	0.514	0.512
Role-emotional	0.760	0.777	0.818	0.819	0.828	0.824
Mental health	0.581	0.604	0.626	0.624	0.628	0.628


\*\*\* *P*-value < 0.001 for all coefficients

## Figures



**Figure 1**

Flow chart of the study cohort



**Figure 2**

Histogram of total scores for psychological resilience (CD-RISC25) (N = 517)