The Older The Better? The Strange Case of Empirical Ground Motion Models in the Near-Source of Moderate-To-Large Magnitude Earthquakes

Roberto Paolucci

Politecnico di Milano

Angela Chiecchio (✉ angela.chiecchio@polimi.it)

Politecnico di Milano <https://orcid.org/0000-0002-4608-8584>

Manuela Vanini

Politecnico di Milano

Research Article

Keywords: empirical ground motion models, near-source conditions, analyses of residuals, strong motion dataset

Posted Date: July 16th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-698487/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Bulletin of Earthquake Engineering on January 17th, 2022. See the published version at <https://doi.org/10.1007/s10518-021-01304-9>.

Abstract

This paper aims at providing a quantitative evaluation of the performance of a set of empirical ground motion models (GMMs), by testing them in a magnitude and distance range ($M_w = 5.5 \div 7.0$ and Joyner-Boore source-to-site distance $R_{jb} \leq 20$ km) which dominates hazard in the highest seismicity areas of Italy for the return periods of upmost interest for seismic design. To this end, we made use of the very recent release of the NESS2.0 dataset (Sgobba et al., 2021), that collects worldwide near-source strong motion records with detailed metadata. After selection of an ample set of GMMs, based on either their application in past seismic hazard assessment (SHA) studies or for their recent introduction, a quantification of between- and within-event residuals of predictions with respect to records was performed, with the final aim of shedding light on the performance of existing GMMs in the near-source of moderate-to-large earthquakes, in view of a proper selection and weighting of GMMs for SHA.

Declarations

Acknowledgments

This work has been partially supported by swissnuclear within the research activity “Development of advanced numerical approaches for earthquake ground motion prediction”, in the framework of the Sigma2 project, and by the Department of Civil Protection within the ReLUIS project WP18 “Normative contributions related to seismic action”. The authors are sincerely grateful to Sreeram Reddy Kotha for providing fruitful suggestions about the interpretations of results, and to Chiara Smerzini for her useful comments. Researchers of INGV are gratefully acknowledged for making the NESS2.0 dataset available.

Conflict of interest

The authors acknowledge that there are no conflicts of interest.

Availability of data and material

The NESS2.0 dataset is freely available at the website <http://ness.mi.ingv.it/> (Last access: June 6th 2021). The records from ITACA were downloaded from the ITalian ACcelerometric Archive, <http://itaca.mi.ingv.it> (Last access: March 12th 2021). The records from SIMBAD were provided by Chiara Smerzini (Politecnico di Milano).

Code availability

All the computations have been done using MATLAB (<http://www.mathworks.com/products/matlab/>), but the authors don't give their consent to share their routines.

Consent for publication

1. Introduction

There is a general consensus that empirical ground motion models (GMMs) are bound to improve their effectiveness and predictive performance as new records become available, although it is also recognized that, in spite of increasing data and more complex parametrizations, a significant reduction in the aleatory variability cannot be achieved (Douglas and Edwards, 2016).

Among the many GMMs available in the literature, quantitative approaches to provide weights in view of their application to seismic hazard assessment (SHA) have recently been applied, starting from the work by Scherbaum et al. (2009), who devised a log-likelihood scoring scheme, based on the calculation of residuals between observed and predicted values. In such ranking procedure of GMMs, it should be underlined that the sample of records should be representative of the conditions of interest, not only in terms of the prevailing tectonic environment, but also in terms of the magnitude and distance (M - R) ranges that mostly dominate SHA at the site. Making reference for example to the highest seismicity regions of Italy (Central-to-Southern Apennines), the disaggregation (Barani et al, 2009) shows that the mean values of magnitude that dominate the 475 years return period hazard range from about 5.5 to 7 (from short to intermediate oscillation periods), with mean distance less than 20 km.

Owing to the relevance of an accurate evaluation of seismic hazard for such regions, in view of a proper planning of risk mitigation strategies, it is reasonable to check the accuracy of predictions of different GMMs in such M - R range, since, as shown in Figure 1, this is not well sampled yet even by the more up-to-date GMMs.

Although this M - R range may not be strictly considered as near-source, we will check the performance of different GMMs with records included in NESS 2.0 (Sgobba et al., 2021, referred to in the following as NESS): a near-source dataset including worldwide records from active shallow crustal regions, with a careful documentation in terms of seismic source and site conditions metadata. Note that in NESS records are selected only based on M - R considerations and not on their impulsive/non-impulsive features.

With reference to NESS records, this paper aims at checking the performance of a set of GMMs, listed in Table 1, selected for their potential use in the framework of SHA studies in Italy, both in terms of median values and standard deviations.

This paper is organized as follows. First, a selection of GMMs is made and a catalogue of the chosen models is given. A general description of the NESS dataset is then provided, together with the introduction of records considered in this work. Subsequently, the main results concerning the analysis of the performance of selected GMMs against NESS are shown. Residuals, separated into their between- and within- event components, are computed with respect to each model. Finally, the standard deviation of the considered models and the ones determined from records are compared.

2. Selection Of Empirical Earthquake Ground Motion Models

Generation of GMMs, or recalibration of old ones, has nowadays become a continuous process, as new data or new regression procedures become available. In the framework of SHA studies in Italy, we selected several GMMs of interest either for their role in the derivation of the official SH maps of Italy (Ambraseys et al., 1996; Sabetta and Pugliese, 1996), or for their specific calibration with near-source records (Ambraseys and Douglas, 2003), or for their calibration in the Italian and European framework (Akkar and Bommer, 2010; Bindi et al., 2011; Akkar et al., 2014a; Bindi et al., 2014; Lanzano et al., 2019; Kotha et al., 2020), or for their worldwide use (Boore et al., 2014; Campbell and Bozorgnia, 2014; Cauzzi et al., 2014).

Table 1 gives an overview of the 12 selected models, in terms of area of calibration of its dataset, magnitude and distance ranges of applicability, distance metrics and intensity measure type. Note that the most recent models (ITA18 and K20) are based on the European Strong Motion dataset (ESM, Luzi et al. 2020), while two of them, ITA14 and AB10, are derived from the European RESORCE dataset (Akkar et al. 2014b). Two of the selected models are derived from only Italian data (SP96 and ITA10). Anyway, all selected models are suitable to be used in the shallow active crustal regions that dominate seismicity of Italy, with few exceptions of volcanic areas and subduction zones offshore Calabria.

Table 1 GMMs selected for the analyses. For each model, reference details, area of calibration of the dataset, distance metric, distance ranges of applicability, earthquake magnitude and intensity measure type are given.

GMM	acronym	Area of calibration	R metric	R [km]	M	IM type (‡)
Ambraseys et al., 1996	AMB96	Europe	R_{jb}	0 - 200	$M_s = 4 - 7.5$	$H_{\max}$
Sabetta and Pugliese, 1996	SP96	Italy	R_{jb}, R_{epi}	0 - 100	4.6 - 6.8	$H_{\max}$
Ambraseys and Douglas, 2003	AD03	Global	R_{jb}	0 - 15	$M_s = 5.8 - 7.8$	$H_{\max}$
Akkar and Bommer, 2010	AB10	Europe - Middle East	R_{jb}	0 - 99	5 - 7.6	HGM
Bindi et al., 2011	ITA10	Italy	R_{jb}	0 - 200	4.1 - 6.9	HGM
Akkar et al., 2014a	ASB14	Europe - Middle East	$R_{jb}, R_{hypo}, R_{epi}$	0 - 200	4.7 - 7.6	HGM
Campbell and Bozorgnia, 2014	CB14	Global	R_{rup}	0 - 300	3 - 7.9	RotD50
Boore et al., 2014	BSSA14	Global	R_{jb}	0 - 400	3 - 7.9	RotD50
Cauzzi et al., 2014	CF14	Global	R_{rup}	0 - 150	4.5 - 7.9	HGM
Bindi et al., 2014	ITA14	Europe - Middle East	R_{jb}, R_{hypo}	0 - 300	4 - 7.6	HGM
Lanzano et al., 2019	ITA18	Europe	R_{jb}, R_{rup}	0 - 200	3.5 - 8	RotD50
Kotha et al., 2020	K20	Europe	R_{jb}	0 - 545	3 - 7.4	RotD50

(‡) Intensity Measures are defined as follows: HGM = horizontal geometric mean; RotD50 = median value of the IM distribution obtained from rotated waveforms (Boore, 2010); $H_{\max}$ = max horizontal component.

Given that the aim of this work is the evaluation of performance of GMMs in the magnitude and distance ranges of engineering relevance in Italy (i.e., M_w from about 5.5 to 7 and distances up to 20 km), Figure 1 highlights, for some of the selected models, the portion of records falling in that range (shown by the rectangle), with respect to the total.

It is clear that the records included in the selected M - R range are by far a minority of the total, especially for the most recent GMMs, despite the richness of their calibration dataset. Moreover, the fraction of the selected records with respect to the total decreases sharply moving towards recent years, because the number of small magnitude - large distance records increase with time much faster than the number of

near-source records. Subsequent Figure 3 shows the observations used in this analysis, within the same ranges of interest.

As a last remark, we point out that, despite the continuous improvement in regression models and metadata, the aleatory variability of the predictions of the GMMs tends to increase, most probably due to the large variability of the available calibration dataset. To support this sentence, Figure 2 illustrates the total standard deviation (s_{ln}) of the selected models as a function of period, showing an increasing trend with time of s_{ln} .

3. Overview Of Ness Near-source Dataset

NESS is a dataset of high-quality metadata and intensity measures of near-source strong-motion records from active shallow crustal regions, with a careful documentation in terms of seismic source (e.g. geometries and rupture mechanisms) and site conditions. The last version of the dataset was released in January 2021 and is available at the website <http://ness.mi.ingv.it/> (Sgobba et al. 2021). In this dataset, records were selected based on $M-R$ criteria (i.e. earthquakes with $M_w \geq 5.5$, hypocentral depth ≤ 40 km and maximum source-to-site distance depending on the fault dimensions through scaling relationships). Figure 3 shows the distribution of data with respect to M_w , R_{jb} , site conditions and fault mechanism.

In the analyses of this study, only a selection of data from NESS was considered (Figure 3, rectangle). The selection has been done according to the following criteria:

- records on ground type B according to EC8 (stiff soil conditions, CEN 2004);
- records with $R_{jb} \leq 20$ km.

Selected data were classified in four M_w classes, centered at 5.5, 6.0, 6.5 and 7.0 (± 0.1), respectively, as shown in Figure 4. The class with the largest number of records is M_w 6.0, with about 70 records. The same figure shows the distribution of the selected data with respect to focal depth as well. Note that most of the records are characterized by a focal depth of about 10 km.

To increase data for the less populated M_w classes (i.e., 5.5 and 6.5) and to have a statistically significant number of samples, some records from other databases have been used, namely: 12 records from the M_w 5.4 Central Italy earthquake of Oct 26, 2016 included in ITACA 3.0 (D'Amico et al. 2020) and 4 records from the M_w 6.5 Jun 2000 Iceland sequence, included in SIMBAD v.06 (Smerzini et al. 2014).

4. Performance Of Selected Gmms Against Near-source Record

The performance of the selected GMMs was tested for the horizontal components of motions, using two ground motion intensity measures: peak ground acceleration (PGA) and spectral acceleration at 1 s (SA1). Predictions have been computed for a set of 4 magnitude classes, centered at 5.5, 6.0, 6.5 and 7.0 (as shown in Figure 4), using the central value for calculations. The source-to-site distance was selected

according to the specific GMM and the corresponding values provided in NESS. For the style of faulting (SoF), the 'unspecified' mechanism option was selected, in order for all the selected GMMs, included those not providing the SoF, to have a proper basis of comparison. For soil conditions, ground type B (or a representative value of $V_{s,30} = 600$ m/s, for the models having $V_{s,30}$ as site proxy) was used. In the case of SP96, the parametrized soil conditions of which cannot be directly associated to the ground type B, a default multiplication factor of 1.2 has been applied to the prediction on rock. In case of additional parameters, suggested default values were used.

For AMB96 and AD03 models, a conversion from M_s to M_w was applied, in agreement with that considered for the Italian seismic hazard map (MPS Working group, 2004). For SP96, as the magnitude scale adopted provides a good correspondence with M_w (Idriss, 1991), no conversion was applied. Finally, note that for each GMM the comparison with records has been carried out considering the corresponding type of predicted intensity measure (see Table 1), with no need of conversion. Figure 5 shows some examples of comparison, for PGA and SA1, between the median predictions for M_w 6.0 and NESS records (dots) with $M_w = 5.9 \div 6.1$; while Figure 6 shows the same comparison using the median prediction for M_w 7.0 and NESS records with $M_w = 6.9 \div 7.1$. Note that, following K20, we have distinguished the following focal depth (D) classes: 'shallow' ($D < 10$ km), 'intermediate' ($D = 10 \div 20$ km) and 'deep' ($D \geq 20$ km), although, as shown in Figure 4, the NESS records are by far mostly populated in the $8 \div 10$ km bin.

5. Calculation Of Residuals For Different Magnitude And Distance Bins For Site Class B

According to their general definition (Strasser et al., 2008) total residuals are the difference between the natural logarithm of records (y_{es}) and the corresponding prediction (μ_{es}), for a given earthquake e at station s , obtained using a specific GMM. In order to analyze the performance of models as a function of either magnitude or distance, total residuals are separated into their between- and within -event components. Following the notation proposed by several authors (e.g., Al Atik et al., 2010; Rodriguez-Marek et al. 2011), the natural logarithm of the observed ground motion y_{es} can be expressed as in Eq.1, where μ_{es} is the median ground motion value predicted by a specific GMM, δB_e is the between-event residual, which corresponds to the average misfit of records from one particular earthquake with respect to the median ground motion model (Eq.2, where NS is the number of station recordings from one particular earthquake e), δW_{es} is the within-event residual defined as the misfit between an individual observation at station s with respect to the event-corrected median estimate (Eq.3). Between- and within-event residuals are modelled using normal distributions and standard deviation τ and ϕ , respectively.

$$y_{es} = \mu_{es} + \delta B_e + \delta W_{es} \quad (1)$$

$$\delta B_e = \frac{1}{NS} \sum_{s=1}^{NS} (y_{es} - \mu_{es}) \quad (2)$$

$$\delta W_{es} = (y_{es} - (\mu_{es} + \delta B_e)) \quad (3)$$

As an example, Figure 7 shows the computed δB_e and δW_{es} for the AMB96 model, together with the 16th, 50th and 84th percentiles of their normal distribution. Note that, as δB_e can identify a systematic misfit of a specific GMM, with reference to a particular magnitude class, a positive δB_e indicates a systematic underprediction of records, while a negative term indicates a systematic overprediction. Figure 7 shows that AMB96 model has an excellent performance for SA1, but also a limited bias in terms of PGA. This holds true for all magnitude classes, despite a positive bump of residuals in PGAs for M_w 6.5. However, as this tendency of underprediction of PGA for M_w 6.5 is apparent for all models (see Figure 8), it may be likely attributed to a specific trend of records for this magnitude class.

The whole set of between-event residuals calculated for all GMMs is shown in Figure 8 and Figure 9, for PGA and SA1, respectively. Note that the AD03 model was excluded from the calculations for M_w 5.5 class, because its range of applicability is beyond $M_s = 5.8$. Instead, the SP96 and ITA10 models were considered also for M_w 7.0, because their calibration dataset includes records of the M_w 6.9 1980 Irpinia earthquake, used for the validation.

In general, between-event residuals have a larger scatter when calculated for SA1 and show different patterns depending on the GMM under consideration. For instance, PGA residuals calculated for ITA18 suggest a systematic underprediction for almost all magnitude classes, with an overall positive mean δB_e up to about +0.6 (in ln units) for M_w 6.5 class (in PGA), that means an underestimation of recorded values by a factor of about 2.

Although the visual inspection of Figure 8 and Figure 9 is sufficient to provide a meaningful overview of the performance of the different GMMs, a quantitative comparison was set up by scoring each model in terms of either the average or the worst absolute values of δB_e (see Table 2). The best scores are for both AD03 and AMB96, the former ranked the best in terms of average δB_e and second in terms of $\max|\delta B_e|$ and viceversa for the latter model.

6. Performance In Terms Of Standard Deviation

The performance check in the previous section was limited to the median prediction values. We aim in this section at checking if the models are suitable to quantify the observed variability of near-source records.

To this end, we computed the standard deviation of the within-event residuals $\delta W_{es,j}$ of NESS records with respect to the j -th GMM, denoted by $\varphi_{NESS,j}$ as follows:

$$\varphi_{NESS,j} = \sqrt{\sum_{s=1}^{NS} \frac{(\ln(y_{es}) - \ln(a_{GMM,j}) - \delta B_{e,j})^2}{NS-1}} = \sqrt{\sum_{s=1}^{NS} \frac{(\delta W_{es,j})^2}{NS-1}} \quad (4)$$

where y_{es} is the observed value at station s from event e (of either PGA or SA1), $a_{GMM,j}$ is the median predicted acceleration from the j -th GMM, $\delta B_{e,j}$ is the between-event residual computed with reference to the same model, for the same event and NS is the number of recordings available for event e .

The $\varphi_{NESS,j}$ values are compared in Figure 10 with the corresponding φ_j that is the standard deviation of the within-event residuals of the j -th GMM. Note that for the older models (AMB96, SP96 and AD03) φ_j was not computed, so the total standard deviation σ_j was used for comparison.

Considering that j is generally slightly lower than s , the performance of both AMB96 and AD03 models (first row of Figure 10) is good, while for SP96 (R_{epi}) the performance is fairly good for PGA, but not for SA1, especially at higher magnitudes. For M_w 7.0 class, as the spectral acceleration SA1 is very close to the corresponding median prediction, the observed variability (φ_{NESS_SP96}) is lower (see Figure 6). For the more recent GMMs (second and third rows of Figure 10), there is a clear trend of $\varphi_{NESS,j}$ values to be substantially lower than φ_j , suggesting that the variability implied by those GMMs tends to overestimate that observed in the near-source records, as expected since the size of the calibration dataset is higher. This is also visually confirmed in Figure 11, showing $\delta W_{es,j}$ for the class with the highest number of records (M_w 6.0), together with the standard deviation of AMB96, SP96, CB14, ITA10, ITA18 and K20 models.

7. Conclusions

In this paper, we provide a quantitative evaluation of the performance of a set of GMMs in the near-source area of moderate-to-large earthquakes ($M_w = 5.5 \div 7.0$ and Joyner-Boore source-to-site distance $R_{jb} \leq 20$ km), selected because it dominates hazard in the high seismicity Italian regions. GMMs were tested taking advantage of the NESS dataset, consisting of worldwide near-source records with high-quality metadata and intensity measures. Through an analysis of residuals extended to short (PGA) and intermediate periods (SA1) intensity measures, we investigated the misfit of the considered GMMs with respect to records and we compared the performance of the different models in order to check for possible systematic bias.

Our results are somehow unexpected, as it turned out that the two best performing GMMs on near-source records in the $5.5 \div 7 M_w$ range and source-to-site distance $R_{jb} \leq 20$ km, both in terms of median values and standard deviation, are the AMB96, dating about 25 years, and the AD03 models, dating about 20 years. While the excellent performance of AD03 was expected, although not granted, because it was specifically devised for near-source conditions and it took advantage of the important sets of records from the 1999 Turkey and Taiwan earthquakes, that of AMB96 was not obvious, also considering that most of the validation dataset does not contain records used for the calibration.

On the contrary, there was no apparent trend of improved performance with more recent and complex models, in most cases tested against records belonging to their calibration dataset. Furthermore, some of the recent models were affected on one side by the largest bias, especially towards underprediction, and, on the other side, by a tendency towards overestimation of within event variability with respect to the observed one.

These results may likely be related to the bias in the calibration sets of newer models, that are growing continuously and very fast in low amplitude motions, while the number of near-source high amplitude records is increasing at a much lower rate. Therefore, the fraction of near-source records over the total calibration set of new GMMs is fastly decreasing as new data become available. As a consequence, the performance of new GMMs for low-magnitude large-distance earthquake ground motions is expected to steadily improve, but a similar improvement is not granted in the near-source region of moderate-to-large earthquakes, that most often govern the seismic hazard assessment for the highest seismicity regions.

As a final remark, it should be pointed out that the NESS dataset itself is likely not sufficiently extended to cover the variety of fault slip distributions, fault geometry details, complex geological configurations that are expected to affect strong ground motion in near-source conditions. This variety is hardly sampled by the available near-source records, the number of which is not expected to grow significantly in the next few years. In these conditions, the most rationale way to progress towards advanced modelling of near-source ground motions seems to take advantage of the increasingly accurate broadband results of physics-based numerical simulations, as a numerical lab providing realistic realizations of future earthquake ground motions in realistic 3D geological configurations. For this purpose, simulated broadband accelerograms may successfully complement the sparse availability of records in near-source conditions, as in the BB-SPEEDset, recently introduced by Paolucci et al. (2021) based on several validated ground motion scenarios from Italian and worldwide earthquakes, or in the subset of CyberShake ground-motion time series, for the Los Angeles area, selected by Baker et al. (2021).

8. References

1. Akkar S, Bommer JJ (2010) Empirical equations for the prediction of PGA, PGV and spectral accelerations in Europe, the Mediterranean region and the Middle East. *Seismol Res Lett* 81 (2), 195–206 (Mar/Apr) <https://doi.org/10.1785/gssrl.81.2.195>

2. Akkar S, Sandkkaya MA, Bommer JJ (2014a) Empirical ground-motion models for point- and extended-source crustal earthquake scenarios in Europe and the Middle East. *Bull Earthq Eng* 12(1):359–387. Doi 10.1007/s10518-013-9461-4
3. Akkar S, Sandikkaya MA, , Senyurt M, Azari AS, Ay B (2014b) Reference database for seismic ground-motion in Europe (RESORCE). *Bull Earthq Eng*. doi:10.1007/s10518-013-9461-4
4. Al Atik L, Abrahamson NA, Bommer JJ, Scherbaum F, Cotton F, Kuehn N (2010) The variability of ground-motion prediction models and its components. *Seismol Res Lett* 81(5):794–801. <https://doi.org/10.1785/gssrl.81.5.794>
5. Ambraseys NN, Simpson KA, Bommer JJ (1996) Prediction of horizontal response spectra in Europe. *Earthq Eng Struct Dyn* 25(4):371–400
6. Ambraseys NN, Douglas J (2003) Near-field horizontal and vertical earthquake ground motions. *Soil Dynamics and Earthquake Engineering*. Volume 23, Issue 1, 1–18. [https://doi.org/10.1016/S0267-7261\(02\)00153-7](https://doi.org/10.1016/S0267-7261(02)00153-7)
7. Baker JW, Rezaeian S, Goulet CA, Luco N, Teng G (2021) A subset of CyberShake ground-motion time series for response-history analysis. *Earthq Spectra* 37:1162–1176. <https://doi.org/10.1177/8755293020981970>
8. Barani S, Spallarossa D, Bazzurro P (2009) Disaggregation of probabilistic ground-motion hazard in Italy. *Bull Seismol Soc Am* 99(5):2638–2661. Doi:10.1785/0120080348
9. Bindi D, Pacor F, Luzi L, Puglia R, Massa M, Ameri G. and Paolucci R (2011) Ground motion prediction equations derived from the Italian strong motion database. *Bull Seismol Soc Am* 9:1899–1920
10. Bindi D, Massa M, Luzi L, Ameri G, Pacor F, Puglia R, Augliera P (2014) Pan – European ground-motion prediction equations for the average horizontal component of PGA, PGV, and 5% - damped PSA at spectral periods up to 3.0s using the RESORSE dataset. *Bull Earthq Eng* 12(1):391–430. Doi 10.1007/s10518-013-9525-5
11. Boore DM (2010) Orientation – independent, nongeometric-mean measures of seismic intensity from two horizontal components of motion. *Bull Seismol Soc Am* 100:1830–1835. <https://doi.org/10.1785/0120090400>
12. Boore DM, Stewart JP, Seyhan E, Atkinson GM (2014) NGA-West2 equations for predicting PGA, PGV, and 5% damped PSA for shallow crustal earthquakes. *Earthq Spectra* 30:1057–1085. <https://doi.org/10.1193/070113EQS184M>
13. Campbell KW, Bozorgnia Y (2014) NGA-West2 ground motion model for the average horizontal components of PGA, PGV, and 5% damped linear acceleration response spectra. *Earthq Spectra* 30:1087–1115. <https://doi.org/10.1193/062913eqs175m>
14. Cauzzi C, Faccioli E, Vanini M, Bianchini A (2014) Updated predictive equations for broadband (0.01–10s) horizontal response spectra and peak ground motions, based on a global dataset of digital acceleration records. *Bull Earthq Eng* 13(6):1587–1612. Doi:10.1007/s10518-014-9685-y
15. Comité Européen de Normalisation (CEN) (2004) Eurocode 8: design of structures for earthquake resistance-Part 1: general rules, seismic actions and rules for buildings. Comité Européen de

16. D'Amico M, Felicetta C, Russo E, Sgobba S, Lanzano G, Pacor F, Luzi L (2020) Italian Accelerometric Archive v 3.1 – Istituto Nazionale di Geofisica e Vulcanologia, Dipartimento della Protezione Civile Nazionale. doi:10.13127/itaca.3.1
17. Douglas J, Edwards B (2016) Recent and future developments in earthquake ground motion estimation. *Earth Sci Rev* 160:203–219. <https://doi.org/10.1016/j.earscirev.2016.07.005>
18. Idriss IM (1991) Selection of earthquake ground motions at rock sites, Report prepared for the Structures Division, Building and Fire Research Laboratory. Department of Civil Engineering, University of California, Davis
19. Kotha SR, Weatherill G, Bindi D et al (2020) A regionally-adaptable ground-motion model for shallow crustal earthquakes in Europe. *Bull Earthquake Eng* 18:4091–4125. <https://doi.org/10.1007/s10518-020-00869-1>
20. Lanzano G, Luzi L, Pacor F, Felicetta C, Puglia R, Sgobba S, D'Amico M (2019) A revised ground-motion prediction model for shallow crustal earthquakes in Italy. *Bull Seismol Soc Am* 109:525–540. <https://doi.org/10.1785/0120180210>
21. Luzi L, Lanzano G, Felicetta C, D'Amico MC, Russo E, Sgobba S, Pacor F, and ORFEUS Working Group 5 (2020) Engineering Strong Motion Database (ESM) (Version 2.0). Istituto Nazionale di Geofisica e Vulcanologia (INGV). <https://doi.org/10.13127/ESM.2>
22. MPS Working Group (2004) Redazione della Mappa di Pericolosità Sismica Prevista dall'Ordinanza PCM del 20 Marzo 2003 n.3274, All. 1, Rapporto Conclusivo per il Dipartimento della Protezione Civile, Istituto Nazionale di Geofisica e Vulcanologia (INGV), Milano-Roma, Italy, Aprile 2004
23. Paolucci R, Smerzini C, Vanini M (2021) BB-SPEEDset: a dataset of broadband near-source earthquake ground motions from physics-based simulations. *Bull. Seismol. Soc. Am.* Accepted for publication
24. Rodriguez-Marek A, Montalva GA, Cotton F, Bonilla F (2011) Analysis of single-station standard deviation using the KiK – net data. *Bull Seismol Soc Am* 101(3):1242–1258. Doi:10.1785/0120100252
25. Sabetta F, Pugliese A (1996) Estimation of response spectra and simulation of nonstationary earthquake ground motions. *Bull Seism Soc Am* 86(2):337–352
26. Scherbaum F, Delavaud E, Riggelsen C (2009) Model selection in seismic hazard analysis: An information-theoretic perspective. *B Seismol Soc Am* 99:3234–3247. Doi:10.1785/0120080347
27. Sgobba S, Felicetta C, Lanzano G, Ramadan F, D'Amico M, Pacor F (2021) NESS2.0: an updated version of the worldwide dataset for calibrating and adjusting ground motion models in near-source. Istituto Nazionale di Geofisica e Vulcanologia (INGV) <https://doi.org/10.13127/NESS.2.0>
28. Smerzini C, Galasso C, Iervolino I, Paolucci R (2014) Ground Motion Record Selection Based on Broadband Spectral Compatibility. *Earthquake Spectra* 30:1427–1448. <https://doi.org/10.1193/052312EQS197M>

29. Strasser FO, Bommer JJ, Abrahamson NA (2008) Truncation of the distribution of ground-motion residuals. J Seismolog 12(1):79–105. <https://doi.org/10.1007/s10950-007-9073-z>

Supplementary

Due to technical limitations, Table 2 is only available as a download in the Supplemental Files section.

Figures

Figure 1

Magnitude-distance distribution of calibration datasets for some of the selected GMMs, namely: AMB96, BSSA14 and K20 (see Table 1 for references). Distance metric is RJB for all plots. The rectangle highlights magnitude and distance ranges used for the selection of the data in this study.

Figure 2

Total standard deviation $\delta \ln$ of selected GMM (used as a function of R_{jb} , when not specified), for unspecified mechanism. Note that for BSSA14 and CB14, $\delta \ln$ depends on M_w , distance and site conditions as well; adopted values are shown.

Figure 3

Mw and Rjb distribution of NESS data with reference to EC8 site class (left panel) or Style of faulting (right panel, Normal, Reverse and Strike-Slip). The rectangle highlights the Mw-R ranges used for the selection of the data in this study.

Figure 4

Number of records used in the analyses with respect to different Mw classes (left) or focal depths (right). Magnitude classes are centered respectively at 5.5, 6.0, 6.5 and 7.0. Note that only records at short distances from the fault (i.e. $R_{jb} \leq 20\text{ km}$) have been considered.

Figure 5

PGA (top) and SA1 (bottom): median predictions calculated for AMB96 and AD03 (R_{jb} and H_{max}); AB10, ITA10, ITA14, ASB14 (R_{jb} and HGM); CB14 and CF14 (R_{rup} and HGM); K20 'shallow' (K20-S), K20 'intermediate' (K20-I), K20 'deep' (K20-D), ITA18, BSSA14 (R_{jb} and $RotD50$) and SP96 (R_{epi} and H_{max}). Predictions are shown for $M_w = 6.0$ together with the corresponding records from NESS, with $M_w = 5.9 \div 6.1$, distinguished by focal depth (grey dots, $D < 10$ km; white dots, $D = 10 \div 20$ km; black dots, $D \geq 20$ km). The comparison between records and GMMs is shown for the corresponding type of predicted intensity measure (see Table 1). The only exception is in the third column (CB14 and CF14), where records are shown in terms of HGM, while CB14 prediction is in terms of $RotD50$.

Figure 6

Same as Figure 5, with GMM predictions for $M_w = 7.0$ and records from NESS with $M_w = 6.9 \div 7.1$.

Figure 7

PGA (top) and SA1 (bottom) between- (δBe) and within-event residuals (δWes) computed for AMB96 model. For δBe four magnitude classes are shown: M_w 5.5 ($M_w = 5.4 \div 5.6$), M_w 6.0 ($M_w = 5.9 \div 6.1$), M_w 6.5 ($M_w = 6.4 \div 6.6$), M_w 7.0 class ($M_w = 6.9 \div 7.1$). For δWes percentiles are computed with bin of distances and residuals are shown for the four magnitude classes. Black dots and lines show the corresponding 16th, 50th and 84th percentiles of the corresponding normal distribution.

Figure 8

PGA: between-event residuals (δBe) computed for all the selected GMMs: SP96 AD03, ITA10, AB10, ASB14, ITA14, BSSA14, CF14, CB14, ITA18, K20. Four magnitude classes are shown: Mw 5.5 (Mw = 5.4÷5.6), Mw 6.0 (Mw = 5.9÷6.1), Mw 6.5 (Mw = 6.4÷6.6), Mw 7.0 class (Mw = 6.9÷7.1). Black dots and lines show the corresponding 16th, 50th and 84th percentiles of the respective normal distribution.

Figure 9

Same as Figure 8 for SA1.

Figure 10

PGA (left) and SA1 (right): Observed standard deviations ϕ_{NESS_j} for some of the GMMs under study, as a function of the four considered magnitude classes, compared to the standard deviations of each model, shown by thick lines. δ is used for the total standard deviation, ϕ for the within event standard deviation (whenever available).

Figure 11

PGA: Within-event residuals (δWes) for M_w 6.0 class, computed using different GMMs, with 16th, 50th and 84th percentiles (with bin of distances) of the corresponding normal distribution. Solid lines represent the total standard deviation (σ), dashed lines the within-event standard deviation (ϕ), when available, of the corresponding GMM.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table2.docx](#)