

Behaviour-directed Interventions for Problematic Person Transfer Situations in Two Dementia Care Dyads: A Single-case Design Study

Hanna Lagerlund

Rehabilitation Unit, Nyköping Municipality

Charlotta Thunborg

Karolinska Institute

Maria Sandborgh (✉ maria.sandborgh@mdh.se)

Mälardalen University

Research Article

Keywords: person transfer situation, single-case design, functional behaviour analysis, dementia, special care unit, physiotherapy

Posted Date: July 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-701428/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Geriatrics on March 29th, 2022. See the published version at <https://doi.org/10.1186/s12877-022-02952-5>.

Abstract

Background:

Persons with dementia living in nursing homes need assistance with moving and transfers; however, caregivers assisting persons with dementia in their daily person transfers report strain-related and complicated transfer-related behavioural problems. The reciprocity of complex dyadic transfer-related behaviours is affected by environmental factors, the health status of the person with dementia and the caregiver's skills and knowledge. The aim of this study was to explore tailored interventions guided by a functional behaviour analysis for problematic person transfer situations in two dementia care dyads.

Methods:

This study was a quasi-experimental single-case study with an A-B design. Tailored interventions were developed in a five-step model for functional behavioural analysis. The study was conducted in a dementia special care unit at a nursing home, and the inclusion criteria were caregivers' experiences of physical strain and/or resistiveness to care, which led to complex transfer-related behaviour. Two care dyads were included. Transfer situations were video-recorded and evaluated with the Dyadic Interaction in Dementia Transfer Assessment Scale, Pain Assessment in Advanced Dementia Scale, and Resistiveness to Care Scale for Dementia of the Alzheimer's Type. The caregiver experience was evaluated with study-specific items addressing caregiver self-efficacy, catastrophizing thoughts, perceived control, and perceived physical strain. Scorings were graphically displayed. The graphs were inspected visually to identify changes in trend, level, latency, and variability. Nonoverlap of all pairs (NAP), including 90% confidence intervals (CIs), was calculated to complement the visual inspection.

Results:

Verbal and nonverbal discomfort decreased in care dyad 1, which mirrored the caregiver changes in adapting their actions to the needs of the person with dementia. High variability was seen in both the intervention and the baseline phases in care dyad 2. In both care dyads, caregiver transfer-related behaviour improved significantly.

Conclusions:

The results indicate that the transfer-related behaviours of the care dyad might be improved through a behaviour-directed intervention tailored to meet the care dyad's needs. The study is a replication of a previous study in a new clinical setting, which in turn strengthens the generalizability for a functional behavioural analysis-guided intervention for problematic transfer situations in dementia care dyads.

Introduction

Persons with dementia (PWD) living in nursing homes and dementia special care units (SCUs) are often vulnerable individuals with complex and advanced needs (Prince, Prina, & Guerchet, 2013). Dementia

reduces cognitive function and impairs walking and balance, which leads to mobility problems and is therefore a common cause of dependency that affects daily living activities (Waite, Broe, Grayson, & Creasey, 2000; Wallace Williams et al., 2005). People in the late stages of dementia living in SCUs require increasing support over time from caregivers (Carpenter, Hastie, Morris, Fries, & Ankri, 2006), including during person transfer situations.

A transfer situation in an SCU is often constructed of one or two caregivers helping a PwD move, such as from sitting to standing or transferring from a bed to a wheelchair. A transfer situation is often complicated and influenced by environmental, psychological, and physical factors and could be seen as a complex care dyadic behaviour involving both the PwD and the professional caregiver (Thunborg, Söderlund, & von Heideken Wågert, 2019).

A care dyad consists of two people: one caregiver and one care receiver (Lyons, Zarit, Sayer, Aline, & Whitlatch, 2002). Most research addressing dementia care dyads concerns a PwD and an informal caregiver (Orsulic-Jeras, Whitlatch, Powers, & Johnson, 2020; Van't Leven et al., 2013). Caring for a PwD is sometimes challenging, and the professional caregiver has a significant impact on the quality of life and the outcomes when supporting optimal functions in the PwD (Anderson & Blair, 2020). What caregivers do and the way they do it affects the PwD and influences the PwD's behaviour in a caring situation (Berg, Hallberg, & Norberg, 1998; Holst, Edberg, & Hallberg, 1999). One example is the resistiveness to care (Belzil & Vézina, 2015), which can also be referred to as a transfer situation (Thunborg et al., 2019). If a PwD displays resistiveness to care by striving backwards, for example, the caregivers might experience physical strain (Wångblad, Ekblad, Wijk, & Ivanoff, 2009), which in turn can result in uncompleted transfers and increased reciprocal struggling by the care dyad (Thunborg, von Heideken Wågert, Söderlund, & Götell, 2012).

The symbiotic relationship between the members of the care dyad, their interrelated behaviour, the functional status and overall health of the PwD and the caregivers' competence and flexibility alongside the environmental factors (e.g., walking and moving aids, and a stressful working environment) could be interpreted as a dynamic triad within social cognitive theory (Bandura, 1986). There is a knowledge gap concerning the complexity of caring for a PwD in transfer situations, especially with regard to interventions that address the reciprocity of the care dyad, environment, and behaviour. Physiotherapy interventions that consider the complexity of dyadic reciprocity in functional behavioural analyses (FBAs) could provide a better understanding of the function of the care dyad's transfer-related behaviour, which in turn could facilitate more effective tailored interventions for problematic dyadic behaviour (Hanley, 2012), e.g., dyadic transfer-related behaviour, as demonstrated in a single case study (SCS) by Thunborg and colleagues (2019). However, systematic and clinical replications of an SCS are needed to increase the generalizability of the findings (Kazdin, 2017).

The aim of this study was to explore tailored interventions guided by an FBA for problematic person transfer situations in two dementia care dyads.

Methods

Design

In this study, a quasi-experimental single-case design was used, with a baseline phase (A) and an intervention phase (B), and tailored interventions were developed. The key characteristics of an SCS are ongoing assessment, baseline assessment, performance stability and different phases (Kazdin, 2017). The design is clinically relevant and features the opportunity to meet the complexity of person transfer-related behavioural problems in care dyads by tailoring and modifying interventions according to the care dyad's changing needs. This study is a replication of a former study by Thunborg et al. (2019); however, it was implemented in another setting.

Setting and participants

The study was conducted in an SCU with 19 PwDs and 13 caregivers giving constant care and support. Caregivers were either licenced practical nurses or nursing assistants. Two care dyads, i.e., the PwD and the caregivers assisting in the person transfer situation, were included. The first author (HL) conducted the recruitment of the care dyads in collaboration with the caregivers, the manager of the care unit and the responsible nurse. The inclusion criteria for the care dyad were that caregivers had reported experience of physical strain when assisting the PwD and/or that the PwD displayed resistiveness to care, which led to difficulties for the PwD in performing any of the following person transfer situations: lying to sitting, sitting to standing, walking, standing to sitting or sitting to lying. PwD who met the selection criteria but were expected to be at risk of being psychologically or physically adversely affected by participating or who were assessed to be in or near a late palliative phase in life were excluded.

All professional caregivers in the SCU were asked to participate. In total, eight gave their written consent to participate. All who consented were included in the study, and they were video recorded at least two times. As all caregivers worked shifts on different schemas, maintaining the same caregiver who was supporting the PwD throughout all video-recorded transfer situations was not possible. A care dyad is therefore defined as a PwD and the caregivers who assist in a given person transfer situation. For participant characteristics, see Table 1.

Table 1
Care dyad characteristics.

Care dyad	Gender	Age	Diagnosis	Time in Special Care Unit
1	woman	83	Alzheimer's disease	2 years
2	woman	78	Alzheimer's disease	2 years
Care dyad	Gender	Age	Professional role	Years working as a caregiver
1	woman (N = 4)	37–58 years	Licensed practical nurses (N = 4)	2–29
2	woman (N = 4)	20–51 years	Licensed practical nurses (N = 3); nursing assistant (N = 1)	2–25

Data collection

Data were collected between April and July 2019. One problematic person transfer situation for each care dyad was identified through semistructured interviews with the caregivers concerning the character of the problematic transfer situation, affecting factors and the functional ability of the PwD. In total, 23 person transfer situations were video-recorded and analysed (see Table 2). The time required for each person transfer situation was measured with a digital timer by observing and calculating the time spent from start to finish.

Dyadic interaction in dementia transfer assessment scale

To assess problematic person transfer situations, the Dyadic Interaction in Dementia Transfer Assessment Scale (DIDTAS) (Thunborg, von Heideken Wågert, Götell, Ivarsson, & Söderlund, 2015) was used. The DIDTAS is an observational assessment scale developed to evaluate problematic person transfer situations when a caregiver assists a PwD in a person transfer situation. The scale contains 17 items: items 1–8 assess the actions of the PwD, and items 9–17 assess the actions of the caregiver. A higher score indicates a more difficult person transfer situation. The intraclass correlation (ICC), which represents the reliability for each of the 17 items, ranges from 0.34 to 0.92 for interrater reliability and from 0.56 to 0.92 for intrarater reliability (Thunborg, Heideken Wågert, Ivarsson, & Söderlund, 2015). See additional file 1 DIDTAS.

Pain assessment in advanced dementia scale

To assess pain during the person transfer situation for the PwD, the Pain Assessment in Advanced Dementia Scale (PAINAID) (Warden, Hurley, & Volicer, 2003) was used. The PAINAID was developed to assess pain in older cognitively impaired individuals who cannot self-report their pain experience. The PAINAID includes five items related to five behaviours. The items include breathing, negative sounds (e.g., crying and moaning), facial expressions, body language and consolability, which are rated from 0 to 2 and summarized based on a total score of 0 to 10. A possible interpretation of the scores is as follows:

1–3 = mild pain; 4–6 = moderate pain; and 7–10 = severe pain. PAINAID shows good conceptual validity and satisfactory interrater validity (Warden et al., 2003).

Resistiveness to care scale-dementia alzheimer type

Resistiveness to care was evaluated with the Resistiveness to Care Scale-Dementia Alzheimer Type (RTC-DAT) (Mahoney et al., 1999), which consists of 13 items concerning the behaviours: turn away, pull away, push away, push/pull, grab object, grab person, adduct, hit/kick, say no, cry, threaten, scream/yell and clench mouth. Each behaviour is observed and rated based on prevalence and severity. RTC-DAT gives a score between 0 and 156, with a lower score representing a small degree of resistiveness to care.

Regarding reliability, internal consistency estimates from 0.82–0.87 and good to excellent kappa values have been demonstrated (Mahoney et al., 1999).

Caregiver’s self-reported ratings

Before each observation, the caregiver completed self-reported ratings regarding their self-efficacy, catastrophizing thoughts and perceived control for the transfer situation in question. Four items addressing self-efficacy were developed in the context of social cognitive theory (Bandura, 1997). Three items were adapted from the Coping Strategies Questionnaire (CSQ) and addressed one statement measuring catastrophizing thoughts and two questions measuring perceived control (Rosenstiel & Keefe, 1983). The item perceived physical strain was adapted from the Patient Transfer Assessment Instrument (PTAI) and measured after the observation with the following question: “How do you experience physical strain during the person transfer situation (0 = not physically strenuous at all, 10 = physically very strenuous)” (Karhula, Rönholm, & Sjögren, 2009). For further details, see additional file 2.

Table 2
Number of video-recorded observations and length of phases

	Number of video-recorded observations		Length of phases in days	
	Baseline (A)	Intervention(B)	Baseline (A)	Intervention((B)
Care dyad 1	6	6	18	29
Care dyad 2	5	6	8	36

Procedural steps in the FBA model

We used a five step FBA-model (see Fig. 1). During the first step, cognitive, functional and/or environmental factors associated with the occurrence (and nonoccurrence) of specific problematic transfer-related behaviour were identified for each care dyad. Sources for information on the person transfer situation were caregiver reports, medical records and next-of-kin information. The first author (HL) observed the person transfer situation without video recording and performed a general physiotherapy assessment (e.g., range of motion, any signs of pain, anxiety and communication

difficulties). During the baseline phase (step 2), the caregivers were instructed to administer care as usual when assisting the PwD in the transfer situation. The transfer situation was video-recorded six (care dyad 1) and five (care dyad 2) times by the first author. The video-recorded environment-related factors, antecedents for the target behaviour, the behaviour itself and the consequences of the behaviour were analysed and rated by all authors. After gathering the baseline data, the DDTAS items of interest were chosen alongside the ratings from the RTC-DAT, PAINAID and caregivers' self-reports, and a problematic transfer-related target behaviour was identified. In step 3, a hypothesis based upon the assessment in steps 1 and 2 describing the target behaviour in sufficient detail was developed. In step 4, the hypothesis was tested by implementing one tailored intervention in each care dyad. In step 5, the intervention effectiveness was monitored, i.e., the caregivers provided rating on the self-report assessment scales, and the person transfers were video-recorded. The caregivers were instructed to administer care according to the tailored intervention when assisting the PwD in the transfer situation. The transfer situation was video-recorded six times in step 5 in each care dyad.

Dyad characteristics and baseline data

Care dyad 1

Step 1. Functional assessment to identify the problem behaviour

The PwD in care dyad 1 was an 83-year-old woman with Alzheimer's disease. She had suffered from neck disabilities of unknown origin most of her adult life and had fibromyalgia since the age of 50. In recent years, she had unspecified back pain and general tremor. She had paratonia in both the upper and lower limbs, and an assessment with the Paratonia Assessment Instrument (PAI) (Hobbelen, Koopmans, Verhey, Habraken, & de Bie, 2008) indicated moderate paratonia (value = 2) on a scale from 0–5, which resulted in involuntary resistance in passive movements. She rarely took the initiative for speech but was able to answer simple spoken questions and was judged by the caregivers to be able to understand the meaning of instructions and conversations to some extent. She was unable to perform a cognitive test; therefore, her cognitive ability was uncertain. The caregivers supposed she had adequate hearing and vision. She used a mobile lift in all daily transfers to the wheelchair. Two caregivers on each side of the PwD's bed took part in the morning care routine when the lifting sling was applied. The caregivers reported that her entire body became tense and she cried out and was perceived to be struggling against the caregivers when they turned her from side-to-side with the help of sliding and draw sheets to apply the lifting sling. The caregivers experienced the transfer situation as difficult and physically heavy, and they also inferred that she did not feel well during the transfer situation. The transfer situation proceeded with other similar transfer situations when changing incontinence protection.

Step 2. Collecting baseline data from multiple sources

During baseline (A), the problem behaviour was best described by the high ratings of DDTAS items 6 and 7, indicating that the PwD expressed verbal and non-verbal discomfort. The high caregiver ratings of

DIDTAS items 9 and 10 indicated a lack of caregiver instructions before the beginning of the transfer and a lack of a clear verbal commands about the transfer. The high ratings of DIDTAS items 11 and 14 indicated that the caregivers did not wait for the PwD to respond during the transfer and that the caregivers did not adapt their actions to facilitate the PwD in the person transfer situation. See appendix DIDTAS. Information about the caregiver's experiences of physical strain was provided through self-reports. The PAINAID (score = 0) and RTC-DAT (score = 0) values indicated that the PwD had no pain and did not display any resistiveness to care in the person transfer situation.

Step 3. Developing the hypothesis

The hypothesis suggested that when the PwD experienced a change in position for which she was unprepared, her unease and paratonia increased, which was expressed through grabbing objects and shouting, and that motor protective reflexes were triggered in combination, with increased muscle tension over the whole body. The caregivers experienced the transfer situation as physically heavy and expressed difficulties in consoling the PwD. Eye contact, maintaining physical contact and clear verbal instructions seemed to decrease anxiety and paratonia. Rapidness during the transfer situation seemed to increase unease and paratonia. The behavioural goals were therefore to decrease the verbal and non-verbal discomfort expressed by the PwD (DIDTAS items 6 and 7) and to increase the adaptation to the PwD during patient transfer by the caregiver (DIDTAS items 9, 10, 11 and 14).

Care dyad 2

Step 1. Functional assessment and problem behaviour identification

The PwD in care dyad 2 was a 78-year-old woman with Alzheimer's disease and no other known diseases. She had severe difficulties sitting down on the toilet, especially in the mornings. When the caregivers assisted physically, she could react with resistiveness and strive in the opposite direction. The nursing staff found that their verbal instructions did not reach the women. The person transfers worked best when she took the initiative herself to step out of bed, but even on these occasions, she could experience problems sitting down on the toilet. The caregivers noticed that when two caregivers assisted the person transfer, the interaction with her was impaired. Therefore, only one caregiver assisted. She had impaired vision and used glasses, although she seldom wore them when she got up from bed and walked to the toilet during the morning routine. The caregivers supposed she had adequate hearing. She communicated with few words and seldom provided adequate responses to direct questions. The caregivers thought she was able to understand the meaning of instructions and conversations to some extent, although the linguistic understanding was difficult to assess. The physiotherapy assessment showed that she had sufficient muscle strength and balance to sit down on the toilet.

Step 2. Collecting baseline data from multiple sources

During baseline (A), the problem behaviour was best described by the high ratings of DIDTAS item 1, indicating that the PwD was not able to remain attentive during the person transfer situation. The high

ratings of DIDTAS item 14 indicated that the caregivers did not adapt their actions to facilitate the PwD in the person transfer situation. Important information for the FBA was also obtained from the PwD's RTC-DAT scores, indicating that the PwD showed resistiveness to care by grabbing the toilet handle and walking away from the caregiver in the transfer situation. A PAINAID score of 0 indicated that the PwD had no pain during the person transfer situation.

Step 3. Developing the hypothesis

The hypothesis was that when the PwD was assisted physically and asked verbally to sit down on the toilet, she had severe difficulties interpreting the instructions and understanding the goal with the person transfer. Verbal instructions could lead to compliance, no action at all or walking away from the toilet. When the caregiver assisted physically, she often reacted with resistiveness and grabbed the toilet handles and did not release them. The caregivers adjusted their instructions, e.g., combined them with visual and auditive cues or used a diverting step, e.g., making the bed and then going back to the toilet. The PwD seemed to follow the instructions better if the caregiver was standing on her right side instead of in front or on her left side. Since the PwD often forgot to use her glasses, she might experience a reduced ability to interpret visual cues. The behavioural goals were therefore to increase the PwD's attentiveness (DIDTAS item 1) and to increase the caregiver's adaption to the PwD in the transfer situation (DIDTAS item 14).

Data management and analysis

The scores for the outcome measures were displayed graphically, and medians for each outcome and phase were calculated. The graphed data points, each representing a person transfer observation, were drawn and connected with trend lines over time and within a given phase. The graphs were inspected visually to identify changes in trend, level, latency and variability (Kazdin, 2017). Nonoverlap of all pairs (NAP), including 90% confidence intervals (CIs), was calculated to complement the visual inspection. NAP is a nonparametric technique for measuring "the percent of nonoverlapping data between baseline and treatment phases" (Parker I. & Vannest, 2009, p 359). The NAP value is equal to the empirical area under the curve (AUC) of a receiver operating characteristic test (Parker I. & Vannest, 2009). To fit the analysis with the NAP, the item scoring for DIDTAS was reversed when calculating the NAP.

Statistical analysis with the Mann-Whitney U-test was used to analyse the time required from start to finish for each person transfer situation. The significance level was set at 0.05. The Paleontological Statistics Software Package for Education and Data Analysis (Hammer, Ø., Harper, D.A.T., Ryan, 2001) was used to perform statistical analyses.

Ethical considerations

The Swedish Ethical Review Authority (dnr 2018/2169- 31) reviewed and approved the study. The rights of the participants were carefully considered and protected in all aspects. The PwD was asked for consent in close connection to each video taken of the person transfer situation. If the PwD was unable to give verbal permission, the person transfer situation was carefully observed by the first author (HL) to

detect any expression of dislike and the caregivers were asked if they detected any inconvenience of the PwD. The video recording was terminated if the PwD demonstrated dislike or aversion. The first author (HL) contacted the next of kin via telephone, provided oral information about the study and the opportunity to ask questions and sent the written study information with the consent form by mail with a prepaid envelope. The next of kin could not make the decision on behalf of the PwD but rather could only state whether they did not oppose the PwD's participation. Thus, written consent was collected by asking "Do you oppose the participation of your next of kin?" The caregivers were provided with oral and written information of the study and asked to provide written and informed consent to participate.

Results

Care dyad 1

Step 4. Intervention planning and hypothesis testing.

Based on the analysis from steps 1–3 of the FBA, a tailored intervention was developed that focused on the nursing staff's problematic transfer-related behaviour, DITAS items 9, 10, 11 and 14. The intervention was a combination of the following components: (1) the caregivers distributed the responsibility during person transfers so that one person maintained contact and interacted with the PwD and the other person pulled the sliding sheet and took care of the practicalities during the transfer; (2) the caregiver maintained eye contact and held the hands of the PwD during the turn, clearly informed what was going to happen and waited for a time between providing the information and performing the person transfer; (3) the turning was divided into several parts, namely, from back-lying to side-lying and from side-lying to back-lying; and (4) the PwD was prepared for movement by being positioned in a natural movement pattern for turning.

Step 5. Monitoring intervention effectiveness and modification

Visual inspection of DITAS items 6 and 7 showed a positive change in the trend. Verbal and nonverbal discomfort decreased in the intervention phase compared to the baseline [DITAS item 6: NAP = 100% (90% CI = 0.429- 1), $p = 0.0039$ and DITAS item 7: NAP = 79% (90 % CI = 0.013- 1)]. The significant improvement in DITAS 6 showed low variability and short latency. The level between the baseline phase and intervention phase changed abruptly in DITAS 7 and DITAS 6. The change in DITAS item 7 was not significant, and high variability at baseline was observed, although stabilization could be seen in the intervention phase. Both DITAS item 6 and DITAS item 7 showed a median change (see Fig. 2).

The caregivers' self-reported ratings showed high self-efficacy, low catastrophizing thoughts and high perceived control for the transfer situation of interest. No change was observed for the caregivers' self-reported ratings between the baseline and intervention phases, for further details, see additional file 2. The variability and median (median baseline = 2.5 and median intervention phase = 1.5) of the caregivers'

self-reported perceived physical strain decreased in the intervention phase, indicating that the caregiver experienced lower physical strain after the implementation of the intervention (see Fig. 3).

The caregivers' performance improved significantly in the intervention phase [DIDTAS 9, 10, 11 and 14: NAP = 100% (90% CI = 0.429- 1), $p = 0.0039$]. Caregiver trends changed positively as shown in DIDTAS items 9, 10, 11 and 14, indicating that the caregiver instructions and adaption of actions to facilitate the PwD improved. The variability for DIDTAS items 9, 10, 11 and 14 was low in the baseline phase and slightly higher in the intervention phase for DIDTAS 9, 11 and 14. The level changed rapidly between the baseline and intervention phases, and the latency was short after implementing the intervention for DIDTAS 9, 10, 11 and 14. The levels for the outcomes, as reflected by DIDTAS items 9, 10, 11 and 14, showed a median change (see Fig. 2).

The time required for the person transfer situation significantly increased (Mann-Whitney U-test, $p = 0.008$) between the baseline and intervention phases. The median increased from 57 seconds at baseline to 137 seconds in the intervention phase.

Care dyad 2

Step 4. Intervention planning and hypothesis testing

According to the hypothesis, both the caregivers and the PwD displayed problematic transfer behaviour, as demonstrated by DIDTAS items 1 and 14, and the intervention to be prioritized was related to environmental factors. The intervention consisted of a combination of the following elements: (1) the PwD was helped to put on glasses before the person transfer began; (2) the left handle on the toilet was folded up before the PwD sat down, and then the left handle was folded down, and if necessary, the caregivers could lower the right handle during the person transfer; and (3) the caregivers provided support from the right side of the PwD.

Step 5. Monitoring intervention effectiveness and modification

In the intervention phase, the PwD was not capable of remaining attentive during the transfer situation, as reflected by an increasing trend in the intervention phase and no significant change in DIDTAS item 1 [DIDTAS item 1: NAP = 73% (90% CI = -0.134- 1)]. DIDTAS item 1 showed no distinct latency or shift in level, and the variability in both the baseline and intervention phases was high. DIDTAS item 1 showed a slight positive change in the median (see Fig. 4).

The PwD was less resistive to care in the intervention phase than the baseline phase, although the change was not significant [RTC-DAT: NAP = 80% (90% CI= -0.001- 1)]. The trend showed increases in both the baseline and intervention phases but at a lower level in the intervention phase. The RTC-DAT values showed a positive change in the median but did not show a distinct latency or shift in level. The variability was high in the baseline phase and intervention phase (see Fig. 4).

High self-efficacy, low catastrophizing thoughts, low perceived physical strain, and high perceived control in the transfer situation were reported by the caregivers. There was no change in self-reported ratings between the baseline and intervention phases, for further details, see additional file 2.

The caregivers significantly improved their performance in the intervention phase [DIDTAS 14: NAP = 88% (90% CI = 0.166- 1], $p = 0.0358$]. The level between the baseline and intervention phases changed abruptly, and the latency was short. There was a change in the median between the baseline and intervention phases, and high variability was observed in both the baseline and the intervention phases (see Fig. 4).

The time required for the person transfer situation between the baseline and intervention phases did not change significantly. The median decreased from 171 seconds (2 minutes and 51 seconds) at baseline to 113 seconds (1 minute and 53 seconds) in the intervention phase.

Discussion

This study explored tailored interventions guided by an FBA for problematic person transfer situations in two dementia care dyads at an SCU. From a social cognitive theoretical perspective, the reciprocity between the individual, environmental and behavioural factors in the transfer situation was clearly illustrated in care dyad 1 but not as evident in care dyad 2.

In care dyad 1, nonverbal discomfort decreased (DIDTAS item 6) and a positive trend in both verbal and nonverbal discomfort (DIDTAS items 6 and 7) was observed. Caregivers' transfer-related behaviour mirrored these changes by providing a clear verbal command (DIDTAS item 11) just before beginning the transfer (DIDTAS item 9), waiting for the PwD to respond (DIDTAS item 11) and adapting to the situation, such as keeping eye contact and holding the hands of the PwD during the transfer (DIDTAS item 14).

The high variability in DIDTAS item 7 in terms of addressing verbal discomfort in the baseline phase was stabilized in the intervention phase. It is well known that PwD behaviour can vary greatly (Wångblad et al., 2009), and stabilization, e.g., transfer-related behaviour, could imply an important improvement from a clinical perspective. Lack of time has been reported by caregivers as a barrier when providing palliative care for people with severe dementia (Midtbust, Alnes Einang, Gjengedal, & Lykkeslet, 2018); however, in care dyad 1, the time for completing the person transfer increased by only approximately one minute. Simultaneously, the caregivers perceived less physical strain during the person transfer. The PwD in care dyad 1 suffered from paratonia, which is a motor behaviour problem prevalent in 90–100% of all PwDs with severe dementia and increases the caregiver burden with time (Souren, Franssen, & Reisberg, 1997). Symptoms of paratonia in person transfers are crucial to address, and the intervention in care dyad 1, including caregivers' verbal and nonverbal communication before initiating the transfer situation and waiting for the PwD to react, shows promising results for PwDs with paratonia.

In care dyad 2, high variability in PwD behaviour was seen in both the intervention and baseline phases. The transfer-related behaviour in the PwD was sustained despite the significant change in the caregivers' behaviour, as reflected by their adaption of actions (DIDTAS item 14) and the inability of the PwD to

remain attentive in the transfer situation (DIDTAS item 1). Possible reasons for the nonsignificant results could be that the tailored intervention did not meet the needs of the PwD. The PwD tended to be less resistive to care in the intervention phase than the baseline phase, although the change was not significant. Reduced resistive behaviour was noted after a behavioural intervention in a previous study (Thunborg et al., 2019), which may indicate that these behaviours can influence behavioural interventions.

In both care dyads 1 and 2, caregiver behaviour improved significantly but still varied in the intervention phase. The variability may be related to the varying level of adherence to the intervention by the individual caregivers, which might reflect the difficulties for caregivers working in SCU to change their accustomed behaviour in their daily work. Caregiver training should focus on more intense and in-depth training to increase the understanding of behavioural change across the development of dementia (Appleton & Pereira, 2019). Genuine professional knowledge concerning the fundamental needs in persons with dementia is recognized as crucial in caregivers for dyadic interactions in dementia care (Jakobsen & Sørli, 2010; Manthorpe et al., 2010), which means that health care professionals need to recognize the perspectives of both parts since the PwD needs to rely on the other's contributions in the interaction (Lichtner et al., 2016).

There are both strengths and limitations in this study. One limitation is that an AB design was used. Unfortunately, the health condition of the PwD made it impossible to add additional phases because the internal validity could have been strengthened by an ABAB design (Kazdin, 2017; Petursdottir & Carr, 2018); however, the design was not feasible, mainly for two reasons. First, it would not have been ethical to end a caregiver behaviour after the intervention phase if it was beneficial for the PwD. Second, it would have been difficult for the caregiver to return to their former behaviour, although one possible change would have been to apply a follow-up phase (Kazdin, 2017). In both care dyads, the intervention implemented was a combination of several components, and it might have been better to divide the intervention into several phases, such as an A-B1-B2-B3 sequence. Considering the need for interventions in both care dyads, such a division would have delayed the possibility for an improved transfer situation for the care dyad.

In care dyad 2, the scarce number of observations and high variability make it difficult to draw causal inferences about the impact of the intervention. More observations during a longer period might have better clarified the needs of the PwD and established a stable baseline. The FBA indicated that environmental factors affected the PwD behaviour in care dyad 2, but these factors were also influenced by personal factors, such as motor and social interaction behaviour (Bandura, 1986). The care dyad's interaction is therefore complex, and the high variability complicated the analyses. It can take time for one factor to influence the others in the triad (Bandura, 1986), and the intervention phase for care dyad 2 may be too short for this to happen. In care dyad 1, the variability in DIDTAS item 7 was high and a stable baseline was not achieved. An extended baseline and additional repeated observations would have been preferable to rule out the possibility that history or maturation could have influenced the change in the dependent variable (Petursdottir & Carr, 2018; Shadish, Cook, & Campbell, 2002). However, the rapid

change of the PwD behaviour in care dyad 1 in the intervention phase together with the short latency in caregivers' behavioural change indicates the impact of the intervention. Thunborg et al. (2019) also demonstrated high variability and a short baseline in a care dyad, thus reflecting the challenges of conducting an intervention in this population. PwDs in the late phase of their dementia disease are frail, and their health condition can rapidly change. Such conditions increase the difficulty of following individuals over time, and in longitudinal dementia studies, dropouts due to death or other reasons commonly reduce the follow-up sample size (Anderson & Blair, 2020). For this reason, we tried to capture as many observations as possible during a short period of time. Unfortunately, only a portion of all the caregivers in the nursing home consented to participate in the study, and due to shift work, it was not possible to conduct and observe the care dyads every day in a row. On the other hand, this study reflects the reality of dementia care and the true conditions for the care dyad.

The results of the visual inspections were strengthened using NAP as a nonparametric statistical complement. Regarding the participants, especially the caregivers, we cannot rule out the possibility that testing could have influenced their actions during the patient transfer situations because they knew they were being filmed (Petursdottir & Carr, 2018; Shadish et al., 2002).

This study is a systematic replication of a former SCS by Thunborg et al. (2019). The present study was conducted in a new setting with regard to the nursing homes, caregivers and municipalities. Additionally, the constellation of the research team for this study was new. Although Thunborg's participation in both studies could be considered a weakness, it aided in the precise operational definitions of DITAS.

Kazdin (2011) emphasized that the selected participants in an SCS should be typical clinical cases. A strength of this study is that the two single cases illustrate two different person transfer situations commonly present in an SCU for PwD. Furthermore, the study was conducted in the participants' natural environment. Every transfer situation was assessed with DITAS, PAINAID and RTC-DAT. The combination of these instruments meant that a behaviour could be evaluated from different aspects. For example, shouting can be interpreted as pain, a way to show resistance to care or an expression of discomfort with words/sounds. By carefully observing each behaviour with all three assessment scales, it was possible to form an idea of what function the behaviour had, which is an essential part in developing an FBA hypothesis (Hanley, 2012).

Conclusions

We could not demonstrate intrasubject replication in this study, but similar to the study by Thunborg et al. (2019), our study demonstrated significant improvements for one care dyad, which presented less discomfort expressed by the person with dementia and simultaneously improved the adaptive behaviour by the caregivers.

The results indicated that the transfer-related behaviours of the care dyad might be improved through a behaviour-directed intervention tailored to meet the care dyad's needs. Further research is needed to

understand how behaviour interventions can be best constructed to meet the complexity of care dyads in problematic person transfer situations.

List Of Abbreviations

Person with Dementia (PwD)

Special Care Unit (SCU)

Single Case Study (SCS)

Functional Behavioural Analysis (FBA)

Dyadic Interaction in Dementia Transfer Assessment Scale (DIDTAS)

Pain Assessment in Advanced Dementia Scale (PAINAID)

Resistiveness to Care Scale- Dementia Alzheimer Type (RTC-DAT)

Confidence Interval (CI)

Nonoverlap of All Pairs (NAP)

Declarations

Ethics approval and consent to participate

The Swedish Ethical Review Authority (dnr 2018/2169–31) reviewed and approved the study. The rights of the participants were carefully considered and protected in all aspects. The PwDs were asked for consent in close connection to each video uptake of the person transfer situation. If a PwD was unable to give verbal permission, the person transfer situation was carefully observed by the first author (HL) to detect any expression of dislike and the caregivers were asked if they detected any inconvenience of the PwD. The video recording was terminated if the PwD demonstrated dislike or aversion. The first author (HL) contacted the next of kin via telephone, provided oral information about the study with the opportunity to ask questions and sent the written study information with the consent form by mail with a prepaid envelope. The next of kin could not make the decision on behalf of the PwD but rather could only state whether they did not oppose the PwD's participation. Thus, written consent was collected by asking "Do you oppose the participation of your next of kin?" The caregivers were provided with oral and written information of the study and asked to provide written and informed consent to participate.

Consent for publication

Written informed consent for publication was obtained from the caregivers and the next of kin in the same procedure as “Consent to participate” (see above).

Availability of data and materials

The datasets generated and analysed during the current study are not publicly available due to ethical restrictions but are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The authors declare that they have no funders.

Authors' contributions

HL, CT and MS all contributed to the study design. HL conducted the data collection. All authors (HL, CT and MS) observed, analysed, rated and interpreted the video-recorded person transfer situations. All authors jointly developed interventions for the care dyads. HL prepared all the figures and tables and drafted the manuscript text. All authors contributed to data interpretation, critical review and amendments of the manuscript, and they also read and approved the final manuscript.

Acknowledgements

Not applicable

Authors' information (optional)

Not applicable

References

1. Anderson, K., & Blair, A. (2020). Why we need to care about the care: A longitudinal study linking the quality of residential dementia care to the residents' quality of life. *Archives of Gerontology and Geriatrics*, 91, 104226. <https://doi.org/10.1016/j.archger.2020.104226>

2. Appleton, K., & Pereira, A. (2019). Behavioural Changes in Dementia and their Impact on Professional Caregivers: A Grounded Theory Approach. *Dementia, 18*(4), 1479–1491.
<https://doi.org/10.1177/1471301217714654>
3. Bandura, A. (1986). *Social foundations of thought and action: a social cognitive theory*. Englewood Cliffs, N.J.: Prentice-Hall.
4. Bandura, A. (1997). *Self-efficacy: the exercise of control*. New York: Basingstoke: W. H. Freeman.
5. Belzil, G., & Vézina, J. (2015). Impact of caregivers' behaviors on resistiveness to care and collaboration in persons with dementia in the context of hygienic care: an interactional perspective. *International Psychogeriatrics, 27*(11), 1861–1873. <https://doi.org/10.1017/S104161021500099X>
6. Berg, A., Hallberg, I. R., & Norberg, A. (1998). Nurses' reflections about dementia care, the patients, the care and themselves in their daily caregiving. *International Journal of Nursing Studies, 35*(5), 271–282. [https://doi.org/10.1016/s0020-7489\(98\)00040-6](https://doi.org/10.1016/s0020-7489(98)00040-6)
7. Carpenter, G. I., Hastie, C. L., Morris, J. N., Fries, B. E., & Ankri, J. (2006). Measuring change in activities of daily living in nursing home residents with moderate to severe cognitive impairment. *BMC Geriatrics, 6*(1), 7. <https://doi.org/10.1186/1471-2318-6-7>
8. Hammer, Ø., Harper, D.A.T., Ryan, P. D. (2001). PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*, pp. 4(1)1-9.
9. Hanley, G. P. (2012). Functional Assessment of Problem Behavior: Dispelling Myths, Overcoming Implementation Obstacles, and Developing New Lore. *Behavior Analysis in Practice, 5*(1), 54–72. <https://doi.org/10.1007/BF03391818>
10. Hobbelen, J. S. M., Koopmans, R. T. C. M., Verhey, F. R. J., Habraken, K. M., & de Bie, R. A. (2008). Diagnosing paratonia in the demented elderly: reliability and validity of the Paratonia Assessment Instrument (PAI). *International Psychogeriatrics, 20*(4), 840–852.
<https://doi.org/10.1017/S1041610207006424>
11. Holst, G., Edberg, A. K., & Hallberg, I. R. (1999). Nurses' narrations and reflections about caring for patients with severe dementia as revealed in systematic clinical supervision sessions. *Journal of Aging Studies, 13*(1), 89–107. [https://doi.org/https://doi.org/10.1016/S0890-4065\(99\)80008-5](https://doi.org/https://doi.org/10.1016/S0890-4065(99)80008-5)
12. Jakobsen, R., & Sørli, V. (2010). Dignity of older people in a nursing home: Narratives of care providers. *Nursing Ethics, 17*(3), 289–300. <https://doi.org/10.1177/0969733009355375>
13. Karhula, K., Rönholm, T., & Sjögren, T. (2009). *A method for evaluating the load of patient transfers* (1st ed.). Tampere: Occupational Safety and Health Administration.
14. Kazdin, A. E. (2017). *Research design in clinical psychology* (5th ed.). Boston: Pearson.
15. Lichtner, V., Dowding, D., Allcock, N., Keady, J., Sampson, E. L., Briggs, M., ... Closs, S. J. (2016). The assessment and management of pain in patients with dementia in hospital settings: A multi-case exploratory study from a decision making perspective. *BMC Health Services Research, 16*(1).
<https://doi.org/10.1186/s12913-016-1690-1>
16. Lyons, K. S., Zarit, S. H., Sayer, Aline, G., & Whitlatch, C. J. (2002). Caregiving as a dyadic process: Perspectives from caregiver and receiver. *The Journals of Gerontology, Series B*(3), P195–P204.

<https://doi.org/10.1093/geronb/57.3.p195>

17. Mahoney, E. K., Hurley, A. C., Volicer, L., Bell, M., Gianotis, P., Hartshorn, M., ... Warden, V. (1999). Development and testing of the Resistiveness to Care Scale. *Research in Nursing & Health, 22*(1), 27–38.
18. Manthorpe, J., Iliffe, S., Samsi, K., Cole, L., Goodman, C., Drennan, V., & Warner, J. (2010). Dementia, dignity and quality of life: Nursing practice and its dilemmas. *International Journal of Older People Nursing, 5*(3), 235–244. <https://doi.org/10.1111/j.1748-3743.2010.00231.x>
19. Midtbust, M. H., Alnes Einang, R., Gjengedal, E., & Lykkeslet, E. (2018). Percieved barriers and facilitators in providing palliative care for people with severe dementia: the healthcare professional's experience. *BMC Health Services Research, 18*:709. <https://doi.org/https://doi.org/a0.1186/s12913-018-3515-x>
20. Orsulic-Jeras, S., Whitlatch, C. J., Powers, S. M., & Johnson, J. (2020). A dyadic perspective on assessment in Alzheimer's dementia: Supporting both care partners across the disease continuum. *Alzheimer's and Dementia: Translational Research and Clinical Interventions, 6*(1). <https://doi.org/10.1002/trc2.12037>
21. Parker I., R., & Vannest, K. (2009). An improved Effect Size for Single-Case Research: Nonoverlap of All Pairs. *Behavior Therapy, 40*, 357–367.
22. Petursdottir, A. I., & Carr, J. E. (2018). Applying the Taxonomy of Validity Threats from Mainstream Research Design to Single-Case Experiments in Applied Behavior Analysis. *Behavior Analysis in Practice, 11*(3), 228–240. <https://doi.org/10.1007/s40617-018-00294-6>
23. Prince, M., Prina, M., & Guerchet, M. (2013). *World Alzheimer Report. Journey of Caring. An Analysis of Long-Term Care for Dementia.*
24. Rosenstiel, A., & Keefe, F. (1983). The use of coping strategies in chronic low back pain patients: Relationship to patient characteristics and current adjustment. *Pain, 17*, 33–44.
25. Shadish, W. R., Cook, T. D., & Campbell, D. T. (2002). *Experimental and Quasi-Experimental Designs for Generalized Causal Inference.* Boston: Houghton Mifflin.
26. Souren, L. E. M., Franssen, E. H., & Reisberg, B. (1997). Neuromotor Changes in Alzheimer's Disease: Implications for Patient Care. *Journal of Geriatric Psychiatry and Neurology, 10*(3), 93–98. <https://doi.org/10.1177/089198879701000301>
27. Thunborg, C., Heideken Wågert, P. von, Ivarsson, A. B., & Söderlund, A. (2015). Inter- and Intra-Rater Reliability of a Newly Developed Assessment Scale: The Dyadic Interaction in Dementia Transfer Assessment Scale (DIDTAS). *Physical & Occupational Therapy in Geriatrics, 33*(4), 279–293. <https://doi.org/10.3109/02703181.2015.1065939>
28. Thunborg, C., Söderlund, A., & von Heideken Wågert, P. (2019). Functional behaviour analysis guided interventions might improve transfer-related behaviour in dementia care dyads: a single case study. *European Journal of Physiotherapy.* <https://doi.org/10.1080/21679169.2018.1549591>
29. Thunborg, C., von Heideken Wågert, P., Götell, E., Ivarsson, A.-B., & Söderlund, A. (2015). Development of a new assessment scale for measuring interaction during staff-assisted transfer of residents in

- dementia special care units. *BMC Geriatrics*, 15(1), 6. <https://doi.org/10.1186/s12877-015-0003-6>
30. Thunborg, C., von Heideken Wågert, P., Söderlund, A., & Götell, E. (2012). Reciprocal struggling in person transfer tasks: Caregivers' experiences in dementia care. *Advances in Physiotherapy VO - 14*, (4), 175. <https://doi.org/10.3109/14038196.2012.725184>
31. Van't Leven, N., Prick, A. E. J. C., Groenewoud, J. G., Roelofs, P. D. D. M., De Lange, J., & Pot, A. M. (2013, October). Dyadic interventions for community-dwelling people with dementia and their family caregivers: A systematic review. *International Psychogeriatrics*. *Int Psychogeriatr*. <https://doi.org/10.1017/S1041610213000860>
32. Waite, L. M., Broe, G. A., Grayson, D. A., & Creasey, H. (2000). Motor function and disability in the dementias. *International Journal of Geriatric Psychiatry*, 15(10), 897–903. [https://doi.org/10.1002/1099-1166\(200010\)15:10<897::AID-GPS215>3.0.CO;2-C](https://doi.org/10.1002/1099-1166(200010)15:10<897::AID-GPS215>3.0.CO;2-C)
33. Wallace Williams, S. W., Williams, C. S., Zimmerman, S., Sloane, P. D., Preisser, J. S., Boustani, M., & Reed, P. S. (2005). Characteristics Associated With Mobility Limitation in Long-Term Care Residents With Dementia. *The Gerontologist*, 45(Supplement 1), 62–67. https://doi.org/10.1093/geront/45.suppl_1.62
34. Wångblad, C., Ekblad, M., Wijk, H., & Ivanoff, S. D. (2009). Experiences of physical strain during person transfer situations in dementia care units. *Scandinavian Journal of Caring Sciences*, 23(4), 644–650. <https://doi.org/10.1111/j.1471-6712.2008.00655.x>
35. Warden, V., Hurley, A. C., & Volicer, L. (2003). Development and Psychometric Evaluation of the Pain Assessment in Advanced Dementia (PAINAD) Scale. *Journal of the American Medical Directors Association*, 4(1), 9–15. <https://doi.org/10.1097/01.JAM.0000043422.31640.F7>

Figures




Figure 1

Five-step model for the functional behavioural analyses (FBAs), adapted from Thunborg et al. 2019, p. 3.




Figure 2

Data points, medians and phases for care dyad 1: DIDTAS item 6, PwD expresses discomfort through body language in the transfer situation; item 7, PwD expresses discomfort through words/sounds in the transfer situation; item 9, caregiver provides instructions for transfer just before beginning transfer; item 10, caregiver provides a clear verbal command about transfer; item 11, transfer request is followed by the caregiver waiting for the PwD to respond; and item 14, caregiver adapts their actions to facilitate the transfer situation of the PwD. A higher score indicated a more difficult person transfer situation.




Figure 3

Caregivers' self-reported ratings of perceived physical strain measured after the observation (where 0 = not physically strenuous at all and 10 = physically very strenuous). Each measurement point on which the box plot is based corresponds to the mean value of each self-reported rating from the two caregivers who participated in the transfer situation at the time of observation. Median phase A = 2.5; median phase B = 1.5.




Figure 4

Data points, medians and phases for care dyad 2. DIDTAS item 1, the PwD is able to remain attentive during the transfer in the transfer situation; and DIDTAS item 14, caregiver adapts their actions to facilitate the transfer situation of the PwD. A higher score indicates a more difficult person transfer situation. For PwD resistiveness to care (RTC-DAT), a higher score indicates more resistiveness to care.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.pdf](#)
- [Additionalfile2.pdf](#)