Cytokine Dysregulation Persists in Childhood Post Neonatal Encephalopathy

Zunera Zareen

University of Dublin Trinity College <https://orcid.org/0000-0003-1531-2034>

Tammy Strickland

University of Dublin Trinity College

Victoria Mc Eneaney

University of Dublin Trinity College

Lynne A. Kelly

University of Dublin Trinity College

Denise McDonald

Tallaght University Hospital

Deirdre Sweetman

National Maternity Hospital

Eleanor J. Molloy (✉ eleanor.molloy@tcd.ie)

<https://orcid.org/0000-0001-6798-2158>

Research article

Keywords: Cytokines; Neonatal encephalopathy; Inflammation; Neurodevelopmental; Hypoxic-ischaemic encephalopathy.

Posted Date: October 23rd, 2019

DOI: <https://doi.org/10.21203/rs.2.16375/v1>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Neurology on March 30th, 2020. See the published version at <https://doi.org/10.1186/s12883-020-01656-w>.

Abstract

Objective: Cytokines are possible mediators of neuroinflammation and associated with adverse outcome in neonatal encephalopathy (NE). Our aim was to explore cytokine response in children with Neonatal Encephalopathy (NE) at school age compared to age-matched controls. **Method:** Follow up at school age, children who had NE and age-matched controls were assessed for their cytokine responses and neurodevelopment outcome. Pro- and anti-inflammatory cytokines in the serum, [Interleukin (IL)-1 α , IL-1 β , IL-2, IL-6, IL-8, IL-18, Tumor necrosis factor (TNF)- α , TNF β , Interferon (IFN)- γ , granulocyte-macrophage colony-stimulating factor (GM-CSF), vascular endothelial growth factor (VEGF), erythropoietin (EPO), IL-10 & IL-1RA] were measured at baseline and in response to in vitro stimulation with lipopolysaccharide (LPS: endotoxin). **Results:** GM-CSF, TNF- β , IL-2 IL-6, IL-8, IL-10, and IL-18 were significantly elevated at school age following NE (n=40) compared to controls (n=37). A significant rise in GM-CSF, IL-8, TNF- α , IL-1 β , & IL-6 were seen in NE group following LPS stimulation. Relative LPS hypo-responsiveness was also noted in children with severe NE with IL-10, VEGF, EPO and TNF- β . Elevated TNF- β was associated with low gross motor scores on assessment at school age. **Conclusion:** School-age children post-NE had significantly altered cytokine responses to endotoxin compared to controls. TNF- β was associated with adverse developmental outcomes. This suggests the inflammatory process may persist into childhood and a longer therapeutic window may be available for immunomodulatory adjunctive therapies.

Background

Neonatal brain injury such as Neonatal encephalopathy (NE) is an important cause of neonatal death and disability such as cerebral palsy. Inflammation combined with Hypoxia-ischemia (HI) play an important pathophysiological role in NE. The injury processes can persist for months and years and a tertiary mechanism of damage has been proposed, which includes inflammation and epigenetic changes, (1) decreased plasticity and reduced number of neurons. Infants with NE have a persistent inflammatory response over the first week of life correlating with the degree of brain injury (2, 3).

Pro-inflammatory cytokine expression within the brain, especially of Interleukin (IL)-1 β and Tumor Necrosis factor (TNF)- α have been demonstrated following perinatal brain damage by pathogen triggers and hypoxic injury both in experimental models and the human neonates (4). Neuroinflammation and IL-6, enhances the growth of neural precursors after neonatal brain injury (5). Pro-inflammatory cytokines activate cytotoxic T cells and natural killer cells, which enhance cellular and tissue damage. This leads to cell proliferation, differentiation and cell death causing white matter changes and long-term neurological damage. (6) (7)

Alternations in cytokines in NE during the neonatal period have been well described. (8) Cytokines including IL-6 and IL-16 in cord blood were associated with severity of neonatal encephalopathy defined by continuous electroencephalography.(9) Elevated Interleukin (IL-6), noted in the first few hours after birth, in infants with NE who received TH was associated with death and abnormal neurodevelopmental outcome at 12 months of age.(10) In addition, levels of IL-6, IL-8, and vascular endothelial growth factor

(VEGF) were also noted to be greater at 6–24 hours in infants with moderate to severe NE compared to mild NE and were associated with abnormal neurological outcomes.(1, 3, 11)

We have previously demonstrated altered VEGF and Epo in this cohort of babies with NE in the first week of life and therefore followed them in childhood to assess immune function. We hypothesised that the inflammatory response persists in childhood and aimed to examine the cytokine profile of school aged children with Neonatal Encephalopathy in comparison to healthy age-matched controls. Understanding this persistent inflammatory mechanism could lead to safe and effective therapies to treat a developmentally disrupted brain long after the initial insult.

Methods

Ethical committee approval was received from the Ethics Committee of the National Maternity Hospital, Dublin and the National Children Hospital, Tallaght, Dublin. Written informed consent was obtained from all parents of children enrolled in the study.

Patient Groups

This study is a follow up of a previous cohort of infants with Neonatal encephalopathy recruited in the neonatal period and performed follow up at school age and the recruitment of these babies as neonates has been previously described (2). The following groups were enrolled:

(1) Control group: serum samples from healthy children age 4-7 years who attended the day ward and were undergoing phlebotomy as part of a day case procedure. These children were born following a full-term normal delivery with normal Apgar scores, neurological examination and postnatal course and had no underlying co-morbidities.

(2) NE group: School age children who were exposed to perinatal asphyxia (PA) according to the criteria of Huang et al (12), were included. Children with NE were classified using the modified Sarnat & Sarnat (13) : (a) NE I: infants with mild encephalopathy, or (b) NE II/III: moderate/severe encephalopathy(2).

Fifteen of 24 children in the NE II/ III group received therapeutic hypothermia (TH) in accordance with the TOBY (Total Body Hypothermia for Neonatal Encephalopathy) criteria (14) and were treated for 72 hours duration. Infants with congenital abnormalities, confirmed sepsis or evidence of maternal substance abuse were not enrolled. (2)

Development Questionnaire

The Ages and Stages Questionnaire was used as a screening tool to assess different domains of development including gross motor (GM), fine motor (FM), problem solving (PS), communication and

personal social skills in children in the control group and children with NE. The age-appropriate development Ages and Stages Questionnaire (ASQ-3) was completed by the parents of school age children in both the NE group and the control group enrolled in the study. The age of the children was in the range of 4-6 years. All domains of development including gross motor, fine motor, problem solving, communication and personal social skills were examined. The parents completed the questionnaire at the clinic visit and any parental concerns were addressed. It took around 10-15 mins for the parent to complete the questionnaire. A detailed history was taken, and neurological examination was carried out at the same visit.

Blood sampling

Peripheral venous samples (1 mL) were collected into vacutainers containing sodium citrate. The sample was split into microtubes and treated with Vehicle (1µl PBS) or Lipopolysaccharide (LPS) (10ng/ml) at 37°C for 1hr. Following incubation, the samples were centrifuged (1500 rpm at 4°C x 10 minutes) and the supernatant removed, aliquoted and stored at -80°C for further analysis.(2)

Multiplex cytokine analysis

Interleukin-1 alpha (IL-1 α), interleukin-1 beta (IL-1 β), interleukin-2 (IL-2), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), interleukin-18 (IL-18), interleukin-1 receptor antagonist (IL-1ra), tumour necrosis factor-alpha (TNF- α), tumour necrosis factor-beta (TNF- β), interferon gamma (IFN- γ), vascular endothelial growth factor (VEGF), erythropoietin (Epo) and granulocyte macrophage/colony-stimulating factor (GM-CSF) were analysed using a multi-spot 96-well, 14 spot custom human serum plasma plate which was customised for our study and selected cytokines by Meso Scale Discovery. The plate was then analysed on the SECTOR Imager and validated (Meso Scale Discovery, Rockville, MD, USA; www.mesoscale.com).2

Statistical Analysis

Statistical analysis was carried out using the PASW statistical package version 24 (www.ibm.com/SPSS_Statistics). Continuous normally distributed data was displayed as means and standard deviations (SDs) and comparisons were made using the independent student-t test. Significance was achieved for values $p < 0.05$. We used ANOVA-one-way analysis of variance for analysis of the cytokine results. Two-way analysis of variance was carried out to compare baseline and stimulated sample results. Spearman correlation was used to assess the association between the cytokines levels and neurodevelopment outcome using ASQ-3 at 5-6 years.

Results

Clinical Characteristics

The study enrolled 77 school age children (age range 4-7 years), including controls (n=40) and children following NE (n= 37). The Sarnat and Sarnat grades (13) were as follows: number of infants with perinatal asphyxia but with no neurological signs (designated as grade 0 for study purposes, n=2); mild NE (grade I, n =11); moderate NE (grade II, n=22); and severe NE (grade III, n=2). Fifteen children received therapeutic hypothermia (TH) in accordance with the TOBY (Total Body Hypothermia for Neonatal Encephalopathy) criteria (14) and were treated for 72 hours duration. Seventeen infants developed clinical seizures in the neonatal period and 13 had an abnormal MRI brain. There were no significant differences between the NE groups 0/I and II/III regarding gestational age, gender or birth weight (Table 1).

Neurodevelopmental Outcome

Twenty-four children with moderate to severe NE (NE II/III) had the following outcomes on clinical review at school age as designated by their clinical teams: Cerebral Palsy with Gross Motor Function Classification System (GMFCS), level III-V (n=5); epilepsy (n=1); speech and language difficulty (n=5); hearing deficit (n=2); functional motor and co-ordination problems (n=3).

The children in the NE group scored poorly in all domains compared to children in the control group, especially in gross & fine motor skills, communication and problem-solving (p value <0.04). The scores for children with mild NE (NE 0/I) though in the normal range, were significantly lower when compared to children in the control group especially for fine motor and problem-solving skills. (NE 0/I v Control; FM: 58 vs 48, p value=0.03; PS: 58 vs 55, p value =0.04, respectively). Similarly, on comparing the NE groups by severity, the NE II/III group scores were significantly low compared to NE 0/I group (NE II/III v NE 0/I: FM: 39 vs 48, p=0.025; PS: 48 vs 55, p value= 0.036). We demonstrated that elevated TNF- β was associated with low gross motor scores on developmental assessment at 5-6 years of age using the ASQ questionnaire (Rho 0.39, p value 0.02).

Cytokines in children with NE versus Controls and Hypo responsiveness to LPS

GM-CSF, TNF- β , IL-2 IL-6, IL-8, IL-18, and TNF- α were significantly higher at school age (p value <0.05) in children with NE compared to children in the control group. In addition, on LPS stimulation a significant rise of cytokines GM-CSF, and IL-8 were seen in children with NE compared to controls (p value <0.05). IL-6, IL-1 β and TNF- α were significantly increased in both groups of children with NE and controls (p value <0.01) on LPS stimulation. (Figure 1&2)

There was no increase in the pro inflammatory cytokines, IL-1 α , IL-1 β , IFN- γ and TNF- α , at baseline in the control group but these rose significantly when stimulated with LPS in vitro. There was no statistically

significant increase in IL-10, IL-1Ra and VEGF in children with NE even on stimulation with LPS. LPS hypo-responsiveness was noted in Epo and TNF- β in children with NE. At baseline levels of both EPO and TNF- β rose slightly but on stimulation with LPS the levels were significantly reduced in children with NE compared to controls. (Figure 1&2).

Children who received Therapeutic Hypothermia (TH) had high levels of GM-CSF and IL-10 compared the control group ($p < 0.05$). However, on stimulation with LPS the levels of GM-CSF and IL-10 were noted to be reduced in the group of children who received TH compared to those without TH. Similarly, decreased response to LPS in cytokines IL-6, IL-8, IL-1 α , IL-1 β , TNF- α and VEGF was demonstrated in children in the TH group compared to children who did not receive TH. (Figure 3 & 4).

Discussion

We found a persistently abnormal cytokine response in children at school age following NE. GM-CSF, TNF- β , IL-2 IL-6, IL-8, IL-10, IL-18, and Epo, were significantly higher in children with NE compared to controls at baseline. GM-CSF and IL-8 rose significantly in children with NE in response to LPS stimulation compared to age-matched controls. Children at school-age following NE also had relative LPS hypo responsiveness in several cytokines demonstrating an altered immune phenotype compared with age-matched controls.

IL-8 was significantly elevated in school children with NE when compared with controls, and after in vitro treatment with an endotoxin (LPS) levels increased further. IL-8 was noted to be high in infants with NE when compared to normal infants and higher in those with severe encephalopathy and abnormal neuroimaging, compared to those with mild NE (3). IL-8 measured before 96 hours of age, to be a potent biomarker predicting death and abnormal neurological outcome in survivors of NE.(8) Elevated IL-8 levels in the group of children with NE, in our study compared to normal children demonstrates a persistently altered immune response at school age.

GM-CSF was also significantly elevated in children with NE compared to the controls. Interestingly increased response was also noted in the neonatal period in the same cohort of children with NE. GM-CSF activates neutrophils and macrophages and stimulates the production of other pro-inflammatory cytokines. GM-CSF production is stimulated by lipopolysaccharide (LPS), tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), and IL-23. GM-CSF has been reported to high in infants with NE who were eligible for TH and in non-survivors at day 1 of life compared to survivors (3). We also noted a similar response at school age in children with NE who underwent TH in the neonatal period. However, GM-CSF crosses the blood brain barrier and may be neuroprotective and reduced brain damage in previous experimental models of stroke.(15)

IL-6 was noted be significantly high on LPS stimulation in children with NE II/III who received TH and had abnormal neurodevelopmental outcome. A study demonstrated that high levels of serum IL-6 at 24-48 hours in a group of infants who received TH, were associated with abnormal MRI brain (16). A meta-analysis identified serum IL-6 and serum IL-1 β measured before 96 hours of age as potential biomarkers

in predicting abnormal outcome in infants with NE (8). Elevated IL-6 in newborn with NE who received TH was associated with death and abnormal neurodevelopmental outcome at 12 months of age (17).

Tumor Necrosis factor (TNF- β) is another pro-inflammatory cytokine which acts on different cells. It promotes the proliferation of fibroblasts and induces the synthesis of GM-CSF, G-CSF, IL-1, and prostaglandin-E2 in fibroblasts (18). We demonstrated high levels of TNF- β in response to LPS stimulation in school age children with NE. We found an association of high TNF- β levels in school age children with NE with low gross motor scores on developmental assessment. Previous studies have demonstrated an association between neonatal serum cytokine levels of IL-8, IL-12, TNF α and TNF- β in extremely low birth weight infants and Cerebral Palsy (19). But there are no studies to our knowledge, showing an association of poor neurodevelopmental outcome and abnormally elevated cytokine levels at school age in children with NE.

Interleukin-10 (IL-10) is an anti-inflammatory cytokine, due to its ability to inhibit macrophage activation. It also has a stimulatory function on B cells and mast cells. Increased IL-10 levels were noted in children with NE compared to controls. High levels of IL-10 were associated with severity of NE and in those who received therapeutic hypothermia (TH). IL-10 also has neuroprotective properties demonstrated in animal studies (20). Correlation of high levels of IL-10 with multiorgan dysfunction & mortality has been demonstrated in children in paediatric intensive care (21). Neonates with sepsis, pneumonia and necrotising enterocolitis demonstrated high levels of IL-10 (22). Increased levels of IL-10 at 72 hrs of age were associated with prolonged NICU stay in neonates with Transposition of great arteries who underwent the arterial switch procedure (23).

Erythropoietin (Epo) is a glycoprotein cytokine secreted by the kidney in response to cellular hypoxia. It stimulates the production of red blood cells from the bone marrow. Apart from being an antioxidant, it also has anti-apoptotic and anti-inflammatory actions. Elevated Epo levels were demonstrated in children with NE compared to controls, especially in response to LPS. Significantly high levels of Epo have been demonstrated in infants with increased severity of NE and who underwent TH at day 2-4 of life. NE grade II/III and death were associated with elevated Epo levels on Day 3 of life (2, 16). High levels of serum Epo may represent markers of severity of encephalopathy and brain injury, this theory can be supported by previous studies reported in preterm infants.(24)

TNF- α is a proinflammatory cytokine which plays a critical role in the local inflammatory response and in initiation of the cytokine cascade (25). TNF- α is an important cytokine involved in sepsis and inflammation and is a potential marker in the diagnosis of early and late onset neonatal sepsis(25, 26). Increased serum level of TNF- α has been reported to be associated with mortality and abnormal neuroimaging in neonates with NE (16). In our study we demonstrated significantly increased TNF- α in response to LPS in school age children in the NE and the control groups.

We have shown in our study that school age children with NE demonstrated vigorous systemic innate immune response compared to children with normal development. Periventricular leukomalacia (PVL) or related inflammation events or both during the perinatal and postnatal period may have a programming

effect, causing altered inflammatory responses in preterm children with CP (27). In their study they demonstrated that preterm school-age children with periventricular leukomalacia (PVL) induced (CP) had significantly higher levels of tumour necrosis factor (TNF α) and elevated TLR4 mRNA in peripheral blood mononuclear cells (PBMC's), in comparison to preterm term school-age control group children.

Lipopolysaccharide (LPS) when introduced in vitro in healthy individuals not only induces the cascade of inflammatory pathways (28) but also initiates a transient refractory state, referred to as LPS hypo-responsiveness or LPS tolerance (29). This LPS tolerance state is associated with decreased capacity of whole blood and peripheral mononuclear cells to produce proinflammatory cytokines in response to LPS stimulation. Decreased levels of proinflammatory cytokines IL-6 and TNF- α in the serum and in the peripheral mononuclear cells (PBMC) supernatant are found in response to LPS stimulation in critically ill patients in ICU in comparison to healthy individuals (30). Similar results were seen in our study, the level of TNF- α rose with LPS stimulation to a lesser degree than in children with NE compared to age-matched controls. In addition, LPS hypo responsiveness was found in children with NE in cytokines Epo and TNF- β . Similarly, children in the TH group demonstrated decreased response to LPS in cytokines IL-6, IL-8, IL-1 α , IL-1 β , TNF- α and VEGF, which correlated with severity of NE.

Conclusions

Neonatal Encephalopathy remains an important cause of mortality and long-term neurodevelopmental impairment including Cerebral Palsy. Hypoxic ischaemic injury induces an inflammatory response involving excessive cytokine production. We demonstrated significant alteration in cytokines in children with NE at school age compared to normal children which supports our hypothesis of a persistent inflammatory response years after hypoxic ischaemic injury. We also showed an association between elevated proinflammatory cytokine TNF- β and poor neurodevelopmental outcome using ASQ-3 questionnaire. Thus demonstrating that proinflammatory cytokines may act as important biomarkers in prediction of neurodevelopmental outcome in children with NE. Persistent inflammatory response seen in children with NE at school age indicates a longer “therapeutic window” may be available for immunomodulatory adjunctive therapies following therapeutic hypothermia.

Abbreviations

Cerebral Palsy (CP), Neonatal Encephalopathy (NE), lipopolysaccharide (LPS), Interleukin (IL), Tumor necrosis factor (TNF), Interferon (IFN), granulocyte-macrophage colony-stimulating factor (GM-CSF), vascular endothelial growth factor (VEGF), erythropoietin (EPO),

Declarations

Ethics Approval & Consent to Participate

Ethical committee approval was received from the Ethics Committee of the National Maternity Hospital, Dublin and the National Children Hospital, Tallaght, Dublin. Written informed consent was obtained from all parents of children enrolled in the study.

Consent for Publication: Not applicable

Availability of Data and Material:

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request

Conflicts of Interest: The authors declare no conflict of interest. The sponsor of the study has no role in the study design, collection, analysis and interpretation of data, writing the paper and in the decision to submit the manuscript for publication.

Funding: Research Funding Committee, National Children’s Hospital Foundation, Tallaght University Hospital, Dublin, Ireland.

Author Contribution: Author involved in the study concept and design; Z.Z, E.M, D.M and D.S. Authors involved in data acquisition and analysis were Z.Z, L.K, V. Mc E, and T.S. Authors involved in drafting texts and figures were Z.Z, T.S and L.K.

Acknowledgement

The study was funded by the National Children Hospital Foundation, Tallaght University Hospital, Dublin, Ireland. There is no commercial party involved. We thank all our patient, parents and nursing staff for their effort and time in this study.

Tables

Table 1: Demographics of infants with NE

Variables		NE 0/1 (n=13)	NE- II/III (n=24)	p- value (95% CI)
GA (wks)a		39.7 (1.43)	40.5 (1.21)	0.06 (-1.4-0.03)
BW (kgs)a		3.48 (0.7)	3.61 (0.61)	0.31 (-4.7-0.13)
Gender, male, n (%) b		9 (76%)	14 (58%)	0.70
Mode of delivery n (%) b	LSCS	7 (53)	11 (45)	0.70
	SVD	3 (23)	7 (29)	0.73
	Inst	3 (23)	6 (25)	0.73
Apgar @1 min c		5(3-6)	2(1-5)	0.003
Apgar@5 mins c		7(5-8)	4(2-7)	0.01
Apgar@10 mins c		7.5 (6-9)	5(3-7)	0.01
TH, n (%) b		0 (0)	15 (62)	<0.001
Seizures, n (%) b		0 (0)	17 (70)	<0.001
MRI- Abnormal, n (%) b		0/0	13 (50)	<0.001
ASQ -GM		56 (4)	48 (9)	0.03

Table 1: Demographics of infants with NE: GA=Gestational age; BW = birth weight; SVD = spontaneous vaginal delivery; inst. = instrumental delivery; TH = therapeutic hypothermia; ASQ= Ages and Stages Questionnaire GM= Gross motor skill score. a) For normally distributed data, mean $\pm$ SD is expressed, and the independent Student-t test was used for comparison, where $p < 0.05$ is significant with 95% confidence intervals (CI). b) For binary variables, the χ^2 test was used for comparison. c) For skewed data, medians and IQRs are expressed and the Mann-Whitney U test was used for comparison;

Table 2 Cytokines levels in control group, NE0/I and NE II/III

Cytokines	Control (n=40) (Mean)	NE (0/I) (Mean)	NE (II/III) (Mean)	p value (a)	P value (b)
IL2	0.54	1.18	1.35	0.02	0.04
IL6	1.27	2.03	2.31	0.05	0.73
IL8	16.77	37.99	77.67	0.11	0.39
TNF α	3.63	4.74	4.69	0.17	0.95
GMCSF	0.24	2.01	0.41	0.29	0.34
IFN γ	15.98	15.33	11.44	0.92	0.29
EPO	90.45	98.07	93.22	0.62	0.63
IL10	0.75	1.68	1.74	0.21	0.95
IL 18	532.57	628.01	683.14	0.16	0.56
IL -IRA	681.91	767.10	649.44	0.68	0.61
IL-1 α	1.60	0.85	2.01	0.12	0.11
IL-1 β	0.62	1.52	1.32	0.20	0.81
TNF β	1.29	3.17	2.63	0.01	0.42
VEGF	106.65	122.57	148.57	0.50	0.68

Table 2: Means and p value using Student t-test. IL= Interleukin, TNF= Tumor necrosis factor, IFN= Interferon, VEGF= Vascular endothelial growth factor, EPO= Erythropoietin, GMCSF= Granulocyte colony stimulating factor. p (a)-p value Control vs NE 0/I and p (b) p value NE0/I vs NEII/III.

References

1. Hagberg H, Gressens P, Mallard C. Inflammation during fetal and neonatal life: implications for neurologic and neuropsychiatric disease in children and adults. *Ann Neurol*. 2012;71(4):444-57.

2. Sweetman DU, Onwuneme C, Watson WR, Murphy JF, Molloy EJ. Perinatal Asphyxia and Erythropoietin and VEGF: Serial Serum and Cerebrospinal Fluid Responses. *Neonatology*. 2017;111(3):253-9.
3. O'Hare FM, Watson RW, O'Neill A, Blanco A, Donoghue V, Molloy EJ. Persistent systemic monocyte and neutrophil activation in neonatal encephalopathy. *J Matern Fetal Neonatal Med*. 2016;29(4):582-9.
4. Dammann O, Leviton A. Inflammatory brain damage in preterm newborns—dry numbers, wet lab, and causal inferences. *Early Hum Dev*. 2004;79(1):1-15.
5. Covey MV, Loporchio D, Buono KD, Levison SW. Opposite effect of inflammation on subventricular zone versus hippocampal precursors in brain injury. *Ann Neurol*. 2011;70(4):616-26.
6. Khwaja O, Volpe JJ. Pathogenesis of cerebral white matter injury of prematurity. *Arch Dis Child Fetal Neonatal Ed*. 2008;93(2):F153-61.
7. Okazaki K, Nishida A, Kato M, Kozawa K, Uga N, Kimura H. Elevation of cytokine concentrations in asphyxiated neonates. *Biol Neonate*. 2006;89(3):183-9.
8. Ramaswamy V, Horton J, Vandermeer B, Buscemi N, Miller S, Yager J. Systematic review of biomarkers of brain injury in term neonatal encephalopathy. *Pediatr Neurol*. 2009;40(3):215-26.
9. Walsh BH, Boylan GB, Livingstone V, Kenny LC, Dempsey EM, Murray DM. Cord blood proteins and multichannel-electroencephalography in hypoxic-ischemic encephalopathy. *Pediatr Crit Care Med*. 2013;14(6):621-30.
10. Jenkins DD, Rollins LG, Perkel JK, Wagner CL, Katikaneni LP, Bass WT, et al. Serum cytokines in a clinical trial of hypothermia for neonatal hypoxic-ischemic encephalopathy. *J Cereb Blood Flow Metab*. 2012;32(10):1888-96.
11. Chalak LF, Sanchez PJ, Adams-Huet B, Laptook AR, Heyne RJ, Rosenfeld CR. Biomarkers for severity of neonatal hypoxic-ischemic encephalopathy and outcomes in newborns receiving hypothermia therapy. *J Pediatr*. 2014;164(3):468-74 e1.
12. Huang CC, Wang ST, Chang YC, Lin KP, Wu PL. Measurement of the urinary lactate:creatinine ratio for the early identification of newborn infants at risk for hypoxic-ischemic encephalopathy. *N Engl J Med*. 1999;341(5):328-35.
13. Sarnat HB, Sarnat MS. Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study. *Arch Neurol*. 1976;33(10):696-705.
14. Azzopardi D, Brocklehurst P, Edwards D, Halliday H, Levene M, Thoresen M, et al. The TOBY Study. Whole body hypothermia for the treatment of perinatal asphyxial encephalopathy: a randomised controlled trial. *BMC Pediatr*. 2008;8:17.
15. Schabitz WR, Kruger C, Pitzer C, Weber D, Laage R, Gassler N, et al. A neuroprotective function for the hematopoietic protein granulocyte-macrophage colony stimulating factor (GM-CSF). *J Cereb Blood Flow Metab*. 2008;28(1):29-43.
16. O'Hare FM, Watson RW, O'Neill A, Segurado R, Sweetman D, Downey P, et al. Serial cytokine alterations and abnormal neuroimaging in newborn infants with encephalopathy. *Acta Paediatr*.

2017;106(4):561-7.

17. Jenkins DD, Lee T, Chiuzan C, Perkel JK, Rollins LG, Wagner CL, et al. Altered circulating leukocytes and their chemokines in a clinical trial of therapeutic hypothermia for neonatal hypoxic ischemic encephalopathy*. *Pediatr Crit Care Med*. 2013;14(8):786-95.
18. Tracey D, Klareskog L, Sasso EH, Salfeld JG, Tak PP. Tumor necrosis factor antagonist mechanisms of action: a comprehensive review. *Pharmacol Ther*. 2008;117(2):244-79.
19. Varner MW, Marshall NE, Rouse DJ, Jablonski KA, Leveno KJ, Reddy UM, et al. The association of cord serum cytokines with neurodevelopmental outcomes. *Am J Perinatol*. 2015;30(2):115-22.
20. Dammann O, O'Shea TM. Cytokines and perinatal brain damage. *Clin Perinatol*. 2008;35(4):643-63, v.
21. Doughty L, Carcillo JA, Kaplan S, Janosky J. The compensatory anti-inflammatory cytokine interleukin 10 response in pediatric sepsis-induced multiple organ failure. *Chest*. 1998;113(6):1625-31.
22. Romagnoli C, Frezza S, Cingolani A, De Luca A, Puopolo M, De Carolis MP, et al. Plasma levels of interleukin-6 and interleukin-10 in preterm neonates evaluated for sepsis. *Eur J Pediatr*. 2001;160(6):345-50.
23. Appachi E, Mossad E, Mee RB, Bokesch P. Perioperative serum interleukins in neonates with hypoplastic left-heart syndrome and transposition of the great arteries. *J Cardiothorac Vasc Anesth*. 2007;21(2):184-90.
24. Korzeniewski SJ, Allred E, Logan JW, Fichorova RN, Engelke S, Kuban KC, et al. Elevated endogenous erythropoietin concentrations are associated with increased risk of brain damage in extremely preterm neonates. *PLoS One*. 2015;10(3):e0115083.
25. Lv B, Huang J, Yuan H, Yan W, Hu G, Wang J. Tumor necrosis factor-alpha as a diagnostic marker for neonatal sepsis: a meta-analysis. *ScientificWorldJournal*. 2014;2014:471463.
26. Machado JR, Soave DF, da Silva MV, de Menezes LB, Etchebehere RM, Monteiro ML, et al. Neonatal sepsis and inflammatory mediators. *Mediators Inflamm*. 2014;2014:269681.
27. Lin CY, Chang YC, Wang ST, Lee TY, Lin CF, Huang CC. Altered inflammatory responses in preterm children with cerebral palsy. *Ann Neurol*. 2010;68(2):204-12.
28. Adrie C, Pinsky MR. The inflammatory balance in human sepsis. *Intensive Care Med*. 2000;26(4):364-75.
29. van der Poll T, Coyle SM, Moldawer LL, Lowry SF. Changes in endotoxin-induced cytokine production by whole blood after in vivo exposure of normal humans to endotoxin. *J Infect Dis*. 1996;174(6):1356-60.
30. Kim M, Kim M, Jeong H, Chae JS, Kim YS, Lee JG, et al. Hyporesponsiveness of natural killer cells and impaired inflammatory responses in critically ill patients. *BMC Immunol*. 2017;18(1):48.

Figures

Figure 1

Figure 1

Cytokine response in school age children with NE compared to control at baseline and on LPS stimulation: Epo, GM-CSF, IFN- γ , IL-10, IL-18, IL-1ra, IL-1 α , IL-1 β and IL-2: Mean serum cytokine concentration expressed in pg/ml, GM-CSF= Granulocyte macrophage colony stimulating factor, IL= Interleukin, IL-1RA= Interleukin1-Receptor antagonist, IL-1 β = Interleukin 1beta, IFN γ = interferon gamma, IL= Interleukin, IL-1 β = Interleukin 1beta, Vehicle or baseline, LPS= Lipopolysaccharide. * p value <0.05 versus Veh/ Control; ** versus Veh/ NE.

Figure 2

Figure 2

Cytokine response in school age children with NE compared to control at baseline and on LPS stimulation: Interleukin IL-6, IL-8, TNF-α, TNF-β and VEGF: Mean serum cytokine concentration expressed in pg/ml, TNF-α= Tumor necrosis factor alpha, VEGF= vascular endothelial growth factor. Veh= Vehicle or baseline, LPS= Lipopolysaccharide. * versus Veh Control p value < 0.05, ** versus veh/NE; *** versus all other parameters.

Figure 3

Figure 3

Cytokine response in children with NE who underwent Therapeutic Hypothermia (TH) vs NE without TH at baseline and on LPS stimulation: Cytokines EPO, GM-CSF, IFN- γ , IL-2, IL-10, IL-18, IL-1ra, IL-1 α , IL-1 β : Cytokine concentration expressed in pg/ml Epo= Erythropoietin, GM-CSF= Granulocyte Macrophage Colony stimulating factor, IL= Interleukin, IFN γ = Interferon gamma, IL-1RA= Interleukin 1 Receptor antagonist, Veh= Vehicle or baseline, LPS= Lipopolysaccharide. * p < 0.05 versus Veh Control

Figure 4

Figure 4

Cytokine response in children with NE who underwent TH vs NE without TH at baseline and on LPS stimulation: Interleukin IL-6, IL-8, TNF-α, TNF-β and VEGF: Cytokine concentration expressed in pg/ml, in children with NE and Therapeutic Hypothermia versus NE without TH. * p<0.05 versus Veh Controls.