State-Selective Electron Capture Cross Sections in Collision Between Be^{4+} and Ground State Hydrogen Atom – Significant Improvement in the Classical Treatment

Iman Ziaeeian

University of Debrecen

Károly Tőkési (✉ tokesi@atomki.hu)

Institute for Nuclear Research

Research Article

Keywords: State-Selective Electron, Collision , Be^{4+} and Ground State Hydrogen Atom , Classical Treatment

Posted Date: July 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-702501/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

K. Tokesi et al.

State-selective electron capture cross sections in collision between Be^{4+} and ground state hydrogen atom significant improvement in the classical treatment

Preprint of Paper to be submitted for publication in Scientific Reports

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 and 2019-2020 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

This document is intended for publication in the open literature. It is made available on the clear understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EUROfusion Programme Management Unit, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK or e-mail Publications.Officer@euro-fusion.org

Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EUROfusion Programme Management Unit, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK or e-mail Publications.Officer@euro-fusion.org

The contents of this preprint and all other EUROfusion Preprints, Reports and Conference Papers are available to view online free at <http://www.euro-fusionscipub.org>. This site has full search facilities and e-mail alert options. In the JET specific papers the diagrams contained within the PDFs on this site are hyperlinked

State-selective electron capture cross sections in collision between Be^{4+} and ground state hydrogen atom – significant improvement in the classical treatment

Iman Ziaeeian^{1,2} & Károly Tókesi¹✉

We present state-selective electron capture cross sections in collision between Be^{4+} and ground state hydrogen atom. The n - and nl -selective electron capture cross sections are calculated by a three-body classical trajectory Monte Carlo method (CTMC) and by a classical simulation schema mimicking quantum features of the collision system. The quantum behavior is taken into account with the correction term in the Hamiltonian as was proposed by Kirschbaum and Wilets (C. L. Kirschbaum, and L. Wilets, Phys. Rev. A **21**, 834 (1980)). Calculations are carried out in the projectile energy range of 1-1000 keV/amu. We found that our model for $\text{Be}^{4+} + \text{H}(1s)$ system remarkably improves the obtained state-selective electron capture cross sections, especially at lower projectile energies. Our results are very close and are in good agreement with the previously obtained quantum-mechanical results. Moreover our model with simplicity can time efficiently carry out simulations where maybe the quantum mechanical ones become complicated, therefore, our model should be an alternative way to calculate accurate cross sections and maybe can replace the quantum-mechanical methods.

¹Institute for Nuclear Research (Atomki), 4026 Debrecen Bem tér 18/c, Hungary. ²Doctoral School of Physics, Faculty of Science and Technology, University of Debrecen, P.O. Box 400, H-4002, Debrecen, Hungary. ✉email: tokesi@atomki.hu

Beryllium is widely used as a first wall element of the fusion reactors¹ because of its unique thermo-physical properties. So, due to wall erosion, Beryllium should be one of main impurity in fusion chamber². The radiative decay of excited impurity ions can be the source for the energy loss of the plasma and can cool the plasma. These radiative decays can be analyzed by the electron capture recombination spectroscopy (CXRS). Therefore, the exact knowledge of electron capture cross sections in collisions between Be ions and hydrogen atoms is essential³. Due to the experimental difficulties, the experimental results for electron capture cross sections in $\text{Be}^{4+} + \text{H}$ collisions are entirely lacking, but those were studied intensively theoretically in the past years. The total electron capture cross sections have been studied using various models and methods such as applying the quantum-mechanical molecular orbital close-coupling (QMOCC)⁴, the atomic orbital close-coupling (AOCC)⁵, the hyper spherical close-coupling (HSCC)⁶ models, using the solution of the time dependent Schrödinger equation (TDSE)⁷, the lattice time dependence Schrödinger equation (LTDSE)⁸, the classical over barrier model (COBM)⁹ and the classical trajectory Monte Carlo method^{10,11}. The partial electron capture cross sections in the collision between Be and hydrogen atom have been also studied using different quantum-mechanical methods such as: QMOCC^{4,12}, AOCC⁵, one-electron diatomic molecule (OEDM)¹³, and boundary corrected continuum intermediate state (BCCIS)¹⁴ models. It is worth noting that all the results have been published for projectile energy below 100 keV/amu. The calculation of the principle quantum number, n , dependent cross sections has been studied by Jorge *et al.*¹⁵ by solving the time-dependent Schrödinger equation with the GridTDSE package (GTDSE) numerically in the broad energy range between 1 keV/amu and 500 keV/amu.

In this work we present the electron capture cross sections into the bound states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collisions. We treat the collision dynamics classically using a three-body classical trajectory Monte Carlo (CTMC) and a three-body quasi classical Monte Carlo (QCTMC) model when the Heisenberg correction term is added to the standard CTMC model via model potential [16-21]. Since there is no experimental data available, our calculated cross sections are compared with the previous theoretical results.

Results

For each collision energies, the calculation of the state selective electron capture cross sections requires to follow 10^7 classical trajectories. At first, we tested three calculation schemes during our simulations since the Heisenberg correlation potential may influenced the obtained results significantly. These are the following: 1) target-centered, where the correction term is taken into account between the target electron the target nucleus, 2) projectile-centered, where the correction term is taken into account between the target electron the projectile. 3) combined one, i.e., target and projectile centered when the correction term is taken into account between target electron and both the target nucleus and projectile.

As an example Fig. 1 shows our CTMC and QCTMC results corresponding to the three calculation schema of the electron capture cross sections into the 4s state of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision as a function of the impact energy. It can be seen that the effects of the correction term at lower energies are significant. While for the case of target-centered, the cross sections at lower incident energies are increasing compared to the standard CTMC results for the case of projectile-centered they are decreasing. The combination of the use of target- and projectile-centered corrections results increases the cross sections and we also obtained good agreement between our QCTMC results and previous full quantum mechanical results in the entire impact energy range. Therefore in followings, for the calculation of the capture cross sections, we will use only the combination scheme.

Physically, due to the Heisenberg constraint, the electron cannot collapse to the target and projectile nucleus in the electron capture channel. To clarify this further, we calculated the electron capture probabilities as a function of the impact parameter.

Figure 1. Electron capture cross sections into the 4s state of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision as a function of the impact energy. Solid red line: present CTMC results, solid black line: present *target-centered* QCTMC results, solid green line: present *projectile-centered* QCTMC results, solid blue line: present *target and projectile centered* QCTMC results, red squares: AOCC results of Fritsch⁵, green circles: QMOCC results of Harel *et al.*⁴, black triangles: BCCIS results of Das *et al.*¹⁴, blue inverse triangle: OEDM results of Errea *et al.*¹³

Figure 2 shows the present CTMC and QCTMC results with the three calculation schemes of the electron capture probabilities into the specific $n = 3, 4$ and $nl = 3d, 4s$ states of the projectile at 10 keV/amu impact energy in $\text{Be}^{4+} + \text{H}(1s)$ as a function of impact parameter. The impact parameter dependent electron capture probabilities, $bP(b)$, were fitted by a Gaussian function. The peak maxima of the Gaussian fitting is also shown in fig. 2. We note that the area under the curves is proportional to the state-selective electron capture cross sections. We found that the probability of electron capture is higher in target-centered QCTMC and lower in projectile-centered QCTMC model compared with the standard CTMC model. This behavior can be understood with the explanation of the acting forces between the interacting particles, $F = -dU/dr$. The attractive force between an electron and both of proton and positive projectile ion, is due to the Coulomb interaction and repulsive force is due to the Heisenberg correction term as follow:

$$F_{\text{Heisenberg}} = -\left(\frac{\xi_H^2}{2\alpha_H r^3} + \frac{rp^4}{\xi_H^2}\right) \exp\left\{\alpha_H \left[1 - \left(\frac{rp}{\xi_H}\right)^4\right]\right\}. \quad (1)$$

The attractive Coulomb force acts between the electron and positively charged, target and projectile, in the same way in all schemes. This force, most of the time of the collision, is much larger than $F_{\text{Heisenberg}}$. On the other hand, in the target-centered scheme, the repulsive force, $F_{\text{Heisenberg}}$, is toward the projectile, but on the contrary, this repulsive force is towards the target in projectile-centered mode. We note that this repulsive force, of course, does not show up in the standard CTMC model. According to the sum of the forces, the electron has the highest attraction to the projectile in the target-centered QCTMC and the least attraction to the projectile in the projectile-centered QCTMC. With this scenario, the case of CTMC is placed

between the above two modes. Therefore, the probability of electron capture in projectile-centered QCTMC, CTMC, and target-centered QCTMC modes increases, respectively.

Another noteworthy point is that the peak maxima in CTMC and QCTMC projectile-centered cases are very close to each other and locate in lower impact parameters. This is also true in QCTMC target-centered and QCTMC combined target- and projectile-centered cases, except that the peak maxima are at higher impact parameters.

Figure 3 shows the present CTMC and QCTMC results of the electron capture cross sections into the $n = 3, 4, 5$ states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision as a function of the impact energy. The present classical results are compared with Fritsch⁵, Harel *et al.*⁴, and Das *et al.*¹⁴, as well. The QCTMC results are higher than the CTMC ones at low and intermediate impact energies. This difference is more significant in $n = 4$ and $n = 5$ states. The best matching between present CTMC and QCTMC is seen at high energies. In $n = 3$ and $n = 5$ states, the present QCTMC results agree well with the available quantum-mechanical approaches such as; QMOCC⁴, AOCC⁵, and BCCIS¹⁴.

The standard statistical error (see Eq. 10) at 1000 keV/amu impact energy is around 4% in CTMC and QCTMC, respectively. Figure 3 also shows the cross sections for higher states where no previous data are available.

Figure 2. Probability for electron capture into $n = 3, 4$ and $nl = 3d, 4s$ states of the projectile (multiplied by impact parameter) in $\text{Be}^{4+} + \text{H}(1s)$, as a function of the impact parameter, at 10 keV/amu impact energy. Dash-dotted red line: present CTMC results, dash black line: present target-centered QCTMC results, solid green line: present projectile-centered QCTMC results, solid blue line: combination of target-projectile-centered.

Figure 4 shows our CTMC and QCTMC results of the electron capture cross sections into 3s, 3p and 3d states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ as a function of the impact. The comparison is made with Fritsch⁵, Harel *et al.*³, Das *et al.*¹⁴, and Errea *et al.*¹³. The QCTMC model significantly improve the cross sections compared to the CTMC at low and intermediate impact energies. Moreover, the unique agreement is obtained between the present QCTMC results and

1) QMOCC results⁴ in the 3s state, 2) OEDM¹³, and BCCIS results¹⁴ in 3p state 3) QMOCC⁴, AOCC⁵, OEDM¹³, and BCCIS¹⁴ results in 3d state of the Be^{3+} at energies lower than 100 keV/amu, respectively.

Figure 3. Electron capture cross sections into the $n = 3, 4, 5$ states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision as a function of the impact energy. Solid red line: present CTMC results, solid blue line: present QCTMC results, red squares: AOCC results of Fritsch⁵, green circles: QMOCC results of Harel *et al.*⁴, black triangles: BCCIS results of Das *et al.*¹⁴. Also, electron capture cross sections into $n = 6, 8, 10$ states are recommended.

Figure 4. Electron capture cross sections into 3s, 3p, and 3d states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision as a function of the impact energy. Solid red line: present CTMC results, solid blue line: present QCTMC results, red squares: AOCC results of Fritsch⁵, green circles: QMOCC results of Harel *et al.*⁴, black triangles: BCCIS results of Das *et al.*¹⁴, blue inverse triangle: OEDM results of Errea *et al.*¹³.

Figure 5 represents our CTMC and QCTMC results of the electron capture cross sections into 4s, 4p, 4d, and 4f states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ as a function of the impact. We have compared the present classical results with the quantum-mechanical approaches such as QMOCC⁴, AOCC⁵, OEDM¹³, and BCCIS¹⁴. The QCTMC model remarkably increases the cross sections compared with the CTMC at low and intermediate energies. The difference between the present CTMC and QCTMC results at low energies gradually increases from 4s to 4f states. It can be seen that our CTMC results have the best agreement with the AOCC results of Fritsch⁵ in 4s and 4p states. Moreover, the QCTMC cross sections in 4s and 4p states have better agreement with the QMOCC results of Harel *et al.*⁴ and the OEDM results of Errea *et al.*¹³. However, both present classical results are in excellent agreement with the BCCIS results of Das *et al.*¹⁴ at intermediate energies. The standard statistical error for 4s, 4p, 4d, and 4f states is around 1.6% in the range of 1-500 keV/amu impact energies. At the same time, for the projectile energy range of 500-1000 keV/amu, the estimated uncertainties around 4%.

Figure 5. Electron capture cross sections into 4s, 4p, 4d, and 4f states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision, as a function of the impact energy. Solid red line: present CTMC results, solid blue line: present QCTMC results, red squares: AOCC results of Fritsch⁵, green circles: QMOCC results of Harel *et al.*⁴, black triangles: BCCIS results of Das *et al.*¹⁴, blue inverse triangle: OEDM results of Errea *et al.*¹³.

Figure 6 shows the present CTMC and QCTMC results of the electron capture cross sections into 5s, 5p, 5d, and 5f states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ as a function of the impact energy. The obtained results are compared with QMOCC⁴, BCCIS¹⁴, and OEDM¹³ methods, as well. According to figure 6, the QCTMC method outstandingly enhances the cross sections compare to the CTMC results at impact energies lower than about 60 keV/amu. Good agreements are obtained between the present QCTMC results with the OEDM results of Errea *et al.*¹³ and the QMOCC results of Harel *et al.*⁴ in 5s, 5d, and 5f states of the projectile. The present CTMC and QCTMC results in all 5l-states agree well with the BCCIS results of Das *et al.*¹⁴ at intermediate energies. The QCTMC and CTMC cross sections are approximately matched at the impact energies greater than 100 keV/amu.

Figure 6. Electron capture cross sections into 5s, 5p, 5d, and 5f states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ collision, as a function of the impact energy. Solid red line: present CTMC results, solid blue line: present QCTMC results, green circles: QMOCC results of Harel *et al.*⁴, black triangles: BCCIS results of Das *et al.*¹⁴, blue inverse triangle: OEDM results of Errea *et al.*¹³

According to Fig. 3-6, the present CTMC and QCTMC results of the electron capture cross sections into specific states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$ are given for several typical impact energies in Table I. As we already mentioned, the QCTMC cross sections are larger compared to CTMC ones at lower incident energies. However as the energy increases, this difference gradually decreases so that at very high energies, this difference is negligible. To explain this behavior physically, we focus on the force between the electron and the hydrogen nucleus. Typically, the net Coulomb force is applied between two bodies, which is inversely related to the square of the distance between them. Heisenberg correction term (see Eq. 6) generates a repulsive force in the opposite direction to the Coulomb force. In this case, the attraction force between the electron and the target's nucleus decreases, increasing the electron's reactivity with the projectile's ion in the electron capture channel. Also, the long- distance of the projectile to the electron practically reduces this repulsive force's effect on the calculations (see Fig. 1).

On the other hand, the passing projectile ion at low energies causes the extension of the interaction time. Therefore, the effect of these factors increases the cross section at low energies in the QCTMC model. Also, the interaction time is shorter at high energies. Furthermore, due to the small Heisenberg repulsive force, the correction term gradually loses its effects; therefore, the CTMC and QCTMC results are approximately the same.

TABLE I. The present CTMC and QCTMC results of the electron capture cross sections into specific states of the projectile in $\text{Be}^{4+} + \text{H}(1s)$.

Energy (keV/amu)	Model	Cross Section (in 10^{-16} cm^2)					
		3s	3d	4s	4d	5s	5d
1	CTMC	2.608	7.175	0.022	0.116	0.001	0.001
	QCTMC	5.698	17.81	0.686	3.370	0.017	0.056
5	CTMC	1.326	11.09	0.135	0.314	0.012	0.019
	QCTMC	3.343	15.74	0.385	3.844	0.061	0.170
10	CTMC	0.681	11.61	0.183	0.571	0.033	0.081
	QCTMC	1.757	15.46	0.276	3.364	0.084	0.316
35	CTMC	0.209	6.780	0.112	1.652	0.062	0.464
	QCTMC	0.288	6.469	0.158	2.275	0.087	1.024
55	CTMC	0.117	3.109	0.077	1.220	0.054	0.623
	QCTMC	0.144	2.787	0.089	1.161	0.055	0.658
70	CTMC	0.081	1.793	0.052	0.797	0.036	0.450
	QCTMC	0.089	1.576	0.062	0.705	0.039	0.401
90	CTMC	0.052	0.879	0.033	0.401	0.022	0.263
	QCTMC	0.040	0.569	0.028	0.405	0.023	0.212
200	CTMC	0.007	0.039	0.005	0.024	0.003	0.015
	QCTMC	0.004	0.039	0.003	0.028	0.002	0.0187

Discussions

The electron capture cross sections into $n = 3, 4, 5, 6, 8, 10$ and $nl = 3l, 4l, 5l$ states of the projectile have been presented in $\text{Be}^{4+} + \text{H}(1s)$ in the framework of CTMC and QCTMC methods. For the determination of the cross sections 10^7 trajectories were calculated for each impact energies. We found that the QCTMC cross sections are higher than the CTMC ones at low energies. We have used the previous AOCC, QMOCC, BCCIS, and OEDM quantum-mechanical approaches for comparison with our present data. Including the potential correction term to mimic the Heisenberg uncertainty principle in the classical Hamiltonian, we have shown that our QCTMC capture cross sections into the projectile states, $n = 3, 5$ and $nl = 3s, 3p, 3d, 4s, 4p, 5s, 5d, 5f$ are in excellent agreement with quantum-mechanical results. We believe that our model, with its simplicity, can be an alternative way to calculate accurate cross sections and maybe can replace the results of the quantum-mechanical models, where the quantum mechanical calculations become complicated.

Methods

The QCTMC model, in principle, takes into account the Heisenberg and Pauli constraints in adding a correction term into the standard original Hamiltonian¹⁶. For hydrogen atom, which has only one electron, the Pauli correction can automatically neglect. Therefore, the quasi classical Hamiltonian consists of correction potential, V_H , inspired by the Heisenberg principles can be written as:

$$H_{QCTMC} = H_0 + V_H, \quad (2)$$

where

$$V_H = \sum_{n=a,b} \sum_{i=1}^N f(r_{ni}, p_{ni}; \xi_H, \alpha_H). \quad (3)$$

a and b denote the nuclei, and the i index the electrons. r and p are the distance and momentum of an electron with respect to a nucleus, which is defined as follows:

$$r_{ab} = r_b - r_a, \quad (4)$$

$$p_{ab} = \frac{m_a p_b - m_b p_a}{m_a + m_b}. \quad (5)$$

The Heisenberg correction function is expressed as¹⁶

$$f(r_{ab}, p_{ab}; \xi_H, \alpha_H) = \frac{\xi_H}{4\alpha_H r_{ab}^2 \mu_{ab}} \exp \left\{ \alpha_H \left[1 - \left(\frac{r_{ab} p_{ab}}{\xi_H} \right)^4 \right] \right\}, \quad (6)$$

where subscripts a and b indicate pairs of particles with reduced mass μ_{ab} . The parameter ξ_H reflects the size of the core while α_H is a hardness parameter. These parameters universally are used where $\alpha_H = 4$ and $\xi_H = 0.9428$, respectively¹⁹⁻²¹. The Heisenberg potential between the target electron and both target core and projectile (p; projectile, e; electron, T; target) are defined as follows:

$$f(\vec{r}_{pe}, \vec{p}_{pe}; \varepsilon_H, \alpha_H) = \frac{\xi_H^2}{4\alpha_H \vec{r}_{pe}^2 \mu_{pe}} \exp \left\{ \alpha_H \left[1 - \left(\frac{\vec{r}_{pe} \vec{p}_{pe}}{\xi_H} \right)^4 \right] \right\}, \quad (7)$$

$$f(\vec{r}_{Te}, \vec{p}_{Te}; \varepsilon_H, \alpha_H) = \frac{\xi_H^2}{4\alpha_H \vec{r}_{Te}^2 \mu_{Te}} \exp \left\{ \alpha_H \left[1 - \left(\frac{\vec{r}_{Te} \vec{p}_{Te}}{\xi_H} \right)^4 \right] \right\}. \quad (8)$$

The total cross sections are computed with the following formula:

$$\sigma = \frac{2\pi b_{max}}{T_N} \sum_j b_j^{(i)}, \quad (9)$$

and the statistical uncertainty of the cross sections is given by:

$$\Delta\sigma = \sigma \left(\frac{T_N - T_N^{(i)}}{T_N T_N^{(i)}} \right)^{1/2}, \quad (10)$$

where T_N is the total number of trajectories calculated for impact parameters less than b_{max} , $T_N^{(i)}$ is the number of trajectories that satisfy the criteria for the corresponding final channels (electron capture), and $b_j^{(i)}$ is the actual impact parameter for the trajectory corresponding electron capture processes.

In the classical approaches, the classical principal (n_c) and the orbital angular momentum (l_c) quantum numbers are defined by

$$n_c = Z_T Z_e \left(\frac{\mu_{Te}}{2U} \right)^{1/2}, \quad (11)$$

$$l_c = \sqrt{m_e [(x\dot{y} - y\dot{x})^2 + (x\dot{z} - z\dot{x})^2 + (y\dot{z} - z\dot{y})^2]}, \quad (12)$$

where μ_{Te} is the reduced mass of the target nucleus and the target electron. x , y , and z are the Cartesian coordinates of the electron relative to the nucleus and $\dot{x}$, $\dot{y}$, and $\dot{z}$ are the corresponding

velocities. The classical values of n_c are “quantized” to a specific level n [22] if they satisfy the relation:

$$[(n-1)(n-1/2)n]^{1/3} \leq n_c \leq [(n+1)(n+1/2)n]^{1/3}. \quad (13)$$

Since l_c is uniformly distributed for a given n level²³, the quantal statistical weights are reproduced by choosing bin sizes such that

$$l \leq \frac{n}{n_c} l_c \leq l+1, \quad (14)$$

where l is the quantum-mechanical orbital angular momentum.

Acknowledgments

This work has been carried out within the framework of the EURO fusion Consortium and has received funding from the Euratom research and training program 2014-2018 and 2019-2020 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

References

1. Pitts, R. *et al.* Physics basis and design of the ITER plasma-facing components. *J. Nucl. Mater.* **415**, S957 (2011).
2. Mattioli, M. *et al.* Laser blow-off injected impurity transport in L mode Tore Supra plasmas. *Nucl. Fusion*. **38**, 1629 (1998).
3. Loarte, A. *et al.* Chapture4: Power and particle control. *Nucl. Fusion*. **47**, S203 (2007).
4. Harel, C. *et al.* Cross sections for electron capture from atomic hydrogen by fully stripped ions in the 0.05-1.00 a.u. impact velocity range. *At. Data. Nucl. Data Tables*. **68**, 279 (1998).
5. Fritsch, W. & Lin, C. D. Atomic-orbital-expansion studies of electron transfer in bare-nucleus $Z(Z=2, 4-8)$ -hydrogen-atom collisions. *Phys. Rev. A*. **29**, 3039 (1984).
6. Le, A. T. *et al.* Hyperspherical close-coupling for charge transfer cross sections in $\text{Si}^{4+} + \text{H}(\text{D})$ and $\text{Be}^{4+} + \text{H}$ collisions at low energies. *J. Phys. B*. **36**, 3281 (2003).
7. Lüdde, H. J. & Dreizler, R. M. Electron capture with He^{2+} , Li^{3+} , Be^{4+} and B^{5+} projectiles from atomic hydrogen. *J. Phys. B*. **15**, 2713 (1982).
8. Minami, T. *et al.* Lattice, time-dependent Schrödinger equation approach for charge transfer in collisions of Be^{4+} with atomic hydrogen. *J. Phys. B*. **39**, 2877 (2006).
9. Sattin, F. Classical overbarrier model to compute charge exchange and ionization between ions and one-optical-electron atoms. *Phys. Rev. A* **62**, 042711 (2000).
10. Illescas C. & Riera, A. Classical study of single-electron capture and ionization processes in $\text{A}^{q+} + (\text{H}, \text{H}_2)$ collisions. *Phys. Rev. A*. **60**, 4546 (1999).
11. Schultz, D. R. *et al.* Inelastic processes in 1-1000 keV/u collisions of Be^{q+} ($q = 2-4$) ions with atomic and molecular hydrogen. *Physica Scripta*. **T62**, 69 (1996).
12. Harel, C. *et al.* Description of ionization in the molecular approach to atomic collisions. *Phys. Rev. A*. **55**, 287 (1997).
13. Errea, L. F. *et al.* Quantal and semiclassical calculations of charge transfer cross sections in $\text{Be}^{4+} + \text{H}$ collisions for impact energies of $2.5 \text{ eV amu}^{-1} < E < 25 \text{ keV amu}^{-1}$. *J. Phys. B*. **31**, 3527 (1998).
14. Das, M. *et al.* Charge-transfer cross sections in collisions of Be^{q+} ($q = 1-4$) and B^{q+} ($q = 1-5$) with ground-state atomic hydrogen. *Phys. Rev. A*. **57**, 3573 (1998).
15. Jorge, A. *et al.* Application of a grid numerical method to calculate state-selective cross sections for electron capture on $\text{Be}^{4+} + \text{H}(1s)$ collisions. *Phys. Rev. A*. **94**, 032707 (2016).
16. Kirschbaun, C. L. & Wilet, L. Classical many-body model for atomic collisions incorporating the Heisenberg and Pauli principles. *Phys. Rev. A*. **21**, 834 (1980).

17. Cohen, J. S. Quasiclassical effective Hamiltonian structure of atoms with $Z=1$ to 38. *Phys. Rev. A.* **51**, 266 (1995).
18. Cohen, J. S. Quasiclassical-trajectory Monte Carlo methods for collisions with two-electron atoms *Phys. Rev. A.* **54**, 573 (1996).
19. Cohen, J. S. Molecular effects on antiproton capture by H_2 and the states of $\overline{p}p$ formed. *Phys. Rev. A.* **56**, 3583 (1997).
20. Cohen, J. S. Extension of quasiclassical effective Hamiltonian structure of atoms through $Z=94$. *Phys. Rev. A.* **57**, 4964 (1998).
21. Cohen, J. S. Multielectron effects in capture of antiprotons and muons by helium and neon. *Phys. Rev. A.* **62**, 022512 (2000).
22. Becker, R. L. & Mackellar, A. Theoretical initial l dependence of ion-Rydberg-atom collision cross sections. *J. Phys. B At. Mol. Opt. Phys.* **17**, 3923 (1984).
23. Tőkési, K. & Mukoyama, K. Theoretical investigation of the ECC peak for charge particles with the CTMC method. *Bull Ins Chem Res Kyoto Univ.* **72**, 62 (1994).