Long noncoding RNA UCA1 facilitates cell proliferation and inhibits apoptosis via activating PI3K/Akt pathway in retinoblastoma

Zhongfang Yuan

Jinan Central Hospital Affiliated to Shandong University

Zhaona Li (✉ lizhaona133@163.com)

The Second People's Hospital of Jinan

Research article

Keywords: lncRNA-UCA1; RB; proliferation; apoptosis; PI3K/Akt pathway

Posted Date: October 23rd, 2019

DOI: <https://doi.org/10.21203/rs.2.16387/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Retinoblastoma (RB) is the most common intraocular malignancy of childhood. This study is aimed to explore the effect of lncRNA-UCA1 on RB and its potential molecular mechanisms.

Methods In our study, the expression of lncRNA-UCA1 was measured by qRT-PCR in both RB tissues and RB HXO-RB44 and Y79 cells. The relationship between lncRNA-UCA1 expression and clinical parameters in RB patients were evaluated. Cell proliferation, cell clones, apoptosis and cell cycle of HXO-RB44 and Y79 cells were measured by cell counting kit-8 (CCK-8) assay, colony formation assay and flow cytometry, respectively. In addition, the expressions of PCNA, Caspase-3, survivin, p16, p21, CDK2, PI3K, p-PI3K, Akt, p-Akt and S6k in HXO-RB44 and Y79 cells were measured by western blot.

Results lncRNA-UCA1 was highly expressed in both RB tissues and RB HXO-RB44 and Y79 cells. Moreover, the expression of lncRNA-UCA1 in RB patients was remarkably correlated with tumor size, optic nerve invasion and pathologic grade. lncRNA-UCA1 markedly facilitated cell proliferation and cell cycle procession, as well as inhibited cell apoptosis in HXO-RB44 and Y79 cells. lncRNA-UCA1 dramatically increased the expression of S6k and the phosphorylation of PI3K and Akt in RB cells. LY294002 (PI3K inhibitor) reversed the effects of lncRNA-UCA1 on RB cell proliferation, apoptosis and cell cycle procession.

Conclusions Our study indicated that lncRNA-UCA1 could promote cell proliferation and cell cycle procession, as well as inhibit cell apoptosis in RB via activating PI3K/Akt pathway.

Background

Retinoblastoma (RB) is the most common intraocular cancer and mainly affects children under 5 years old [1, 2]. The incidence of RB is about 1/15000 all over the world and has no gender predilection [3]. Although the therapeutic techniques have greatly improved, the survival rate of RB patients still remains poor, mainly due to its insensitivity to chemotherapy and radiotherapy [4]. Therefore, it is urgent to investigate new diagnostic markers and therapeutic targets for RB treatment.

Long noncoding RNAs (lncRNAs) are defined as RNA transcripts longer than 200 nts that have no protein-encoding ability [5]. Accumulating evidence has demonstrated that lncRNAs play important roles in the development and progression of cancers including RB [6]. More and more researches have confirmed that lncRNAs participate in numerous biological activities, such as cell proliferation, migration, invasion and cell cycle arrest [7, 8]. For example, lncRNA BDNF-AS has been reported to suppress RB cell proliferation, migration and cell cycle transition [9]. Su et al. [10] have indicated that lncRNA BANC1 could suppress cell proliferation, migration and invasion in RB. Dong et al. [11] have confirmed that lncRNA-HOTAIR could promote cell proliferation and invasion via regulating Notch pathway in RB. It is reported that lncRNA-H19 inhibited RB cell proliferation, migration and invasion, and promoted cell apoptosis [12]. However, the effect of lncRNA-UCA1 was still fully unclear in RB.

The phosphatidylinositol 3'-kinase/protein kinase B (PI3K/Akt) pathway is reported to play a crucial part in a variety of malignancies [13–15]. In addition, some lncRNAs may exert their functions by regulating PI3K/Akt pathway in cancer [5]. Previous researches have confirmed that lncRNA-UCA1 could promote osteosarcoma metastasis by activating PI3K/Akt pathway [16]. Li et al. [17] have indicated that lncRNA-UCA1 could facilitate cell proliferation and suppress apoptosis through regulating PI3K/Akt pathway in gastric cancer. Therefore, we investigate whether lncRNA-UCA1 could affect RB via regulating PI3K/Akt pathway.

In this research, we explored the functions of lncRNA-UCA1 on cell proliferation and apoptosis in RB, and its related molecular mechanisms. Our results indicated that lncRNA-UCA1 could facilitate cell proliferation and inhibit apoptosis through activating PI3K/Akt pathway in RB. Our results indicate that lncRNA-UCA1 may serve as a new therapeutic target for RB.

Methods

Clinical samples

Total 42 fresh RB specimens and the paired adjacent normal retina specimens from the patients of RB were provided by the Ophthalmology Department of our hospital. None of the patients received anti-tumor treatment, chemotherapy and radiotherapy before surgery. In addition, the characteristics of RB patients were presented in Table 1. This research was permitted by the Ethics Committee of our hospital. All patients or their guardians have signed informed consent.

Cell cultures

Two RB cell lines (HXO-RB44 and Y79) and normal retinal vascular endothelial cell line (ACBRI-181) were supplied by American Type Culture Collection (ATCC, USA). The cell lines were cultured in RPMI1640 medium (Gibco, USA) containing 10% FBS and 1% penicillin/streptomycin. All cells were cultured in a humidified incubator with 5% CO₂ at 37°C.

Cell transfection assay

The pcDNA3.1-UCA1, UCA1 siRNA and their corresponding negative control (pcDNA3.1-Mock and UCA1 siRNA control) were obtained from Thermo Fisher, USA. LY294002 (PI3K inhibitor) was purchased from Sigma, USA. The pcDNA3.1-UCA1, pcDNA3.1-Mock, UCA1 siRNA and UCA1 siRNA control were separately transfected into HXO-RB44 and Y79 cells by Lipofectamine[®] 3000 Reagent (Invitrogen, USA). The transfected HXO-RB44 and Y79 cells were randomly divided into six groups: BLANK group (no-treated group), pcDNA-UCA1 group (transfected with pcDNA3.1-UCA1), pcDNA-Mock group (transfected with pcDNA3.1-Mock), si-UCA1 group (transfected with UCA1 siRNA), si-NC group (transfected with UCA1

siRNA control) and pcDNA-UCA1 + LY294002 group (transfected with pcDNA-UCA1 and treated with 50 μ M LY294002). Finally, all cells were cultured in 37°C for 48 hours.

Cell counting kit–8 assay

The cell proliferation abilities of HXO-RB44 and Y79 cells were tested by cell counting kit–8 (CCK–8) kit (Sigma, USA). In brief, the transfected HXO-RB44 and Y79 cells (1×10^4 cells/well) were planted into 96-well plates. At different time points of 24, 48, 72 and 96 h, 10 μ L CCK–8 was added into each well. Subsequently, the cells were then cultured at room temperature for 1 h. Finally, the absorbance was measured at 450 nm by a microplate reader (Bio-Rad, USA).

Colony formation assay

The transfected HXO-RB44 and Y79 cells were respectively seeded into 6-well plates at a density of 300 cells per well and cultivated for 14 d. After fixed in 4% cold formaldehyde for 30 min, the colonies were stained with 0.1% crystal violet (Beyotime, China) for 20 min. Finally, the colonies were photographed and counted under an optical microscope (Olympus, Tokyo, Japan).

Cell cycle analyses

The transfected HXO-RB44 and Y79 cells were harvested, washed with PBS, fixed in pre-cold 70% (v/v) ethanol for 2 h at -20°C . Subsequently, the cells were stained with Muse™ Cell Cycle Reagent (Merck Millipore, USA) in dark for 30 min. Finally, the cells were observed and analyzed by MUSE™ flow cytometry (Merck Millipore, USA).

Apoptosis assay

The apoptosis of HXO-RB44 and Y79 cells was determined by Annexin V-FITC apoptosis detection kit (Invitrogen, USA). In brief, the transfected HXO-RB44 and Y79 cells were collected and resuspended in Binding buffer. Then, the transfected HXO-RB44 and Y79 cells were stained by using Annexin V-FITC and propidium iodide (PI) for 15 min in a dark room. Subsequently, the apoptotic cells were observed using MUSE™ flow cytometer.

Real-Time fluorogenic PCR assays

Total RNA from RB tissue, adjacent normal retina tissue, HXO-RB44 cells, Y79 cells and ACBRI–181 cells was extracted using TRIZOL (Invitrogen, USA). Then, cDNA was synthesized from total RNA using Revert Aid First Strand cDNA Synthesis Kit (Thermo, USA). Subsequently, the qRT-PCR was performed using

SYBR green qPCR Master Mix (Thermo Scientific, USA) according to the manufacturer's protocol. Primers used in this study were as follows: lncRNA-UCA1 F: 5'-CCACACCCAAAACAAAAATCT-3', R: 5'-TCCCAAGCCCTCTAACAAACA-3'; GAPDH F: 5'-TGTTTCGTCATGGGTGTGAAC-3', R: 5'-ATGGCATGGACTGTGGTCAT-3'.

Western blot analysis

The total proteins were extracted by RIPA buffer (Beyotime, Jiangsu, China). 50 µg of protein samples were separated by 10% SDS-PAGE and then transferred onto nitrocellulose membrane. Subsequently, the membranes were blocked with 5% milk in TBST for 2 h, followed by overnight incubations at 4°C with the primary antibody (PI3K, 1:1000; Akt, 1:1000; PCNA, 1:1000; GAPDH, 1:1000, Sino Biological, USA. p-PI3K, 1:500; p-Akt, 1:500; S6K, 1:1000, Abcam, USA. survivin, 1:1000; CDK2, 1:1000; Caspase-3, 1:1000; p16, 1:1000; p21, 1:1000, Cell Signal, USA). Subsequently, the membranes were incubated with the peroxidase-labeled secondary antibody (anti-rabbit IgG, 1:5000, I5381MSDS, Sigma, USA) for 1 h. Finally, the protein bands were detected by an enhanced chemiluminescence (ECL) kit (Thermo Fisher, USA).

Statistical analysis

All statistical analyses were performed using SPSS 22.0 Statistical Software (Chicago, IL). The results were presented as the mean ± SD. Statistical significance was tested using the one-way ANOVA or Student's t-test. $P < 0.05$ was considered to be statistically significant.

Results

The expression of lncRNA-UCA1 is upregulated and associated with clinical parameters in RB

The results of qRT-PCR showed that lncRNA-UCA1 expression in RB tissue was markedly higher than that in paracancerous tissue ($P < 0.01$) (*Fig 1A*). Similarly, lncRNA-UCA1 expression was remarkably increased in HXO-RB44 and Y79 cells compared with ACBRI-181 cells ($P < 0.01$) (*Fig 1B*). All the results suggested that lncRNA-UCA1 was highly expressed in RB. We also found that the expression of lncRNA-UCA1 was remarkably correlated with tumor size ($P < 0.01$), optic nerve invasion ($P < 0.01$) and pathologic grade ($P < 0.01$) (*Table 1*). Meanwhile, the expression of lncRNA-UCA1 had no significantly correlation with age and gender in the patients with RB ($P > 0.05$) (*Table 1*). In addition, the expression of lncRNA-UCA1 in transfected HXO-RB44 and Y79 cells was detected by qRT-PCR. The results showed that lncRNA-UCA1 expression of HXO-RB44 and Y79 cells was significantly increased in pcDNA-UCA1 group compared with BLANK and pcDNA-Mock group ($P < 0.01$) (*Fig 1C*). Meanwhile, when compared with BLANK and si-NC group, lncRNA-UCA1 expression of HXO-RB44 and Y79 cells was markedly decreased in si-UCA1 group ($P < 0.01$) (*Fig 1C*), suggesting that the transfection was successful.

lncRNA-UCA1 promotes cell proliferation in HXO-RB44 and Y79 cells

The results of CCK-8 assay showed that the proliferation ability of HXO-RB44 cells was markedly decreased in si-UCA1 group compared with BLANK and si-NC group at 48 h ($P < 0.05$), 72 h and 96 h ($P < 0.01$) (*Fig 2A*). When compared with BLANK and pcDNA-Mock group, the proliferation ability of Y79 cells was remarkably increased in pcDNA-UCA1 group at 48 h ($P < 0.05$), 72 h and 96 h ($P < 0.01$) (*Fig 2A*). Furthermore, colony formation assay also confirmed that silencing lncRNA-UCA1 expression markedly decreased the cell clones number of HXO-RB44 cells ($P < 0.01$) (*Fig 2B*), while lncRNA-UCA1 overexpression significantly increased the cell clones number of Y79 cells ($P < 0.01$) (*Fig 2B*). To further confirm the function of lncRNA-UCA1 on RB cell proliferation, we also measured the expression of PCNA in both HXO-RB44 and Y79 cells. The expression of PCNA of HXO-RB44 cells was dramatically decreased in si-UCA1 group compared with BLANK and si-NC group ($P < 0.01$) (*Fig 2C*). When compared with BLANK and pcDNA-Mock group, the expression of PCNA of Y79 cells was markedly increased in pcDNA-UCA1 group ($P < 0.01$) (*Fig 2C*). All the results indicated that lncRNA-UCA1 could promote cell proliferation in RB cells.

lncRNA-UCA1 inhibits cell apoptosis in HXO-RB44 and Y79 cells

As shown in *Fig 3A*, silencing lncRNA-UCA1 expression dramatically increased HXO-RB44 cells apoptosis ($P < 0.01$) (*Fig 3A*). However, lncRNA-UCA1 overexpression significantly decreased the apoptosis of Y79 cells ($P < 0.01$) (*Fig 3A*). To further confirm the effect of lncRNA-UCA1 on RB cell apoptosis, we detected the expression of Caspase-3 and survivin in both HXO-RB44 and Y79 cells. Silencing lncRNA-UCA1 expression remarkably increased Caspase-3 expression and decreased survivin expression in HXO-RB44 cells ($P < 0.01$) (*Fig 3B*). On the contrary, lncRNA-UCA1 overexpression markedly decreased Caspase-3 expression and increased survivin expression in Y79 cells ($P < 0.01$) (*Fig 3B*). The results above suggested that lncRNA-UCA1 could inhibit cell apoptosis in RB cells.

lncRNA-UCA1 promotes cell cycle procession in HXO-RB44 and Y79 cells

As shown in *Fig 4A*, silencing lncRNA-UCA1 in HXO-RB44 cells dramatically increased the ratio of G0/G1 phase cells ($P < 0.01$), and notably reduced the ratio of S ($P < 0.01$) and G2/M phase cells ($P < 0.01$). Meanwhile, lncRNA-UCA1 overexpression in Y79 cells dramatically decreased the ratio of G0/G1 phase cells ($P < 0.01$), and markedly increased the ratio of S ($P < 0.01$) and G2/M phase cells ($P < 0.01$). This suggests that lncRNA-UCA1 could arrest RB cells at the G0/G1 phase. To further confirm the effect of lncRNA-UCA1 on RB cell cycle, we measured the expression of p16, p21 and CDK2 in both HXO-RB44 and

Y79 cells. Our results indicated that silencing lncRNA-UCA1 significantly increased p16 and p21 expression, as well as decreased CDK2 expression in HXO-RB44 cells ($P < 0.01$) (*Fig 4B*). Nevertheless, lncRNA-UCA1 overexpression dramatically decreased p16 and p21 expression, and increased CDK2 expression in Y79 cells ($P < 0.01$) (*Fig 4B*). All the results indicated that lncRNA-UCA1 could promote cell cycle procession in RB cells.

lncRNA-UCA1 activates PI3K/Akt signaling pathway in HXO-RB44 and Y79 cells

The expression of p-PI3K, p-Akt and S6K in HXO-RB44 cells in si-UCA1 group was dramatically lower than that in BLANK and si-NC group ($P < 0.01$) (*Fig 5*). Meanwhile, the expressions of BLANK and si-NC group in Y79 cells were remarkably increased in pcDNA-UCA1 group compared with BLANK and pcDNA-Mock group ($P < 0.01$) (*Fig 5*), suggesting that lncRNA-UCA1 could activate PI3K/Akt signaling pathway in HXO-RB44 and Y79 cells.

lncRNA-UCA1 promotes cell proliferation and cell cycle procession and inhibits apoptosis via activating PI3K/Akt signaling pathway in RB

To further confirm whether lncRNA-UCA1 exerted its function on RB through activating PI3K/Akt signaling pathway, we performed a variety of experiments in Y79 cells. The results of CCK-8 showed that Y79 cells proliferation was markedly promoted in pcDNA-UCA1 group compared with BLANK and pcDNA-Mock group at 48 ($P < 0.05$), 72 ($P < 0.01$) and 96 h ($P < 0.01$) (*Fig 6A*). LY294002 markedly reversed the function of lncRNA-UCA1 on cell proliferation ($P < 0.01$) (*Fig 6A*). Colony formation assay results also confirmed that the cell clones number of Y79 cells was dramatically increased in pcDNA-UCA1 group compared with BLANK and pcDNA-Mock group ($P < 0.01$) (*Fig 6B*). Meanwhile, when compared with pcDNA-UCA1 group, the number of Y79 cells clones was notably decreased in pcDNA-UCA1 + LY294002 group ($P < 0.01$) (*Fig 6B*). As shown in *Fig 6C*, Y79 cells apoptosis was dramatically lower in pcDNA-UCA1 group than that in BLANK and pcDNA-Mock group ($P < 0.01$). When compared with pcDNA-UCA1 group, the apoptosis of Y79 cells was remarkably increased in pcDNA-UCA1 + LY294002 group ($P < 0.01$). The results of western blot showed that Caspase-3, p16 and p21 expressions of Y79 cells in pcDNA-UCA1 group were markedly lower than those in BLANK and pcDNA-Mock group ($P < 0.01$) (*Fig 6D and E*), while survivin and CDK2 expressions were significantly higher ($P < 0.01$) (*Fig 6D and E*). When compared with pcDNA-UCA1 group, Caspase-3, p16 and p21 expressions of Y79 cells were remarkably higher in pcDNA-UCA1 + LY294002 group ($P < 0.01$), while survivin and CDK2 expressions were significantly lower ($P < 0.01$) (*Fig 6D and E*). All the results indicated that lncRNA-UCA1 could promote cell proliferation and cell cycle procession, as well as inhibit apoptosis via activating PI3K/Akt pathway in RB.

Discussion

RB, the most common intraocular malignancy, is diagnosed in approximately 8000 children all over the world each year. [18] In developing countries, the survival rate of RB patients is still below 70%. [19] It is urgent to investigate new and effective molecular targets for the treatment of RB. [20, 21] In this study, we confirmed that lncRNA-UCA1 could facilitate cell proliferation and inhibit apoptosis through activating PI3K/Akt pathway in RB.

More and more evidences have indicated that lncRNAs play a carcinogenic or anticancer role in various malignant tumors, such as RB. In addition, lncRNAs are reported to be abnormally expressed in tumors. Previous researches have confirmed that the expression of lncRNA-UCA1 is upregulated in gastric cancer [16], ovarian cancer [22] and lung cancer. [23] Furthermore, Sun et al. [24] have reported that lncRNA-UCA1 is highly expressed in laryngeal squamous cell carcinoma. In the present study, we assessed the expression of lncRNA-UCA1 in both RB tissues and RB HXO-RB44 and Y79 cells, and found that lncRNA-UCA1 was highly expressed in both RB tissues and RB HXO-RB44 and Y79 cells. Furthermore, we also found lncRNA-UCA1 expression was associated with some clinical parameters of RB patients, such as tumor size, optic nerve invasion and pathologic grade. Accumulating evidences have confirmed that lncRNAs participate in the regulation of cell proliferation, apoptosis, migration and cell cycle in various diseases [7, 25]. For example, lncRNA CCAT1 could promote cell proliferation, migration and invasion, as well as suppress cell apoptosis of RB cells by modulating miR-218-5p. [26] Li et al. [27] have indicated that lncRNA 00152 could promote cell proliferation and migration, and suppress apoptosis in RB. lncRNA H19 could suppress RB cell proliferation and promote cell cycle arrest and apoptosis through counteracting miR-17-92 cluster. [28] In this study, lncRNA-UCA1 promoted cell proliferation and cell cycle procession, and inhibited cell apoptosis in RB, indicating that lncRNA-UCA1 may serve as a potential prognostic biomarker for RB.

More and more researches have suggested that PI3K/Akt signaling pathway affects various biological behaviour in the process of malignancies including cell proliferation, migration, invasion, apoptosis and cell cycle. [29] S6K is a downstream target molecule of PI3K/Akt pathway. Once PI3K/Akt pathway is activated, S6K may be activated and then regulate important process of cell biology. [17] Moreover, previous findings have indicated that lncRNAs could participate in cell biological behaviour by modulating PI3K/AKT pathway in many kinds of cancers. For instance, lncRNA-UCA1 has been reported to regulate cell cycle progression through PI3K signaling pathway in bladder carcinoma cells. [30] A study of Li et al. [17] has indicated that lncRNA-UCA1 significantly promotes cell proliferation and suppresses apoptosis by activating PI3K-Akt-mTOR signaling pathway in gastric cancer. Ma et al. [16] have suggested that lncRNA-UCA1 could promote osteosarcoma metastasis through the activation of PI3K/AKT signaling pathway. In the present study, silencing lncRNA-UCA1 significantly decreased the phosphorylation of PI3K and Akt, as well as the expression of S6k in HXO-RB44 cells, while lncRNA-UCA1 overexpression dramatically promoted the phosphorylation of PI3K and Akt, and S6k expression in Y79 cells, suggesting that lncRNA-UCA1 could activate PI3K/Akt signaling pathway in RB cells. In addition, we also confirmed that LY294002 reversed the function of lncRNA-UCA1 on RB cell proliferation, apoptosis

and cell cycle procession. All results indicated that lncRNA-UCA1 could promote cell proliferation and cell cycle procession, as well as inhibit apoptosis via activating PI3K/Akt pathway in RB.

Conclusions

In conclusion, our work confirmed that lncRNA-UCA1 expression was dramatically upregulated in both RB tissue and RB HXO-RB44 and Y79 cells. In addition, we found that lncRNA-UCA1 could promote cell proliferation and cell cycle procession, as well as inhibit apoptosis via activating PI3K/Akt pathway in RB. Our research provides an innovatively regulatory mechanism about lncRNA-UCA1 in RB and point a new way to RB treatment.

Abbreviations

retinoblastoma (RB)

Long noncoding RNAs (lncRNAs)

American Type Culture Collection (ATCC)

Declarations

Ethics approval and consent to participate: This study was conducted after obtaining local ethical committee approval of The Second People's Hospital of Jinan and written informed consent from the patients.

Consent for publication: Not applicable.

Availability of data and material: All data generated or analyzed during this study are included in this published article [and its supplementary information files].

Competing interests: The authors declare that they have no competing interests.

Funding: Not applicable.

Authors' contributions:

ZFY: Substantial contributions to conception and design, data acquisition, drafting the article, final approval of the version to be published;

ZNL: Drafting the article or critically revising it for important intellectual content.

Acknowledgements: Not applicable.

References

- [1]Mahoney MC, Burnett WS, Majerovics A and Tanenbaum H. The epidemiology of ophthalmic malignancies in New York State. *Ophthalmology* 1990; 97: 1143–1147.
- [2]Dimaras H, Dimba EA and Gallie BL. Challenging the global retinoblastoma survival disparity through a collaborative research effort. *Br J Ophthalmol* 2010; 94: 1415–1416.
- [3]Rao R and Honavar SG. Retinoblastoma. *Indian J Pediatr* 2017; 84: 937–944.
- [4]Kivela T. The epidemiological challenge of the most frequent eye cancer: retinoblastoma, an issue of birth and death. *Br J Ophthalmol* 2009; 93: 1129–1131.
- [5]Peng WX, Koirala P and Mo YY. LncRNA-mediated regulation of cell signaling in cancer. *Oncogene* 2017; 36: 5661–5667.
- [6]Wang L, Yang D, Tian R and Zhang H. NEAT1 promotes retinoblastoma progression via modulating miR-124. *J Cell Biochem* 2019;
- [7]Jiang C, Li X, Zhao H and Liu H. Long non-coding RNAs: potential new biomarkers for predicting tumor invasion and metastasis. *Mol Cancer* 2016; 15: 62.
- [8]Chen L, Dzakah EE and Shan G. Targetable long non-coding RNAs in cancer treatments. *Cancer Lett* 2018; 418: 119–124.
- [9]Shang W, Yang Y, Zhang J and Wu Q. Long noncoding RNA BDNF-AS is a potential biomarker and regulates cancer development in human retinoblastoma. *Biochem Biophys Res Commun* 2018; 497: 1142–1148.
- [10]Su S, Gao J, Wang T, Wang J, Li H and Wang Z. Long non-coding RNA BANCR regulates growth and metastasis and is associated with poor prognosis in retinoblastoma. *Tumour Biol* 2015; 36: 7205–7211.
- [11]Dong C, Liu S, Lv Y, Zhang C, Gao H, Tan L and Wang H. Long non-coding RNA HOTAIR regulates proliferation and invasion via activating Notch signalling pathway in retinoblastoma. *J Biosci* 2016; 41: 677–687.
- [12]Qi D, Wang M and Yu F. Knockdown of lncRNA-H19 inhibits cell viability, migration and invasion while promotes apoptosis via microRNA-143/RUNX2 axis in retinoblastoma. *Biomed Pharmacother* 2019; 109: 798–805.
- [13]Zhang Y, Kwok-Shing Ng P, Kucherlapati M, Chen F, Liu Y, Tsang YH, de Velasco G, Jeong KJ, Akbani R, Hadjipanayis A, Pantazi A, Bristow CA, Lee E, Mahadeshwar HS, Tang J, Zhang J, Yang L, Seth S, Lee S, Ren X, Song X, Sun H, Seidman J, Luquette LJ, Xi R, Chin L, Protopopov A, Westbrook TF, Shelley CS, Choueiri TK, Ittmann M, Van Waes C, Weinstein JN, Liang H, Henske EP, Godwin AK, Park PJ, Kucherlapati R, Scott KL, Mills GB, Kwiatkowski DJ and Creighton CJ. A Pan-Cancer Proteogenomic Atlas of PI3K/AKT/mTOR Pathway Alterations. *Cancer Cell* 2017; 31: 820–832 e823.

- [14]Zeng L, Liao Q, Zou Z, Wen Y, Wang J, Liu C, He Q, Weng N, Zeng J, Tang H, Fang R, Lei Z, Tang Z, Yang X and Cui S. Long Non-Coding RNA XLOC_006753 Promotes the Development of Multidrug Resistance in Gastric Cancer Cells Through the PI3K/AKT/mTOR Signaling Pathway. *Cell Physiol Biochem* 2018; 51: 1221–1236.
- [15]Xu S, Sui S, Zhang J, Bai N, Shi Q, Zhang G, Gao S, You Z, Zhan C, Liu F and Pang D. Downregulation of long noncoding RNA MALAT1 induces epithelial-to-mesenchymal transition via the PI3K-AKT pathway in breast cancer. *Int J Clin Exp Pathol* 2015; 8: 4881–4891.
- [16]Ma H, Su R, Feng H, Guo Y and Su G. Long noncoding RNA UCA1 promotes osteosarcoma metastasis through CREB1-mediated epithelial-mesenchymal transition and activating PI3K/AKT/mTOR pathway. *J Bone Oncol* 2019; 16: 100228.
- [17]Li C, Liang G, Yang S, Sui J, Yao W, Shen X, Zhang Y, Peng H, Hong W, Xu S, Wu W, Ye Y, Zhang Z, Zhang W, Yin L and Pu Y. Dysregulated lncRNA-UCA1 contributes to the progression of gastric cancer through regulation of the PI3K-Akt-mTOR signaling pathway. *Oncotarget* 2017; 8: 93476–93491.
- [18]Soliman SE, Racher H, Zhang C, MacDonald H and Gallie BL. Genetics and Molecular Diagnostics in Retinoblastoma—An Update. *Asia Pac J Ophthalmol (Phila)* 2017; 6: 197–207.
- [19]Fabian ID, Onadim Z, Karaa E, Duncan C, Chowdhury T, Scheimberg I, Ohnuma SI, Reddy MA and Sagoo MS. The management of retinoblastoma. *Oncogene* 2018; 37: 1551–1560.
- [20]Ma C, Shi X, Zhu Q, Li Q, Liu Y, Yao Y and Song Y. The growth arrest-specific transcript 5 (GAS5): a pivotal tumor suppressor long noncoding RNA in human cancers. *Tumour Biol* 2016; 37: 1437–1444.
- [21]Weiss M, Plass C and Gerhauser C. Role of lncRNAs in prostate cancer development and progression. *Biol Chem* 2014; 395: 1275–1290.
- [22]Zhang L, Cao X, Zhang L, Zhang X, Sheng H and Tao K. UCA1 overexpression predicts clinical outcome of patients with ovarian cancer receiving adjuvant chemotherapy. *Cancer Chemotherapy & Pharmacology* 2016; 77: 629–634.
- [23]Nie W, Ge HJ, Yang XQ, Sun X, Huang H, Tao X, Chen WS and Li B. LncRNA- UCA1 exerts oncogenic functions in non-small cell lung cancer by targeting miR-193a-3p. *Cancer Letters* 2016; 371: 99–106.
- [24]Sun S, Gong C and Yuan K. LncRNA UCA1 promotes cell proliferation, invasion and migration of laryngeal squamous cell carcinoma cells by activating Wnt/beta-catenin signaling pathway. *Exp Ther Med* 2019; 17: 1182–1189.
- [25]Gao D, Lv AE, Li HP, Han DH and Zhang YP. LncRNA MALAT-1 Elevates HMGB1 to Promote Autophagy Resulting in Inhibition of Tumor Cell Apoptosis in Multiple Myeloma. *J Cell Biochem* 2017; 118: 3341–3348.

[26]Zhang H, Zhong J, Bian Z, Fang X, Peng Y and Hu Y. Long non-coding RNA CCAT1 promotes human retinoblastoma SO-RB50 and Y79 cells through negative regulation of miR-218-5p. *Biomed Pharmacother* 2017; 87: 683–691.

[27]Li S, Wen D, Che S, Cui Z, Sun Y, Ren H and Hao J. Knockdown of long noncoding RNA 00152 (LINC00152) inhibits human retinoblastoma progression. *Onco Targets Ther* 2018; 11: 3215–3223.

[28]Zhang A, Shang W, Nie Q, Li T and Li S. Long non-coding RNA H19 suppresses retinoblastoma progression via counteracting miR-17-92 cluster. *J Cell Biochem* 2018; 119: 3497–3509.

[29]Chen S, Fisher RC, Signs S, Molina LA, Shenoy AK, Lopez MC, Baker HV, Koomen JM, Chen Y, Gittleman H, Barnholtz-Sloan J, Berg A, Appelman HD and Huang EH. Inhibition of PI3K/Akt/mTOR signaling in PI3KR2-overexpressing colon cancer stem cells reduces tumor growth due to apoptosis. *Oncotarget* 2017; 8: 50476–50488.

[30]Yang C, Li X, Wang Y, Zhao L and Chen W. Long non-coding RNA UCA1 regulated cell cycle distribution via CREB through PI3-K dependent pathway in bladder carcinoma cells. *Gene* 2012; 496: 8–16.

Tables

Table 1 Association between lncRNA-UCA1 expression and clinical parameters in RB patients

clinical parameters	n	UCA1 expression	P value
Age (years)			
≤ 2	22	2.4 ± 0.268	0.7679
> 2	20	2.4 ± 0.300	
Gender			
Male	20	2.4 ± 0.300	0.8395
Female	22	2.4 ± 0.299	
Tumor size			
≤ 10 mm	22	2.6 ± 0.100	0.0074**
> 10 mm	20	2.2 ± 0.105	
optic nerve invasion			
Yes	25	2.6 ± 0.101	0.0015**
No	17	2.1 ± 0.108	
pathologic grade			
Poorly differentiated	20	2.2 ± 0.101	0.0078**
Well differentiated	22	2.6 ± 0.102	

Figures

Figure 1

The expression of lncRNA-UCA1 in both RB tissues and RB HXO-RB44 and Y79 cells. (A) The expression of lncRNA-UCA1 in RB tissues and adjacent tissue. (B) The expression of lncRNA-UCA1 in ACBRI-181, HXO-RB44 and Y79 cells. (C) The expression of lncRNA-UCA1 in transfected HXO-RB44 and Y79 cells. **P < 0.01, vs. paracancerous tissue group (A); **P < 0.01, vs. ACBRI-181 group (B); **P < 0.01, vs. BLANK and pcDNA-Mock group; &&P < 0.01, vs. BLANK and si-NC group (C).

Figure 2

lncRNA-UCA1 promoted cell proliferation in HXO-RB44 and Y79 cells. (A) The cell proliferation of transfected HXO-RB44 and Y79 cells. (B) The cell clones number of transfected HXO-RB44 and Y79 cells. (C) The expression of PCNA in transfected HXO-RB44 and Y79 cells. **P < 0.01, vs. BLANK and pcDNA-Mock group; &&P < 0.01, vs. BLANK and si-NC group.

Figure 3

lncRNA-UCA1 inhibited cell apoptosis in HXO-RB44 and Y79 cells. (A) The apoptosis of transfected HXO-RB44 and Y79 cells. (B) The expressions of Caspase-3 and survivin in transfected HXO-RB44 and Y79 cells. ^{**} $P < 0.01$, vs. BLANK and pcDNA-Mock group; ^{&&} $P < 0.01$, vs. BLANK and si-NC group.

Figure 4

lncRNA-UCA1 promoted cell cycle procession in HXO-RB44 and Y79 cells. (A) The cell cycle distribution in transfected HXO-RB44 and Y79 cells. (B) The expressions of p16, p21 and CDK2 in transfected HXO-RB44 and Y79 cells. ** $P < 0.01$, vs. BLANK and pcDNA-Mock group; && $P < 0.01$, vs. BLANK and si-NC group.

Figure 5

lncRNA-UCA1 activated PI3K/Akt signaling pathway in HXO-RB44 and Y79 cells. $**P < 0.01$, vs. BLANK and pcDNA-Mock group; $\&\&P < 0.01$, vs. BLANK and si-NC group.

Figure 6

lncRNA-UCA1 promoted RB cell proliferation and cell cycle procession and inhibited cell apoptosis via activating PI3K/Akt pathway. (A) The cell proliferation of transfected Y79 cells. (B) The cell clone number of transfected Y79 cells. (C) The cell apoptosis of transfected Y79 cells. (D) The expressions of Caspase-3 and survivin in transfected Y79 cells. (E) The expressions of p16, p21 and CDK2 in transfected Y79 cells. $**P < 0.01$, vs. BLANK and pcDNA-Mock group; $\&\&P < 0.01$, vs. pcDNA-UCA1 group.