Laminoplasty with selective fusion at unstable segment versus laminectomy with fusion for multilevel cervical myelopathy: A retrospective comparative study

Lin Du

Henan Provincial People's Hospital, People's Hospital of Zhengzhou University

Yanzheng Gao

Department of Spine Surgery, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University

Changqing Zhao

Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine

Tangjun Zhou

Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine

Haijun Tian

Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine

Kai Zhang

Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine

Jie Zhao (✉ sh9hzhaojie@163.com)

Shanghai Jiao Tong University School of Medicine

Research article

Keywords: Myelopathy, Segmental instability, Laminectomy and fusion, Laminoplasty, Selective fusion

Posted Date: October 23rd, 2019

DOI: <https://doi.org/10.21203/rs.2.16395/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Purpose The aim of this study was to compare laminoplasty with selective fixation (LPSF) versus laminectomy with fusion (LCF) in patients with multilevel cervical myelopathy accompanied by segmental instability. **Methods** A retrospective cohort study was conducted by reviewing data from 63 patients who underwent LPSF (n=30) or LCF (n=33). Cervical alignment, range of motion (ROM), neurologic status and axial symptom severity pre-operation, 3-days after operation, and at the final follow-up (minimum 24 months) were measured and compared between groups. **Results** Postoperation, patients in the LPSF group lost $31.1 \pm 17.3\%$ of cervical lordosis and $43.2 \pm 10.9\%$ cervical ROM while patients in the LCF group lost $5.7 \pm 8.2\%$ and $67.9 \pm 15.5\%$, respectively. Both LPSF and LCF groups significantly improved neurologic status and axial symptom severity at the final follow-up with similar between-group results ($P > 0.05$). Blood loss, operation time, hospital stay, and medical cost in the LPSF group were significantly less than in the LCF group ($P < 0.05$). The incidence of negative events were statistically similar between groups. **Conclusion** According to our results, LPSF was effective in maintaining the stability of the cervical spine with less sacrifice of mobility and surgical trauma for multilevel myelopathy with segmental instability compared to LCF.

Background

Multilevel degenerative changes and ossification of the posterior longitudinal ligament (OPLL) in the cervical spine leading to myelopathy has been studied extensively. Posterior decompression, including laminoplasty and laminectomy with fusion (LCF), are common procedures for multilevel spinal cord decompression.

Multilevel myelopathy often coexists with segmental instability, a degenerative change frequently seen in the elderly population. Segmental instability is a risk factor for the progression of osteophytic bone spurs and development of myelopathy [1,2]. Laminoplasty is not an appropriate choice for this type of patient, because it can aggravate pre-existing segmental spinal instability which predisposes a patient to loss of cervical lordosis that may progress to kyphosis [3,4]. Cervical kyphosis prevents indirect decompression and is associated with delayed neurologic deterioration and axial neck pain [5]. Therefore, laminectomy and fusion are commonly recommended for this group of patients. Despite posterior instrumentation to stabilize unstable segments and prevent post-laminectomy kyphosis, it has been recognized that without the protection of the lamina, adhesion formation may occur and lead to neurologic deterioration [6]. In addition, decrease in cervical range of motion (ROM) and higher incidence rate of C5 motor palsies [7] have been reported after LCF procedures.

Laminoplasty with selective fixation at the unstable segments (LPSF) was first reported by Yu et al for the treatment of OPLL associated with segmental instability [8, 9]. This new technique has the potential to provide a stable environment for spinal cord recovery and prevents the progression of kyphosis and ossification, but also preserves the cervical ROM. However, there is a paucity of evidence-based comparisons between this promising new technology and standard LCF surgery. The current study

addresses this gap, comparing outcomes between LPSF and LCF surgery for patients with identical operative indications in the hands of a single surgeon. The purpose of the present retrospective cohort study was to compare radiological and clinical outcomes of LPSF with LCF in the treatment of patients with multilevel cervical myelopathy accompanying segmental instability.

Methods

Patient selection

We reviewed medical records and radiological data of 255 patients who were diagnosed with cervical myelopathy and underwent posterior decompression surgery at our hospital between March 2010 and March 2015 (Fig 1).

Only patients with segmental instability were included in the study, which was confirmed by radiological findings. The radiological criterion was defined as more than 3.5 mm horizontal displacement of one vertebra in relation to an adjacent vertebra or more than 11° change of segmental lordotic angle on pre-operation cervical dynamic lateral view radiography according to White and Panjabi standards [10].

Patients with cervical kyphosis alignment, a previous history of cervical spine surgery, decompression extended to C2 or T1, or loss to follow-up within 2 years after surgery were excluded from the study. The final study sample consisted of 63 patients with 30 patients in the LPSF group (Fig 2a and b) and 33 patients in the LCF group (Fig 2c and d). All patients included in the study showed symptoms such hand clumsiness, gait instability, loss of coordination, as well as spinal cord compression confirmed by magnetic resonance imaging (MRI) and computed tomography (CT) that could explain their symptoms.

Surgical technique

For all patients, posterior decompression started at C3 and terminated at C6 or C7 and surgery was performed by the senior author (J.Z.). After induction of general anaesthesia, the patient was positioned prone in a prefabricated plaster cast. The unstable segment was confirmed by radiological findings. If LPSF was planned, the open side was placed on the side with the greater degree of symptoms. A laminoplasty plate was employed to support the lamina in a decompression position (Centerpiece, Medtronic Sofamor Danek, USA). Unilateral pedicle screw fixations were performed at the unstable segments. For these, the pedicle screws were inserted at the pedicle opposite to the side of dominance of the vertebral artery, regardless of whether the side was the open or hinge side. Autologous bone grafts were placed over the unstable segments on the hinge side. If LCF was scheduled, en bloc resection of the laminae and instrumentation with pedicle screw or lateral mass screw on the decompression segments were performed. For all patients, poly-axial screws combination with 3.5 mm titanium rods were used for posterior instrumentation (Vertex, Medtronic Sofamor Danek, USA). Postoperatively, all patients were

taught to wear a Philadelphia collar for comfort and were encouraged to perform early neck muscle exercises.

Radiographic evaluation

We evaluated the cervical alignment in upright standard lateral view radiographs obtained preoperatively, 3-days after operation, and at the final follow-up. Cervical sagittal alignment was assessed by the C2-7 Cobb angle. The C2-C7 Cobb angle was defined as the angle between the line parallel to the inferior endplate of C2 and C7 (Fig 3a). Straight alignment was defined as fewer than 4° of either the kyphotic or lordotic angulations. Postoperative loss of cervical lordosis (LCL) was calculated with the following formula: $LCL (\%) = (3\text{-day C2-7 Cobb angle} - \text{final C2-7 Cobb angle}) / 3\text{-day C2-7 Cobb angle} \times 100\%$ [11].

Dynamic lateral views of the cervical spine taken before surgery and at the final follow-up were used to evaluate cervical ROM changes. We defined the cervical ROM as the sum of C2-7 Cobb angle measured on flexion-extension lateral radiographs (Fig 3b and c) [12]. Loss of ROM (%) was calculated as $(\text{preoperative ROM} - \text{postoperative ROM}) / \text{preoperative ROM} \times 100\%$. Measurements are performed using tools provided by the SYNAPSE Picture Archiving and Communication Systems. To evaluate and minimize interrater and intrarater difference, all radiologic images were evaluated 3 times by 2 of the authors (LD and CZ) independently and the mean value was used for analysis.

Clinical evaluation

Clinical evaluation was made preoperatively and at the final follow-up. Neurological status was evaluated using the Japanese Orthopaedic Association (JOA) disability scale and axial symptom severity was quantified by the Visual Analogue Scale (VAS) and Neck Disability Index (NDI) [13]. The neurological recovery rate was calculated based on the following formula: $\text{recovery rate} = (\text{postoperative JOA score} - \text{preoperative JOA score}) / (17 - \text{preoperative JOA score}) \times 100\%$ [14]. Data on surgical parameters including duration, bleeding amount, length of hospital stay, and medical costs were also collected. Postoperative complications, including wound infection, instrumentation failure, cerebrospinal fluid (CSF) leak and nerve root palsy were recorded in detail.

Statistical analysis

All continuous variables are reported as the mean $\pm$ standard deviation (SD) and categorical variables as frequencies and percentages. The Fisher's exact test or chi-square test was used for comparison of categorical variables. For comparison of continuous variables between groups, we used independent sample t test if the assumptions of equal variance and the Gaussian distribution were met. Otherwise, a

nonparametric Mann-Whitney test was used. To determine statistical significance of changes between preoperative and follow-up parameters in each group, we used paired t tests. Statistical analyses were performed using SPSS version 22.0 for Windows (SPSS, Inc., Chicago, IL, USA). p values < 0.05 based on a 2-sided hypothesis test were considered significant.

Results

Baseline results

The general characteristics of patients are shown in Table 1. Of the 63 patients included in this study, the mean follow-up period was 29.0 ± 3.0 months in the LPSF group and 30.1 ± 3.7 in the LCF group. No significant difference was detected among the groups in baseline demographic including sex ($p=0.597$), age ($p=0.073$), symptomatic duration ($p=0.085$), follow-up period ($p=0.240$), BMI ($p=0.189$), Diabetes mellitus ($p=0.614$), smoking ($p=0.182$), and diagnosis ($p=0.556$) (Table 1).

Radiological analysis

The mean pre-operation C2-7 Cobb angle was $17.5 \pm 7.7^\circ$ in the LCF group and $17.4 \pm 8.0^\circ$ in the LPSF group. No patients with kyphosis cervical alignment were included in either group and the difference was not statically significant between the groups. After operation, both groups showed a slight increase in C2-7 Cobb angle. At the final follow-up the cervical alignment decreased by 5.7% to $18.4 \pm 6.9^\circ$ in the LCF group and by 31.1% to $13.4 \pm 7.7^\circ$ in the LPSF group. The change in sagittal alignment between the LCF and LPSF groups was statically different at the final follow-up ($P < 0.001$) (Table 2).

Preoperatively, the cervical ROM of the LPSF and LCF was $32.7 \pm 8.8^\circ$ and $33.7 \pm 8.4^\circ$, respectively. At the final follow-up, both groups showed a significant reduction in ROM, however the magnitude of reduction was much larger in the LCF group $67.9 \pm 15.5\%$ than in the LPSF group $43.2 \pm 10.9\%$ ($p < 0.001$) (Table 2).

Clinical analysis

There were no significant differences in preoperative self-reported clinical data between the 2 groups. Both groups (LPSF and LCF) showed significant improvement in their JOA, NDI and VAS scores postoperatively. The JOA score improved from 8.9 ± 2.3 (preoperative) to 14.1 ± 2.5 (postoperative) in the LPSF group and from 9.3 ± 2.5 (preoperative) to 14.5 ± 2.0 (postoperative) in the LCF group. The recovery rates of the JOA scores at the final follow-up was $66.8 \pm 25.7\%$ in the LPSF group and $66.1 \pm 26.9\%$ in the LCF group. No significant difference was detected in the final JOA score and JOA recovery rate between the 2 groups. Comparison of VAS and NDI score at the final follow-up showed that axial pain was not significantly different between the 2 groups (VAS: $p=0.722$, NDI: $p=0.432$) (Table 3).

In the LCF compared with LPSF group, there were longer operating times, greater amount of blood loss, longer hospital stays, and higher medical costs, and the differences were statistically significant. There were 3 superficial incision infections in the LCF group and 2 in the LPSF group but the difference was not statistically significant. There was no significant difference in the incidence of CSF leakage either. No instrumentation failure or reoperation was reported in this group of patients (Table 3).

Discussion

The purpose of the present study was to compare radiological and clinical outcomes of LPSF with LCF in the treatment of patients with multilevel cervical myelopathy accompanying segmental instability. Our results suggest that LPSF was effective in maintaining the stability of the cervical spine with less sacrifice of mobility and surgical trauma for multilevel myelopathy with segmental instability compared to LCF.

Well maintained cervical alignment is needed for indirect spinal cord decompression. It is intuitive that loss of cervical lordosis is especially relevant to the LPSF group but not to the LCF group. This is manifested within the present study where the cervical lordosis was well maintained in the LCF group. Meanwhile, in the LPSF group there was a 31.1% decreased of cervical lordosis and 5 patients developed straight alignment at the final follow-up. This result is similar to previous reports of cervical lordosis loss after laminoplasty surgery alone [4]. Considering it is possible that LPSF may result in loss of lordosis and even progression to kyphosis, LPSF should only be recommended for patients with a well-aligned cervical spine.

Restricted cervical motion after operation is one of the patient concerns before cervical spinal operations. Preserving cervical movement is an advantage of laminoplasty over laminectomy and fusion. With strong metal instrumentation, LCF is a powerful tool to correct mal-aligned cervical spines and maintain cervical lordosis, but much of the cervical spine mobility is sacrificed. There is also a reduction of cervical ROM after laminoplasty alone, assumed to be caused by spontaneous laminar fusion [15]. Therefore, it is meaningful to investigate the effect of additional selective fusion after laminoplasty and its effect on cervical ROM. In this study, we found both groups experienced a significant decrease in ROM, but the decrease was considerably less in the LPSF than in the LCF group. This result indicates that LPSF provides the benefit of preserving physiological cervical motion.

Preoperative cervical kyphosis is not suitable for cervical laminoplasty and loss of cervical lordosis is correlated with delayed neurological function deterioration [16]. In this study, although LPSF was less powerful than LCF in preventing loss of cervical lordosis, this effect did not result in inferior neurological recovery. We found that both LPSF and LCF significantly improved the JOA score and no significant difference was detected in the recovery rates between the 2 groups. These results were similar to previous reports on the comparison between LPSF and LCF [17].

Axial symptoms, including neck and shoulder pain and shoulder muscle spasm, occur frequently in patients with cervical spondylosis and have a significant negative impact on quality of life [18]. The axial

symptoms are generally considered to originate from degenerated intervertebral discs or facet joints and are reported to worsen or newly emerge after laminoplasty [19]. Segmental instability is a significant risk factor for neck pain after laminoplasty [20, 21] and postoperative cervical ROM restriction could alleviate segmental instability and concomitant axial symptoms [21, 22]. In this study, both LPSF and LCF patients showed significant improvement in VAS and NDI scores at the final follow-up and the improvement between the 2 groups was similar. These findings indicate that selective fusion is effective for relieving axial symptoms after laminoplasty due to stabilization at the unstable segments and/or the restriction of cervical ROM after LPSF.

Extensive posterior structure resection and instrumentation will increase surgical time, blood loss and medical expenditure. As shown in this study, LPSF resulted in less surgical time and blood loss, indicating a relatively less invasive technique. Because implant costs are a big part of medical expenditure for patients in China, LPSF was more cost effective than LCF. We noticed shorten hospital stays may have also decreased the expenditure of patients who underwent LPSF surgery.

LPSF preserved the lamina, which may restrict excessive spinal cord posterior drift at the level of C5 and decreased mechanical tethering of C5 nerve root [23]. However, this study did not show a lower incidence of C5 nerve root palsies in the LPSF group. wound infection, instrumentation failure, and CSF leakage were similar between the groups. These data shows that LPSF is at least as safe as traditional LCF surgery.

There are a few limitations to consider in the present study. We acknowledge the grouping was not prospectively randomized and controlled. Nearly all the LCFs were performed in the first two years, and LPSFs in the last four years of the study period. This variation was based on new techniques for less invasive and effective treatment of patients, but could have introduced bias. However, the demographic and baseline outcomes between the 2 groups were similar. We believe that the well-matched grouping was the major strength of this study. Considering the limitation of the current study, the efficiency of LPSF needs to be determined by further long-term clinical observation.

Conclusions

In conclusion, our results suggest that LPSF produces comparable clinical outcomes with LCF surgery in the treatment of patients with OPLL accompanied by segmental instability. LPSF was found to be superior to LCF in preserving cervical ROM and decreasing surgical trauma and medical cost.

Abbreviations

CSF: Cerebrospinal fluid; CT: Computer tomography; JOA: Japanese orthopaedic association; LCF: Laminectomy with fusion; LCL: Loss of cervical lordosis; LPSF: Laminoplasty with selective fixation; MRI: Magnetic resonance imaging; NDI: Neck disability index; OPLL: Ossification of the posterior longitudinal ligament; ROM: Range of motion; VAS: Visual analogue scale.

Declarations

Acknowledgements

We would like to thank Editage (www.editage.cn) for English language editing.

Authors' contributions

JZ performed all the surgery, and designed this study. CQZ are the main assistants in operation, and were major contributor in data collection. HJT and KZ contributed to design of the study, obtained ethical permission, performed data analysis, interpretation of results. LD and YZG wrote the main manuscript text and TJZ prepared the figures. All authors reviewed the manuscript and approved the final manuscript.

Funding

This study was partly supported by the National Natural Science Foundation of China (grant no. 81071503, 81572168, 81871790); 2016 Henan Science and Technology Research Project (162102310018); 2018 National and Provincial Co-Construction Project of Henan Provincial Medical Science and Technology Research (SBGJ2018076) and Henan Provincial Zhongyuan Thousand Talents Plan.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

This research involving human data have been performed in accordance with

the Declaration of Helsinki and have been approved by The Ethics Committee of Clinical Medicine Research, Shanghai Ninth People's Hospital. Because of its retrospective design, this study was exempt from obtaining informed consent for patients and all of the clinical and radiological data were collected and analysed anonymously.

Consent for publication

Written, informed consent was obtained for participants, for the publication of identifying images or other personal or clinical details of participants that compromise anonymity.

Competing interests

The authors declare that they have non-financial competing interests.

Author details

¹Department of Spine Surgery, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, 7 Weiwu Road, Zhengzhou, Henan, 450000, People's Republic of China.

²Shanghai Key Laboratory of Orthopedic Implants, Department of Orthopedics, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, 639 Zhizaoju Road, Shanghai, 200011, People's Republic of China.

References

- [1] Iwasaki K, Furukawa KI, Tanno M, et al. Uni-axial cyclic stretch induces Cbfa1 expression in spinal ligament cells derived from the patients with ossification of the posterior longitudinal ligament. *Calcif Tissue Int* . 2004;74:448-457.
- [2] Azuma Y, Kato Y, Taguchi T. Etiology of cervical myelopathy induced by ossification of the posterior longitudinal ligament: determining the responsible level of OPLL myelopathy by correlating static compression and dynamic factors. *J Spinal Disord Tech*. 2010;23:166-169.
- [3] van Geest S, de Vormer AM, Arts MP, et al. Long-term follow-up of clinical and radiological outcome after cervical laminectomy. *Eur Spine J*. 2015;24:S229-S235.
- [4] Cao J, Zhang J, Yang D, et al. Multivariate analysis of factors associated with kyphotic deformity after laminoplasty in cervical spondylotic myelopathy patients without preoperative kyphotic alignment. *Sci Rep*. 2017;7: 43443.
- [5] Tang JA, Scheer JK, Smith JS, et al. The impact of standing regional cervical sagittal alignment on outcomes in posterior cervical fusion surgery. *Neurosurgery*. 2015;76: S14-S21
- [6] Yonenobu K, Hosono N, Iwasaki M, et al. Neurologic complications of surgery for cervical compression myelopathy. *Spine (Phila Pa 1976)*. 1991;16:1277-1282.
- [7] Dai L, Ni B, Yuan W, et al. Radiculopathy after laminectomy for cervical compression myelopathy. *J Bone Joint Surg Br*. 1998;80:846-849.
- [8] Chen Y, Wang X, Chen D, et al. Posterior Hybrid Technique for Ossification of the Posterior Longitudinal Ligament Associated With Segmental Instability in the Cervical Spine. *J Spinal Disord Tech*. 2014;27:240-244.
- [9] Chen Y, Chen D, Wang X, et al. Significance of segmental instability in cervical ossification of the posterior longitudinal ligament and treated by a posterior hybrid technique. *Arch Orthop Trauma Surg*. 2013;133:171-177.

- [10] Vernon H, Mior S. The Neck Disability Index: a study of reliability and validity. *J Manipulative Physiol Ther.* 1991;14(7):409-415.
- [11] Kim TH, Lee SY, Kim YC, et al. T1 slope as a predictor of kyphotic alignment change after laminoplasty in patients with cervical myelopathy. *Spine (Phila Pa 1976).* 2013;38: E992-E997.
- [12] Zhang JT, Li JQ, Niu RJ, et al. Predictors of cervical lordosis loss after laminoplasty in patients with cervical spondylotic myelopathy. *Eur Spine J.* 2017;26:1205-1210.
- [13] Chang H, Kim C, Choi BW. Selective laminectomy for cervical spondylotic myelopathy a comparative analysis with laminoplasty technique. *Arch Orthop Trauma Surg.* 2017;137: 611-616.
- [14] Su N, Fei Q, Wang BQ, et al. long-term outcomes and prognostic analysis of modified open-door laminoplasty with lateral mass screw fusion in treatment of cervical spondylotic myelopathy. *Ther Clin Risk Manag.* 2016;12:1329-1337.
- [15] Hyun SJ, Rhim SC, Roh SW, et al. The Time Course of Range of Motion Loss After Cervical Laminoplasty: A Prospective Study With Minimum Two-Year Follow-up. *Spine (Phila Pa 1976).* 2009;34(11):1134-1139
- [16] Du W, Wang L, Shen Y, et al. Long-term impacts of different posterior operations on curvature, neurological recovery and axial symptoms for multilevel cervical degenerative myelopathy. *Eur Spine J.* 2013;22(7):1594-1602.
- [17] Blizzard DJ, Caputo AM, Sheets CZ, et al. Laminoplasty versus laminectomy with fusion for the treatment of spondylotic cervical myelopathy: short-term follow-up. *Eur Spine J.* 2017;26(1):85-93.
- [18] Kimura A, Endo T, Inoue H, et al. Impact of axial neck pain on quality of life after laminoplasty. *Spine (Phila Pa 1976).* 2015;40(24): E1292-E1298.
- [19] Wang M, Luo XJ, Deng QX, et al. Prevalence of axial symptoms after posterior cervical decompression: a meta-analysis. *Eur Spine J.* 2016;25(7): 2302-2310.
- [20] Oichi T. Cervical anterolisthesis: A predictor of poor neurological outcomes in cervical spondylotic myelopathy patients after cervical laminoplasty. *Spine (Phila Pa 1976).* 2016;41(8): E467-E473.
- [21] Kimura A, Shiraishi Y, Inoue H, et al. Predictors of Persistent Axial Neck Pain After Cervical Laminoplasty. *Spine (Phila Pa 1976).* 2018;43(1):10-15.
- [22] Nakashima H, Kato F, Yukawa Y, et al. Comparative effectiveness of open-door laminoplasty versus French-door laminoplasty in cervical compressive myelopathy. *Spine (Phila Pa 1976).* 2014;39(8): 642-647.

[23] Sodeyama T, Goto S, Mochizuki M, et al. Effect of decompression enlargement laminoplasty for posterior shifting of the spinal cord. Spine (Phila Pa 1976). 1999;24(15):1527-1531.

Tables

Table 1. Baseline patient demographics

Variables	Group LPSF (n=30)	Group LCF (n=33)	<i>p</i> value
Gender			0.597
Male (%)	21(70.0)	20(60.6)	
Female (%)	9(30.0)	13(39.4)	
Age (years)	61.1± 7.7	64.6± 7.4	0.073
BMI (kg/m ²)	23.2±4.2	24.5±3.9	0.189
Smoker (%)	7(23.3)	13(39.4)	0.182
Diabetes mellitus	7(23.3)	6(18.2)	0.614
Symptomatic duration (months)	39.1±10.9	43.4±8.9	0.085
Follow-up period (months)	29.0±3.0	30.1±3.7	0.240
Diagnosis			0.556
CSM	19	24	
OPLL	11	9	

LPSF=laminoplasty and selective fixation at the unstable segment; LCF=laminectomy and fusion; BMI=body mass index; CSM=cervical spondylotic myelopathy; OPLL=ossification of the posterior longitudinal ligament

Table 2. Radiographic outcomes

Variables	Group LPSF (n=30)	Group LCF (n=33)	<i>p</i> value
C2-7 Cobb angle			
Preoperative (°)	17.4±8.0	17.5±7.7	0.941
3-day Postoperative (°)	18.1±8.2	19.3±6.0	0.528
Final (°)	13.4±7.7	18.4±6.9	0.013
loss of cervical lordosis (%)	31.1±17.3	5.7±8.2	<0.001
Cervical ROM			
Preoperative (°)	32.7±8.8	33.7±8.4	0.647
Final (°)	19.2±8.2	10.1±3.7	<0.001
loss of ROM (%)	43.2±10.9	67.9±15.5	<0.001

LPSF=laminoplasty and selective fixation at the unstable segment; LCF=laminectomy and fusion; ROM= range of motion

Table 3. Clinical outcomes

Variables	Group LPSF (n=30)	Group LCF (n=33)	P value
JOA score			
Preoperative	8.9±2.3	9.3±2.5	0.481
Final	14.1±2.5	14.5±2.0	0.579
Recovery rate (%)	66.8±25.7	66.1±26.9	0.968
VAS score			
Preoperative	5.0±2.2	4.2±1.9	0.148
Final	1.6±1.1	1.7±1.4	0.722
NDI score			
Preoperative	17.3±9.5	18.7±6.9	0.495
Final	10.9±5.6	11.9±5.1	0.432
Blood loss (ml)	313.8±88.4	387.5±127.8	0.011
Operating time (min)	126.3±27.1	145.4±38.6	0.028
Length of stay (days)	5.8±1.3	7.1±1.5	<0.001
Complications (n)			
Nerve root palsy	1	8	0.018
Superficial wound infection	2	3	1.000
CSF leakage	1	2	1.000
Instrumentation failure	0	0	NA
Reoperation	0	0	NA
Cost (RMB)	71.9±11.2	78.6±9.4	0.013

JOA=Japanese Orthopaedic Association; VAS= visual analogue scale; NDI = neck disability index; CSF= cerebrospinal fluid; RMB=Chinese Yuan; NA= Not Applicable

Figures

Figure 1

The flowchart of patient inclusion and exclusion.

Figure 2

Postoperative cervical radiography of patients treated with laminoplasty and fixation at the unstable segment (a and b) and laminectomy and fusion (c and d).

Figure 3

The C2-C7 Cobb (α) angle was defined as the angle between the line parallel to the inferior endplate of C2 and C7 (a). The cervical range of motion (ROM) was defined as the sum of C2-7 Cobb angles measured on flexion-extension lateral radiographs (b and c). $ROM = \beta_1 + \beta_2$.