The continuity of prime numbers can lead to even continuity(The ultimate)

Xie Ling (✉ 29997609@qq.com)

Cosmic Mathematical Physics

Research Article

Keywords: prime even continuity, Bertrand Chebyshev theorem, Ascending and descending, Extreme law

Posted Date: July 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-713902/v5>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

The continuity of prime numbers can lead to even continuity(The ultimate)

Author: Xie Ling

Abstract : n continuous prime numbers can combine a group of continuous even numbers. If an adjacent prime number is followed, the even number will continue.

For example, if we take prime number **3**, we can get even number **6**. If we follow an adjacent prime number **5**, we can get even numbers by using **3** and **5**: **6, 8** and **10**.

If a group of continuous prime numbers **3,5,7,11,... P**, we can get a group of continuous even numbers **6,8,10,12,..., 2n**. Then if an adjacent prime number q is followed, the original group of even numbers **6,8,10,12,..., 2n** will be finitely extended to **2 (n + 1)** or more adjacent even numbers. My purpose is to prove that the continuity of prime numbers will lead to even continuity as long as **2 (n + 1)** can be extended.

If the continuity of even numbers is discontinuous, it violates the Bertrand Chebyshev theorem of prime numbers.

Because there are infinitely many prime numbers: **3, 5, 7, 11,...**

We can get infinitely many continuous even numbers: **6,8,10,12,...**

Key words: prime even continuity; Bertrand Chebyshev theorem; Ascending and descending; Extreme law; Mathematical complete induction

1 Preface

The whole proof begins with the ascent

Rule 1: the minimum odd prime number is **3**

Rule 2: add two odd prime numbers.(any combination of two odd primes).

Rule 3: odd prime number can be quoted repeatedly.

Rule 4: meet the previous provisions, and all prime combination to the maximum.

Rule 5: $P_a + P_b = 2S$, and $P_b + P_a = 2S$, Delete one and leave only one.

I can quote the minimum odd prime number **3**

The results are as follows

$$3+3=6$$

5 + 1 = 6 (it is stipulated that **1** is not a prime number, which is deleted because it violates regulation 2)

$\therefore$ The unique formula: **3 + 3 = 6** .(comply with Rule 4: all prime numbers are quoted to the maximum).

According to the rules, the prime number 3 uses the maximum combination, and can only get:

$$3 + 3 = 6$$

How to make **6** continuous to **8**?

If we take the adjacent prime number P greater than 3, we can get $P-3 \geq 2$ (a)

From $3+3=6$, $\rightarrow 3+(3+2)=6+2$

If $(3+2)=P$, the proposition holds when the even number is 8

Extreme law $\rightarrow$ Assumption: $(3+2) \neq p$

$\therefore p-3 \neq 2$.

$\therefore$ (a): $\{p-3 \geq 2, p-3 \neq 2\} \rightarrow (p-3 > 2) \rightarrow (p-3 \geq 4)$

$\therefore p > 2 \times 3$

$\therefore p > 2 \times 3 > 3$

The results of Bertrand Chebyshev theorem are as follows:

$(2 \times 3) > (\text{prime number } p_1) > 3$.

$\therefore (p > 2 \times 3 > 3) \rightarrow (p > p_1 > 3) \rightarrow (P \text{ and } 3 \text{ are adjacent prime numbers})$ contradiction. Negation hypothesis.

$\therefore (3+2)=p$, $\therefore 3+p=8$.

$\therefore 6 \rightarrow 8$

The proof process uses the known prime number, and the sieve method is not used in the derivation process,

Prime number theorem and Bertrand Chebyshev theorem.

By quoting the prime number theorem, we get a finite number of continuous prime numbers, and then generate even numbers according to the requirements of this manuscript.

Use the extreme rule again ψ , Force even numbers to continue.

2 nouns and definitions, citing theorems

ϕ_1 Definition of prime number: prime number refers to the natural number with no other factors except 1 and itself in the natural number greater than 1.

ϕ_2 Extreme law : A may or may not be true. What conclusion can we get if we only prove that A is not true.

$\{A \mid A=x, A=y\}, (A=x) \rightarrow (QED)$. Take: $A \neq x$, only prove the $A=y$ conclusion.

ϕ_3 [References cited 1] Bertrand Chebyshev theorem: if the integer $n > 3$, then there is at least one prime P , which conforms to $n < p < 2n-2$. Another slightly weaker argument is: for all integers n greater than 1, there is at least one prime P , which conforms to $n < p < 2n$.

ϕ_4 In this paper, the even number generation rules are as follows

- ① Only odd primes are allowed as elements.
- ② Only two prime numbers can be added.(any combination of two odd primes).
- ③ Two prime numbers can be used repeatedly: $(3+3)$, or $(3+5)$, or $(P+P)$.
- ④ meet the previous provisions, and all prime combination to the maximum.
- ⑤ Take only one of (p_a+p_b) and (p_b+p_a) .

ϕ_5 [References cited 2] The theorem of infinite number of primes:

the n bit after each prime can always find another prime.

For example, **3** is followed by **5** and **13** is followed by **17**; There must be an adjacent prime p_i after the prime P .

ϕ_6 Here we only discuss the following cases: prime number sequence and even number sequence

$(\phi_5) \rightarrow$ Prime number sequence: **3, 5, 7, 11, 13, 17, 19, 23, ...**

Even number sequence: **6, 8, 10, 12, 14, 16, 18, 20, ...**

Explain ϕ_6 in today's words: the prime number in the prime number sequence discussed in this paper is adjacent and continuous, and the first number is **3**.

An even number in an even number sequence is contiguous and the first number is **6**.

ϕ_7 Remember: all the primes I'll talk about below refer to odd primes (excluding **2**).

Prime symbol P , different primes use $P_1, P_2, P_3, \dots, P_x$

3 Logical argument

Θ_1 Goldbach conjecture : $3 \leq \forall n \in \mathbb{N}, 2n = p_x + p_y$. set up $\Delta_1: p_x \geq p_y$

Θ_2 Theorem: the continuity of prime numbers leads to the continuity of even numbers

In mathematical language :

Known: $\{3, 5, 7, 11, 13, \dots, p_2, p_1, p_0\} \in (\text{prime})$, The next neighbor of p_1 is p_0 ,

$3 < 5 < 7 < 11 < 13 < \dots p_2 < p_1 < p_0$,

$\{6, 8, 10, 12, 14, 16, \dots, 2n\} \in (\text{continuous even number})$.

If : $\{3, 5, 7, 11, 13, \dots, p_2, p_1\} \rightarrow \{6, 8, 10, 12, 14, 16, \dots, 2n\}$.

inevitable: $\{3, 5, 7, 11, 13, \dots, p_2, p_1, p_0\} \rightarrow \{6, 8, 10, 12, 14, 16, \dots, 2n, 2(n+1)\}$.

prove:

Humans use computers to calculate a finite number of even numbers: $\{6, 8, 10, 12, 14, \dots, 2n\}$

every even number satisfies Θ_1 .

The computer process is finite $\{6, 8, 10, 12, 14, \dots, 2n\}$

It is not logically proved that any even number greater than **4** satisfies Θ_1 .

$(\phi_5 + \phi_6)$ Take odd prime sequence: **3, 5, 7, 11, 13, 17, 19, 23, ...**

Take the minimum prime number **3** from the front of the prime sequence,

A_1 $\{\phi_1 + \phi_4 + \{3\}\} \rightarrow$:

$\{3+3=6\}$ It is recorded as A_1

$\rightarrow 6$

According to the rule, 3 can only get $\{3+3=6\}$

Nonexistence: $5+p=6$

【 Because 1 in $5+1=6$ is not defined as a prime number. If 1 is defined as a prime number, this paper will come to the same conclusion 】

Note: $5 \notin \{3\}$.

Prime number 3, limit is used according to ϕ_4 , cannot be: $6 \rightarrow 8$.

$\{\phi_1 + \phi_4 + \{3\}\} \rightarrow$: Quoting prime number 3 can only get even number 6.

If you want to: $6 \rightarrow 8$, you must add an adjacent prime number 5.

A_2 $\{\phi_1 + \phi_4 + \{3,5\}\} \rightarrow$:

$3+3=6$

$3+5=8$

$5+5=3+7=10$ $\therefore \phi_4 \{(5,5),(3,7)\} \in 10$ $\therefore \{\phi_1 + \phi_4 + \{3,5,7\}\} \rightarrow$

$7+5=12$

$7+7=11+3=14$ $\therefore \phi_4 \{(7,7),(3,11)\} \in 14$ $\therefore \{\phi_1 + \phi_4 + \{3,5,7,11\}\} \rightarrow$

$11+5=13+3=16$ $\therefore \phi_4 \{(11,5),(13,3)\} \in 16$ $\therefore \{\phi_1 + \phi_4 + \{3,5,7,11,13\}\} \rightarrow$

$11+7=13+5=18$.

$13+7=17+3=20$. There is a new prime number 17 in continuity.

Note: $\{(13,7),(17,3)\} \in 20$. even numbers $\rightarrow :6,8,10,12,14,16,18,20$.

Prime numbers are continuous, and there is a new prime number 17.

even numbers $\rightarrow :6,8,10,12,14,16,18,20$.

Note the key point: A_1 broken, increase the adjacent prime number 5 to have A_2

From $A_1 \rightarrow A_2 \rightarrow$ is it always infinite? Or will it stop?

Here's the wonderful thing:

(analysis I) :

Always Unlimited: $A_1 \rightarrow A_2 \rightarrow \dots$

There are: $(\phi_5 + \phi_6) \Rightarrow \{ 3, 5, 7, 11, 13, 17, 19, 23, \dots$

Get: 6, 8, 10, 12, 14, 16, 20, 22, ...

Conclusion : Θ_2 (QED).

$(\phi_2) \rightarrow$ Stop at A_n , cannot continue.

(analysis II): Stop at A_n , not to be continued.

Stop at A_n : $A_1 \rightarrow A_2 \rightarrow \dots \rightarrow A_n$

There are: 3, 5, 7, 11, 13, 17, 19, 23, ..., p_2 , p_1

Get: 6, 8, 10, 12, 14, 16, 20, 22, ..., $2(n-1)$, $2n$

$$\Theta_3 : \{\phi_1 + \phi_4 + \{3, 5, 7, 11, \dots, p_2, p_1\}\} \Rightarrow \{6, 8, 10, 12, \dots, 2(n-1), 2n\}$$

$$\Theta_4 : \{\phi_1 + \phi_4 + \{3, 5, 7, 11, \dots, p_2, p_1\}\} \neq \Rightarrow 2(n+1)$$

(1): { Let: prime p satisfy: $p \notin \{3, 5, 7, 11, \dots, p_2, p_1\}$, $p \in \forall$. Θ_4 : $p \notin A_n$, $p_1 < p$. }

Take the prime number that is greater than p_1 and adjacent to p_1 as p_0 ,

$$\therefore (1) \rightarrow \{p_0 + 3 \neq 2n, p_0 + 3 \leq 2n\}$$

$$\therefore p_0 + 3 > 2n$$

$$\therefore p_0 + 2 \geq 2n \quad \because (\text{odd}) \neq (\text{even})$$

$$\therefore p_0 + 2 > 2n \rightarrow p_0 + 1 \geq 2n \quad (W)$$

Starting from (Analysis II)

The principle of mathematical complete induction: it is correct in the front, until A_n .

Take the continuous prime number (3, 5, 7, 11, ..., p_2 , p_1) from small to large.

$$A_n \quad \{\phi_1 + \phi_4 + \{3, 5, 7, 11, \dots, p_2, p_1\}\} \rightarrow :$$

$$\{3+3=6$$

$$5+3=8$$

$$7+3=5+5=10. \text{ set up: } (7+3=5+5) \text{ sequence: } 7 > 5$$

$$7+5=12$$

$$11+3=7+7=14. \text{ set up } \Delta_2: (11+3=7+7) \text{ sequence: } 11 > 7$$

$$13+3=11+5=16. \text{ set up } \Delta_2: (13+3=11+5) \text{ sequence: } 13 > 11$$

$$13+5=11+7=18. \text{ set up } \Delta_2: (13+5=11+7) \text{ sequence: } 13 > 11$$

$$17+3=13+7=20. \text{ set up } \Delta_2: (17+3=13+7) \text{ sequence: } 17 > 13$$

$$19+3=17+5=11+11=22. \text{ set up } \Delta_2: (19+3=17+5=11+11) \text{ sequence: } 19 > 17 > 11$$

$$19+5=17+7=13+11=24. \text{ set up } \Delta_2: (19+5=17+7=13+11) \text{ sequence: } 19 > 17 > 13$$

.....

$$p_e + p_f = 2(n-2)$$

$$p_c + p_d = 2(n-1)$$

$$p_1 + p_b = 2n \quad \text{It is recorded as } A_n$$

$$\rightarrow : 6, 8, 10, 12, 14, 16, 18, \dots, 2(n-1), 2n.$$

See A_n prime numbers of the same color: $3 \rightarrow 5 \rightarrow 7 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow 19 \dots \rightarrow p_1$

α line: $3 \rightarrow 5 \rightarrow 7 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow 19 \rightarrow \dots \rightarrow p_1$

Record it as continuous rise: $3 \rightarrow 5 \rightarrow 7$ and $13 \rightarrow 17 \rightarrow 19$. Mold record: $P \rightarrow P + 2a$

Record as continuous platform: $7 \rightarrow 7$. And: $13 \rightarrow 13$, Mold record: $P \rightarrow P$.

Confirm: ($\rightarrow$) superior subordinate relationship, (lower) $\rightarrow$ (upper).

From what we know now (α) The line is: (lower) $\leq$ (upper).

See: $3 \rightarrow 5$ is: (lower) $<$ (upper). $7 \rightarrow 7$ is: (lower) = (upper).

What man knows: in A_n α When we reach the last p_1 , all of them conform to:

(lower) $\leq$ (upper).

α Route (limited) ascending and descending in turn:

$\therefore A_1 \rightarrow A_n$ is a finite number: $3 \rightarrow 5 \rightarrow 7 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow 19 \dots \rightarrow p_1$

$\therefore A_1 \leftarrow A_n$ is a finite number: $3 \leftarrow 5 \leftarrow 7 \leftarrow 7 \leftarrow 11 \leftarrow 13 \leftarrow 13 \leftarrow 17 \leftarrow 19 \leftarrow 19 \dots \leftarrow p_1$

Regulation (flat form): $p_x \rightarrow p_x$

A_n is simplified as B_n .

$$\{3+3=6$$

$$5+3=8$$

$$7+3=10$$

$$7+5=12$$

$$11+3=14$$

$$13+3=16$$

$$13+5=18$$

$$17+3=20$$

$$19+3=22$$

$$19+5=24$$

$$23+3=26$$

.....

$$p_e + p_f = 2(n-2)$$

$$p_c + p_d = 2(n-1)$$

$$p_a + p_b = 2n \quad \text{It is recorded as } B_n$$

Add the number to B_n and change to: C_n

$$\{3+2(n-2)+3=2(n+1)$$

$$5+2(n-3)+3=2(n+1)$$

$$7+2(n+4)+3=2(n+1)$$

$$7+2(n+5)+5=2(n+1)$$

$$11+2(n-6)+3=2(n+1)$$

.....

$$p_e+6+p_f=2(n+1)$$

$$p_c+4+p_d=2(n+1)$$

$$p_1+2+p_b=2(n+1)\} \quad \text{It is recorded as } C_n$$

$$\beta \text{ line: } 2 \rightarrow 4 \rightarrow 6 \rightarrow 8 \rightarrow \dots \rightarrow 2(n-6) \rightarrow 2(n-5) \rightarrow 2(n-4) \rightarrow 2(n-3) \rightarrow 2(n-2)$$

$$\alpha \text{ line: } 3 \rightarrow 5 \rightarrow 7 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow 19 \rightarrow \dots \rightarrow p_1$$

α line: Any two adjacent prime numbers satisfy: (lower) $\leq$ (upper).

$$\alpha \text{ line is simplified to } \kappa \text{ line: } 3 \rightarrow 5 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow \dots \rightarrow p_2 \rightarrow p_1$$

α line has (T station), (T station): $7 \rightarrow 7, 13 \rightarrow 13, p \rightarrow p$.

According to (κ line), we get:

$$\{p_1-p_2=2a_1$$

$$p_2-p_3=2a_2$$

$$p_3-p_4=2a_3$$

.....

$$7-5=2a_n$$

$$5-3=2a_{(n+1)}\} \quad \text{is recorded as: } D_n$$

Theorem(ω_1) :

Known:

$$\{ A_n, B_n, C_n, D_n, \phi_2, \quad \text{The first prime after } p_1 \text{ is } p_0, p_0 > p_1,$$

$$\beta \text{ line: } 2 \rightarrow 4 \rightarrow 6 \rightarrow 8 \rightarrow \dots \rightarrow 2(n-6) \rightarrow 2(n-5) \rightarrow 2(n-4) \rightarrow 2(n-3) \rightarrow 2(n-2)$$

$$\alpha \text{ line: } 3 \rightarrow 5 \rightarrow 7 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow 19 \rightarrow \dots \rightarrow p_1$$

$$\kappa \text{ line: } 3 \rightarrow 5 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow \dots \rightarrow p_2 \rightarrow p_1 \}$$

There must be: $p_0 - p_1 > 2(n-2)$

Proof:

$\therefore \alpha$ line: Any two adjacent prime numbers satisfy: (lower) $\leq$ (upper).

$$\therefore p_0 > p_1$$

$$\therefore p_0 - p_1 \geq 2$$

$$(p_0 - p_1 \geq 2) \rightarrow \{p_0 - p_1 = 2, p_0 - p_1 > 2\}. \quad (2)$$

$$\text{Take: } p_0 - p_1 = 2 \quad (3)$$

$$\text{Substituting (3) into } C_n: p_1 + 2 + p_b = 2(n+1)$$

$$\text{The result is: } p_0 + p_b = 2(n+1)$$

$$\therefore \Theta_2 \text{ (QED).}$$

$$\{\phi_2\} \rightarrow \text{It negates (3) and obtains: } p_0 - p_1 \neq 2$$

$$\therefore (2) \rightarrow (p_0 - p_1 > 2)$$

$$\text{Get: } \{p_0 - p_1 = 4, p_0 - p_1 > 4\}. \quad (4)$$

$$p_0 - p_1 = 4 \quad (5)$$

$$\text{If } C_n \text{ and } p_1 \text{ are the same } p_c \text{ (T station), then: } p_1 = p_c$$

$$\text{In } C_n, \text{ we get the following results}$$

$$p_1 + 4 + p_d = 2(n+1)$$

$$\text{Substituting (5), we get: } p_0 + p_d = 2(n+1)$$

$$\therefore \Theta_2 \text{ (QED).}$$

$$\{\phi_2\} \text{ negates (5), The result is: } p_0 - p_1 \neq 4.$$

$$\text{With the same logic, all the results of the same (T station) are obtained: } p_0 - p_1 > 2a_1 \quad (6)$$

$$\text{If } p_c \text{ and } p_1 \text{ in } C_n \text{ are not in the same (T station).}$$

$$\text{In (6), } a_1 = 1.. \text{ Go straight to the next (T station). I'm not sure how big } a_1 \text{ is. The natural number } a_1 \text{ is to be determined.}$$

$$\text{(T station) in } C_n \text{ and } A_n,$$

$$\text{If: } p_1 - p_2 = 2a_1. \text{ No (T station): } p_1 - p_2 = 2.$$

$$A_n \text{ schematic diagram:}$$

.....

$$p_2 + p_d = 2(n-1)$$

$$p_1 + p_b = 2n \text{ It is recorded as } A_n$$

$$\text{If } p_1 - p_2 = 2a_1, \text{ there is (t station) } a_1 = 2, p_1 - p_2 = 4.$$

A_n schematic diagram:

.....

$$p_1 + p_d = 2(n-1)$$

$$p_1 + p_b = 2n \} \text{ It is recorded as } A_n$$

If $p_1 - p_2 = 2a_1$, there is (t station) $a_1=3$, $p_1 - p_2=6$.

A_n schematic diagram:

.....

$$p_1 + p_f = 2(n-2)$$

$$p_1 + p_d = 2(n-1)$$

$$p_1 + p_b = 2n \} \text{ It is recorded as } A_n$$

If: $p_1 - p_2 = 2a_1$. $p_2 - p_3 = 2a_2$. No (T station): $p_1 - p_2 = 2$, $p_2 - p_3 = 2$.

A_n schematic diagram:

.....

$$p_3 + p_f = 2(n-2)$$

$$p_2 + p_d = 2(n-1)$$

$$p_1 + p_b = 2n \} \text{ It is recorded as } A_n$$

The explanation of the schematic diagram is over.

$$\text{In } C_n : p_2 + p_s = 2n - 2a_1 - 2 \quad (7)$$

$$\therefore \text{ Adjacent to (7): } p_1 + p_v = 2n - 2a_1 \quad (8)$$

$$\text{From (6): } p_0 - p_1 = 2a_1 + 2 \quad (9)$$

$$\text{In } D_n : p_1 - p_2 = 2a_1 \quad (10)$$

$$(9) \rightarrow \{ p_0 - (2n - 2a_1 - p_v) = 2a_1 + 2 \}$$

The results show that: $p_0 + p_v = 2n + 2 \therefore \Theta_2$ (QED).

$\{ \phi_2 \} \rightarrow$ It negates (9), The results showed that $p_0 - p_1 > 2a_1 + 2$

$$\text{From (6): } p_0 - p_1 = 2a_1 + 4 \quad (11)$$

Similarly, it negates the (11) formula,

The results showed that $p_0 - p_1 > 2a_1 + 4$

Similarly, the negative (11) form,

The results are as follows: $p_0 - p_1 > 2a_1 + 2a_2 + 2$

$$\therefore D_n + (\beta \text{ line: } 2(n-2))$$

∴ Conclusion: Descending

$(p_0 > p_1) \rightarrow$

$$\left\{ \begin{array}{l} (p_0 - p_1 \geq 2) \rightarrow p_0 - p_1 \neq 2, \therefore p_0 - p_1 > 2 \\ (p_0 - p_1 \geq 4) \rightarrow p_0 - p_1 \neq 4, \therefore p_0 - p_1 > 4 \\ (p_0 - p_1 \geq 6) \rightarrow p_0 - p_1 \neq 6, \therefore p_0 - p_1 > 6 \\ \dots\dots\dots \\ (p_0 - p_1 \geq 2(n-3)) \rightarrow p_0 - p_1 \neq 2(n-3), \therefore p_0 - p_1 > 2(n-3) \\ (p_0 - p_1 \geq 2(n-2)) \rightarrow p_0 - p_1 \neq 2(n-2), \therefore p_0 - p_1 > 2(n-2) \end{array} \right\} \rightarrow p_0 - p_1 > 2(n-2)$$

By analogy, we get: $p_0 - p_1 > 2(n-2)$ Note: $C_n \rightarrow \beta$ line: $2(n-2)$

$(\omega_1)(QED).$

$$(\omega_1) \rightarrow (p_0 - p_1 > 2(n-2)) \rightarrow (p_0 - p_1 > 2n-4 = p_1 + p_b - 4)$$

$$\therefore p_0 > (2p_1 + p_b - 4)$$

$$\therefore p_0 \geq (2p_1 + p_b - 3)$$

$$\because (\text{the smallest prime in } \kappa \text{ is } 3) \rightarrow p_b \geq 3 \therefore (p_0 \geq 2p_1 + p_b - 3) \rightarrow p_0 \geq 2p_1$$

$$\because (\text{odd}) \neq (\text{even}). \therefore p_0 > 2p_1$$

It is proved that: $p_0 > 2p_1$

$$\therefore p_0 > 2p_1 > p_1$$

$$\phi_3 \rightarrow p_0 > 2p_1 > p_g > p_1$$

Get: $(p_0 > p_g > p_1) \rightarrow$ And p_0, p_1 are adjacent, contradictory.

It is proved that it is wrong to quote "Extreme law ϕ_2 " in the process.

$$\text{Get: } p_x + p_y = 2(n+1)$$

$$\therefore p_x + p_y - 2 = 2n \quad (12)$$

$$(W) + (12) : p_0 + 3 \geq p_x + p_y \quad (13)$$

$$\because (\text{the smallest prime in } \kappa \text{ is } 3)$$

$$\therefore p_y \geq 3$$

$$\therefore (13) \rightarrow p_0 \geq p_x \quad (14)$$

$$\therefore \left\{ \Theta_4 : \{ \phi_4 + \{3, 5, 7, 11, \dots, p_2, p_1\} \neq \Delta 2(n+1) = p_i + p_{ii} \} \right. \quad (15)$$

$$\therefore \{ (1) + (15) \rightarrow p_x \notin \{3,5,7,11,\dots,p_2,p_1\} \rightarrow p_x > p_1 \} \quad (16)$$

【reason:

Hypothesis: $p_x \in \{3,5,7,11,\dots,p_2,p_1\}$

$$\because p_x \geq p_y \rightarrow \{p_x, p_y\} \in \{3,5,7,11,\dots,p_2,p_1\}$$

$$\{3,5,7,11,\dots,p_2,p_1\} \rightarrow p_x + p_y = 2(n+1) \quad \text{Contradiction with (15)}$$

$$\therefore p_x \notin \{3,5,7,11,\dots,p_2,p_1\} \rightarrow p_x > p_1 \quad \text{】}$$

$$\{ (14) + (16) \} \rightarrow p_0 \geq p_x > p_1 \quad (17)$$

Because $p_0 > p_1$, and the prime number: p_0 and p_1 adjacent.

$$\{ p_0 \text{ and } p_1 \text{ adjacent.} + (17) \} \rightarrow \therefore p_x = p_0$$

It is proved that: ($p_x = p_0$), get: new (α line).

new (α line): $3 \rightarrow 5 \rightarrow 7 \rightarrow 11 \rightarrow 13 \rightarrow 17 \rightarrow 19 \rightarrow \dots \rightarrow p_1 \rightarrow p_0$

$\therefore \alpha$ line: Any two adjacent prime numbers satisfy: (lower) $\leq$ (upper).

New (α line) and (α line) are not contradictory. (lower) $\leq$ (upper).

At this time, it is completely proved that: Θ_2 (QED).

Θ_1 and Θ_2 are equivalent: Θ_2 (QED) $\rightarrow \Theta_1$ (QED).

4 Conclusion

Θ_1 and Θ_2 are equivalent: Θ_2 (QED) $\rightarrow \Theta_1$ (QED).

The new α -line logic extends the old one. Θ_1, Θ_2 (QED).

Complete the mathematical complete induction:

$$\{A_1 (\kappa \text{ line: } 3) + (\alpha \text{ line}) (\text{lower}) \leq (\text{upper})\} \Theta_1 \text{ (QED)}.$$

$$(A_n (\kappa \text{ line: } 3,5,7,\dots,p_1) + (\alpha \text{ line}) (\text{lower}) \leq (\text{upper})) \Theta_1 \text{ (QED)}.$$

$$(A_{n+1} (\kappa \text{ line: } 3,5,7,\dots,p_1,p_0) + (\alpha \text{ line}) (\text{lower}) \leq (\text{upper})) \Theta_1 \text{ (QED)}.$$

5 Statement:

1. There is no relevant data in my manuscript.
2. I cited references in my manuscript.
3. There is no conflict of interest.

6

References cited:

1 Bertrand Chebyshev theorem: if the integer $n > 3$, then there is at least one prime P , which conforms to $n < p < 2n-2$. Another slightly weaker argument is: for all integers n greater than 1, there is at least one prime P , which conforms to $n < p < 2n$.

https://www.researchgate.net/publication/228592091_A_Generalization_of_Erdos's_Proof_of_Bertrand-Chebyshev_Theorem

2 The theorem of infinite number of primes: the n bit after each prime can always find another prime.

https://www.researchgate.net/publication/266044680_THE_NUMBER_OF_PRIMES_IS_INFINITE