

Determinants of low breastfeeding self-efficacy amongst mothers of children aged less than six months old: Results from the BADUTA Study in East Java, Indonesia

Christiana R Titaley (✉ christiana_rialine@yahoo.com)

Faculty of Medicine, Pattimura University <https://orcid.org/0000-0002-3023-003X>

Michael Dibley

Sydney School of Public Health, The University of Sydney

Iwan Ariawan

Center for Health Research Universitas Indonesia

Anifatun Mu'asyaroh

Center for Health Research Universitas Indonesia

Ashraful Alam

Sydney School of Public Health, The University of Sydney

Rita Damayanti

Center for Health Research, Universitas Indonesia

Do Thanh Tran

National Institute of Nutrition

Elaine Ferguson

London School of Hygiene and Tropical Medicine

Htet Kyaw

Southeast Asian Ministers of Education Organization, Regional Center for Food and Nutrition

Mu Li

Sydney School of Public Health, The University of Sydney

Aang Sutrisna

Global Alliance for Improved Nutrition

Umi Fahmida

Southeast Asian Ministers of Education Organization, Regional Center for Food and Nutrition

Research

Keywords: breastfeeding, self-efficacy, children aged <6 months, BADUTA study, Malang District, Sidoarjo District, Indonesia

Posted Date: November 11th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-71492/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 19th, 2021. See the published version at <https://doi.org/10.1186/s13006-021-00357-5>.

Abstract

Background Despite the increasing rate of exclusive breastfeeding in Indonesia, there is still a need for supportive interventions. Breastfeeding self-efficacy of mothers is a key factor positively associated with optimum breastfeeding practices. Our analysis aims to assess the determinants of low breastfeeding self-efficacy amongst a sample of women with children aged under six months in Malang and Sidoarjo Districts, East Java, Indonesia.

Methods We used information from 1210 mothers of children aged <6 months recruited in the BADUTA study conducted in 2015-2016 in Malang and Sidoarjo Districts. The outcome variable in this analysis was mothers' self-efficacy for breastfeeding using the 14 statements in the Breastfeeding Self Efficacy-Short Form. We evaluated 17 potential predictors of breastfeeding self-efficacy, organized into six sub-groups of variables: (1) context/demographic; (2) household factors; (3) maternal characteristics; (4) child characteristics; (5) breastfeeding practices; and (6) antenatal and delivery care. Logistic regression analyses were employed to examine factors associated with mothers' self-efficacy with breastfeeding.

Results More than half of the women in this study had a low level of self-efficacy. One of the factors associated with low breastfeeding self-efficacy found in this study was problems encountered by mothers related to breastfeeding. Mothers who had problems with breastfeeding not related to illness (aOR=3.27, 95%CI: 2.45-4.36) or problems related to both illness and non-illness conditions (aOR=3.57, 95%CI: 1.37-9.33) had higher odds of low breastfeeding self-efficacy than those who did not have any problems. Compared to mothers who completed university education, there was a significantly higher odds of low breastfeeding self-efficacy in mothers who completed primary school or lower (aOR=1.88, 95%CI: 1.16-3.05); completed junior high school (aOR=2.27; 95%CI: 1.42-3.63); and completed senior high school (aOR=1.94, 95%CI: 1.29-2.91). Other significant predictors of low breastfeeding self-efficacy were mothers not exposed to any breastfeeding interventions (aOR=1.87, 95%CI: 1.09-3.22); working outside the house (aOR=1.69, 95% CI: 1.23-2.32); not obtaining any advice on breastfeeding (aOR=1.40, 95%CI: 1.08-1.82); with low knowledge of breastfeeding (aOR=1.38, 95%CI: 1.03-1.84); and delivered by Caesarean section (aOR=1.34, 95%CI: 1.05-1.70).

Conclusions Multipronged breastfeeding education programs and support are required to improve women's self-efficacy with breastfeeding. Improved access to breastfeeding counsellors, active support for mothers following caesarean delivery, and increased supporting facilities at workplaces are essential to improve self-efficacy with breastfeeding.

Introduction

The importance of breastfeeding for both babies and mothers has been widely acknowledged [1]. For infants, exclusive breastfeeding is strongly recommended in their first six months of life to provide the ideal nutrition for optimal growth and development [2]. After this period, infants should continue to receive breast milk along with appropriate complementary feeding until reaching at least two years of age

[3]. The short-term and long-term effects of breastfeeding for babies include the short-term effects of reductions in the risk of diarrhea and respiratory infections [4], the long-term effects of protection against overweight and obesity, as well as a positive effect on intelligence [5]. For mothers, breastfeeding helps to increase child spacing, and reduce the risk of mastitis, postpartum haemorrhage, depression, as well as ovarian and breast cancer [3, 6, 7].

In Indonesia, the rate of exclusive breastfeeding is increasing, and contributing to this change is the Decree of the Minister of Health of the Republic of Indonesia No.450/MENKES/SK/IV/2004 concerning exclusive breastfeeding for infants, which contains the ten steps towards successful breastfeeding [8]. Subsequently, the Indonesian Government issued Regulation No. 33 Year 2012, declaring that exclusive breastfeeding fulfilled the baby's right to get the best source of food until the age of six months [9]. However, despite the improvement in breastfeeding practices, supportive interventions are still required. Based on the last two Indonesia Demographic and Health Survey data, the national rate of exclusive breastfeeding has increased from 32.4% in 2007 [10] to 41.5 % in 2012 [11] and to 52.0% in 2017 [11]. However, the 2018 Basic Health Survey from the Ministry of Health reported the national rate of exclusive breastfeeding amongst infants aged 0-5 months was 74.5%, ranging from 72.7% in urban areas to 76.6% in rural areas [12].

There is a range of different factors associated with breastfeeding practices, including the intention to breastfeed, maternal age, maternal education, smoking status, economic status, knowledge of breastfeeding, advice from health professionals, problems encountered with breastfeeding, or child's birth weight [13]. One of the factors also reported positively associated with breastfeeding is the mothers' breastfeeding self-efficacy. Self-efficacy, an element of the social cognitive theory of Bandura [14], is a predictor of health-related behaviours [15]. Self-efficacy consists of two components: (1) the outcome expectancy or the belief that a given behaviour will produce a particular outcome, and (2) self-efficacy expectancy or an individual's conviction that they can successfully perform certain tasks or behaviours to produce the desired outcome [14, 15]. We defined breastfeeding self-efficacy as the mothers' beliefs and confidence in their ability to successfully breastfeed their infants [16]. Breastfeeding self-efficacy is an important predictor of the duration [17] as well as the exclusivity of breastfeeding a [18, 19]. Thus, the assessment of mothers' breastfeeding self-efficacy will help in identifying those women who need more support for breastfeeding during the postnatal period [17].

The Global Alliance for Improved Nutrition (GAIN) and the University of Sydney, in collaboration with the Centre for Health Research, Universitas Indonesia (CHR-UI), SEAMEO-RECFON, and the London School Hygiene and Tropical Medicine, conducted the BADUTA Study, an impact evaluation of the BADUTA Program, in 2015 to 2017 in East Java, Indonesia [20]. Our analysis used the cross-sectional surveys conducted for the BADUTA Program evaluation. It aimed to examine the factors associated with low breastfeeding self-efficacy amongst the sample of women with children aged under six months in Malang and Sidoarjo Districts, East Java, Indonesia.

Methods

Data source and study sites

This analysis used data derived from the BADUTA study conducted in 2015-2016 in Sidoarjo and Malang District of East Java, Indonesia. The Ministry of Health, Republic of Indonesia identified Malang and Sidoarjo Districts as the study sites for evaluating the BADUTA program since they represented peri-urban and rural areas of East Java Province. In both districts, we selected six sub-districts to conduct the trial. The sub-districts in Sidoarjo District were Tulangan, Wonoayu, Sidoarjo, Prambon, Taman, and Krian; and in Malang District were Dampit, Turen, Tumpang, Poncokusumo, Gondanglegi, and Jabang.

We have presented detailed information about the BADUTA study protocol elsewhere [20]. We used data for this analysis from two independent cross-sectional surveys conducted in 2015 at the beginning, and in 2017 at the end of the project. To assess breastfeeding self-efficacy amongst mothers, we only used information collected from mothers of children less than six months of age.

Background information on study sites

East Java Province is one of the provinces in Indonesia located in Java Island, and the capital is Surabaya City, the second-largest city in Indonesia. The total population of East Java is approximately 37 million people, the second-most populous province in the country [21]. Malang District, located in the centre-south region of East Java Province, has an estimated total population in 2017 of 2,576,596 people [22]. Most of the people were working as labourers or private employees (37.63%) [22]. Sidoarjo District, located north of Surabaya City, has an estimated total population in 2017 of 2,207,600 people [23].

Study design and samples of the study

We conducted an observational epidemiological study to examine factors associated with low breastfeeding self-efficacy. For our analysis, we combined the data from the baseline and endline cross-sectional surveys, for both the intervention and comparison groups in the BADUTA study.

The sampling design in this trial used a three-stage cluster sampling procedure. At the initial stage, in each of the 12 subdistricts selected, 10 villages were selected using the probability proportionate to size sampling method. Next, we selected two sub-villages from each chosen village using simple random sampling method. Finally, in each of the selected sub-villages, we conducted a mini census to list all children aged <2 years, and using the listing as a sampling frame we selected 12 children aged <2 years and their mothers using simple random sampling.

In the baseline survey of the BADUTA study, the sample size for children under two years old was 2435 children; while in the endline survey, the sample for children under two years old was 2740 [20]. For this analysis, we only used information from 1210 women with children under six months (575 from the baseline and 635 from the endline survey).

Survey instruments and field personnel

The field team carried out house-to-house interviews using pre-tested and structured questionnaires. The information collected in this study included socio-economic and demographic characteristics; infant feeding practices as well as the intention of the mother to breastfeed and self-efficacy for breastfeeding of the mothers; child morbidity, reported by mother/caregiver; as well as contact with the health system and exposure to the interventions. Information about the mothers' self-efficacy for breastfeeding was collected using the Breastfeeding Self-Efficacy Scale-Short Form questionnaire developed by Dennis [15], a 14-item instrument aimed at measuring breastfeeding confidence.

At the baseline, we established eight fieldwork teams in each district. However, in the endline study, we established ten fieldwork teams to shorten the duration of data collection. For interviews, each team consisted of one field coordinator, one assistant field coordinator, and ten enumerators. In total, there were 10 field coordinators, 130 interviewers, and 20 nurses or midwives recruited [20]. The nurses and midwives conducted blood samples collection and took anthropometric measurements.

Before data collection, all field workers attended a one-week training program to standardize the enumerators, and their coordinators, with the questionnaire, sampling methodology, as well as interview techniques. The training covered different aspects of the study, i.e., an overview of the BADUTA study, the use of CommCare application; household listing and data collection procedures; explanations of study instruments (listing forms and questionnaires); quality controls for data collection; as well as a field plan. The training program included a two-day tryout to allow all training participants to practice the household listing and interviews using CommCare application. A pre and post-test was also carried out before and after the training sessions, respectively. Enumerators with low post-test results were monitored closely by field coordinators and supervisors, particularly at the beginning of data collection, to ensure their ability and quality to conduct all fieldwork activities.

Data were collected electronically on hand-held devices using CommCare system developed by Dimagi [24]. Information was recorded on structured, error detecting forms on tablets and then dispatched directly to a server for cleaning and merging. Field supervisors and a data manager monitored the data quality regularly.

Outcome variable

The outcome variable in this analysis was mothers' self-efficacy for breastfeeding as a binary variable (low or high self-efficacy on breastfeeding). We defined breastfeeding self-efficacy as the mothers' beliefs and confidence in their ability to breastfeed their infants successfully. Information about the mothers' self-efficacy for breastfeeding was collected using the Breastfeeding Self-Efficacy Scale-Short Form [15]. For each of the 14 statements, we asked the mothers to give a score from 1 to 5 that offered a range of answer options from "strongly disagree" to "strongly agree," respectively. We added all the scores to calculate the total score. As in other studies, we based the classification of breastfeeding self-efficacy on the median of the total score [25, 26]. Previous studies supported using either the mean and or the median as the cut-off point to categorize low and high breastfeeding self-efficacy [26-28]. We classified

mothers scoring less than the median as having a low self-efficacy on breastfeeding, and those scoring equal to or above the median as having a high self-efficacy.

Potential predictors

The potential predictors were selected using the analytical framework shown in Figure 1. In total, there were 17 potential predictors of breastfeeding self-efficacy included in the analyses, categorized into six sub-groups: (1) context/demographic variables; (2) household characteristics; (3) maternal characteristics; (4) child characteristics; (5) breastfeeding characteristics; and (6) antenatal and delivery care characteristics.

In the group of contextual and intervention variables, we constructed a composite variable indicating the total number of interventions from 13 variables representing breastfeeding-related interventions in the BADUTA study. Those 13 interventions were: (1) discussing breastfeeding with cadres on a home visit during pregnancy; (2) discussing exclusive breastfeeding in pregnant women's class during pregnancy; (3) did not receive any free formula milk after delivery (part of the Baby Friendly Hospital Initiative); (4) discussing breastfeeding with a village facilitator during pregnancy; (5) watching a breastfeeding-related video shown by the village facilitator during pregnancy; (6) discussing the topic of breastfeeding in emo-demo sessions; (7) receiving mobile phone messages on early initiation of breastfeeding; (8) receiving mobile phone messages on the benefits of colostrum; (9) receiving mobile phone messages on exclusive breastfeeding; (10) receiving mobile phone messages on exclusive breastfeeding problems and how to handle them; (11) receive breastfeeding counseling by midwives during pregnancy; (12) receive breastfeeding counseling by cadres during pregnancy; and (13) watching TV commercials about breastfeeding. For each question, we scored the answers "1 (one)" if the mothers answered "yes," and scored "0 (zero)" if answered otherwise. We then summed all the scores to obtain a total intervention score. We then categorized the total intervention score for each individual into "no intervention" (total score = 0); "one intervention" (total score = 1); "two interventions" (total score = 2), and "three or more intervention" (total score is ≥ 3). Finally we calculated the total intervention score for all women from both the intervention and comparison groups included in this analysis. Our purpose was to assess the impact of any breastfeeding intervention, whether from the study interventions or routine programs on breastfeeding self-efficacy. We have documented a detailed explanation of the interventions in the BADUTA study elsewhere [20].

In household characteristics, we constructed the household wealth index variable using Principal Component Analysis (PCA) [29] from an inventory of the household's facilities and assets. These items included ownership of electricity, drinking water, toilet facility, type of toilet facility, fecal final disposal and ownership of bicycle, television, water heater, 12kg of LPG, fridge, and car. We ranked households by this index and classified them into five quintiles, i.e., poorest, poor, middle, rich, and richest categories of households.

In the breastfeeding knowledge and experience group, we developed one composite variable to represent mothers' level of knowledge about breastfeeding. We constructed this variable from five questions: (1)

the best food or liquid to be provided to children aged < 6 months; (2) the duration for exclusively breastfeeding a child; (3) the duration a child should receive breast milk; (4) the benefits of giving breast milk to children; and (5) the time a child should receive complementary feeding. For each question, score "1 (one)" was assigned to all correct answers, and "0 (zero)" for all incorrect answers. We summed all the scores to get the total knowledge score, and we calculated the median value. Two categories of knowledge were developed: (1) high level of knowledge, for those whose total knowledge score was greater or equal to the median, and (2) a low level of knowledge for those whose total knowledge score was less than the median. To test if previous experience with feeding infants influenced breastfeeding self-efficacy, we also used an indicator for previous live births as we did not specifically ask the mothers about their earlier breastfeeding experiences. For the variable of problems with breastfeeding, we categorized mothers into four groups: (1) Mothers who did not experience any problems with breastfeeding; (2) Mothers who had breastfeeding problems not related to illness; (3) Mothers who had breastfeeding problems related to illness or anatomical conditions; and (4) Mothers who had both types of problems. We categorized mothers who mentioned their breastmilk was insufficient, or they could not express it, or the infant refused breastfeeds as "breastfeeding problems not related to illness. We categorized mothers with problems due to swollen breasts/mastitis, sore nipples, or flat/embedded/large nipples as a "problem related to illness/anatomical conditions". We categorized mothers reporting both types of breastfeeding problems as "mothers who had both types of problems.

Data analysis

To examine the characteristics of all variables (outcome variables and potential predictors) used in the analysis we used contingency tables. We then applied logistic regression analyses to determine factors associated with all outcome variables using ORs (odds ratios) as the estimated measures of association. We used Stata survey commands (svyset) to adjust for the clustering from the cluster randomization. All estimates presented in this analysis considered the complex sample design.

In the first step of logistic regression, we used bivariate analyses to assess the relationship between outcome variables and their potential predictors independently. In the second step, we performed multivariate analyses using a backward elimination method to remove all variables not significantly related to the study outcome, with a significance level of 0.05. Two variables selected a priori and retained in the final model regardless of the significance level, were: (1) Period of the survey (baseline or endline) and (2) the fulfilment of the minimum requirement of four antenatal care visits by trimester (met or did not met). In the final model, we obtained the adjusted ORs (aOR) and 95% confidence intervals (95% CIs) for all variables in the model.

In multivariate analysis, we used problems of breastfeeding and the number of breastfeeding interventions as composite variables. After obtaining the final model (Model #1), we developed the second model by replacing problems of breastfeeding with each type of breastfeeding problem (Model #2). We also developed the third model by replacing the breastfeeding intervention variable with all the individual exposure to intervention indicators (Model #3). We then retained the other variables in the final

model of Model #1 in Model #2 and Model #3. We used Stata/MP software (version 13.1; Stata Corp) for all analyses.

Collaborating institutions

This study was conducted by an International Research Consortium that comprised of experienced researchers from the University of Sydney (Australia), the London School of Hygiene and Tropical Medicine (LSHTM) (United Kingdom), the Center for Health Research Universitas Indonesia (CHR-UI), the Indonesia Nutrition Foundation for Food Fortification (KFI), and the Southeast Asian Ministers of Education Organization (SEAMEO), Regional Center for Food and Nutrition (RECFON).

Results

Table 1 shows the frequency distribution of mothers with children under six months by households, mothers, child characteristics, breastfeeding experience and knowledge, antenatal and delivery care services, as well as for women with low breastfeeding self-efficacy. Approximately three-quarters of the mothers were aged 20-34 years, and the majority were housewives. Slightly more than half reported ever receiving some advice on breastfeeding. Almost 31% of mothers had problems with breastfeeding: 16.9% of mothers had problems not related to illness; 11.3% had problems related to illness or anatomical condition and 2.5% had both types of problems. Around 25% of mothers had a low level of knowledge regarding breastfeeding. The characteristics of women with low breastfeeding self-efficacy were similar to all the women in the study except more worked outside their house, more had low levels of knowledge about breastfeeding, and more had experienced problems with breastfeeding. From the six knowledge components used in this analysis, the lowest percentage of correct answers was regarding the time to start complementary feeding (71.7%), followed by the minimum duration children should receive breast milk (74.3%) (Figure 2).

Table 1. Frequency distribution of mothers with children under six months old by households, mothers, child characteristics, breastfeeding experience, and knowledge, antenatal and delivery care services as well as breastfeeding self-efficacy status, The BADUTA Study in East Java, Indonesia, 2015-2016

Variables	Total Sample		Breastfeeding Self-Efficacy ¹		
	n	%	Low (%)	High (%)	<i>p</i>
Household characteristics					
Household wealth index					
Poorest	243	20.1	54.7	45.3	<i>0.086</i>
Poor	281	23.2	54.5	45.5	
Middle	236	19.5	59.8	40.2	
Rich	268	22.2	61.9	38.1	
Richest	182	15.0	50.0	50.0	
Maternal characteristics					
Maternal age					
≤19 years	85	7.0	64.7	35.3	<i>0.280</i>
20-34 years	913	75.5	56.1	43.3	
35+ years	212	17.5	55.2	44.8	
Maternal education					
University/Academy	154	12.7	44.2	55.8	<i>0.008</i>
Completed senior high school	495	40.9	57.2	42.8	
Completed junior high school	334	27.6	60.5	39.5	
No school/incomplete primary/completed primary school	227	18.8	57.7	42.3	
Maternal occupation					
Housewife	963	79.6	54.9	45.1	<i>0.027</i>
Working outside the house	247	20.4	62.8	37.2	
Number of children still alive					
1	534	44.1	56.6	43.4	<i>0.993</i>
2	477	39.4	56.2	43.8	
3	156	12.9	57.1	42.9	
4+	43	3.6	58.1	41.9	
Previous live birth					
None	520	43.0	56.4	43.6	<i>0.911</i>
Any	690	57.0	56.7	43.3	
Antenatal, delivery care					
Minimum antenatal care visits²					
Completed (4+ visits)	858	71.6	54.6	45.4	<i>0.057</i>
Incomplete (<4 visits)	340	28.4	60.6	39.4	
Mode of delivery					
Normal	821	67.9	55.2	44.8	<i>0.168</i>

Variables	Total Sample		Breastfeeding Self-Efficacy ¹		
	n	%	Low (%)	High (%)	p
Caesarea	389	32.2	59.4	40.6	
Birth attendant					
General Practitioner/OBGYN	483	39.9	57.1	42.3	0.772
Midwife/nurse	708	58.5	55.9	44.1	
Traditional birth attendant/family/friend	19	1.6	63.2	36.8	
Child's characteristics					
Sex of the child					
Male	582	48.7	56.0	44.0	0.792
Female	613	51.3	56.8	43.2	
Birth weight according to monitoring card					
Larger than average	275	22.8	62.6	37.4	0.062
Average	831	69.0	54.5	45.5	
Smaller than average	99	8.2	54.6	45.4	
Breastfeeding experience and knowledge					
Ever received any advice on breastfeeding					
Yes	638	52.7	52.7	47.3	0.004
No	572	47.3	60.8	39.2	
Knowledge about breastfeeding					
High ³	909	75.1	53.8	46.2	0.001
Low ⁴	301	24.9	64.8	35.2	
Problems with breastfeeding					
None	838	69.3	50.8	49.2	<0.001
Not related to illness	204	16.9	76.5	23.5	
Related to illness/anatomical condition	137	11.3	56.9	43.1	
Both types of problems	30	2.5	76.7	23.3	

Note:

¹Low breastfeeding self-efficacy was mothers whose total self-efficacy score was less than the median value. ²Minimum Antenatal Care refers to the recommendation of at least four antenatal visits, i.e., once in trimester one to three, and twice in trimester three. ³High level of knowledge was mothers whose total knowledge score was greater than, or equal to the median knowledge score value. ⁴Low level of knowledge was mothers whose total knowledge score was less than the median knowledge score value

Table 2 presents the distribution of 1210 (575 from the baseline and 635 from the endline surveys) mothers of children aged less than six months who were interviewed in the BADUTA study by contextual, intervention, and breastfeeding characteristics against low self-efficacy status. Based on the interventions received, 40% received at least one intervention on breastfeeding, but less than 8% received

three or more interventions. The low percentage of mothers receiving BADUTA interventions is due to the pooled dataset from baseline and endline surveys. At the endline survey, around 33% of women in the comparison group did not receive any breastfeeding interventions, while 67% received at least one type of intervention (47% received one; 17% received two, and 4% received three or more). In the intervention group, 13% of women did not receive any breastfeeding interventions, and 87% received at least one intervention (36% received one; 24% received two, and 27% received three or more interventions).

Table 2. Frequency distribution of mothers with children under six months by contextual and intervention characteristics and breastfeeding and self-efficacy status, The BADUTA Study in East Java, Indonesia, 2015-2016

Variable	Total Sample		Breastfeeding Self-Efficacy ¹		
	n	%	Low (%)	High (%)	p
Exposure to interventions					
Exposed to intervention ²	286	23.6	49.7	50.3	0.007
Not exposed to intervention ³	924	76.4	58.7	41.3	
Number of breastfeeding interventions mothers were exposed to					
Three or more interventions	91	7.5	38.5	61.5	<0.001
Two interventions	127	10.5	48.8	51.2	
One intervention	266	22.0	56.4	43.6	
No interventions	726	60.0	60.2	39.8	
Period					
Baseline	575	47.5	60.5	39.5	0.008
Endline	635	52.5	52.9	47.1	
Ever visited by village facilitator at home					
Yes	22	1.8	36.4	63.6	0.054
Ever attended "emo-demo."					
Yes	50	4.1	44.0	56.0	0.068
Ever received SMS Bunda messages					
Yes	20	1.7	35.0	65.0	0.050
Ever visited by cadre at home					
Yes	17	1.4	29.4	70.6	0.023
Ever attended pregnancy class					
Yes	40	3.3	40.0	60.0	0.032
Roomed in with baby after delivery					
Yes	363	30.0	48.2	51.8	<0.001
Ever received counselling by a midwife					
Yes	188	15.5	52.1	47.9	0.185
Ever received counselling by a cadre					
Yes	24	2.0	37.5	62.5	0.058
Ever seen "Rumpi Sehat" TV commercials					
Yes	123	10.2	43.9	56.1	0.003

Note:

¹Low self-efficacy was mothers whose total self-efficacy score was less than the median value. ²Exposed to intervention refers to respondents living in the intervention sub-districts at the endline survey. ³Not exposed to intervention referred to all respondents from the baseline survey and those living in the control sub-districts at the endline survey.

Over half the women (56%) in the study had low breastfeeding self-efficacy. The median score of breastfeeding self-efficacy in mothers with low efficacy was 35 (SD: 6.33) and in mothers with high self-

efficacy was 43 (SD=4.31). As shown in Figure 3, there was a significant difference in the percentage of women with low breastfeeding self-efficacy related to how they were feeding their infants under six months of age ($p<0.001$). There was no significant difference between the percentage of mothers with low breastfeeding self-efficacy who exclusively or predominantly breastfed their infants. However, the percentage of mothers with low breastfeeding self-efficacy was significantly higher among women who were exclusively or predominantly breastfeeding compared with women feeding with breast milk and formula, or with breast milk and solids/semi-solids, or not breastfeeding at all (Figure 3).

Figure 4 presents the distribution of the respondents' answers to each component of the Breastfeeding Self-Efficacy Scale. Overall more than 50% of the mothers answered either agree or strongly agree to all components of the Breastfeeding Self-Efficacy Scale score. We found the highest percentage of agreement in mothers who felt able to tell when their baby finished breastfeeding (80%), and the lowest was in mothers who felt they could deal with breastfeeding being time-consuming. As shown in Table 1, low breastfeeding self-efficacy was highest amongst women with lower levels of education, of younger age, and working outside the house. Low breastfeeding efficacy declined as the number of breastfeeding interventions the women received increased.

Table 3 presents the results of the analysis of factors associated with low breastfeeding self-efficacy. One of the factors that were significantly related to low mothers' breastfeeding self-efficacy was problems encountered related to breastfeeding. There was no association between mothers who had breastfeeding problems related to illness or anatomical conditions and low breastfeeding self-efficacy (aOR=1.36, 95%CI: 0.95-1.93, $p=0.089$). However, there was a significant association between mothers who had problems not related to illness (aOR=3.27, 95% CI: 2.45-4.36, $p<0.001$), or both types of problems (aOR=3.57, 95%CI: 1.37-9.33, $p=0.010$) and low breastfeeding self-efficacy. When the 'problems with breastfeeding' variable was replaced in the final model by each type of mothers' breastfeeding problem, we found an increased odds for low breastfeeding self-efficacy for mothers with a swollen breast (aOR=3.10, 95%CI: 1.33-7.21, $p=0.009$); with flat nipples (aOR=2.75, 95%CI: 1.29-5.87, $p=0.009$); whose infants refused breastfeeds (aOR=5.45, 95%CI: 1.64-18.13, $p=0.006$); and who perceived their breastmilk is not enough (aOR=2.80, 95%CI: 2.04-3.84, $p<0.001$) (Table S1).

The level of mothers' education was also significantly associated with low breastfeeding self-efficacy. A higher odds was found in mothers who completed primary school or lower (aOR=1.88, 95%CI: 1.16-3.05, $p=0.011$); completed junior high school (aOR=2.27; 95%CI: 1.42-3.63, $p=0.001$), and completed senior high school (aOR=1.94, 95%CI: 1.29-2.91, $p=0.002$), compared to mothers who completed university education.

As expected, the likelihood of having low breastfeeding self-efficacy reduced as the number of breastfeeding interventions received by mothers increased. The odds of low breastfeeding self-efficacy significantly increased in mothers not exposed to any breastfeeding interventions (aOR=1.87, 95%CI: 1.09-3.22, $p=0.024$). In the analysis shown in the supplementary table (Table S2), we replaced the total number of interventions by an indicator of exposure for each specific intervention. It is interesting to see

that in general, there was no significant association between each intervention and self-efficacy, except for mothers visited by a cadre at home to talk about breastfeeding (aOR=0.28, 95%CI: 0.09-0.85, p=0.025).

Moreover, we found an increased odds of low breastfeeding self-efficacy in mothers working outside the house (aOR=1.69, 95%CI: 1.23-2.32, p=0.001); who had never received any breastfeeding advice (aOR=1.40, 95%CI: 1.08-1.82, p=0.013); and who had low knowledge of breastfeeding (aOR=1.38, 95%CI: 1.03-1.84, p=0.029). The odds of low breastfeeding self-efficacy was also higher in mothers who delivered by a Caesarean section (aOR=1.34, 95%CI: 1.05-1.70, p=0.017) than those with normal delivery.

Table 3. Factors associated with low breastfeeding self-efficacy amongst mothers of children under six months old, The BADUTA Study in East Java, Indonesia, 2015-2016

Variable	Univariate				Multivariate ¹				
	aOR	95% CI		<i>p</i>	aOR	95% CI		<i>p</i>	
Contextual and intervention characteristics									
Exposure to intervention									
Exposed to intervention ²	1.00								
Not exposed to intervention ³	1.44	1.13	1.83	0.003					
Number of breastfeeding interventions mothers exposed to									
Three or more interventions	1.00				1.00				
Two interventions	1.53	0.83	2.81	0.175	1.47	0.74	2.89	0.269	
One intervention	2.07	1.16	3.69	0.014	1.81	0.98	3.35	0.059	
No interventions	2.42	1.41	4.15	0.001	1.87	1.09	3.22	0.024	
Period									
Baseline	1.00				1.00				
Endline	0.73	0.55	0.97	0.032	0.89	0.61	1.28	0.517	
Household characteristics									
Household wealth index									
Poorest	1.00								
Poor	0.99	0.70	1.39	0.948					
Middle	1.23	0.86	1.75	0.257					
Rich	1.35	0.93	1.95	0.118					
Richest	0.83	0.52	1.32	0.426					
Mother's characteristics									
Maternal age									
≤19 years	1.00								
20-34 years	0.70	0.41	1.17	0.172					
35+ years	0.67	0.40	1.14	0.138					
Maternal education									
University/Academy	1.00				1.00				
Completed senior high school	1.69	1.19	2.39	0.003	1.94	1.29	2.91	0.002	
Completed junior high school	1.94	1.31	2.86	0.001	2.27	1.42	3.63	0.001	

Variable	Univariate				Multivariate ¹			
	aOR	95% CI		<i>p</i>	aOR	95% CI		<i>p</i>
No school/incomplete primary/completed primary school	1.73	1.12	2.67	<i>0.014</i>	1.88	1.16	3.05	<i>0.011</i>
Maternal occupation								
Housework	1.00				1.00			
Working outside the house	1.38	1.06	1.81	<i>0.018</i>	1.69	1.23	2.32	<i>0.001</i>
Number of children still alive								
1	1.00							
2	0.99	0.77	1.26	<i>0.904</i>				
3	1.02	0.69	1.52	<i>0.920</i>				
4+	1.07	0.55	2.07	<i>0.847</i>				
Previous live birth								
None	1.00							
Any	1.01	0.82	1.26	<i>0.906</i>				
Antenatal and delivery care								
Minimum antenatal care visits⁴								
Completed (4+ visits)	1.00				1.00			
Incomplete (<4 visits)	1.28	0.98	1.68	<i>0.073</i>	1.23	0.92	1.65	<i>0.166</i>
Mode of delivery								
Normal	1.00				1.00			
Caesarean	1.19	0.96	1.47	<i>0.108</i>	1.34	1.05	1.70	<i>0.017</i>
Birth attendant								
General practitioner/OBGYN	1.00							
Midwife/nurse	0.95	0.78	1.165	<i>0.631</i>				
Traditional birth attendant/family/friend	1.29	0.54	3.082	<i>0.572</i>				
Child's characteristics								
Sex of the child								
Male	1.00							
Female	1.03	0.82	1.30	<i>0.791</i>				

Variable	Univariate			Multivariate ¹			
	aOR	95% CI	<i>p</i>	aOR	95% CI	<i>p</i>	
Birth weight from monitoring card							
Larger than average	1.00						
Average	0.72	0.51	1.00			0.053	
Smaller than average	0.72	0.44	1.17			0.183	
Breastfeeding knowledge & experience							
Ever received any breastfeeding advice							
Yes	1.00			1.00			
No	1.40	1.09	1.78	1.40	1.08	1.82	0.013
Knowledge about breastfeeding							
High level ⁵	1.00			1.00			
Low level ⁶	1.58	1.19	2.10	1.38	1.03	1.84	0.029
Problems with breastfeeding							
None	1.00			1.00			
Not related to illness	3.14	2.34	4.23	3.27	2.45	4.36	<0.001
Related to illness/anatomical condition	1.28	0.91	1.80	1.36	0.95	1.93	0.089
Both types of problems	3.18	1.29	7.81	3.57	1.37	9.33	0.010

Note:

¹Multivariate logistic regression using the backward elimination method to select significant predictors of low breastfeeding self-efficacy. The variable for the minimum requirement of four antenatal care visits by trimester was selected a priori to be retained in the final model regardless of its significance level. ²Exposed to intervention refers to respondents living in the intervention sub-districts at the endline survey ³Not exposed to intervention referred to all respondents from the baseline survey and those living in the control sub-districts at the endline survey. ⁴Minimum antenatal care refers to the recommendation of at least four antenatal visits, i.e., once in trimester one to three, and twice in trimester three. ⁵High level of knowledge was mothers whose total knowledge score was greater than, or equal to the median knowledge score value. ⁶Low level of knowledge was mothers whose total knowledge score was less than the median knowledge score value.

Discussion

Main findings

Our study found that more than half of the women with children under six months had a low level of breastfeeding self-efficacy. It was also confirmed by the exclusive breastfeeding status of mothers, as a significantly higher percentage of mothers with high breastfeeding self-efficacy exclusive breastfed their babies than those with low breastfeeding self-efficacy. The significant predictors of low breastfeeding self-efficacy included low education level, working outside the house, never receiving any advice on breastfeeding, low knowledge about breastfeeding, and problems encountered with breastfeeding. Mothers who had a caesarean section delivery also had a low self-efficacy for breastfeeding. As expected, the odds of having low self-efficacy were lower as the number of breastfeeding interventions received increased. The findings of this study should assist decision-makers and program managers in designing and implementing supportive interventions to increase mothers' self-efficacy with breastfeeding. Increased self-efficacy will promote and accelerate the improvement of breastfeeding practices to the benefit of both mothers and their children.

The role of knowledge and education in breastfeeding self-efficacy

We confirmed the relationship between knowledge and breastfeeding self-efficacy in our study. Firstly, mothers with a low level of breastfeeding knowledge were more likely to have lower self-efficacy than those with a high level of knowledge. Secondly, we found an association between low self-efficacy with lower educational attainments. The level of education mirrors mothers' level of knowledge and awareness, not only about breastfeeding but also health in general, the possibility of having more exposure to health-related information, and comprehension about health information received. A study has reported that mothers with secondary and high school education are more likely to have a higher self-efficacy score than those graduating from university [30]. The difference was assumed due to the mother's occupation, as highly educated mothers were more likely to have a job and work outside the house. However, in our analysis, we adjusted the association between education or knowledge and breastfeeding self-efficacy for the maternal occupation to remove the confounding of the effects by occupation.

Education-based interventions and support to improve breastfeeding self-efficacy

The important role of knowledge and education found in our study indicates the importance of promoting strategies to enhance mothers' and other family members' awareness of breastfeeding. Previous literature has reported the benefits of both prenatal [31-34] and postnatal education and the support mothers received for breastfeeding [35, 36]. It is also important to consider the type, timing, setting, and frequency of education interventions [33]. A systematic review highlighted the importance of support-based initiatives during postnatal care through interaction with lactation experts [37]. The use of

combined settings (health facility and community) for health education, not solely in hospitals or community, was reported to be more beneficial than education merely in a health facility [33, 38].

Furthermore, we found that mothers who had never received any advice on breastfeeding were more likely to have lower breastfeeding self-efficacy than those who had ever received any advice. Interactive and face-to-face education, in addition to the consistent delivery of breastfeeding messages, will increase self-efficacy. Previous literature revealed that combined individual and group counselling is more effective than individual or group counselling only [38]. Multiple intervention contacts, rather than only a single contact intervention, have more favourable outcomes [33, 39]. Thereby, health workers should effectively use every contact opportunity with mothers and other family members, from antenatal to postnatal period, to improve mothers' awareness and self-efficacy with breastfeeding.

One review found that peer support interventions were also effective in promoting optimum breastfeeding [39], suggesting an opportunity to encourage health-volunteers and mothers, who have successfully breastfed, to support other mothers breastfeeding their children. This analysis found that mothers with breastfeeding problems had a lower self-efficacy than those who never experienced any problems. The problems included those not related to illness, such as mothers whose infants refused to breastfeed, and mothers who perceived their breastmilk was not enough; or problems related to illness or anatomical conditions such as swollen breasts or fat nipples. Peer support will assist first-time mothers to build their confidence in breastfeeding through experience sharing sessions. It will also help support mothers to overcome their breastfeeding challenges and problems.

Other intervention channels for providing education and support are the use of telephone contacts for discussion or counselling or production of promotional materials such as leaflets, flip charts, DVDs, or workbooks. An experimental study in Brazil found that mothers who received educational interventions, enhanced by a flip chart with illustrations of breastfeeding themes, showed increased breastfeeding self-efficacy [35]. The use of checklists, pamphlets, and audiovisual materials on breastfeeding among recently delivered mothers in hospitals also improved the mothers' self-efficacy for breastfeeding [36].

Mothers who delivered by Caesarean section in this study were more likely to have low breastfeeding self-efficacy. This finding was similar to other studies where women who had delivered by Caesarean section were less likely to breastfeed or to delay breastfeeding initiation [40, 41]. The problems related to lactogenesis due to abdominal surgery or stress response, as a result of delivery complications, contributed to increased difficulty with or early cessation of breastfeeding [40, 42]. Consequently, interventions targeting mothers, both with planned or emergency caesarean section, are required. Support and counselling programs by health workers and lactation counsellors, including postpartum home visits, might be beneficial for these women.

Previous research reported the critical role of family support to help mothers to breastfeed their children [43]. Women who received praise from other family members had higher scores of breastfeeding self-efficacy than those receiving praise from friends [44]. Thus, the involvement of family members,

particularly husbands in educational programs, for example, during prenatal classes, is vital for promoting optimal breastfeeding practices.

The odds of high self-efficacy with breastfeeding amongst women in this study increased as the number of breastfeeding interventions received by the mothers increased. Many of BADUTA interventions aimed to strengthen the existing health system, which was also available to mothers in the comparison group. Consequently, in the endline survey, we found a high percentage of women from the comparison group also received breastfeeding interventions. These breastfeeding interventions heavily relied on awareness-raising and education strategies to improve the community's knowledge and skills in breastfeeding. The combined effect of different interventions might reflect a dose-response effect of increasing odds of higher breastfeeding self-efficacy with an increasing number of breastfeeding intervention services received by mothers [45, 46].

To ensure the quality of support and educational interventions received by mothers, interventions to increase knowledge and skills of health workers, lactation counsellors, community health workers (cadres), particularly concerning their counselling skills, are essential. Trained birth attendants, whose services are still widely used by the community in some settings for mother and childcare, could be trained by health workers to promote breastfeeding amongst pregnant and recently delivered mothers.

The role of maternal occupation on breastfeeding self-efficacy

We found a low level of breastfeeding self-efficacy in women working outside their homes. Secondary analysis using nationally representative survey data for Indonesia found that a mother working outside her home was a barrier for optimal breastfeeding practices [47]. A strong reason for this could be the short duration of maternity leave in Indonesia. In the formal sector, only 1.5 months of maternity leave is given before and after delivery [48], while in the informal sector, this regulation is often not fully applied. Although the Act of the Republic of Indonesia stated that female workers still breastfeeding their children should be given appropriate opportunities to breastfeed even during working hours [48], supportive breastfeeding facilities are limited or even unavailable. Advocacy, in addition to the development of supportive policy and regulations, to ensure the availability of lactation space and breastfeeding breaks in the workplace is therefore crucial for women working outside their homes [46].

Strengths and limitations

This study has several strengths. It has a large sample size, giving adequate power to analyze the role of different predictors on breastfeeding self-efficacy. It is the first study in Indonesia to examine self-efficacy with breastfeeding in mothers of children aged less than six months. Some limitations are worth noting when interpreting the results. We based our results on the mothers' recall ability, and we did not validate the information provided by respondents. Another limitation was how mothers responded to breastfeeding self-efficacy questions will be influenced by their current feeding practices, particularly amongst those who decided not to breastfeed their infants. There are other possible determinants of mothers' self-efficacy with breastfeeding that we did not analyze because the variables were not

available in the dataset, such as the mothers' previous breastfeeding experience or the level of family support. To partially address previous breastfeeding experience, we examined the association of having had a previous live birth on the odds of low breastfeeding self-efficacy in mothers, and we found no association between them.

Conclusions

Overall, our study found maternal education, knowledge of breastfeeding, occupation, and mode of delivery as predictors of mothers' breastfeeding self-efficacy. Consequently, multipronged breastfeeding education and support are required to increase women's self-efficacy with breastfeeding that will accelerate appropriate breastfeeding practices despite the ongoing improvements in breastfeeding in Indonesia. Program managers should consider the use of combined settings, individual and group counselling with multiple contact opportunities. Effective education strategies and support programs targeting not only mothers but also other family members are required.

Additionally, efforts to ensure the availability and access to breastfeeding counsellors or peer counsellors will help mothers to increase their confidence in breastfeeding. Training of cadres and traditional birth attendants to provide necessary counselling on breastfeeding will be beneficial to provide continuous support to women. Furthermore, home visits, including those women who delivered with caesarean section, will ensure mothers have ongoing support and overcome challenges or barriers in breastfeeding. Interventions to increase the availability of supporting facilities in the workplace are also required to enhance optimum breastfeeding practices amongst women working outside their homes.

Declarations

Ethics approval and consent to participate

We obtained ethics approval from the Faculty of Public Health, Universitas Indonesia (323/H2.F10/PPM.00.02/2016) and Human Research Ethics Committee of the University of Sydney, Australia (Protocol number: 2015/115). We also secured a research clearance from the Ministry of Internal Affairs at the central level, the Office of National Unity, and from the Community Protection branch at the provincial and district levels. We took written informed consent from all respondents before the interview. All individuals included in this research are 18 years or older.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Funding

This paper is based on the evaluation of phase one (2014-2017) of the Global Alliance for Improved Nutrition (GAIN) BADUTA Program in Sidoarjo and Malang districts of East Java. The study was funded from a grant by Kingdom of the Netherlands to GAIN [activity number:400000062], who contracted a consortium led by the University of Sydney to conduct an independent impact evaluation. Responsibility for the contents and for the opinions expressed rests solely with the authors; publication does not constitute an endorsement by the Netherlands Ministries of Foreign Affairs.

Author Contributions

CRT and MJD designed the study; CRT, MJD, IA and AM performed data analyses; CRT and MJD wrote the original draft of the manuscript; AA, RD, TDT, EF, HK, LM, AA, UF provided data analysis advice; MJD, IA, AA, RD, TDT, EF, HK, LM, AA, UF reviewed the manuscript. All authors read and approved the manuscript.

Acknowledgments

The authors extend thanks to the women and children who participated in the evaluation of phase one (2014-2017) of the BADUTA program. We also thank the following organizations that contributed to the success of the BADUTA program: Directorate of Community Nutrition, Ministry of Health, Indonesia, East Java Provincial Health Office, District Health Offices of Sidoarjo and Malang, Yayasan Sayangi Tunas Cilik, Save the Children, Paramitra Foundation, and PT Holland for Water (Nazava). Also, we thank GAIN staff both past and present for their support of phase one of the BADUTA program evaluation, including Lynnette Neufeld and Alison Tumilowicz, for their contributions to the design of the evaluation. We also thank those who contributed to the program evaluation, including the enumerators, fieldwork teams, the staff, and students from local health and nutrition academies (Poltelkes Kemenkes Malang and Akademi Gizi Surabaya) for their enthusiastic support of the field activities.

References

1. Zeng C, Guo X, Long J, Kuchenbaecker KB, Droit A, Michailidou K, Ghoussaini M, Kar S, Freeman A, Hopper JL *et al*. **Identification of independent association signals and putative functional variants for breast cancer risk through fine-scale mapping of the 12p11 locus.** *Breast Cancer Res* 2016, **18**(1):64.

2. Kramer MS, Kakuma R: **The optimal duration of exclusive breastfeeding: a systematic review.** *Advances in experimental medicine and biology* 2004, **554**:63-77.
3. World Health Organization: **Infant and young child feeding: Model Chapter for textbooks for medical students and allied health professionals.** Geneva: World Health Organization; 2009.
4. Horta BL, Victora CG: **Short-term effects of breastfeeding: systematic review on the benefits of breastfeeding on diarrhoea and pneumonia mortality** In.: World Health Organization; 2013.
5. Horta BL, Victora CG: **Long-term effects of breastfeeding: a systematic review.** In.: World Health Organization; 2013.
6. Gartner LM, Morton J, Lawrence RA, Naylor AJ, O'Hare D, Schanler RJ, Eidelman AI: **Breastfeeding and the use of human milk.** *Pediatrics* 2005, **115**(2):496-506.
7. Ystrom E: **Breastfeeding cessation and symptoms of anxiety and depression: a longitudinal cohort study.** *BMC Pregnancy and Childbirth* 2012, **12**(1):36.
8. Ministry of Health Republic of Indonesia: **Ministry of Health Republic of Indonesia Decree No. 450/MENKES/SK/IV/2004 on Exclusive Breastfeeding to Infants in Indonesia [Pemberian Air Susu Ibu (ASI) secara eksklusif pada bayi di Indonesia].** In. Edited by Indonesia MoHRo; 2004.
9. President of the Republic of Indonesia: **Government Regulation No 33 Year 2012 on Exclusive Breastfeeding [Pemberian Air Susu Ibu Eksklusif].** 2012.
10. Statistics Indonesia, National Family Planning Coordinating Board, Ministry of Health Republic of Indonesia, ORC Macro: **Indonesia Demographic and Health Survey 2007.** In. Calverton, Maryland: BPS and ORC Macro; 2008.
11. Statistics Indonesia, National Family Planning Coordinating Board, Ministry of Health Republic of Indonesia, Macro O: **Indonesia Demographic and Health Survey 2017.** In. Calverton, Maryland: BPS and ORC Macro; 2017.
12. National Institute of Health Research and Development MoH, Republic of Indonesia: **Basic Health Research 2018: National Report 2018.** In. Jakarta: National Institute of Health Research and Development, Ministry of Health, Republic of Indonesia; 2018.
13. Forster DA, McLachlan HL, Lumley J: **Factors associated with breastfeeding at six months postpartum in a group of Australian women.** *Int Breastfeed J* 2006, **1**:18.
14. Bandura A: **Self-efficacy: toward a unifying theory of behavioral change.** *Psychological review* 1977, **84**(2):191-215.
15. Dennis CL: **The breastfeeding self-efficacy scale: psychometric assessment of the short form.** *J Obstet Gynecol Neonatal Nurs* 2003, **32**(6):734-744.
16. Dennis CL: **Theoretical underpinnings of breastfeeding confidence: a self-efficacy framework.** *Journal of human lactation : official journal of International Lactation Consultant Association* 1999, **15**(3):195-201.
17. Baghurst P, Pincombe J, Peat B, Henderson A, Reddin E, Antoniou G: **Breastfeeding self-efficacy and other determinants of the duration of breastfeeding in a cohort of first-time mothers in Adelaide,**

- Australia.** *Midwifery* 2007, **23**(4):382-391.
18. Glassman ME, McKearney K, Saslaw M, Sirota DR: **Impact of breastfeeding self-efficacy and sociocultural factors on early breastfeeding in an urban, predominantly Dominican community.** *Breastfeeding medicine : the official journal of the Academy of Breastfeeding Medicine* 2014, **9**(6):301-307.
 19. Semenic S, Loiselle C, Gottlieb L: **Predictors of the duration of exclusive breastfeeding among first-time mothers.** *Research in nursing & health* 2008, **31**(5):428-441.
 20. Dibley MJ, Alam A, Fahmida U, Ariawan I, Titaley CR, Htet MK, Damayanti R, Li M, Sutrisna A, Ferguson E: **The BADUTA study: a protocol for a cluster randomized controlled trial of of a package of behaviour change interventions to improve maternal and child nutrition in East Java Indonesia (unpublished manuscript).** 2020.
 21. **Population by age group and sex in East Java Province. 2010 [Penduduk Menurut Kelompok Umur dan Jenis Kelamin Provinsi Jawa Timur. 2010]** [<http://sp2010.bps.go.id/index.php/site/tabel?tid=336&wid=3500000000&lang=id>]
 22. BPS-Statistics of Malang District: **Malang District in Figures 2018.** Malang, East Java: BPS-Statistics of Malang District; 2018.
 23. BPS-Statistics of Sidoarjo District: **Sidoarjo District in Figures 2018.** Sidoarjo, East Java BPS-Statistics of Sidoarjo District; 2018.
 24. **Commcare: The world's most powerful mobile data collection platform** [<https://www.dimagi.com/commcare/>]
 25. Keemer F: **Breastfeeding self-efficacy of women using second-line strategies for healthy term infants in the first week postpartum: an Australian observational study.** *International breastfeeding journal* 2013, **8**(1):18-18.
 26. Poorshaban F, Pakseresht S, Bostani Khalesi Z, KazemNejad Leili E: **Factors Associated with Breastfeeding Self-Efficacy of Mothers Within 6 Weeks of Delivery** *Journal of Holistic Nursing and Midwifery* 2017, **27**(1):27-34.
 27. Awaliyah SN, Rachmawati IN, Rahmah H: **Breastfeeding self-efficacy as a dominant factor affecting maternal breastfeeding satisfaction.** *BMC Nursing* 2019, **18**(1):30.
 28. Nankumbi J, Mukama AA, Ngabirano TD: **Predictors of breastfeeding self-efficacy among women attending an urban postnatal clinic, Uganda.** *Nurs Open* 2019, **6**(3):765-771.
 29. Filmer D, Pritchett LH: **Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India.** *Demography* 2001, **38**(1):115-132.
 30. (NLIS) WHO-NLIS: **Country profile indicators: interpretation guide.** Geneva: World Health Organization; 2010.
 31. Otsuka K, Taguri M, Dennis CL, Wakutani K, Awano M, Yamaguchi T, Jimba M: **Effectiveness of a breastfeeding self-efficacy intervention: do hospital practices make a difference?** *Maternal and child health journal* 2014, **18**(1):296-306.

32. Chan MY, Ip WY, Choi KC: **The effect of a self-efficacy-based educational programme on maternal breastfeeding self-efficacy, breast feeding duration and exclusive breastfeeding rates: A longitudinal study.** *Midwifery* 2016, **36**:92-98.
33. Brockway M, Benzies K, Hayden KA: **Interventions to Improve Breastfeeding Self-Efficacy and Resultant Breastfeeding Rates: A Systematic Review and Meta-Analysis.** *Journal of human lactation : official journal of International Lactation Consultant Association* 2017, **33**(3):486-499.
34. Ansari S, Abedi P, Hasanpoor S, Bani S: **The Effect of Interventional Program on Breastfeeding Self-Efficacy and Duration of Exclusive Breastfeeding in Pregnant Women in Ahvaz, Iran.** *International Scholarly Research Notices* 2014, **2014**:6.
35. Dodt RC, Joventino ES, Aquino PS, Almeida PC, Ximenes LB: **An experimental study of an educational intervention to promote maternal self-efficacy in breastfeeding.** *Revista latino-americana de enfermagem* 2015, **23**(4):725-732.
36. Awano M, Shimada K: **Development and evaluation of a self care program on breastfeeding in Japan: A quasi-experimental study.** *Int Breastfeed J* 2010, **5**:9.
37. Skouteris H, Nagle C, Fowler M, Kent B, Sahota P, Morris H: **Interventions designed to promote exclusive breastfeeding in high-income countries: a systematic review.** *Breastfeeding medicine : the official journal of the Academy of Breastfeeding Medicine* 2014, **9**(3):113-127.
38. Haroon S, Das JK, Salam RA, Imdad A, Bhutta ZA: **Breastfeeding promotion interventions and breastfeeding practices: a systematic review.** *BMC public health* 2013, **13** Suppl 3:S20.
39. Jolly K, Ingram L, Khan KS, Deeks JJ, Freemantle N, MacArthur C: **Systematic review of peer support for breastfeeding continuation: metaregression analysis of the effect of setting, intensity, and timing.** *BMJ (Clinical research ed)* 2012, **344**:d8287.
40. Hobbs AJ, Mannion CA, McDonald SW, Brockway M, Tough SC: **The impact of caesarean section on breastfeeding initiation, duration and difficulties in the first four months postpartum.** *BMC Pregnancy and Childbirth* 2016, **16**(1):90.
41. Regan J, Thompson A, DeFranco E: **The Influence of Mode of Delivery on Breastfeeding Initiation in Women with a Prior Cesarean Delivery: A Population-Based Study.** *Breastfeeding Medicine* 2013, **8**(2):181-186.
42. Brown A, Jordan S: **Impact of birth complications on breastfeeding duration: an internet survey.** *J Adv Nurs* 2013, **69**(4):828-839.
43. Ratnasari D, Paramashanti BA, Hadi H, Yugistyowati A, Astiti D, Nurhayati E: **Family support and exclusive breastfeeding among Yogyakarta mothers in employment.** *Asia Pacific journal of clinical nutrition* 2017, **26**(Suppl 1):S31-s35.
44. Kingston D, Dennis C-L, Sword W: **Exploring Breastfeeding Self-efficacy.** *The Journal of perinatal & neonatal nursing* 2007, **21**(3):207-215.
45. Balkam JA, Cadwell K, Fein SB: **Effect of components of a workplace lactation program on breastfeeding duration among employees of a public-sector employer.** *Maternal and child health journal* 2011, **15**(5):677-683.

46. Dinour LM, Szaro JM: **Employer-Based Programs to Support Breastfeeding Among Working Mothers: A Systematic Review.** *Breastfeeding Medicine* 2017, **12**(3):131-141.
47. Titaley CR, Loh PC, Prasetyo S, Ariawan I, Shankar AH: **Socio-economic factors and use of maternal health services are associated with delayed initiation and non-exclusive breastfeeding in Indonesia: secondary analysis of Indonesia Demographic and Health Surveys 2002/2003 and 2007.** *Asia Pacific journal of clinical nutrition* 2014, **23**(1):91-104.
48. Saunders RP, Evans MH, Joshi P: **Developing a process-evaluation plan for assessing health promotion program implementation: a how-to guide.** *Health promotion practice* 2005, **6**(2):134-147.

Supplementary Tables

Table S1. Factors associated with low breastfeeding self-efficacy amongst mothers of children under six months old, including specific breastfeeding problems, The BADUTA Study in East Java, Indonesia, 2015-2016

Variable	Univariate			Multivariate ¹				
	aOR	95% CI		<i>p</i>	aOR	95% CI		<i>p</i>
Contextual and intervention characteristics								
Exposure to intervention								
Exposed to intervention ²	1.00							
Not exposed to intervention ³	1.44	1.13	1.83	0.003				
Number of breastfeeding interventions mothers exposed to								
Three or more interventions	1.00				1.00			
Two interventions	1.53	0.83	2.81	0.175	1.45	0.71	2.95	0.302
One intervention	2.07	1.16	3.69	0.014	1.79	0.97	3.29	0.062
No interventions	2.42	1.41	4.15	0.001	1.82	1.05	3.13	0.032
Period								
Baseline	1.00				1.00			
Endline	0.73	0.55	0.97	0.032	0.89	0.62	1.27	0.516
Household characteristics								
Household wealth index								
Poorest	1.00							
Poor	0.99	0.70	1.39	0.948				
Middle	1.23	0.86	1.75	0.257				
Rich	1.35	0.93	1.95	0.118				
Richest	0.83	0.52	1.32	0.426				
Mother's characteristics								
Maternal age								
≤19 years	1.00							
20-34 years	0.70	0.41	1.17	0.172				
35+ years	0.67	0.40	1.14	0.138				
Maternal education								
University/Academy	1.00				1.00			
Completed senior high school	1.69	1.19	2.39	0.003	2.05	1.36	3.08	0.001
Completed junior high school	1.94	1.31	2.86	0.001	2.36	1.48	3.78	<0.001
No school/incomplete primary/completed primary school	1.73	1.12	2.67	0.014	1.97	1.21	3.21	0.007
Maternal occupation								
Housework	1.00				1.00			
Working outside the house	1.38	1.06	1.81	0.018	1.68	1.22	2.31	0.002

Variable	Univariate			Multivariate ¹		
	aOR	95% CI	<i>p</i>	aOR	95% CI	<i>p</i>
Number of children still alive						
1	1.00					
2	0.99	0.77 1.26	<i>0.904</i>			
3	1.02	0.69 1.52	<i>0.920</i>			
4+	1.07	0.55 2.07	<i>0.847</i>			
Previous live birth						
None	1.00					
Any	1.01	0.82 1.26	<i>0.906</i>			
Antenatal and delivery care						
Minimum antenatal care visits ⁴						
Completed (4+ visits)	1.00			1.00		
Incomplete (<4 visits)	1.28	0.98 1.68	<i>0.073</i>	1.27	0.95 1.71	<i>0.112</i>
Mode of delivery						
Normal	1.00			1.00		
Caesarean	1.19	0.96 1.47	<i>0.108</i>	1.34	1.06 1.70	<i>0.017</i>
Birth attendant						
General practitioner/OBGYN	1.00					
Midwife/nurse	0.95	0.78 1.165	<i>0.631</i>			
Traditional birth attendant/family/friend	1.29	0.54 3.082	<i>0.572</i>			
Child's characteristics						
Sex of the child						
Male	1.00					
Female	1.03	0.82 1.30	<i>0.791</i>			
Birth weight from monitoring card						
Larger than average	1.00					
Average	0.72	0.51 1.00	<i>0.053</i>			
Smaller than average	0.72	0.44 1.17	<i>0.183</i>			
Breastfeeding knowledge & experience						
Ever received any breastfeeding advice						
Yes	1.00			1.00		
No	1.40	1.09 1.78	<i>0.007</i>	1.43	1.10 1.86	<i>0.009</i>

Variable	Univariate			Multivariate ¹		
	aOR	95% CI	<i>p</i>	aOR	95% CI	<i>p</i>
Knowledge about breastfeeding						
High level ⁵	1.00			1.00		
Low level ⁶	1.58	1.19 2.10	0.002	1.35	1.01 1.81	0.045
Problems with breastfeeding						
Swollen breast						
No	1.00			1.00		
Yes	2.35	1.03 5.37	0.043	3.10	1.33 7.21	0.009
Sore nipples						
No	1.00			1.00		
Yes	0.78	0.53 1.16	0.217	0.79	0.50 1.24	0.305
Flat nipples						
No	1.00			1.00		
Yes	2.87	1.42 5.81	0.004	2.75	1.29 5.87	0.009
Infants refused breastfeeds						
No	1.00			1.00		
Yes	7.18	2.16 23.79	0.001	5.45	1.64 18.13	0.006
Perceived breastmilk is not enough						
No	1.00			1.00		
Yes	2.75	2.03 3.74	<0.001	2.80	2.04 3.84	<0.001

Note:

¹Multivariate logistic regression using the backward elimination method to select significant predictors of low breastfeeding self-efficacy. The variable for the minimum requirement of four antenatal care visits by trimester was selected a priori to be retained in the final model regardless of its significance level. ²"Exposed to intervention," refers to respondents living in the intervention sub-districts at the endline survey ³Not exposed to intervention referred to all respondents from the baseline survey and those living in the control sub-districts at the endline survey. ⁴Minimum Antenatal Care refers to the recommendation of at least four antenatal visits, i.e., once in trimester one to three, and twice in trimester three. ⁵High level of knowledge was mothers whose total knowledge score was greater than, or equal to the median knowledge score value. ⁶Low level of knowledge was mothers whose total knowledge score was less than the median knowledge score value.

Table S2: Factors associated with low breastfeeding self-efficacy amongst mothers of children under six months old including specific breastfeeding interventions, The BADUTA Study in East Java, Indonesia, 2015-2016

Variable	Univariate				Multivariate ¹			
	aOR	95% CI		<i>p</i>	aOR	95% CI		<i>p</i>
Contextual and intervention characteristics								
Exposure to intervention								
Exposed to intervention ²	1.00							
Not exposed to intervention ³	1.44	1.13	1.83	0.003				
Period								
Baseline	1.00				1.00			
Endline	0.73	0.55	0.97	0.032	0.97	0.65	1.43	0.867
Ever visited by village facilitator at home								
No	1.00				1.00			
Yes	0.43	0.17	1.12	0.084	0.97	0.34	2.75	0.948
Ever attended "emo-demo"								
No	1.00				1.00			
Yes	0.59	0.35	1.00	0.049	1.17	0.61	2.23	0.635
Ever received SMS Bunda messages								
No	1.00				1.00			
Yes	0.41	0.15	1.14	0.087	0.56	0.16	1.94	0.360
Ever visited by cadre at home								
No	1.00				1.00			
Yes	0.32	0.12	0.84	0.022	0.28	0.09	0.85	0.025
Ever attended pregnancy class								
No	1.00				1.00			
Yes	0.50	0.21	1.21	0.125	0.87	0.33	2.25	0.767
Roomed in with baby after delivery								
No	1.00				1.00			
Yes	0.62	0.49	0.78	0.000	0.75	0.56	1.01	0.063
Ever received counselling by a midwife								
No	1.00				1.00			
Yes	0.81	0.59	1.11	0.185	1.17	0.83	1.65	0.371
Ever received counselling by a cadre								
No	1.00				1.00			
Yes	0.45	0.22	0.95	0.035	0.74	0.33	1.65	0.463
Ever seen "Rumpi Sehat" TV commercials								
No	1.00				1.00			
Yes	0.57	0.38	0.84	0.005	0.72	0.42	1.22	0.216
Household characteristics								
Household wealth index								

Variable	Univariate				Multivariate ¹			
	aOR	95% CI		<i>p</i>	aOR	95% CI		<i>p</i>
Poorest	1.00							
Poor	0.99	0.70	1.39	<i>0.948</i>				
Middle	1.23	0.86	1.75	<i>0.257</i>				
Rich	1.35	0.93	1.95	<i>0.118</i>				
Richest	0.83	0.52	1.32	<i>0.426</i>				
Mother's characteristics								
Maternal age								
≤19 years	1.00							
20-34 years	0.70	0.41	1.17	<i>0.172</i>				
35+ years	0.67	0.40	1.14	<i>0.138</i>				
Maternal education								
University/Academy	1.00				1.00			
Completed senior high school	1.69	1.19	2.39	<i>0.003</i>	1.92	1.28	2.87	<i>0.002</i>
Completed junior high school	1.94	1.31	2.86	<i>0.001</i>	2.17	1.39	3.37	<i>0.001</i>
No school/incomplete primary/completed primary school	1.73	1.12	2.67	<i>0.014</i>	1.85	1.16	2.97	<i>0.011</i>
Maternal occupation								
Housework	1.00				1.00			
Working outside the house	1.38	1.06	1.81	<i>0.018</i>	1.72	1.25	2.36	<i>0.001</i>
Number of children still alive								
1	1.00							
2	0.99	0.77	1.26	<i>0.904</i>				
3	1.02	0.69	1.52	<i>0.920</i>				
4+	1.07	0.55	2.07	<i>0.847</i>				
Previous live birth								
None	1.00							
Any	1.01	0.82	1.26	<i>0.906</i>				
Antenatal and delivery care								
Minimum antenatal care visits ⁴								
Completed (4+ visits)	1.00				1.00			
Incomplete (<4 visits)	1.28	0.98	1.68	<i>0.073</i>	1.21	0.90	1.64	<i>0.209</i>
Mode of delivery								
Normal	1.00				1.00			
Caesarean	1.19	0.96	1.47	<i>0.108</i>	1.37	1.08	1.74	<i>0.011</i>

Variable	Univariate				Multivariate ¹			
	aOR	95% CI		<i>p</i>	aOR	95% CI		<i>p</i>
Birth attendant								
General practitioner/OBGYN	1.00							
Midwife/nurse	0.95	0.78	1.165	0.631				
Traditional birth attendant/family/friend	1.29	0.54	3.082	0.572				
Child's characteristics								
Sex of the child								
Male	1.00							
Female	1.03	0.82	1.30	0.791				
Birth weight from monitoring card								
Larger than average	1.00							
Average	0.72	0.51	1.00	0.053				
Smaller than average	0.72	0.44	1.17	0.183				
Breastfeeding knowledge & experience								
Ever received any breastfeeding advice								
Yes	1.00				1.00			
No	1.40	1.09	1.78	0.007	1.47	1.11	1.93	0.006
Knowledge about breastfeeding								
High level ⁵	1.00				1.00			
Low level ⁶	1.58	1.19	2.10	0.002	1.39	1.04	1.87	0.029
Problems with breastfeeding								
None	1.00				1.00			
Not related to illness	3.14	2.34	4.23	<0.001	3.40	2.54	4.56	<0.001
Related to illness/anatomical condition	1.28	0.91	1.80	0.158	1.35	0.95	1.93	0.098
Both types of problems	3.18	1.29	7.81	0.012	3.73	1.33	10.43	0.012

Note:

¹Multivariate logistic regression using the backward elimination method to select significant predictors of low breastfeeding self-efficacy. The variable for the minimum requirement of four antenatal care visits by trimester was selected a priori to be retained in the final model regardless of its significance level. ²Exposed to intervention refers to respondents living in the intervention sub-districts at the endline survey ³Not exposed to intervention refers to all respondents from the baseline survey and those living in the control sub-districts at the endline survey. ⁴Minimum antenatal care refers to the recommendation of at least four antenatal visits, i.e., once in trimester one to three, and twice in trimester three. ⁵High level of knowledge was mothers whose total knowledge score was greater than, or equal to the median knowledge score

value. ⁶Low level of knowledge was mothers whose total knowledge score was less than the median knowledge score value.

Figures




Figure 1

The analytical framework used to examine factors associated with mothers' low self efficacy on breastfeeding, The BADUTA Study in East Java, Indonesia, 2015-2016




Figure 2

Distribution of mothers of children under six months by their answers to each breastfeeding knowledge component, The BADUTA Study in East Java, Indonesia, 2015-2016.




Figure 3

Distribution of mothers with low breastfeeding self-efficacy by the feeding patterns of their infants under six months of age, The BADUTA Study in East Java, Indonesia, 2015-2016. Note: A chi-square statistic was performed to examine the distribution of infants who were exclusively breastfed based on mothers'

self-efficacy level. The bars represent 95% CI. Svy commands were used to adjust for the cluster sampling design.




Figure 4

Distribution of mothers of children under six months by their answers to each component of breastfeeding self-efficacy short form (BSES-SF) The BADUTA Study in East Java, Indonesia, 2015-2016
 Note: For each question, a score of 1 to 5 was assigned to the responses from "strongly disagree" to "strongly agree." BF=breastfeeding