

Effectiveness of a Community Health Worker(CHW) training in monitoring and care of patients with Chronic Obstructive Pulmonary Disease(COPD) in rural Gujarat, India.

Ashish Gupta

K M Patel Institute of Physiotherapy

Ajay Phatak

H M Patel Center for Medical Care and Education

Meha Patel

K M Patel Institute of Physiotherapy

Neha Das

Pramukhswami Medical College

Nirav Vaghela

K M Patel Institute of Physiotherapy

Harihara Prakash

K M Patel Institute of Physiotherapy

Shyamsundar J Raithatha (✉ sundar.shyamsundar@gmail.com)

H M Patel Center For Medical Care and Education <https://orcid.org/0000-0002-5617-5737>

Research article

Keywords: Pulmonary Disease, Chronic Obstructive, Community Health Workers, Primary Health Care, Learning

Posted Date: October 28th, 2019

DOI: <https://doi.org/10.21203/rs.2.16515/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background- COPD is the second leading contributor to the disease burden of India. The current COPD burden cannot be managed effectively just through a physician based approach. Community Health Workers(CHWs) can play an effective role in making COPD care accessible and effective. Findings of an assessment of a training program for CHWs on COPD are reported here.

Methods- 90 CHWs were exposed to a series of 5 training sessions designed and administered jointly by a team of public health experts and physiotherapists in the year 2017-18. Topics covered were basic clinical aspects of COPD, monitoring a patient with COPD and basic aspects of pulmonary rehabilitation. The assessment comprised 12 MCQs and short questions, 7 video exercises, 2 case vignettes and 5 skill assessments through Objectively Structured Clinical Examination(OSCE). Mean percentage scores were calculated for each domain of assessment to make it comparable.

Results 70 CHWs with a mean age of 42.2 years completed all the trainings and underwent the assessment. Mean percentage score(SD) for knowledge was 62%(16.3). In OSCE assessment, scores were best in sputum clearance technique demonstration(92.1%) and the least in dyspnoea relieving positions(59.2%). The CHWs had difficulties in identifying signs of respiratory distress(score – 55.1%). No statistically significant association was observed between performance scores and their sociodemographic profile.

Conclusion The results were encouraging and the program may be pilot tested in a Government setting.

Background

Non-communicable Diseases(NCDs) contribute to 61.2 % of deaths and 55.4% of Disability Adjusted Life Years(DALYs) lost in India.¹ As per the Global Burden of Diseases, Injuries and Risk factors study(GBD) 2016, COPD is the second leading cause of disease burden in India after ischaemic heart diseases, contributing to 8.7% and 4.8% of the total deaths and DALYs respectively. There are an estimated 55-3 million COPD cases in India with a crude prevalence of 4.2%.²

COPD can be managed effectively through a multidimensional approach involving pharmacotherapy, exercise therapy, diet counselling and psychosocial support. Such an approach leads to an improvement in dyspnoea, exercise capacity and quality of life.^{3,4} Although guidelines for management of COPD are available for more than a decade⁵, a recent study indicated that the practice pattern is significantly different among physicians.⁶ In hospital settings, a nurse led model of COPD management model has been proposed.⁷ However, the evidence on effectiveness of such models is not promising.⁸ In a resource constrained setting like India, the availability of chest physicians and other health professionals is not adequate to meet the high burden of the disease. In order to address this burden, there is a strong need to integrate COPD care in the existing primary health care system through a task sharing approach.⁹ To strengthen the General Practitioners in managing COPD and to standardize the management pathways, a

capacity enhancing model has been designed in India which is a welcome step.¹⁰ However simultaneously other members of the health workforce also need to be empowered to deliver COPD prevention and care services in an integrated fashion. Community Health Workers supported by mid-level health workers and primary care physicians can form an appropriate care delivery system to manage COPD in the community. Such a model has been proposed in Nepal which focuses mainly on integration of health system components.¹¹

Abhay Bang successfully demonstrated utility of CHWs in Home Based Newborn Care and the model was replicated in other parts of India as well as in other developing economies.¹² Based on the learnings from that model, CHWs (known as Accredited Social Health Activist (ASHA)) have become a back bone of the Indian primary health care system. Government of India through the Comprehensive Primary Health Care (CPHC) initiative aims to provide NCD prevention and care services through the network of Health and Wellness Centres across the country where ASHA workers have an important role to play.¹³ Under this initiative, trained ASHA workers can form an important part of the service delivery framework for COPD prevention and care. SPARSH (Shree Krishna Hospital Program for Advancement of Rural and Social Health) is an initiative of Charutar Arogya Mandal to create a rural integrated model of NCD prevention and care. The initiative is implemented through a linkage between CHWs known as Village Health Workers (VHWs), Mobile Health Teams (MHTs), Extension Centres and a tertiary care teaching hospital. SPARSH is implemented in 150 villages across 3 districts of Gujarat located in western India. A training program was implemented under SPARSH to empower its CHWs in providing COPD prevention and care services at the community level. It was evaluated through a competency assessment focusing on cognitive and psycho-motor domains of the CHWs. This paper reports the findings of the assessment.

Methods

It was an educational intervention without a control group. Since the CHWs had never received a training in COPD in the past, baseline assessment of their knowledge and skills on COPD was not relevant. In the year 2017-2018, 90 CHWs of SPARSH were invited for the training through a series of five sessions conducted at monthly intervals. The training program was designed and delivered by a team of public health professionals and physiotherapists. The sessions were conducted in vernacular language (Gujarati or Hindi) through power-point presentations, demonstrations, hands on practice sessions and screening of videos.

The topics covered were basic clinical aspects of COPD (risk factors, clinical features, diagnosis and treatment), monitoring a patient with COPD and basic aspects of pulmonary rehabilitation (dyspnea relieving positions, sputum clearance technique and chest mobility exercises). The concepts were explained in simple language through interactive power point slides and screening of video clips. While teaching a skill, the skill was first demonstrated live and/or through a video, followed by practicing the skill in pairs on each other in small groups of 5-10 under guidance of a facilitator. They were also given an opportunity to practice the same in presence of the program staff during camps organized in their

respective villages. At the end of the training, CHWs were given a reading booklet providing information about the sessions. Two refresher sessions were conducted for all CHWs.

At the end of the training program they were assessed in various domains through different techniques(**Table 1**). The Multiple Choice Questions (MCQs), short questions and different checklists were prepared in consultation with chest physicians and consensually validated. CHWs were assessed at each OSCE(Objectively Structure Clinical Examination) station by a trained assessor(not involved in training sessions) using a check-list of steps(**Table 2**). The assessors (postgraduate students in physiotherapy) were sensitized to the entire training program conducted for the CHWs and the latter's role in the community. They were also oriented to the scoring pattern at the OSCE station. A pilot examination was carried out on 10 field supervisors and paramedical staff working in the program to standardize the assessment process.

Table 1: Various domains and assessment techniques for the CHWs

No	Domain	Assessment technique	No. of questions	Marks (% weightage)
1	Basic clinical knowledge about COPD	Multiple choice questions(MCQs) about COPD	12 MCQs	20 marks (19.6%)
2	Ability to identify signs of respiratory distress	Identifying the signs by videos	Five videos	10 marks (9.8%)
3	Ability to assess severity of breathlessness through the Medical Research Council(MRC) scale ²²	Applying the MRC scale by case scenarios.	Two scenarios	14 marks (13.7%)
4	Counting respiratory rate	Counting respiratory rate through a video showing breathing movements for one minute	Two videos	8 marks (7.8%)
5	The ability to use and document PEFR (Peak Expiratory Flow Rate)	OSCE	One - Station Ten steps	10 marks (9.8%)
6	The ability to demonstrate basic breathing and general mobility exercises to COPD patients.	OSCE	Two Station with two skills at each station, each skill was of seven steps	40 marks (39.2%)

Table 2: Checklist of steps for the three stations with marks for each step shown in parenthesis

Station 1- Peak flow measurement	Station 2 - Pursed lip breathing and Dyspnoea relieving positions	Station 3 - Sputum clearance technique and thoracic mobility exercises
<p>1. Set the dial to zero (1)</p> <p>2. CHW should hold the peak flow meter correctly with both hands without placing any finger on the dial (1)</p> <p>3. Sit up straight or stand (1)</p> <p>4. Take as deep a breathe as possible (1)</p> <p>5. Put the mouth piece of the PEFR machine in the mouth and seal it between the lips (1)</p> <p>6. Exhale as hard and fast as possible. (1)</p> <p>7. Record the reading (1)</p> <p>8. The technique should be performed 3 times (1)</p> <p>9. Ask the CHW the procedure to clean the PEFR machine</p> <p>9a She should mention about asking the patient to wash the mouth piece with soap and water (1)</p> <p>9b She should mention about washing the mouth piece with clean water and spirit (1)</p>	<p>Station 2a - Demonstration of pursed lip breathing</p> <p>1. Inspiration through nose (3)</p> <p>2. Pursing the lips (4)</p> <p>3. Expiration through pursed lips (3)</p> <p>Station 2b - Demonstrating the dyspnoea relieving positions</p> <p>1. Standing position with wall support and performing pursed lip breathing (2.5)</p> <p>2. In sitting position, laying the head on the table and turning the face on one side and performing pursed lip breathing (2.5)</p> <p>3. In sitting position placing hands on the thighs and forward bending and performing pursed lip breathing (2.5)</p> <p>4. In side lying position, resting on 3-4 pillows and performing pursed lip breathing (2.5)</p>	<p>Station 3a - Sputum clearance technique</p> <p>1. CHW performs the procedure for humidification (2.5)</p> <p>2. Demonstrates first position for huffing (2.5)</p> <p>3. Demonstrates second position for huffing (2.5)</p> <p>4. Demonstrates third position for huffing (2.5)</p> <p>Station 3b - Thoracic mobility exercises</p> <p>1. Demonstrates 1 exercise (3)</p> <p>2. Demonstrates 2nd exercise (3)</p> <p>3. Demonstrates 3rd exercise (4)</p>

Data Analysis: Descriptive statistics[Mean (SD) and Frequency (%)] were used to depict the characteristics of the study population such as years of schooling, duration of work experience and involvement in other community work (ASHA/Anganwadi Worker(AWW)/Working with an other NGO). The raw scores in each domain were transformed into percentage for easy understanding. Chi Square test/t-test was applied to determine association between the CHWs performance scores with their age, education and their involvement in other community work (ASHA/AW/NGO) at univariate level depending on nature of the variables involved. Simple regression analysis was performed to find out important baseline characteristics that influence performance of CHWs. Ethical clearance for the study was obtained from the Institutional Ethics Committee of the H M Patel Center for Medical Care and Education, Karamsad (Approval no. - IEC/HMPCMCE/87/Faculty/6/ Dated 15/11/2017.)

Results

A total of 90 CHWs village health workers were invited for the training. Eleven health workers could not complete the training program due to various personal reasons whereas nine health workers were not available during evaluation process. Thus, the analysis included 70 health workers. The mean(SD)[IQR] age of the participants was 42.2(11.2)[17, 75] years. Most of them (71.4%) had 8-12 years of education and had involvement in some other community work) (57.1%). They were associated with SPARSH for an average of 3.5 years.

The mean percent score(SD) on basic knowledge of COPD was 62.0%(16.3). Sputum clearance 92.1% (18.9), using PEFr 82.1%(13.3)%, pursed lip breathing 81.3%(27.1) were best demonstrated by the trainees whereas thoracic mobility exercises 66.6%(33.2) and Dyspnea relieving positions 59.2%(25.6) needed the most improvement. **(Table 3)**

Most participants 56(80%) could correctly count respiratory rate from the two videos. Most of them 51(73%) and 54(77%) applied MRC scale satisfactorily in two scenarios provided. However, signs of respiratory distress were identified correctly by only 31(44%), 17(24%), 33(47%), 43(61%) and 15(21%) participants from the five videos respectively.

No statistically significant association was observed between performance of the participants (theory as well as skills) and their sociodemographic profile like age, experience, education level etc.

Table 3: Performance of participants in knowledge test and skills

Sr. No.	Particulars	Mean(SD) % Score *
Skill performance - OSCE		
1	Sputum clearance techniques	92.14(18.93)
2	Peak flow meter performance	82.14(13.26)
3	Pursed lip breathing	81.29(27.12)
4	Thoracic Mobility	66.64(33.20)
5	Dyspnoea relieving positions	59.21(25.57)
Skill performance - Video Exercises		
6	Respiratory rate counting	85.36(27.08)
7	Identifying signs of respiratory distress	55.14(19.47)
Applying MRC scale through case vignettes		
8	Case scenario score	63.83(23.95)
Knowledge score		
9	MCQs and short questions	62.00(16.26)

Discussion

Seventy CHWs were thus assessed for their knowledge and skills related to COPD prevention and care after undergoing an intensive training program. It is probably the first such rigorous attempt to train CHWs in multiple cognitive and psychomotor domains related to prevention and care of COPD in the country. The results were encouraging and have implications for addressing the COPD burden through the Government primary health care system. Their performance in demonstration of peak flow meter use, pursed lip breathing and sputum clearance technique was good. Thoracic mobility exercises and dyspnoea relieving positions needed some improvement. Majority of them could count the respiratory rate appropriately while there were some issues in identifying the signs of respiratory distress. Mean percentage score for the knowledge domain was more than 60%. There was no significant difference in scores with respect to experience of working with SPARSH, schooling and involvement in work with other organizations.

There is considerable evidence on effectiveness of CHW trainings for maternal and child health and communicable diseases in low and middle income countries(LMICs).¹⁴ However there is relatively less literature available on the role of CHWs in addressing the burden of NCDs in low and middle income countries, inspite of NCDs being the major disease burden. The evidence, whatever available, is mainly focused on addressing cardiovascular diseases and its risk factors such as diabetes, hypertension and tobacco.¹⁵⁻¹⁷ Evidence on the role of CHWs in chronic respiratory diseases is hardly available. In a systematic review for effectiveness of CHW trainings for cardiovascular diseases, post test scores in

most of the studies were found to be 70–80% irrespective of the pre-test scores.¹⁸ In our study the post test knowledge score was found to be 64%. In one of the studies from the review, it was found that CHWs preferred interactive trainings, hands on experience and case scenarios over didactic training.¹⁹ Our training also comprised of a combination of demonstrations, videos, case scenario discussions and hands on exercises in small groups with minimal didactic sessions.

As per the Kirkpatrick model, a training program needs to be evaluated at four levels.²⁰ Through this paper we are reporting the findings of the assessment of the CHW performance post-training which reflects the second level of the model - learning level. However a more comprehensive evaluation would require an assessment of the reaction of the CHWs to the trainings, their change in behaviours and the health outcomes. Also for the CHW to deliver the interventions based on the knowledge and skills gained by them, other program level interventions such as supervision and feedback, materials management and information systems are required to be in place.

Conclusions

Considering the encouraging results obtained in this paper, this intervention should be pilot tested in a Government setting with ASHA workers. It will provide valuable learnings in scaling up the COPD prevention and care services across the network of 1.5 lakh Health and Wellness centres across the country under the Comprehensive Primary Health Care(CPHC) component of the Ayushman Bharat initiative.²¹

Abbreviations

ASHA- Accredited Social Health Activist

AW - Anganwadi

AWW - Anganwadi Worker

CHW- Community Health Worker

COPD- Chronic Obstructive Pulmonary Disease

CPHC - Comprehensive Primary Health Care

DALYs- Disability Adjusted Life Years

GBD- Global Burden of Diseases

IQR- Inter Quartile Range

LMICs - low and middle income countries

MCQs -Multiple choice questions

MHTs- Mobile Health Teams

MRC -Medical Research Council

NCDs- Non-communicable Diseases

NGO-Non-Governmental Organization

OSCE- Objective Structure Clinical Examination

PEFR- Peak Expiratory Flow Rate

SPARSH - Shree Krishna Hospital Program for Advancement of Rural and Social Health

VHWs- Village Health Workers

Declarations

1. Ethical clearance for the study was obtained from the Institutional Ethics Committee of the H M Patel Center for Medical Care and Education, Karamsad (Approval no. - IEC/HMPCMCE/87/Faculty/6/ Dated 15/11/2017.) Since this study was undertaken as a part of a routine training assessment by the training team and the data was not shared with any third party, waiver of written consent was granted by the institutional ethics committee.
2. Consent for publication – Not applicable
3. Availability of data and materials – Available on request to the corresponding author over email
4. Competing interests - The authors declare that they have no competing interests
5. Funding - The training program was funded through a service delivery program grant from GMM Pfaudler , Shamdasani Foundation, Hongkong and Tata trusts. The funders had no role in execution and publication of this study.
6. Authors' contributions - Brief statement of the contribution of each author -
 - AG – Designing and implementing the training program and the assessment and preparing the manuscript
 - AP – Designing the assessment, data analysis and preparing the manuscript
 - MP – Conducting the training and assessments, data entry
 - ND – Conducting the training, data cleaning
 - NV – Guiding data collection
 - HP – Conceptualizing the study
 - SR – Conceptualising the study, designing the training module, preparing the analysis plan and writing the manuscript
 - All authors have read and approved the manuscript
7. Acknowledgement – Not applicable

References

1. ICMR, PHFI, IHME. India: Health of the nation's states - The India State-level Disease Burden Initiative. New Delhi, India [Internet]. 2017 [cited 2019 Jun 4]. Available from: http://www.healthdata.org/sites/default/files/files/2017_India_State-Level_Disease_Burden_Initiative_-_Full_Report%5B1%5D.pdf
2. India State-Level Disease Burden Initiative CRD Collaborators. The burden of chronic respiratory diseases and their heterogeneity across the states of India: the Global Burden of Disease Study 1990–2016. *Lancet Glob Health*. 2018;6(12):e1363–74.
3. Lacasse Y, Martin S, Lasserson TJ, Goldstein RS. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. A Cochrane systematic review. *Eur Medicophysica*. 2007 Dec;43(4):475–85.

4. Van Wetering CR, Hoogendoorn M, Mol SJM, Rutten-van Mólken MPMH, Schols AM. Short- and long-term efficacy of a community-based COPD management programme in less advanced COPD: a randomised controlled trial. *Thorax*. 2010 Jan;65(1):7–13.
5. Bellamy D, Bouchard J, Henrichsen S, Johansson G, Langhammer A, Reid J, et al. International Primary Care Respiratory Group (IPCRG) Guidelines: management of chronic obstructive pulmonary disease (COPD). *Prim Care Respir J J Gen Pract Airw Group*. 2006 Feb;15(1):48–57.
6. Wang DY, Cho S-H, Lin H-C, Ghoshal AG, Bin Abdul Muttalif AR, Thanaviratnanich S, et al. Practice Patterns for Chronic Respiratory Diseases in the Asia-Pacific Region: A Cross-Sectional Observational Study. *Int Arch Allergy Immunol*. 2018;177(1):69–79.
7. Fletcher MJ, Dahl BH. Expanding nurse practice in COPD: is it key to providing high quality, effective and safe patient care? *Prim Care Respir J J Gen Pract Airw Group*. 2013 Jun;22(2):230–3.
8. Jolly K, Sidhu MS, Bates E, Majothi S, Sitch A, Bayliss S, et al. Systematic review of the effectiveness of community-based self-management interventions among primary care COPD patients. *NPJ Prim Care Respir Med*. 2018 23;28(1):44.
9. Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness: the chronic care model, Part 2. *JAMA*. 2002 Oct 16;288(15):1909–14.
10. Bhalla S, Sinha SK, Jain S, Gupta P, Kumar P, Chandwani H, et al. Improving primary care physicians' capacity: A pan India initiative on management of chronic obstructive pulmonary disease and asthma. *Lung India Off Organ Indian Chest Soc*. 2018 Oct;35(5):452–3.
11. Kumar A, Schwarz D, Acharya B, Agrawal P, Aryal A, Choudhury N, et al. Designing and implementing an integrated non-communicable disease primary care intervention in rural Nepal. *BMJ Glob Health*. 2019;4(2):e001343.
12. Bang AT, Bang RA, Baitule SB, Reddy MH, Deshmukh MD. Effect of home-based neonatal care and management of sepsis on neonatal mortality: field trial in rural India. *Lancet Lond Engl*. 1999 Dec 4;354(9194):1955–61.
13. Ved RR, Gupta G, Singh S. India's health and wellness centres: realizing universal health coverage through comprehensive primary health care. *WHO South-East Asia J Public Health*. 2019 Apr;8(1):18–20.
14. WHO, Global health workforce alliance. Global Experience of Community Health Workers for Delivery of Health Related Millennium Development Goals: A Systematic Review, Country Case Studies, and Recommendations for Integration into National Health Systems [Internet]. 2010 [cited 2019 Jun 18]. Available from: www.who.int/workforcealliance/knowledge/publications/CHW_FullReport_2010.pdf
15. Hill J, Peer N, Oldenburg B, Kengne AP. Roles, responsibilities and characteristics of lay community health workers involved in diabetes prevention programmes: A systematic review. *PloS One*. 2017;12(12):e0189069.
16. Jacob V, Chattopadhyay SK, Hopkins DP, Reynolds JA, Xiong KZ, Jones CD, et al. Economics of Community Health Workers for Chronic Disease: Findings From Community Guide Systematic Reviews. *Am J Prev Med*. 2019 Mar;56(3):e95–106.

17. Jeet G, Thakur JS, Prinja S, Singh M. Community health workers for non-communicable diseases prevention and control in developing countries: Evidence and implications. *PloS One*. 2017;12(7):e0180640.
18. Abdel-All M, Putica B, Praveen D, Abimbola S, Joshi R. Effectiveness of community health worker training programmes for cardiovascular disease management in low-income and middle-income countries: a systematic review. *BMJ Open* [Internet]. 2017 Nov 3;7(11).
19. Abrahams-Gessel S, Denman CA, Montano CM, Gaziano TA, Levitt N, Rivera-Andrade A, et al. The training and fieldwork experiences of community health workers conducting population-based, noninvasive screening for CVD in LMIC. *Glob Heart*. 2015 Mar;10(1):45–54.
20. The Kirkpatrick Model [Internet]. [cited 2019 Jul 22]. Available from: <https://www.kirkpatrickpartners.com/Our-Philosophy/The-Kirkpatrick-Model>
21. Year Ender 2018: Ministry of Health and Family Welfare [Internet]. [cited 2019 Jul 30]. Available from: <http://pib.nic.in/newsite/PrintRelease.aspx?relid = 187429>
22. Stenton C. The MRC breathlessness scale. *Occup Med*. 2008 May 1;58(3):226–7.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [titlepage.docx](#)