Transcriptome based identification and validation of heat stress transcription factors in wheat progenitor species *Aegilops speltoides*

Sushmita Seni

Punjab Agricultural University

Satinder Kaur (✉ satinder.biotech@pau.edu)

Punjab Agricultural University

Palvi Malik

Punjab Agricultural University

Inderjit Singh Yadav

Punjab Agricultural University

Parul Sirohi

Indian Institute of Technology Roorkee

Harsh Chauhan

Indian Institute of Technology Roorkee

Amandeep Kaur

Punjab Agricultural University

Parveen Chhuneja

Punjab Agricultural University

Research Article

Keywords: Wild wheat, *Aegilops speltoides*, heat stress, heat stress transcription factors (HSFs), RNA-Seq, Transcriptome, qRT-PCR

Posted Date: August 2nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-721527/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on November 11th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-01596-6>.

Abstract

Wheat, one of the major cereal crops worldwide, get adversely affected by rising global temperature. We have identified the diploid B genome progenitor of wheat, *Aegilops speltoides* (SS), as a potential donor for heat stress tolerance. Therefore, the objective of the present work was to study the total transcriptome profile of *Ae. speltoides* accession pau3809 and compare with that of tetraploid and hexaploid wheat cultivars PDW274 and PBW725, respectively. The comparative transcriptome was utilized to identify and validate heat stress transcription factors (HSFs), the key genes involved in imparting heat stress tolerance. Transcriptome analysis led to the identification of a total of 74K, 68K, and 76K genes in AS3809, PDW274, and PBW725, respectively. There was a high uniformity of GO profiles under the biological, molecular, and cellular functions across the three wheat transcriptomes, suggesting the conservation of gene function. Twelve HSFs with the highest FPKM value in the *Ae. speltoides* transcriptome data were selected and six of these HSFs namely *HSFA3*, *HSFA5*, *HSFA9*, *HSFB2a*, *HSFC1b*, and *HSFB2b* were validated with qRT PCR. These six HSFs were identified as an important component of thermotolerance in *Ae. speltoides* as evident from their comparative higher expression under heat stress.

Introduction

Wheat is the world's second most important source of nutritional energy and nutrients, after rice. Despite being the world's second-largest wheat producer, yields have been decreasing for several years, notably in India's central regions¹. Climate change on a global scale, particularly the current anomalous temperature regimes, poses a serious threat to the agricultural environment. Given future climatic uncertainty and resource limits, providing food and nutritional security for an expanding population will be a huge challenge².

With an increase in ambient temperature since the turn of the century, global climate models project a rise in mean ambient temperatures ranging from 1.8°C to 5.8°C by the end of the century³. Heat stress has the most adverse effect on productivity as wheat flowers at cool temperatures between 0 - 5°C. Although heat stress affects wheat growth to varying degrees at different phenological stages, its' effects are more prominent on reproductive development (also called terminal heat stress) than on vegetative growth as an increase in temperature during grain filling leads to a reduction in grain filling duration, which directly impacts grain number and yield⁴. Previous studies⁵ have estimated grain filling duration and grain yield to decrease by 5% and 3-4%, respectively, with a per degree increase in temperature between 18 and 22 °C during grain filling. Based on this calculation, losses up to 50% in yield potential on exposure to 32–38 °C during the critical grain formation period have been estimated⁶.

Wheat belongs to the family Poaceae, tribe *Triticeae* and is placed in the genus *Triticum*. The genus *Triticum* comprises of diverse types of cultivated and wild species of wheat. The genome size of bread wheat (*Triticum aestivum* L.), a cultivated allohexaploid species is about 16000 Mb and contains three different genomes (AABBDD), while durum is a cultivated tetraploid species (AABB) of approximately 12000 Mb. The B genome of the bread wheat and durum wheat has been derived from a closely related diploid species of wild wheat *Aegilops speltoides* (SS), but this probably happened only once or a few times and involved only a few progenitors. Consequently, the potential genetic diversity of *Ae. speltoides* have not been represented well in bread and durum wheat germplasm⁷.

The importance of wild species, particularly, *Aegilops* species in breeding programs is documented by various studies where they have reported improvement for drought tolerance⁸, heat tolerance^{9,10} salinity¹¹, resistance to several pests and diseases such as rusts^{12–14} and complex traits such as yield⁷. Therefore, besides being a progenitor of bread wheat, *Ae. speltoides* has established itself as a valuable genetic resource for important agronomic traits. Also, the ability of this wild progenitor to adapt to different climatic zones including drought/heat environments, different disease hot spots, and nutrient-poor areas has enhanced its value as a valuable breeding resource.

Heat stress transcription factors (HSFs) regulate core aspects of the heat stress response¹⁵. Several heat-inducible genes known as "heat shock genes" (HSGs) are up-regulated in response to heat stress and encode heat shock proteins (HSPs) which protect intracellular proteins from denaturation¹⁶. The presence of conserved heat shock elements (HSEs) in the promoter region of HSGs initiates transcription in response to heat. Overexpression of HSFs turns on almost all HSGs thus, protecting against heat stress. Plant HSFs are a complex gene family having an important role in the modulation of transcription during heat stress¹⁷. Based on details of their oligomerization domains, plant HSFs have been distinguished into three classes, i.e., class A, class B, and class C HSFs¹⁵. Only A-type HSFs show transactivation property due to the presence of a C-terminal AHA type activation domain, while B-type HSFs have been proposed as either co-activators or repressors. Multiple copies of HSFs have been reported in plants: 21 in Arabidopsis, 24 in tomato, 25 in pepper, and 56 in wheat^{18–21}.

Previous studies have highlighted the importance of HSFs in wheat under various abiotic stress conditions, particularly heat stress^{21–23} but knowledge about the roles of HSFs in wheat wild species is still inadequate. A seed preferential HSF; *TaHSA2d* now annotated as *TaHsfA6b* was cloned for the first time in wheat²³ and overexpression of *TaHSA2d* in Arabidopsis leading to increased thermotolerance was reported. In response to heat stress, the induction of *HSFA2*, *HSFA6* in wheat²¹ and high expression of *TaHsfC2a* gene during grain filling and transient induction in the leaves by heat stress has been reported²⁴. In transgenic wheat overexpressing *TaHsfC2a-B*, thermo-protectant genes- *TaHSP70d* and *TaGalSyn* were found to be upregulated.

We have a collection of about 160 accessions of *Ae. speltoides* and preliminary investigations led to the identification of *Ae. speltoides* as a source of heat stress tolerance as evident by its stay-green trait, normal pollen fertility, and normal seed setting at temperatures as high as 39° C⁶. In the present study, we compared comparative gene expression of HSFs among heat-tolerant *Ae. speltoides* accession pau 3809 (AS3809) and two heat susceptible cultivars, PBW725 (hexaploid, bread wheat) and PDW274 (tetraploid, durum wheat) from transcriptome assemblies and validated the selected HSF genes. This analysis provides fundamental insights into HSFs and their variants available in the wild wheat relative *Ae. speltoides*, which could be utilized in wheat improvement programmes for heat stress tolerance.

Results

Transcriptomic data filtering and *de novo* assembly

A comprehensive overview of the transcriptome in three different wheat ploidies yielded 16, 21, and 18 million paired-end reads for AS3809, PBW725, and PDW274, respectively. A summary of statistics of *de novo*

assemblies is given in Table 1. After removing adapter sequences, ambiguous and low-quality reads, the high-quality clean reads (Phred score ≥ 30) led to the assembly of AS3809 with 113,157 trinity transcripts corresponding to 74,946 genes, 121,466 and 109,827 trinity transcripts from PBW725 and PDW274, respectively corresponding to 76,404 and 68,873 genes. N50 values of 1409, 1398, and 1004 bp for AS3809, PBW725, and PDW274 respectively, and average GC content ranging from 47.57 to 48.22%, indicated the good quality of all the assemblies, suitable for further annotation (Table 1).

Assessment of transcriptome assembly completeness

BUSCO analysis revealed that the majority of the Liliopsida core genes were successfully recovered in the AS3809 assembly (Table 2). Specifically, of the 3278 single-copy orthologs searched, 74.3% of core genes were completely recovered in AS3809 as compared to 65.4% and 56.7% in PBW725 and PDW274, respectively. Around 12.1%, 14%, and 16% of genes in AS3809, PBW725, and PDW274, respectively were partially recovered in fragmented form. Only 13.6% of genes were missing from AS3809 assembly as compared to 16.1% and 19.4% missing rate for hexaploid and tetraploid wheat. Given the high quality of the dataset, recovery for both 'complete-single copy' and 'complete-duplicated' BUSCOs was considerably higher for AS3809 assembly as compared to PBW725 and PDW274. Therefore, the AS3809 transcriptome was used as the standard for extracting transcripts related to heat stress tolerance.

Annotation and Gene Ontology analysis

Blast2GO annotation based on the BLASTX results led to functional characterization of 55838, 50077, and 49872 transcripts in AS3809, PBW725, and PDW274, respectively. All the assemblies exhibited a diverse range of GO, suggesting that biological process, molecular function, and cellular component were all well represented (Fig. 1). These three main categories were further divided into 53 GO functional subcategories uniformly across the three transcriptomes. In all the assemblies, genes encoding cellular and metabolic processes along with the response to stimuli were significantly represented in the biological process category while under the molecular function category, the percentage of genes coding for binding and catalytic activities was higher. Within the cellular components, proteins related to cell and cellular parts were expressing more, followed by the cellular membrane and organelles.

Functional terms were also assigned to the trinity transcripts from each wheat assembly using the MapMan Mercator. MapMan employs a basic hierarchical tree structure of terms called "bins" to represent biological contexts and concepts. MapMan annotation resulted in the classification of the transcripts into different bins, each bin representing a different group as photosynthesis, carbohydrate metabolism, glycolysis, gluconeogenesis, oxidative phosphorylation, TCA, cell wall synthesis, lipid and amino acid metabolism, biotic and abiotic stress response, etc. The abiotic stress bin revealed that the majority of the active transcripts were from redox state, respiratory, cell wall, MAPK, defense genes, PR proteins, heat shock proteins, heat shock transcription factors, auxin, hormone signaling, etc. in all three assemblies. The PfamScan categorized transcripts into domains, families, and repeats with all maximum in AS3809 and minimum in PDW274. The annotation statistics obtained for the three assemblies are shown in Table 3.

Identification of orthologous genes

Ortholog transcript detection, based on the OrthoMCL program, demonstrated considerable overlap in transcripts sequences across all three assemblies. Over 39% of the transcripts were identified as putative orthologs between AS3809 and PBW725 assemblies while PDW274 and AS3809 assemblies shared around 37% of the transcripts as putative orthologs. The annotation results from BLAST2GO, MapMan, and PfamScan of the ortholog transcripts were similar suggesting the authenticity of identified ortholog.

Identification of putative heat shock transcription factor (HSF) genes

The three different classes of HSF i.e. A, B, and C were annotated/identified from the transcripts. In total, 37 HSF transcripts were retrieved from AS3809 assembly along with their orthologs from PDW274 and PBW725. Of these, 12 HSFs viz. *HSFA1b*, *HSFA2b*, *HSFA4b*, *HSFA3*, *HSFA5*, *HSFA6b*, *HSFA9*, *HSFB1*, *HSFB3*, *HSFB2a*, *HSFB2b* and *HSFC1b* showing maximum FPKM values were chosen for further analysis. All these 12 HSFs had higher expression in AS3809, followed by PDW274 and the least expression value in PBW725 (Fig. 2). The selected HSF genes have high sequence similarity (95-98%) among three wheat species and possessed HSF DNA binding domain as evident from Pfam analysis. Gene Ontology from Blast2GO and PfamScan annotation of these HSF transcripts are given in Table 4.

In silico expression analysis of HSF genes (tissue-specific and under abiotic stress)

BLASTn search of selected 12 HSF transcripts with the wheatEXP database suggested expression levels of all the HSF genes to be significantly higher in leaf (Z71) tissues. At the basal level, the expression of *HSFA6b* was found to be the highest, followed by *HSFB2b* and *HSFC1b* (Fig. 3). Similarly, at 1 hour of heat stress, the expression of *HSFA6b* was the highest, followed by *HSFB3*, *HSFB1*, and *HSFB2b*, indicating these genes play regulatory functions and be induced in plants within 1 hour of heat stress. At 6 hrs of heat stress, the expression of *HSFB2b* was found the highest amongst all 12 HSFs. But in the expression database, the expression values recorded under prolonged heat stress (6 h) were less than those observed under 1 h of heat stress.

Validation of the expression of HSF genes

Of the 12 selected HSFs, only eight could be validated through qRT-PCR, as four (*HSFA1b*, *HSFA2b*, *HSFA6b*, and *HSFB3*) failed to amplify in two or more selected wheat species (Fig. 4). After 1h of heat stress on seedlings, *HSFA3*, *HSFA5*, *HSFA9*, *HSFB2a*, and *HSFC1b* exhibited the highest expression in the wild wheat species AS3809 while two HSF genes, *HSFA4b* and *HSFB1* exhibited high expression in cultivated wheat, PDW274, and PBW725 respectively. *HSFB2b* amplified in control but did not show expression after heat stress.

Discussion

Heat stress is one of the most serious constraints that limits wheat productivity. The mechanism of heat stress tolerance is a complicated phenomenon governed by numerous genes which cause a variety of physiological and biochemical changes. Wild species always fascinate scientists for having a repertoire of exotic genes/alleles which have almost vanished in cultivated wheat due to domestication and breeding^{8,10,12-14}. Various studies have highlighted the importance of *Ae. speltooides*, *Ae. tauschii* and *Ae. geniculata* being a valuable genetic resource for enhancing thermotolerance^{9,25,26}. Previous studies^{6,27} reported *Ae. speltooides* to

be highly heat-tolerant which can be utilized to enhance the thermotolerance of wheat²⁶. Therefore, our study was designed to compare the whole transcriptome of *Ae. speltoides* with the cultivated tetraploid and hexaploid wheat and study the expression profiling of selected HSFs among three ploidies.

Transcriptome data analysis of wild and cultivated wheat

De novo transcriptome assembly was performed for all three transcriptomes using Trinity, to avoid any bias arising due to the non-availability of reference genomes of *Ae. speltoides* and *T. durum*. Regardless of the huge difference in the genome size (4.9–7 GB)^{28–30} and the number of genes (50,000–1,00,000)^{29,31} among the wild diploid, cultivated tetraploid and hexaploid wheat, the total number of expressed transcripts obtained were 113.1K in AS3809 which is just marginally less than hexaploid wheat PBW725 having 121.4K and tetraploid PDW274 with 109.8K transcripts. The disproportionate trends between genome size and gene expression have been reported in polyploid species owing to the momentous number of repetitive sequences in the non-expressing heterochromatin region of higher ploidies as compared to diploid relatives. Furthermore, loss of some genes/functional alleles during artificial selection and domestication seems to play a direct role in the observed trend^{32,33}.

The N50 values obtained for the three transcriptome assemblies were found to be appropriate for downstream analysis^{34,35}. BUSCO analysis-based completeness assessment of the three transcriptomes revealed the highest recovery of conserved single-copy orthologs in the diploid wild followed by hexaploid and tetraploid wheat, indicating good coverage and high quality of AS3809, relative to higher ploidies, demonstrating loss of genes owing to the domestication process as well as polyploidy events³⁶. Recovery reported for the assemblies was enough for identification of single-copy orthologs³⁷ thus the current assemblies of the cultivated as well as wild species of wheat will supplement the published resources for wheat. BLAST2GO and MapMan based comprehensive annotation revealed that GO representations for all three categories of molecular function, biological processes, and cellular component were successfully captured representing GO profiles with the highest proportions of mapped GO terms³⁸.

Identification and validation of HSFs

AS3809 shared 39% and 37% putative orthologs with PBW725 and PDW274 respectively. The transcriptome data have been analyzed for differential expression of putative HSF gene candidates for heat stress tolerance. HSFs play an imperative role in acquired thermotolerance as they are the terminal components of the signal transduction chain mediating the activation of genes responsive to heat stress¹⁶. The 12 important HSF genes under study had the highest FPKM in wild wheat indicating higher expression in AS3809 as compared to cultivated durum and hexaploid wheat. These genes are primarily involved in the rapid synthesis and accumulation of heat shock proteins (HSPs), which not only act as molecular chaperones shielding proteins from thermal aggregation but are also involved in various aspects of proteins homeostasis, such as protein translocation and degradation³⁹.

Of the 12 important HSF genes with highest expression in AS3809, eight genes viz *HSFA3*, *HSFA4b*, *HSFA5*, *HSFA9*, *HSFB1*, *HSFB2a*, *HSFB2b* and *HSFC1b* could be validated by qRT-PCR, (Fig. 4). Of these 6 genes *HSFA3*, *HSFA5*, *HSFA9*, *HSFB2a*, *HSFB2b*, and *HSFC1b* showed comparative highest expression in the wild AS3809 as

proven in comparative transcriptome data. Five of these HSF genes were also validated from the WheatExp database from which *HSFB2b*, exhibited the highest expression at basal level as well as after 1h and 6 h of heat stress, *HSFA6b*, at basal level and after 1 hr of heat stress, *HSFB1* & *HSFB3* after 1h of heat stress and *HSFC1b* having higher expression at a basal level only. All HSF genes in the wheatEXP database revealed that the expression values recorded under prolonged heat stress of 6 hr were less than those observed under 1 h of heat stress suggesting the HSF gene family be quickly induced within 1 h of heat stress, activating downstream pathways for thermotolerance as compared to 6 h of heat stress, thus allowing a quick and elevated response system of plants to combat heat stress. *HSFC1b* and *HSFB2b* might have a key regulatory role in thermotolerance in AS3809. Higher expression of *HSFC1b* in transcriptome data and qRT-PCR and wheatExp database results indicating this to be important HSF in AS3809. *HSFB2b* could be another important candidate though it expressed only under control conditions in AS3809 and with a decreased expression upon 1 h of heat stress. The early heat-responsive nature of *HSFA6b* and *HSFB2b* was observed to be upregulated within 10 min of heat stress onset in barley and *HSFA6b* also showed considerable tolerance to salinity and drought stresses^{23,40,41}. Decrease in expression of *TaHsfB2b* under 1.5 h of heat stress²¹ as compared to control conditions suggests that this gene performs well under non-stress conditions but its transcript level decreases after heat stress. *FaHsfC1b* gene from *Festuca arundinacea* in *Arabidopsis thaliana*, plays a positive role in heat stress by upregulating heat-protective genes⁴². Upregulation of *HsfC1* genes in rice (a diploid species) by heat and oxidative stresses or a combination of these stresses⁴³ corroborating with our findings from this experiment.

HSFA3, *HSFA5*, *HSFA9*, *HSFB2a* genes with higher expression in qRT-PCR results were also an important part of the thermotolerance profile of AS3809 though we could not find their expression in the wheatExp database. The data in the wheatExp database is from hexaploid wheat cultivar Chinese spring and TAM107 and the candidate HSFs specific to the S genome of *Ae. speltoides* could not be represented. Overexpression of homologous DREB2A from *Zea mays* in *Arabidopsis thaliana* led to induction of HS-related genes, including *HSFA3*⁴⁴. Downregulation of *HSFA5* in hexaploid and tetraploid wheat under heat stress has also been reported²¹. A9 of class A HSFs and subclass B3, B5 of class B HSFs is absent in monocot species like wheat and rice²⁰. Contrastingly in the current investigation, *HSFA9* transcript was found in all three assemblies and is also validated by real-time PCR reaction. The results revealed the highest upregulation of this gene in AS3809 followed by PBW725 with a marginal difference. However, the role of *HSFA9* in regulating HSP expression during seed development has been demonstrated in *Arabidopsis* and sunflower⁴⁵ though no report of seedling expression has been known. The overexpression of wheat *TaHsfB2a* in transgenic *Arabidopsis* leads to enhanced thermo and freeze tolerance⁴⁶.

Two of the candidate HSFs, *HSFB1*, and *HSFA4b* exhibited the highest expression under heat stress in cultivated wheat, PDW274, and PBW725, respectively indicating the species-specific expression. Subclass B1 has been reported to be highly heat-inducible and its overexpression activated the expression promoter-driven reporter genes under optimal conditions²¹.

Differences in expression profiling of diploid, tetraploid, and hexaploid wheat

Significant differences were observed in the transcriptome expression profiling of allopolyploid wheat species, *T. durum*, and *T. aestivum* as compared to one of the wild diploid progenitors, *Ae. speltoides*. The highest

expression of heat stress transcription factor genes was also observed in the diploid wild wheat, in contrast to the cultivated polyploids. Similarly, recent comparative studies⁴⁷ focused on the expression analysis of gene pairs in the syntenic regions between hexaploid wheat chromosome 3DL and its progenitor 3L arm of *Aegilops tauschii* demonstrated 60% decreased gene expression in 70% of gene pairs in hexaploid context of the 3DL genes. Such a reduction in gene expression has been attributed to altered interactions between transcription factors and nucleosomes on the introduction of new genomes leading to alteration in chromosome accessibility. Another study based on analyzing models of homeolog expression patterns has also demonstrated inter-genome interactions to play a key role in altered gene expression⁴⁸. Thus, polyploidization events bring about changes in mode of gene action that varies from additive, non-additive and dominant type in addition to the epigenetic and change in transposon activity. The expression changes observed in allopolyploids and the wild progenitor may be due to gene repression, genetic dominance, sub-functionalization, and novel activation in the former relative to the latter. The HSF genes that have been identified and validated in current study suggest these genes play positive roles in regulating thermotolerance, especially in the wild wheat. These genes are of breeding importance as their introgression into cultivars will improve their thermotolerance and thus avoiding the yield penalty due to heat stress, ultimately leading to better productivity.

Conclusion

We generated important genomic resource of transcriptome representing differential gene expression among three ploidies. Study highlights the conservation of gene expression across the three species as well loss of functional alleles and inter genome interactions during domestication and polyploidy as major reason behind reduction of expression in higher ploidies in contrast to their diploid relatives. We report the eight important HSFs in the comparative RNA-Seq analysis of three wheat species of diploid *Ae. speltooides*, tetraploid *T. durum* PDW274, and hexaploid *T. aestivum* PBW725, of which six were validated as potential novel candidates for thermotolerance from *Ae. speltooides*. Transcript profiling of HSFs under basal and heat stress conditions revealed a good consistency between the expression levels of the eight genes analyzed by qRT-PCR and their transcript levels detected using RNA-Seq in all the three wheat species. Our results encourage the exploitation of novel alleles from *Ae. speltooides* and other wild relatives for broadening the narrow genetic base of cultivated wheat.

Materials And Methods

Plant material

Plant material included diploid wild wheat *Ae. speltooides* accession pau3809, a heat-tolerant accession identified in Punjab Agricultural University, and the two cultivated wheat varieties, tetraploid *T. durum* cv. PDW274 and hexaploid *T. aestivum* cv. PBW725 which are susceptible to heat stress. The seeds of the PDW274 and PBW725 have been developed and maintained by Punjab Agricultural University (PAU), Ludhiana, India. The seeds of *Aegilops speltooides* accession pau3809 were procured from the Weizmann Institute of Science, Rehovot, maintained by PAU and have also been deposited in National Bureau of Plant Genetic Resources (NBPGR), New Delhi, India. Since the plant material has been maintained by PAU, Ludhiana, India,

permissions regarding the collection of seed specimen was not required. The three wheat genotypes under study were sown in the field in three replications.

RNA extraction and Illumina Sequencing

The 10-12 days old seedlings of the three selected wheat lines were taken and immediately snap-frozen in liquid nitrogen and stored at -80°C for further use. Seedlings from three different plants were pooled to form one biological replicate. Total RNA from two biological replicates was extracted from the frozen seedling tissue samples using RNA isolation kit (Qiagen) as per manufacturer protocols. The RNA concentration and purity were determined by Nanodrop™ 1000 Spectrophotometer (Thermo Scientific). Only high-quality RNA samples with OD 260/280 ranging from 1.8 to 2.2 and RIN (RNA integrity number) values ranging from 7.4 to 10.0 were used to construct the RNA-Seq library. Six separate libraries consisting of two biological replicates of the three genotypes were prepared and outsourced for transcriptome sequencing. Indexed TruSeq libraries were prepared for the six RNA samples and 100bp paired-end sequencing was performed using a HiSeq™ 2000 platform.

Transcriptomic data filtering and *de novo* assembly

The quality of reads was assessed using the FastQC toolkit⁴⁹. The adaptor sequences and low-quality bases (phred score < 30) were removed from the raw reads using the Trimmomatic tool⁵⁰. *De novo* transcriptome assembly was performed for all the three transcriptomes using Trinity software version 2.8.4⁵¹. For each of the three *de novo* assemblies, the transcript abundance was computed by RSEM which estimated the number of RNA-Seq fragments corresponding to each Trinity transcript, including normalized expression values as FPKM (fragments per kilo base of target transcript length per million reads mapped). The assembled transcripts from each assembly were also searched for sequence homology by performing standalone BLASTX with e-value of 1e-5 against publicly available NR (NCBI non-redundant protein sequences) database. A schematic representation of the *de novo* transcriptome reconstruction and analysis pipeline is shown in Fig. 5.

Assessment of transcriptome assembly completeness

The comprehensive and quantitative level of completeness of transcriptome assemblies was assessed by comparing the three assembled transcript sets to a set of highly conserved single-copy orthologs using the BUSCO (Benchmarking Universal Single-Copy Orthologs) v2 pipeline⁵². This compared the assembled transcripts to the predefined set of 3278 Liliopsida single-copy orthologs from the OrthoDB v10 database. The number of complete, duplicated, fragmented, and missing BUSCOs were obtained in each of the three assemblies.

Prediction of protein-coding regions and domains

TransDecoder v5.5.0 (<http://transdecoder.github.io/>) was used to identify all likely coding regions in the three assemblies, and the single best open reading frame (ORF) per transcript was selected as per the TransDecoder pipeline (-single best orf). The protein sequence file from TransDecoder was used to predict the protein structure domains and families of all transcripts by searching against the Protein family (Pfam) database using the HMM-based tool PfamScan⁵³.

Functional annotation of whole transcriptome assemblies

The identified transcripts in each of the assemblies were annotated based on corresponding homologs identified from the BLASTX program against the NCBI protein database 'NR' at an e-value of $1e^{-5}$. Based on the NR annotations, the Blast2GO program was used to obtain Gene Ontology (GO) annotations for genes in biological process, molecular function, and cellular component category for each assembly separately⁵⁴. Mercator v3.6 (www.plabipd.de/portal/mercator-sequence-annotation) was used to assign functional terms to nucleotide sequences using the MapMan 'BIN' ontology. MapMan was used for functional analysis and visualization⁵⁵. The comparative GO terms for molecular function, biological process, and cellular component for each of the three assemblies were plotted using wego⁵⁶.

Identification of orthologous genes

OrthoMCL program was used for constructing orthologous groups amongst the three wheat assemblies, using the Markov Cluster algorithm to identify (putative) orthologs and paralogs³¹. OrthoMCL was conducted for all pair-wise comparisons among the three assemblies. The output of OrthoMCL was used to determine the number of overlapping (shared across species) transcripts across the three assemblies. Further, the transcripts that were annotated/identified as heat stress transcription factors (HSFs) belonging to classes A, B, and C⁵⁷ were retrieved from transcriptome assembly of AS3809 along with their corresponding orthologs from PBW725 and PDW274. Blast2sequence and multiple sequence alignment with ClustalX were performed to see the extent of homology between HSFs of different wheat species.

In silico expression analysis of genes

The expression patterns of selected HSF transcripts were investigated using wheat transcriptome data from the wheatExp database. This database comprises RNA-Seq datasets derived from five different tissues (spike, root, leaf, grain, and stem) of hexaploid bread wheat variety Chinese Spring each sampled at three different developmental stages⁵⁸. Another dataset consists of one-week old seedlings of hexaploid wheat variety TAM107 treated with heat (40°C) for 1 h and 6 h⁵⁹. The 12 selected HSF transcripts were BLASTn searched with the wheat transcriptome database with an expected cut-off of $1e^{-5}$.

Validation of HSF genes by Quantitative Real-Time PCR (qRT-PCR)

The seeds of *Ae. speltooides* acc. pau3809, *T. aestivum* cv. PBW725 and *T. durum* cv. PDW274 were sown in germination trays in a growth chamber maintained at 22°C /15°C Day: night temperature, 16:8 h photoperiod. Then heat stress was given to the 10 days old seedlings at 35°C for 1 hour and immediately after stress RNA from control and heat-stressed plants was extracted for performing qRT-PCR. Primers for the 12 selected HSFs were generated using Primer Express software v3.0.1 following the default parameters and are listed in Supplementary Table 1. Since there was considerable homology (98%) in the active coding region of the HSF genes across the three wheat species, we used the same primers for all three species. *TaActin* gene⁶⁰ was used as the internal control. The relative gene expression levels for RT-qPCR data were calculated using the $2^{-\Delta\Delta Ct}$ method⁶¹. All reactions were conducted in triplicate.

Declarations

Author Contributions

SK and PC - conceived and designed research, reviewed the manuscript, SK - planned the experiments, SS- conducted all analysis of transcriptome data, identification of HSF and primer designing, ISY- helped in analysis of transcriptome data, HC and PS - designed and conducted the real-time experiment. SK, SS, PM, and AK- compiled the work and produced the final draft of the manuscript

Declarations

Ethics approval: NA

Consent to participate: NA

Competing interests: The authors declare no competing interests

Availability of data and material: NA

References

1. Akter, N. & Rafiqul Islam, M. Heat stress effects and management in wheat. A review. *Agron. Sustain. Dev.* **37**, 37 (2017).
2. Arora, S., Cheema, J., Poland, J., Uauy, C. & Chhuneja, P. Genome-Wide Association Mapping of Grain Micronutrients Concentration in *Aegilops tauschii*. *Front. Plant Sci.* **10**, 54 (2019).
3. *Climate change 2007: the physical science basis: contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. (Cambridge University Press, 2007).
4. Wollenweber, B., Porter, J. R. & Schellberg, J. Lack of Interaction between Extreme High-Temperature Events at Vegetative and Reproductive Growth Stages in Wheat. *Journal of Agronomy and Crop Science* **189**, 142–150 (2003).
5. Gupta, P. K., Balyan, H. S., Gahlaut, V. & Kulwal, P. L. Phenotyping, Genetic Dissection, and Breeding for Drought and Heat Tolerance in Common Wheat: Status and Prospects. in *Plant Breeding Reviews* (ed. Janick, J.) 85–168 (John Wiley & Sons, Inc., 2012). doi:10.1002/9781118358566.ch2.
6. Awlachev, Z. T., Singh, R., Kaur, S., Bains, N. S. & Chhuneja, P. Transfer and mapping of the heat tolerance component traits of *Aegilops speltoides* in tetraploid wheat *Triticum durum*. *Mol Breeding* **36**, 78 (2016).
7. Li, J., Wan, H. & Yang, W. Synthetic hexaploid wheat enhances variation and adaptive evolution of bread wheat in breeding processes. *Jnl of Sytematics Evolution* **52**, 735–742 (2014).
8. Farooq, S. Co-Existence of Salt and Drought Tolerance in Triticeae. *Hereditas* **135**, 205–210 (2004).
9. Waines, J. High Temperature Stress in Wild Wheats and Spring Wheats. *Functional Plant Biol.* **21**, 705 (1994).

10. Jafarzadeh, J. *et al.* Breeding Value of Primary Synthetic Wheat Genotypes for Grain Yield. *PLoS ONE* **11**, e0162860 (2016).
11. Colmer, T. D., Flowers, T. J. & Munns, R. Use of wild relatives to improve salt tolerance in wheat. *Journal of Experimental Botany* **57**, 1059–1078 (2006).
12. Vikas, V. K. *et al.* Stem and leaf rust resistance in wild relatives of wheat with D genome (*Aegilops* spp.). *Genet Resour Crop Evol* **61**, 861–874 (2014).
13. Huang, M. *et al.* Genetic analysis of heading date in winter and spring wheat. *Euphytica* **214**, (2018).
14. Olivera, P. D., Rouse, M. N. & Jin, Y. Identification of New Sources of Resistance to Wheat Stem Rust in *Aegilops* spp. in the Tertiary Genepool of Wheat. *Front. Plant Sci.* **9**, 1719 (2018).
15. von Koskull-Döring, P., Scharf, K.-D. & Nover, L. The diversity of plant heat stress transcription factors. *Trends in Plant Science* **12**, 452–457 (2007).
16. Baniwal, S. K. *et al.* Heat stress response in plants: a complex game with chaperones and more than twenty heat stress transcription factors. *J. Biosci.* **29**, 471–487 (2004).
17. Nover, L., Bharti, K. & Scharf, K.-D. Arabidopsis and the heat stress transcription factor world: how many heat stress transcription factors do we need? *Cell Stress* **17** (2001).
18. Fragkostefanakis, S., Röth, S., Schleiff, E. & Scharf, K.-D. Prospects of engineering thermotolerance in crops through modulation of heat stress transcription factor and heat shock protein networks: Hsfs and Hsps for improvement of crop thermotolerance. *Plant Cell Environ* **38**, 1881–1895 (2015).
19. Guo, M. *et al.* Genome-wide analysis, expression profile of heat shock factor gene family (CaHsfs) and characterisation of CaHsfA2 in pepper (*Capsicum annuum* L.). *BMC Plant Biol* **15**, 151 (2015).
20. Scharf, K.-D., Berberich, T., Ebersberger, I. & Nover, L. The plant heat stress transcription factor (Hsf) family: Structure, function and evolution. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **1819**, 104–119 (2012).
21. Xue, G.-P., Sadat, S., Drenth, J. & McIntyre, C. L. The heat shock factor family from *Triticum aestivum* in response to heat and other major abiotic stresses and their role in regulation of heat shock protein genes. *EXBOTJ* **65**, 539–557 (2014).
22. Agarwal, P. & Khurana, P. Functional characterization of HSFs from wheat in response to heat and other abiotic stress conditions. *Funct Integr Genomics* **19**, 497–513 (2019).
23. Chauhan, H., Khurana, N., Agarwal, P., Khurana, J. P. & Khurana, P. A Seed Preferential Heat Shock Transcription Factor from Wheat Provides Abiotic Stress Tolerance and Yield Enhancement in Transgenic Arabidopsis under Heat Stress Environment. *PLoS ONE* **8**, e79577 (2013).
24. Hu, Q. *et al.* Meiotic Chromosome Association 1 Interacts with TOP3α and Regulates Meiotic Recombination in Rice. *The Plant Cell* **29**, 1697–1708 (2017).

25. Zaharieva, M., Gaulin, E., Havaux, M., Acevedo, E. & Monneveux, P. Drought and Heat Responses in the Wild Wheat Relative *Aegilops geniculata* Roth: Potential Interest for Wheat Improvement. *Crop Sci.* **41**, 1321–1329 (2001).
26. Pradhan, G. P., Prasad, P. V. V., Fritz, A. K., Kirkham, M. B. & Gill, B. S. High Temperature Tolerance in *Aegilops* Species and Its Potential Transfer to Wheat. *Crop Sci.* **52**, 292–304 (2012).
27. Jakhu, P. *et al.* Cloning, expression analysis and In silico characterization of HSP101: a potential player conferring heat stress in *Aegilops speltoides* (Tausch) Gren. *Physiol Mol Biol Plants* **27**, 1205–1218 (2021).
28. Salse, J. *et al.* New insights into the origin of the B genome of hexaploid wheat: Evolutionary relationships at the SPA genomic region with the S genome of the diploid relative *Aegilops speltoides*. *BMC Genomics* **9**, 555 (2008).
29. Borrill, P., Adamski, N. & Uauy, C. Genomics as the key to unlocking the polyploid potential of wheat. *New Phytol* **208**, 1008–1022 (2015).
30. Ruban, A. S. & Badaeva, E. D. Evolution of the S-Genomes in Triticum-Aegilops Alliance: Evidences From Chromosome Analysis. *Front. Plant Sci.* **9**, 1756 (2018).
31. El Baidouri, M. *et al.* Reconciling the evolutionary origin of bread wheat (*Triticum aestivum*). *New Phytol* **213**, 1477–1486 (2017).
32. Appels, R. *et al.* Shifting the limits in wheat research and breeding using a fully annotated reference genome. *Science* **361**, (2018).
33. Miki, Y. *et al.* Origin of wheat B-genome chromosomes inferred from RNA sequencing analysis of leaf transcripts from section Sitopsis species of *Aegilops*. *DNA Research* **26**, 171–182 (2019).
34. Liu, X. *et al.* The rice enhancer of zeste [E(z)] genes SDG711 and SDG718 are respectively involved in long day and short day signaling to mediate the accurate photoperiod control of flowering time. *Frontiers in Plant Science* **5**, (2014).
35. Yadav, I. S. *et al.* Comparative Temporal Transcriptome Profiling of Wheat near Isogenic Line Carrying Lr57 under Compatible and Incompatible Interactions. *Front. Plant Sci.* **7**, (2016).
36. Chen, Z. J. & Ni, Z. Mechanisms of genomic rearrangements and gene expression changes in plant polyploids. *Bioessays* **28**, 240–252 (2006).
37. Kumar Kushwaha, S. *et al.* Differential Gene Expression Analysis of Wheat Breeding Lines Reveal Molecular Insights in Yellow Rust Resistance under Field Conditions. *Agronomy* **10**, 1888 (2020).
38. Kumar, R. R. *et al.* Harnessing Next Generation Sequencing in Climate Change: RNA-Seq Analysis of Heat Stress-Responsive Genes in Wheat (*Triticum aestivum* L.). *OMICS: A Journal of Integrative Biology* **19**, 632–647 (2015).

39. Schleiff, E. & Becker, T. Common ground for protein translocation: access control for mitochondria and chloroplasts. *Nat Rev Mol Cell Biol* **12**, 48–59 (2011).
40. Xue, G.-P., Drenth, J. & McIntyre, C. L. TaHsfA6f is a transcriptional activator that regulates a suite of heat stress protection genes in wheat (*Triticum aestivum* L.) including previously unknown Hsf targets. *Journal of Experimental Botany* **66**, 1025–1039 (2015).
41. Mishra, S. K. *et al.* Genome-wide identification, phylogeny and expression analysis of HSF gene family in barley during abiotic stress response and reproductive development. *Plant Gene* **23**, 100231 (2020).
42. Zhuang, L. *et al.* Characterization and Functional Analysis of FaHsfC1b from *Festuca arundinacea* Conferring Heat Tolerance in Arabidopsis. *IJMS* **19**, 2702 (2018).
43. Mittal, D., Madhyastha, D. A. & Grover, A. Gene expression analysis in response to low and high temperature and oxidative stresses in rice: Combination of stresses evokes different transcriptional changes as against stresses applied individually. *Plant Science* **197**, 102–113 (2012).
44. Qin, F. *et al.* Regulation and functional analysis of ZmDREB2A in response to drought and heat stresses in *Zea mays* L: ZmDREB2A in drought and heat stress response. *The Plant Journal* **50**, 54–69 (2007).
45. Kotak, S. *et al.* Complexity of the heat stress response in plants. *Current Opinion in Plant Biology* **10**, 310–316 (2007).
46. Zhang, S. *et al.* Overexpression of TaHSF3 in Transgenic Arabidopsis Enhances Tolerance to Extreme Temperatures. *Plant Mol Biol Rep* **31**, 688–697 (2013).
47. Lu, F.-H. *et al.* Reduced chromatin accessibility underlies gene expression differences in homologous chromosome arms of diploid *Aegilops tauschii* and hexaploid wheat. *GigaScience* **9**, giaa070 (2020).
48. Hu, G. & Wendel, J. F. *Cis – trans* controls and regulatory novelty accompanying allopolyploidization. *New Phytol* **221**, 1691–1700 (2019).
49. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol* **11**, R106 (2010).
50. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).
51. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc* **8**, 1494–1512 (2013).
52. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212 (2015).
53. Finn, R. D. *et al.* Pfam: the protein families database. *Nucl. Acids Res.* **42**, D222–D230 (2014).

54. Conesa, A. & Götz, S. Blast2GO: A Comprehensive Suite for Functional Analysis in Plant Genomics. *International Journal of Plant Genomics* **2008**, 1–12 (2008).

55. Thimm, O. *et al.* mapman: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *The Plant Journal* **37**, 914–939 (2004).

56. Ye, J. *et al.* WEGO: a web tool for plotting GO annotations. *Nucleic Acids Research* **34**, W293–W297 (2006).

57. Doring, P. *et al.* The Role of AHA Motifs in the Activator Function of Tomato Heat Stress Transcription Factors HsfA1 and HsfA2. *The Plant Cell* **12**, 265 (2000).

58. Choulet, F. *et al.* Structural and functional partitioning of bread wheat chromosome 3B. *Science* **345**, 1249721–1249721 (2014).

59. Liu, Z. *et al.* Temporal transcriptome profiling reveals expression partitioning of homeologous genes contributing to heat and drought acclimation in wheat (*Triticum aestivum* L.). *BMC Plant Biol* **15**, 152 (2015).

60. Risk, J. M. *et al.* The wheat *Lr34* gene provides resistance against multiple fungal pathogens in barley. *Plant Biotechnol J* **11**, 847–854 (2013).

61. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2– $\Delta\Delta$ CT Method. *Methods* **25**, 402–408 (2001).

Tables

Table 1: Summary of paired-end sequencing data (2 x 100 bp) and *de novo* transcriptome assembly statistics for *Ae. speltoides* acc. pau3809, *T. aestivum* cv. PBW725 and *T. durum* cv. PDW274

Feature	<i>Aegilops speltoides</i> acc. pau3809	<i>Triticum aestivum</i> cv. PBW725	<i>Triticum durum</i> cv. PDW274
Total number of raw paired-end reads	16,991,059	21,158,755	18,808,576
Paired-end reads obtained after filtering	13,397,450 (78.85%)	16,480,554 (77.89%)	13,824,303 (73.50%)
Number of trinity transcripts	113,157	121,466	109,827
Number of trinity ‘genes’	74,946	76,404	68,873
GC content (%)	47.57	48.11	48.22
Contig N50 (bp)	1409	1398	1004
Average contig length (bp)	809.01	850.16	796.23

Table 2: Summary of the complete, duplicated, fragmented and missing orthologs inferred from BUSCO searched against the 3278 single-copy orthologs for Liliopsida

BUSCO statistics	<i>Aegilops speltoides</i> acc. pau 3809 (AS3809)	<i>Triticum aestivum</i> (PBW725)	<i>Triticum durum</i> (PDW274)
Complete BUSCOs (C)	2436 (74.3%)	2144 (65.4%)	1859 (56.7%)
Complete and single-copy BUSCOs (S)	1534 (46.8%)	1373 (41.9%)	1272 (38.8%)
Complete and duplicated BUSCOs (D)	901 (27.5%)	754 (23.0%)	679 (20.7%)
Fragmented BUSCOs (F)	396 (12.1%)	459 (14.0%)	524 (16.0%)
Missing BUSCOs (M)	445 (13.6%)	528 (16.1%)	636 (19.4%)

Table 3: Summary of transcripts with significant BLAST2GO alignments and annotation using annotation tools of MapMan Mercator and PfamScan

Annotation	<i>Ae. speltoides</i> acc. pau3809	<i>T. aestivum</i> cv. PBW725	<i>T. durum</i> cv. PDW274
BLAST2GO	55,838	50,077	49,872
MapMan mercator annotations			
Total mapped transcripts	103,683	118,104	99,565
Transcript specific for abiotic stress	772	823	689
Pfam Scan based on HMM			
Number of Domains	26,386	24,992	19,321
Number of Families	19,748	18,820	16,242
Number of Repeats	5,127	4,239	3,403

Table 4: List of Trinity IDs of 12 HSFs selected from *Aegilops speltoides* acc. pau3809 transcriptome assembly with maximum FPKM value along with their GO annotation

S. No.	AS3809 transcript ID	HSF subtype	Length (bp)	FPKM value	GO annotation	PfamScan annotation
1.	TRINITY_DN7915_c1_g2_i3	HSFA2B	688	15.51	AHB61248.1heat shock factor HsfA2b	HSF_DNA-bind (Domain)
2.	TRINITY_DN1997_c0_g4_i2	HSFA1B	2257	237.36	AHZ44763.1heat shock factor A1b	HSF_DNA-bind (Domain)
3.	TRINITY_DN7915_c1_g1_i1	HSFA4B	2357	58.73	XP_020165982.1heat stress transcription factor A-4b	HSF_DNA-bind (Domain)
4.	TRINITY_DN49987_c0_g1_i1	HSFA3	660	55	XP_020146599.1heat stress transcription factor A-3-like	HSF_DNA-bind (Domain)
5.	TRINITY_DN16884_c0_g1_i1	HSFA5	1562	56	XP_020200706.1heat stress transcription factor A-5	HSF_DNA-bind (Domain)
6.	TRINITY_DN4362_c0_g1_i10	HSFA6B	2640	77	AHZ44769.1heat shock factor A6b	HSF_DNA-bind (Domain)
7.	TRINITY_DN3035_c0_g1_i2	HSFA9	1521	62	XP_020148069.1heat stress transcription factor A-9-like	HSF_DNA-bind (Domain)
8.	TRINITY_DN4144_c0_g1_i2	HSFB1	1280	71.4	XP_020177659.1heat stress transcription factor B-1-like	HSF_DNA-bind (Domain)
9.	TRINITY_DN4144_c0_g1_i5	HSFB3	2471	60.23	BAJ96942.1predicted protein	HSF_DNA-bind (Domain)
10.	TRINITY_DN4527_c0_g1_i1	HSFB2A	1628	72	AJR10106.1heat-responsive transcription factor	HSF_DNA-bind (Domain)
11.	TRINITY_DN4527_c0_g2_i1	HSFB2B	1800	88	XP_020163127.1heat stress transcription factor B-2b-like	HSF_DNA-bind (Domain)
12.	TRINITY_DN32883_c0_g1_i1	HSFC1B	786	49	XP_020179067.1heat stress transcription factor C-1b-like	HSF_DNA-bind (Domain)

Figures

(a)

(b)

(c)

Figure 1

Proportion of gene ontology annotations made using WEGO for transcripts of (a) *Aegilops speltoides* acc. pau3809, (b) *T. aestivum* cv. PBW725 and (c) *T. durum* cv. PDW274 representing cellular component, molecular function, and biological process.

Figure 2

Heat map representing the different classes of HSF transcripts having a similar function in AS3809, PBW725, and PDW274 (red: maximum expression, black: median, green: minimum expression).

Figure 4

Expression analysis of HSF genes in response to control and high-temperature stress in three species of wheat (AS3809, PBW725 and PDW274). Change in transcript abundance of HSF genes in 10 days old seedlings of wheat species of different ploidy levels subjected to heat stress at 35°C for 1 h. (a) HSFA3 (b) HSFA4b (c) HSFA5 (d) HSFA9 (e) HSFB1 (f) HSFB2a (g) HSFB2b (h) HSFC1b. The expression level at control (AS3809) i.e. 0 h of treatment was normalized as 1.0. The results shown are the means $\pm$ SE of three independent experiments.

Figure 5

Schematic representation of de novo transcriptome reconstruction and analysis pipeline used to generate whole transcriptome assemblies for *Aegilops speltoides* acc. pau3809, *Triticum aestivum* cv. PBW725 and *Triticum durum* cv. PDW274.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [July26Supplementary.docx](#)