Comparison of Surface Wind Speed and Wind Speed Profiles in the Taklimakan Desert

Liu Xinchun

Institute of Desert Meteorology

kang yongde (✉ kyd0115@yeah.net)

Xi an university of technology

Chen Hongna

Institute of Desert Meteorology

Lu Hui

Guangxi Normal University

Research Article

Keywords: Desert hinterland, Surface layer, Diurnal variation, Profile

Posted Date: July 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-723326/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Comparison of Surface Wind Speed and Wind Speed Profiles in the Taklimakan Desert

Liu Xinchun¹, Kang Yongde^{2*}, Chen Hongna³, Lu Hui⁴

1. Institute of Desert Meteorology, CMA, Taklimakan Desert Meteorology Field Experiment Station of CMA, Xinjiang Laboratory of Tree Ring Ecology, Key Laboratory of Tree-ring Physical and Chemical Research of China Meteorological Administration, Urumqi 830002, China; liuxch@idm.cn;

2. State Key Laboratory of Eco-hydraulics in Northwest Arid Region of China, School of Water Resources and Hydroelectric Engineering, Xi'an University of Technology, Xi'an 710048, China; kyd0115@yeah.net

3. Department, Urumqi Environmental Monitoring Center, Urumqi 830001, China; liux-inchun2001@163.com

4. Key Laboratory of Ecology of Rare and Endangered Species and Environmental Protection (Guangxi Normal University), Ministry of Education, Guilin 541004, China; College of Environment and Resources, Guangxi Normal University, Guilin 541004, China

Abstract:

Near-surface (10 m) wind speed (NWS) plays a crucial role in many areas, including the hydrological cycle, wind energy production, and the dispersion of air pollution. Based on wind speed data from Tazhong and the northern margins of the Taklimakan Desert in Xiaotang in spring, summer, autumn, and winter of 2014 and 2015, statistical methods were applied to determine the characteristics of the diurnal changes in wind speed near the ground and the differences in the wind speed profiles between the two sites. The average wind speed on a sunny day increased slowly with height during the day and rapidly at night. At heights below 4 m the wind speed during the day was higher than at night, whereas at 10 m the wind speed was lower during the day than at night. The semi-empirical theory and Monin-Obukhov (M-O) similarity theory were used to fit the NWS profile in the hinterland of the Tazhong Desert. A logarithmic law was applied to the neutral stratification wind speed profile, and an exponential fitting correlation was used for non-neutral stratification. The more unstable the stratification, the smaller the n . Using M-O similarity theory, the "linear to tens of" law was applied to the near-neutral stratification. According to the measured data, the distribution of ϕ_m with stability was obtained. The γ_m was obtained when the near-surface stratum was stable in the hinterland of Tazhong Desert and the β_m was obtained when it was unstable. In summer, γ_m and β_m were 5.84 and 15.1, respectively, while in winter, γ_m and β_m were 1.9 and 27.1, respectively.

Keywords: Desert hinterland; Surface layer; Diurnal variation; Profile

1. Introduction

Near-surface (10 m) wind speed (NWS) and temperature are important parameters for studying atmospheric dynamics and climate change. Research on wind speed and temperature changes will improve our understanding of atmospheric circulation, leading to better climate analyses and predictions. The NWS is one of the key variables in climate research. Changes in the NWS have

significant implications for human society and the natural environment (Pryor et al., 2006). The intensification of NWS may aggravate soil erosion, resulting in more severe sandstorms (Alizadeh-Chooabari et al., 2014; Wang et al., 2017). Over the past decade, scientists have conducted studies to determine the trends in wind speed at the height of 10 m near the surface (McVicar et al., 2012). Globally, wind speeds have recovered slightly since the 2010s, particularly in Central and East Asia (Kim and Paik, 2015; Dunn et al., 2016; Azorin-Molina et al., 2017a; Ming, H, 2019). The observed upward trend in the global ocean surface has revealed the complexity of decades of wind speed changes (Wentz et al., 2007; Tokinaga and Xie, 2011; Young et al., 2011). In addition, variations in wind speed above and below the boundary layer suggest many uncertainties behind the stationary phenomenon. As a product of the atmospheric boundary layer, near-surface wind is transient in nature and is affected by topography and boundary layer processes (Mahrt, 2009; Belušić and Güttler, 2010; Güttler and Belušić, 2012).

The stability of the near-surface layer has a significant relationship with the near-surface wind. Previous studies have shown that near-surface stability was comprehensively affected by surface and boundary layer processes. In a stable stratified boundary layer environment, the net downward flow of warm air leads to an increase in surface temperature. In the daytime, when the rate of decline is negative, the net downward transport of cold air will cause a ground cooling effect.

Studies of the atmospheric boundary layer in arid regions are an important part of climate research in arid regions (Li et al., 2017). Similarity theory is one of the most important analysis and research methods in atmospheric boundary layer meteorology. After establishing the relatively mature Monin-Obukhov (M-O) similarity theory for describing the atmospheric near-surface layer, researchers have attempted to develop a similar similarity theory for the whole boundary layer. The atmospheric boundary layer is also the main characteristic quantity for assessing turbulent mixing, vertical disturbance, convective transmission, cloud belts, atmospheric pollutant diffusion, and analyzing atmospheric environmental capacity (Therry and Lacarrere, 1983; Holtslag and Boville, 1993; Hong and Pan, 1996; Beyrich, 1997; Collier et al., 2005). Earlier studies of the boundary layer mainly focused on the near-surface layer. The development of turbulence theory and technological developments in measuring atmospheric processes have promoted research on the atmospheric boundary layer (Monin and Obukhov, 1954; Johnson, B.D.2021). The structure of the atmospheric boundary layer and its evolution have significant diurnal characteristics. After sunrise, solar radiation heats the ground. The increase in the heat flux generated by the near-surface layer strengthens turbulent mixing, the height of the atmospheric boundary layer increases, and the heat content of the boundary layer also increases. It mixes uniformly and the wind speed, temperature, and specific humidity change little with height. This is referred to as the mixed layer. After sunset, long-wave radiation is emitted from the ground, which cools down, resulting in a weakening of turbulent transport. The near-surface layer forms a stable boundary layer, and the upper mixed layer rises from the ground. During this uplift, the atmospheric turbulence characteristics are significantly weakened, but the distribution of meteorological elements in the mixed layer during the day are still maintained. This layer is obviously different from the upper free atmosphere, and is called the residual layer. Because there is often an inversion layer on the top of the atmospheric boundary layer, the upward development of turbulent mixing is inhibited, and therefore a boundary is formed between the atmospheric boundary layer and the free atmosphere. Due to the differences in the thermal properties of the topography and underlying surface, the height of the atmospheric boundary layer presents obvious spatial variation characteristics. Although the Taklimakan Desert in China

and the Pearl River Delta have similar frequencies of boundary layer occurrence, the average atmospheric boundary layer height in the Taklimakan Desert is significantly higher (Zhang, 2017). Therefore, in different locations and under different weather backgrounds, the height of the boundary layer can be very different.

With the completion of the ecological protection barrier along the highway of the Taklimakan Desert and the restored green spaces of the oil base in the hinterland of the Taklimakan Desert, the nature of the regional underlying surface has changed, resulting in changes in surface wind speed. In the context of climate change, the Taklimakan Desert climate is essentially a complex of basin and desert climates, resulting in an extreme arid continental climate.

This study used wind speed data collected by the gradient system of the Taklimakan Desert Tazhong and Xiaotang 10 m meteorological observation towers to analyze the characteristics of the wind speed diurnal variation in the surface layer at different observation stations in the Taklimakan Desert from August 2014 to May 2015. A comparison of the exchange rates between different wind speed profile models in the Taklimakan Desert provided observational data for the development of near-surface atmospheric observation experiments and atmospheric environment research in the Taklimakan Desert. This will be useful for understanding the factors controlling the microclimate of the farmland around the Tarim Basin, improving the near-surface model of the soil-vegetation-atmosphere system, and better understanding the local agrometeorology. The level of technology in the area was of great significance in the study.

2. Materials and methods

The Taklimakan Desert, located between the Tianshan and Kunlun Mountains, is a closed inland basin. While the east of the desert is bounded by the Gobi Desert, the other three sides are surrounded by mountains, including the southern Tibetan Plateau, the Pamir Plateau in the southwest, and the Tianshan Mountains in the north and northwest. Due to this special terrain and geographic location, it is difficult for ocean currents to penetrate the Tarim Basin. the Taklimakan Desert therefore has a typical continental climate, with an abundance of light and heat, little precipitation, strong sunshine, sandy weather, and large temperature differences between day and night, and the different seasons. In this study, two typical observation stations were selected in Xiaotang and Tazhong (Fig. 1). Xiaotang observatory (40°48.126' N, 84°18.211' E, altitude 912 m) is located on the northern edge of the Taklimakan Desert, about 40 km from the Tarim River. It is a typical desert hinterland-desert-oasis transition zone. The underlying surface is flat sandy land, with ancient river bed in some areas, and there is no vegetation. The station is located on the south bank of an ancient river bed. An area of mobile dunes is located about 250 m to the south, and mainly consists of crescent dunes and compound crescent dune chains. The dune landform belongs to the ancient Tarim River alluvial-flood plain, and is the intersection of two desert areas. The area has an inland warm temperate desert climate, which is subject to drought and little rain. The annual average wind speed is $2.5 \text{ m}\cdot\text{s}^{-1}$, with a maximum in spring and summer, and a minimum in winter. The change of air temperature at Xiaotang station is similar to that of wind speed, with the maximum in June–July and the minimum in December–January. The change of wind speed and air temperature has an obvious synchronization. Most (80%) sandstorms occur in spring and summer, with the lowest frequency in winter. The annual average wind speed is 2.8 m/s, and the average temperature is 11.2°C. Tazhong is located in the center of the hinterland of the Taklimakan Desert (83°39' E, 38°58' N), with the highest temperature in summer of 46.0°C, and the lowest temperature in winter of -

25.0°C. The average temperature of the four seasons is 13.6°C, the average annual precipitation is about 25.9 mm, and the evaporation is about 3,812.3 mm. The regional climate is abnormal, the vegetation coverage is extremely low, and only cold and drought-resistant shrubs (e.g., *Haloxylon ammodendron*) can survive. The peak period of dust storms is from March to August every year. The frequency of dust storms at 12 m above the ground is about 500 per year. The normal average wind speed is 2.5 m·s⁻¹, and the sand activity intensity index reaches a maximum value of about 8000 each year. The prevailing wind direction is northeast and northwest.

Figure 1. Map of the study area.

In this study, a windsonic two-dimensional ultrasonic wind speed and direction sensor (1590-PK-020, Campbell Scientific, Logan, UT, USA) and temperature and humidity sensor (1590-PK-020, Campbell Scientific) installed on meteorological observation towers were used to conduct parallel comparative experimental observations of microclimate elements in Tazhong and Xiaotang from January 1, 2014 to December 31, 2015 (**Fig. 2**). The wind speed and direction sensor had a starting wind speed of 0.01 m/s; precision of wind speed $\pm 2\%$; range of 0–60 m/s and 0–359°; and resolution of 0.01 m/s and 1°. The range of the temperature sensor was -80–60 °C, the accuracy was $\pm 0.17^\circ\text{C}$, and the resolution was 0.1°C. The wind speed, temperature, and relative humidity data used in the study were obtained at the height of 10 m. The data used were subjected to quality control, including the synchronous calibration of the two observation points, the logical extremum of observation data, and a time consistency check. In this study, the four seasons were classed as spring (March to May), summer (June to August), autumn (September to November), and winter (December to February). January, April, July, and October were the representative months of winter, spring, summer, and autumn, respectively.

Figure 2. The 10 m observation meteorological towers in Tazhong and Xiaotang.

3. Results

3.1 Diurnal variations of NWS

Figures 3 (a), (b), (c), and (d) show the diurnal variations of mean wind speed in summer, autumn, winter, and spring. Using the ground meteorological data, typical sunny days in each seasonally representative month were selected to determine an average wind speed per hour. The sunny days in summer were August 27, 28, and 31, 2014, the sunny days in autumn were October 3, 10, and 11, 2014, the sunny days in winter were January 4, 6, and 7, 2015, and the sunny days in spring were April 12, 13, and 15, 2015. Based on the average daily variation of the surface layer height in the four seasons, it was found that the daily variation of wind speed in Tazhong and Xiaotang on sunny days had two characteristics. First, there were two peaks and two low values in the daily variation of wind speed. The two peaks occurred at night and in the daytime, and the two low values occurred in the morning and evening, respectively. However, the magnitude and occurrence of wind speed were different in the different seasons. Second, there were two distinct forms of change in the lower and upper layers. In the lower layer, the daytime wind speed was greater than in the night, in the upper layer the nighttime wind speed was greater than during the daytime, and the middle layer was stable, but there were differences in each season

3.1.1 Diurnal variations of NWS in summer

The diurnal variations of surface wind speed in Tazhong and Xiaotang in summer are shown in **Fig. 3 (a)**. The wind speed at different altitudes from 0.5 to 10 m displayed an increasing trend. The maximum wind speeds in each layer in Tazhong and Xiaotang were similar, with values of 5.286, 5.525, 7.479, 8.030, and 8.905 m/s at 0.5, 1, 2, 4, and 10 m, respectively. From 0.5 to 2 m, the highest wind speed occurred at 3:00 at night and 15:30 in the daytime, and the lowest wind speed occurred at 0:30 and 23:30 at night. At 4 and 10 m, the wind speed in Xiaotang was larger than that in Tazhong at 0:30–13:30, and the wind speed difference was about 1–2 m/s. However, the situation at 13:30–23:30 was the complete opposite, with the wind speed in Tazhong being greater than that in Xiaotang. At 0.5–2 m, the situation between the two sites at 17:30–23:30 was also the opposite. Comparing the diurnal variation curves of wind speed at different altitudes, it was found that the

wind speed increased continuously from 0.5 to 1 m. The wind speed at 0.5 m was below 5 m/s, while the wind speed at 2 m was always above 4 m/s.

After sunrise (07:00) in the morning, due to the increasing solar altitude angle and the strengthening of turbulence exchange, the wind speed in each height layer increased rapidly. The average increase of wind speed in each height layer from 0.5 to 10 m was about 0.9 m/s per hour, until 10:00. However, the wind speed at Xiaotang and Tazhong in the 0.5 and 1 m layers displayed a sharp downward trend and then increased slightly until 16:30. At 2, 4, and 10 m, the wind speed increased relatively quickly until 19:00, at which point the wind speed began to decline sharply, leading to a diurnal difference in wind speed of about 6 m/s. After 18:00, with the decreasing solar altitude angle, the ground radiation balance decreased rapidly, the turbulence weakened, and the wind speed decreased rapidly. The decreasing trend in NWS was most obvious below 4 m, with the average wind speed decreasing by 1 m/s per hour. Due to the gradual formation of a near-surface inversion, the turbulence was further weakened, and the wind speed below 10 m continued to decrease, reaching a minimum at 21:00, with an average wind speed of less than 1 m/s from 21:00 to 22:00. This may be because the valley between the large sand ridges alongside the tower was associated with downhill winds at this time.

3.1.2 Diurnal variations of NWS in autumn

Figure 3 (b) shows the diurnal variations of wind speed at 10 m height in Xiaotang and Tazhong in autumn, with the results being similar to those in summer. Overall, the diurnal variation of wind speed in Tazhong was greater than that in Xiaotang, especially in the 2, 4, and 10 m height layers from 09:30 to 23:30. In both Xiaotang and Tazhong, at 12:30 in the autumn the wind speed was high in the daytime at low levels and low in the nighttime at high levels. Compared with summer, the average wind speed in autumn was slightly lower. At 2, 4, and 10 m, the highest wind speed occurred at 03:30 at night and 09:30 in the daytime, with values of 7.9 and 8.79 m/s, respectively. The lowest wind speed occurred at 06:00, with values of 4.8 and 2.3 m/s, respectively. The difference between the maximum and minimum daily wind speed at 10 m was 3.5 m/s. The time at which the maximum wind speed occurred in Xiaotang was the opposite of that in Tazhong, i.e., the time of the minimum wind speed in Tazhong was the time of maximum wind speed in Xiaotang. The highest wind speeds at 10 m in Xiaotang occurred at 01:00 and 08:30 pm in the daytime, with values of 5.81 and 6.9 m/s, respectively. The lowest wind speeds occurred at 03:00 and 23:30, with values of 2.1 and 1.8 m/s, respectively. The average daily maximum wind speed was only 2.8 m/s larger than the minimum wind speed, and the daily variation of the low-level wind speed was much reduced compared with that in the higher layer.

At 08:00, the NWS within 10 m in Tazhong and Xiaotang decreased rapidly, and at 15:30, the wind speed in each altitude layer decreased significantly. The decrease in Tazhong was greater than that in Xiaotang, with a difference of about 4 m/s, which was different from the pattern observed in the summer. In Tazhong, the wind speeds remained high but the wind speed curves at all five levels in the autumn daytime were more closely arranged than those in Xiaotang, indicating that the wind speed difference in the surface layer within 10 m during the autumn daytime was smaller than that in summer. The wind speed difference between the highest and lowest layers in autumn was 1–2.2 m/s, which was about 0.5 m/s smaller than that in summer.

3.1.3 Diurnal variations of NWS in winter

As shown in Fig.3 (c), the diurnal variation of wind speed in the 10 m layer of Tazhong and Xiaotang in winter was generally small, and the wind speed in the whole 10 m layer was less than

3 m/s. In the Xiaotang area, below 4 m from 00:30 to 09:00 the wind speed was maintained at about 0.2 m/s. At 09:30, the wind speed began to increase, until at 23:30 it reached about 1.5 m/s. The wind speed trends in Tazhong and Xiaotang followed the completely opposite trend. From 00:30 to 09:00 the wind speed at 0.5 m was greater than in the other layers, with a maximum of about 1.8 m/s. In winter, the average wind speed in the daytime was larger than that in the nighttime above 2 m. The highest wind speeds in Tazhong and Xiaotang generally occurred at 13:00 and 19:30 in the daytime, and the lowest wind speeds occurred at 21:00 and 06:00, respectively. The highest wind speeds at 10 m were 2.5 and 2.9 m/s, and the lowest wind speeds were 0.2 and 0.02 m/s, respectively. The average wind speeds at night in Tazhong and Xiaotang from 0.5 to 1 m were higher than in the daytime. The difference between the maximum and minimum wind speed at 10 m was only 1.3 m/s, which was smaller than in summer and autumn. After 10:30 in winter, the wind speed at each altitude increased slowly. This increase was small and occurred about 2 h later than in summer. After 10:00, the wind speed began to increase at 2 m, and then began to decrease until 16:30. This feature also occurred at 09:00 in autumn at 2 m.

3.1.4 Diurnal variations of NWS in spring

Figure 3 (d) shows the diurnal variations of surface wind speed in Tazhong and Xiaotang in spring. Similarly, to summer, the wind speed was high in daytime and low at night within 10 m. The wind speed in each layer in Tazhong from 09:30 was obviously larger than in Xiaotang, with the maximum difference in each layer being about 1–3.3 m/s. From the morning of the previous evening to 09:30 the next day, the trends in wind speed changes in Tazhong and Xiaotang were similar, with a stable transition and wind speed difference of about 0.9 m/s. The difference between the maximum and minimum daily average wind speed at each height in the tower was 5.3–6.1 m/s. The difference between the maximum and minimum daily average wind speed in each layer in Xiaotang was 2–3.1 m/s. The diurnal variations of wind speed in the 0.5–2 and 2–4 m layers were similar. From 09:00 to 12:30 in the morning, with the increase in the solar altitude angle, the turbulence exchange was enhanced and the wind speed increased rapidly. In the next 2 hours, the wind speed at 1 m increased from 0.5 to 4.1 m/s, at 2 m it increased from 0.3 to 4.5 m/s, at 4 m it increased from 0.1 to 5.8 m/s, and at 10 m it increased from 0.5 to 6.1 m/s. After 18:30, the wind speed at all altitudes showed a downward trend, although this was less obvious than that in summer and autumn. From midnight to sunrise, the NWS within 10 m was much smaller than that in summer, and was similar to that in winter.

251

252

253

254

255

Figure 3. Diurnal variations of NWS in Tazhong and Xiaotang.

3.2 Comparison of the different wind speed profiles

A curve fitting of the variation in wind speed with height was applied to enable an understanding the near-surface meteorological characteristics in the hinterland of the Taklimakan Desert, as well as an in-depth understanding of the desert wind-sand movement law, and the development and utilization of wind energy resources. It is important to note that in the subsurface the wind direction does not change with height, and therefore the effect of Coriolis force in the subsurface can be ignored. The most commonly used methods for fitting the wind speed profile are the logarithmic law and the exponential law model. Based on a semi-empirical theory, an improved model of the variation of wind speed with height was established according to the M-O similarity theory.

3.2.1 Wind speed profile model under a semi-empirical theory

3.2.1.1 Wind speed profile under neutral stratification

Thermal turbulence does not develop in neutral stratified air, with turbulent motion being completely dependent on the role of dynamic factors. Turbulent motion in the near-surface layer is very similar to the simulated turbulence in a wind tunnel. The relationship $l=kz$ between the mixing length l and the vertical height z from the ground in a wind tunnel experiment was introduced into a near-surface layer under neutral stratification, where k is the Karman constant (generally 0.4).

The relationship between shear stress and mixing length, and the average wind speed $\bar{u}$ was substituted into $l=kz$:

$$\tau = \rho \kappa^2 z^2 \left(\frac{\partial \bar{u}}{\partial z} \right)^2 \quad (1)$$

By introducing the friction velocity u_* , it was considered that $u_* = \sqrt{\frac{\tau}{\rho}}$ did not change with height in the near-surface layer, and Eq. (1) was rewritten to Eq. (2):

$$\frac{\partial \bar{u}}{\partial z} = \frac{u_*}{kz} \quad (2)$$

The pairwise (2) integral is:

$$\bar{u} = \frac{u_*}{k} \ln z + A \quad (3)$$

A is an integral constant. Many experiments have found that the height at which the average wind speed is zero typically occurs at a height z from the ground that is referred to as z_0 .

If boundary conditions are applied: $z = z_0$ 时, $\bar{u} = 0$, Eq. (3) can be written as:

$$\bar{u} = \frac{u_*}{k} \ln \frac{z}{z_0} \quad (4)$$

Equation (4) expresses a typical NWS profile under neutral stratification as a logarithmic wind speed profile.

Figure 4. Average wind speed and logarithmic law curve fitting under neutral stratification in the middle of Tazhong tower.

Figure 4 shows the average wind speed and logarithmic fitting under neutral stratification in the middle of the Tazhong tower, taking the y axis as $\ln z$. It can be seen from the diagram that the near strata in the tower are neutral layers and are very suitable for expressing a logarithmic distribution, with large correlation coefficients. The time period of the neutral stratification in the atmosphere was consistent with the transition period of the temperature profile.

3.2.1.2 Wind velocity profile under non-neutral stratification

Under non-neutral stratification, the wind speed profile will deviate from the logarithmic law, and can be expressed by a simple exponential law in the form of:

$$\bar{u} = \bar{u}_1 \left(\frac{z}{z_1} \right)^n \quad (5)$$

where $\bar{u}$ is the average wind speed at height z and n is the stability parameter, which is expressed as a positive fraction, $0 < n < 1$

Figure 5. Fitting of the mean wind speed and exponential law under non-neutral stratification in the middle of Tazhong tower.

Figure 5 shows the fitting of the wind speed profile under non-neutral stratification in Tazhong on August 25, 2014 with the exponential law. The fitting results were very good, and the correlation coefficients were all above 0.98. The stability parameters of each period were fitted. The stability parameter n was smaller in the daytime than in the night, changing from 0.145 to 0.140 from 10:00 to 14:00 pm, indicating that the more unstable the stratification was, the smaller the n value was. After 20:00, the atmosphere gradually became stable with an n value of 0.284. With the gradual formation of a nighttime inversion layer, the n value increased, tending to 1.

3.2.2 The wind speed profile model established by M-O similarity theory

Under uniform and steady conditions, in the atmospheric layer where the wind direction was not significantly deflected with height, according to M-O similarity theory, the dimensionless wind speed gradient in the near-surface layer can be expressed as:

$$\frac{kz}{u_*} \frac{\partial u}{\partial z} = \varphi_M \left(\frac{Z}{L} \right) \quad (6)$$

where k is the Kaman constant (generally 0.4), u_* is the local friction velocity, and φ_M is a general function. Under neutral stratification, $\varphi_M = 1$. According to the Businger-Nyer relationship, under non-neutral stratification the following expression applies:

$$\varphi_M\left(\frac{Z}{L}\right) = 1 + \beta_m \frac{Z}{L}, \frac{Z}{L} \leq 0 \quad (7)$$

$$\varphi_M\left(\frac{Z}{L}\right) = (1 - \gamma_m \frac{Z}{L})^{\frac{1}{4}}, \frac{Z}{L} \geq 0 \quad (8)$$

where L is the M-O stability constant, and β_m and γ_m are empirical constants determined by measured values. Due to the different data sources, the values of the empirical constants are different.

3.2.1.1 The wind speed profile under near-neutral stratification

When the atmosphere is close to neutral stratification, $\left|\frac{Z}{L}\right| \ll 1$, [Please insert the correct symbol and finish the sentence with a full stop rather than a comma] At this time, the general function is $\varphi_M\left(\frac{Z}{L}\right) \rightarrow 1$. At the point where 0 is reached the formula is expanded according to the Taylor series and then the second order term is omitted:

$$\varphi_M\left(\frac{Z}{L}\right) \cong 1 + \beta \frac{Z}{L} \quad (9)$$

$$\frac{\partial \bar{u}}{\partial z} = \frac{u_*}{kz} \varphi_M\left(\frac{Z}{L}\right) \quad (10)$$

Equation (10) is a two-sided integral, with boundary conditions, $z = z_0$ 时, $\bar{u} = 0$, [Please finish the sentence with a full stop rather than a comma] Equation (11) is the expression of the wind speed profile under near-neutral stratification:

$$\bar{u} = \frac{u_*}{k} \left[\ln \frac{z}{z_0} + \beta \frac{(z - z_0)}{L} \right] \quad (11)$$

Equation (11) adds only one highly linear term to the logarithmic contour, and is therefore referred to as the logarithmic + linear contour. Under neutral stratification, because $L \rightarrow \infty$, the logarithmic + linear profile can be simplified to a logarithmic profile.

341

342

343

344

345 **Figure 6.** Pairwise linear fitting of the average wind speed under near-neutral stratification in the
 346 middle of Tazhong tower.

347

348 **Figure 6** shows the wind speed profile fitted by the logarithmic dozens of linear laws [This
349 phrase requires more explanation. The term is not widely used in the literature. Is it the “linear to tens of’
350 term that is used in the abstract or something else? Please consider adding further text to explain this
351 term.] under near-neutral stratification in the Tazhong area. Generally, the atmospheric stratification
352 was characterized by near-neutral conditions around sunrise and sunset, and the fitting results were
353 also consistent with this period. The correlation coefficient from the fitting was large, indicating that
354 the wind speed profile in a near-neutral atmosphere of weak stability and weak instability conditions
355 in Tazhong satisfied the logarithmic dozens of linear laws.

356 3.2.1.2 The wind speed profile under stable stratification

357 In general, the wind profile theory under stable stratification is not satisfactory. The M-O
358 similarity model is:

$$359 \frac{kz}{u_*} \frac{\partial u}{\partial z} = 1 + \gamma_m \frac{Z}{L} \quad (12)$$

360 Although stable stratification inhibits turbulence and the use of the M-O similarity model is
361 limited, many NWS gradient observations have confirmed the reliability of the M-O model. For the
362 low-level atmosphere above the constant flux layer, Yokoyama et al. verified that their generalized
363 M-O similarity theory could be applied to the surface layer, and proposed a first-order approximate
364 expression of u_* that varied with height:

$$365 u_* = u_0 \left(1 - \frac{z}{h}\right)^n = u_{0*} \zeta^n \quad (13)$$

366 where h is the atmospheric boundary layer thickness and u_{0*} is the friction velocity at the
367 ground surface $\zeta = 1 - \frac{z}{h}$.

368 According to [Yamamoto \(1997\)](#), in the layer where the wind direction has no obvious monotonic
369 deflection with height, if the influence of Coriolis force is ignored, only the change of u with height
370 should be considered. The differential equation obtained by Eqs. (12) and (13) is as follows:

$$371 \frac{kz}{u_*} \frac{\partial u}{\partial z} = (1 + \gamma_m \frac{z}{L}) \left(1 - \frac{z}{h}\right)^n \quad (14)$$

372 3.2.1.3 The wind speed profile under stable stratification

373 Under unstable conditions, the concept of M-O similarity can be extended. Therefore, in this
374 study u was still regarded as a constant from z_0 to the z integral,

$$375 u = \frac{u_*}{k} \left[\ln \frac{z}{z_0} - \varphi_m \left(\frac{z}{L} \right) + \varphi_m \left(\frac{z_0}{L} \right) \right] \quad (15)$$

376 In which, $\varphi_m = \ln \left[\frac{1+x^2}{2} \left(\frac{1+x}{2} \right)^2 \right] - 2 \arctan x + \frac{\pi}{2}$, $x = \left(1 - \beta_m \frac{z}{L} \right)^{0.25} = \varphi_m^{-1}$.

377 The φ_m value for the measured data at the tower height of 10 m was calculated using Eq. (6),

378 and then the empirical constants γ_m and β_m in the tower were fitted according to the relationship

379 between Eq. (7) φ_m and $\frac{z}{L}$.

380 **Figure 7** shows the distribution of φ_m with stability under stable and unstable summer

381 conditions in the middle of the tower. The parameters γ_m and β_m for the surface strata in the

382 middle of the tower in summer were fitted by Eq. (7) and were 5.84 and 15.1.

383

Figure 7. The relationship between surface strata φ_m and $\frac{z}{L}$ at Tazhong tower in summer.

Using the same treatment as in summer, the relationship between near-surface strata φ_m and

$\frac{z}{L}$ in the hinterland of Tazhong Desert was also fitted in winter, and the measured parameters γ_m

and β_m were obtained. **Figure 8** shows the relationship. The surface layer parameters γ_m and

β_m in the middle of the tower in winter were obtained by fitting and had values of 1.9 and 27.1,

respectively.

391

392

Figure 8. The relationship between surface strata φ_m and $\frac{z}{L}$ at Tazhong tower in winter.

4. Conclusion and discussion

Vegetation in desert areas can improve the local microclimate. Desert plants regulate the microclimate in different ways, such as reducing surface wind speed, cooling, and humidification. Many restored green spaces have been constructed in the Tarim Basin oilfield. Compared with Xiaotang, wind speed, temperature, and other meteorological indicators were different in the Tarim Basin, mainly due to the inverse temperature and humidity conditions in winter. The main causes of this phenomenon were the high daytime temperature and low relative humidity of the desert. The water vapor from diffusion at night migrated to the oasis-desert transition zone and increased its relative humidity; thus, forming a circulation mechanism. Vegetation and shelter forest reduced the solar radiation reaching the surface and the emission of long-wave radiation from the ground. The airflow into the canopy was blocked by vegetation, weakening the intensity of heat exchange, resulting in temperature differences between shelter forests, transition zones, and desert areas. Protective forests and vegetation reduced diurnal temperature variations, keeping temperatures relatively stable. In the hinterland of the Taklimakan Desert, due to the continuous expansion of restored green spaces and the enhancement of transpiration from restored vegetation, the relative humidity, temperature, surface temperature, and ground temperature of the near-surface layer of restored green spaces inevitably decreased; thus, changing the hydrothermal conditions of the desert. Due to the influence of vegetation, the increase in ground moisture reduced the surface reflectance, increased the absorption of surface radiation, increased the radiation balance value, and slowed down the temperature change. In the growing season, restored green spaces reduced the surface wind speed, maintain water and soil conditions, and reduced the extent of erosion caused by wind-sand disaster weather.

By analyzing the diurnal variations of the mean wind speed and temperature on sunny days and the characteristics of different wind speed profiles in the middle of observation towers at Tazhong and Xiaotang in summer, autumn, winter, and spring of 2014 and 2015, the following conclusions were reached.

In the four seasons, the diurnal variation of average wind speed on sunny days at different heights in the surface layer of Tazhong and Xiaotang had two characteristics. First, there were two peaks and two low values in the diurnal variation of wind speed. The two peaks occurred at night and in the daytime, and the two low values occurred in the morning and evening, respectively. However, the wind speed varied and the time of occurrence of maximum values differed among the seasons. Second, there were two distinct forms of change in the lower and upper layers. In the lower layer, the daytime wind speed was greater than in the night, in the upper layer the nighttime wind speed was greater than during the daytime, and the middle layer was stable, but there were differences in each season the diurnal variation of wind speed in autumn and spring was larger in Tazhong than in Xiaotang, whereas the variation was similar in summer and winter.

When a semi-empirical theory was applied, under neutral stratification there was a high correlation coefficient when a logarithmic law was used for the fitting, and the timing of the occurrence of a neutral atmosphere was consistent with the transition time of the temperature profile. Under non-neutral stratification, the wind speed profile deviated from the logarithmic law, and the

fitting was better for an exponential law. For the M-O wind speed profile model, when the atmosphere was close to a neutral stratification and when the general function $\varphi_M\left(\frac{z}{L}\right) \rightarrow 1$ approached 1, the near-neutral stratification was fitted by a “logarithmic + linear” profile, and the results obtained were ideal. The γ_m , and β_m values under stable and unstable stratification in the hinterland of the Tazhong Desert were 5.84 and 15.1 in summer and 1.9 and 27.1 in winter, respectively.

Author Contributions: Conceptualization, Liu Xinchun. and Kang Yongde; data curation, Chen Hongna; formal analysis, Lu Hui.; funding acquisition, Chen Hongna.; investigation, Kang Yongde.; project administration, Liu Xinchun; software, Kang Yongde.; supervision, Lu Hui.; Validation, Chen Hongna.; visualization, Liu Xinchun.; Writing-original draft, Liu Xinchun.; Writing - review & editing, Kang Yongde. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the “National Natural Science Foundation of China (41905009)” and “Tianshan Youth Talents (Xinjiang)” Plan Project (2019Q037), the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (grant no.2019QZKK010206) and General scientific research project of Xinjiang Meteorological Bureau (MS201907).

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Alizadeh-Choobari, O., P. Zawar-Reza, and A. Sturman, 2014. The wind of 120 days and dust storm activity over the Sistan Basin. *Atmos. Res.*, 143, 328–341.
- Azorin-Molina C, Vicente-Serrano SM, et al. 2017c. Assessing the impact of measurement time interval when calculating wind speed means and trends under the stilling phenomenon. *Int. J. Climatol.* 37(1): 480–492.
- Belušić, D., and Güttler, G. 2010. Can mesoscale models reproduce meandering motions. *Q. J. R. Meteorol. Soc.* 136 (648), 553–565.
- Beyrich F. 1997. Mixing height estimation from sodar data- a critical discussion. *Atmospheric Environment*, 31(23):3941–3953.
- Collier C G, Davies F, Bozier K E, et al. 2005. Dual-Doppler lidar measurements for improving dispersion models, *Bulletin of the American Meteorological Society*, 86(6):825–838
- Dunn RJH, Azorin-Molina C, Mears CA, et al. 2016. State of the climate in 2016. *Bull. Am. Meteorol. Soc.* 97(8): S38–S40.
- Güttler, I., and Belušić, D. 2012. The nature of small-scale non-turbulent variability in a mesoscale model. *Atmos. Sci. Lett.* 13 (3), 169–173.
- Holtstlag A A M, Boville B A. 1993. Local versus nonlocal boundary-layer diffusion in a global climate model. *Journal of Climate*, 6 (10): 1825–1842.
- Johnson, B.D.; Williams, M.D.; Shepherd, J.M. 2021. Urbanization and Winter Precipitation: A

Case Study Analysis of Land Surface Sensitivity. *Atmosphere*, 12, 805.

Kim J, Paik K. 2015. Recent recovery of surface wind speed after decadal decrease: a focus on South Korea. *Clim. Dyn.* 45(5): 1699–1712.

Li Zhaoguo, Lyu Shihua, et al. 2017. Effect of a cold dry air incursion on atmospheric boundary layer processes over a high-altitude lake in the Tibetan Plateau. *Atmospheric Research*, 12(4):185–190.

Mahrt, L. 2009. Variability and maintenance of turbulence in the very stable boundary layer. *Boundary-Layer Meteorol.* 135 (1), 1–18.

McVicar TR, Roderick ML, Donohue RJ, et al. 2012. Global review and synthesis of trends in observed terrestrial near-surface wind speeds: implications for evaporation. *J. Hydrol.* 416–417:182–205.

Monin A S, Obukhov A M. 1954. Basic laws of turbulent mixing in the surface layer of the atmosphere. *Tr Akad Nauk SSSR Geofiz Inst*, 24(151):163–187.

Ming, H.; Wei, M.; Wang, M. 2019. Quantitative Detection of Dust Storms with the Millimeter Wave Radar in the Taklimakan Desert. *Atmosphere*, 10, 511.

Pryor et al, J. T. Schoof, and R. J. Barthelmie, 2006. Winds of change Projections of near-surface winds under climate change scenarios. *Geophys. Res. Lett.*, 33, L 11072.

Therry G, Lacarrere P. 1983. Improving the eddy kinetic energy model for planetary boundary layer description. *Boundary-layer Meteorology*, 25(1):63–88.

Tokunaga H, Xie SP. 2011. Wave and Anemometer-based Sea-surface wind (WASWind) for climate change analysis. *J. Clim.* 24(1):267–285.

Wang, R., and Coauthors, 2017. Variation of strong dust storms events in northern China during 1978–2007. *Atmos. Res.*, 183, 166–172.

Wentz FJ, Ricciardulli L, Hilburn K, Mears C. 2007. How much more rain will global warming bring *Science* 317(5835): 233–235.

Young IR, Zieger S, Babanin A V. 2011. Global trends in wind speed and wave height. *Science* 332(6028): 451–455.

Yamamoto. S. et al. 1979. Observational study on the turbulent structure of the atmospheric boundary layer under stable conditions. *Journal Meteor Soc. Japan.* 57:423–431

Zhang W C, Guo J P, Miao Y C, et al, 2017. On the summertime planetary boundary layer with different thermodynamic stability in China: A radiosonde perspective. *Journal of Climate*, 31(4):1451–1465.