Correlation Analysis Between Preoperative Systemic Immune Inflammation Index and Prognosis of Gastric Carcinoma Patients: Based on Propensity Score Matching Method

Zhaojun Xu

Qinghai University <https://orcid.org/0000-0001-9275-9652>

Xiaobin Chen

Qinghai University

Juan An

Qinghai University

Jiaqi Yuan

Qinghai University

Shuyun Jiang

Qinghai University

Tao Liu

Qinghai University

Baojia Cai

Qinghai University

Cheng Wang

Qinghai University

Xiaoming Ma (✉ hilton007@126.com)

Qinghai University Affiliated Hospital <https://orcid.org/0000-0003-4539-7438>

Research

Keywords: gastric carcinoma, systemic immune inflammation index, prognosis, overall survival, propensity score matching method

Posted Date: July 26th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-725912/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: To explore the correlation between the preoperative systemic immune inflammation index (SII) and the prognosis of patients with gastric carcinoma (GC).

Methods: The clinical data of 771 GC patients surgically treated in the Department of Gastrointestinal Surgery, Qinghai University Affiliated Hospital from June 2010 to June 2015 were retrospectively analyzed, and their preoperative SII was calculated. The optimal cut-off value of preoperative SII was determined using the receiver operating characteristic (ROC) curve, the confounding factors between the two groups were eliminated using the propensity score matching (PSM) method, and the correlation between preoperative SII and clinicopathological characteristics was assessed by chi-square test. Moreover, the overall survival was calculated using Kaplan-Meier method, the survival curve was plotted, and log-rank test was performed for the significance analysis between the curves. Univariate and multivariate analyses were also conducted using the Cox proportional hazards model.

Results: It was determined by the ROC curve that the optimal cut-off value of preoperative SII was 489.52, based on which 771 GC patients were divided into high SII (H-SII) group and low SII (L-SII) group, followed by PSM in the two groups. The results of Kaplan-Meier analysis showed that before and after PSM, the postoperative 1-, 3- and 5-year survival rates in L-SII group were superior to those in H-SII group, and the overall survival rate had a statistically significant difference between the two groups ($P < 0.05$). Before PSM, preoperative SII [hazard ratio (HR) = 2.707, 95% confidence interval (CI): 2.074-3.533, $P < 0.001$] was an independent risk factor for the prognosis of GC patients. After 1:1 PSM, preoperative SII (HR=2.669, 95%CI: 1.881-3.788, $P < 0.001$) was still an independent risk factor for the prognosis of GC patients.

Conclusions: Preoperative SII is an independent risk factor for the prognosis of GC patients. The increase in preoperative SII in peripheral blood indicates a worse prognosis.

Introduction

Gastric carcinoma (GC) is a common digestive tract malignancy. In 2018, there were up to 782,000 deaths of GC, making it the third cause of death in malignancies^[1]. China is a GC-prone country, and the new cases and deaths account for 44.1% and 49.9%, respectively, of the total globally, with an age-standardized 5-year survival rate of 27.4%. At the same time, GC patients have insidious early symptoms and the therapeutic effect is poor, leading to a poor prognosis^[2]. Therefore, researchers have been constantly exploring the new simple, economical and accurate prognostic evaluation index similar to TNM stage currently, so as to better guide the clinical treatment^[3].

According to related studies, both inflammatory response and immune response are closely related to the occurrence and development of tumors^[4, 5]. Tumor-induced inflammatory response can cause corresponding changes in the blood neutrophil (NE), lymphocyte (LY) and platelet (PLT) counts^[6, 7, 8]. On this basis, some studies have tried to discover new potential biomarkers related to the prognosis. For example, the preoperative systemic immune inflammation index (SII) calculated based on NE, LY and PLT is correlated with the prognosis of breast cancer, liver cancer and pancreatic cancer^[9, 10, 11, 12]. However, there are currently few studies on the correlation between SII and prognosis of GC. In the present study, therefore, the value of SII for the prognosis of GC patients was explored using propensity score matching (PSM), so as to provide references for clinical treatment.

Methods

This study was agreed by the Institutional Research Ethics Board of Qinghai University Affiliated Hospital and obtain the informed consent of all subjects themselves or their families. All methods were performed in accordance with the Declaration of Helsinki, and this study did not involve human or animal tissue or blood samples, and all patients signed the written informed consent before surgery.

General data

The clinical data of GC patients surgically treated in the Department of Gastrointestinal Surgery, Qinghai University Affiliated Hospital from June 2010 to June 2015 were retrospectively analyzed. The treatment methods included distal gastrectomy (DG), proximal gastrectomy (PG) and total gastrectomy (TG). Inclusion criteria were as follows: (1) patients pathologically diagnosed with gastric mucosal adenocarcinoma after operation, (2) those without undergoing radiotherapy, chemotherapy and biotherapy before operation, (3) those without acute/chronic inflammation before operation, (4) those undergoing standard D2 gastrectomy, (5) those without other severe concomitant diseases and with good organ function, and (6) those without medical contraindications that seriously affect anesthesia and operation. Exclusion criteria were as follows: (1) patients with missing medical data or lost to follow-up, (2) those who refused to undergo operation, (3) those complicated with blood system diseases or other tumors before operation, (4) those with severe uncontrolled recurrent infections, or other severe uncontrolled concomitant diseases, or (5) those who require immunosuppressive therapy due to organ transplantation. The pathological staging was based on the 8th edition UICC/AJCC staging criteria ^[13], and the pathological diagnosis and classification of GC were based on the *Japanese Gastric Cancer Treatment Guidelines (5th edition)* ^[14].

Analysis methods

SII was calculated using the blood routine test results first time after admission ($SII = PLT \times NE/LY$). According to the optimal cut-off value of SII, the patients were grouped, and the correlation between SII and clinicopathological factors of patients was analyzed. The patients were also divided into non-elderly group (<60 years old) and elderly group (≥ 60 years old). The correlation between SII and patients' age and its influence on prognosis were analyzed.

PSM

This was a retrospective study, the data were not strictly randomized and there were many potential influencing factors for SII in clinic, so various confounding factors might be distributed in GC patients with different SII. To eliminate the influence of confounding factors on the prognosis, 1:1 PSM was performed for patients in high SII (H-SII) group and low SII (L-SII) group. Other covariates had no statistically significant difference ($P > 0.05$).

Follow-up

All patients were regularly followed up after operation every 3-6 months by outpatient clinic, message, telephone, e-mail and network communication tools. Overall survival (OS) was defined as the duration from definite diagnosis to death or end of follow-up (February 2021 or patients' death).

Statistical analysis

SPSS 26.0, R 3.6.1 and GraphPad Prism 8.0 were used for statistical analysis. Enumeration data were expressed as n (%), and χ^2 test was performed for intergroup comparison. The optimal cut-off value of SII was determined using the receiver operating characteristic (ROC) curve. The confounding factors in data were eliminated using PSM, and the matching precision was set to 0.02. OS was calculated using Kaplan-Meier method, the survival curve was plotted, and log-rank test was performed for the difference between groups. Univariate and multivariate analyses were also conducted using the Cox proportional hazards model. The hazard ratio (HR) and the corresponding 95% confidence interval (95% CI) were calculated. $P < 0.05$ was considered to be statistically significant.

Results

General data

A total of 771 patients met the screening criteria, including 165 females (21.40%) and 606 males (78.60%), with a median age of 59.30 years old (Fig. 1). All patients were pathologically diagnosed with gastric mucosal adenocarcinoma after operation, and none of them died of postoperative complications. Among all patients, 297 cases (38.52%) underwent DG, 250 cases (32.43%) underwent PG, and 224 cases (29.05%) underwent TG. Standard D2 lymph node dissection was conducted for all patients. All of the 771 patients were followed up until February 6, 2021, with a median follow-up time of 46 months (95%CI: 43.986–48.014).

Optimal Cut-off Value Of Sii

The ROC curve of SII was plotted, and its AUC and Youden index were 0.721 (95%CI: 0.681–0.761) and 0.384, respectively. The corresponding optimal cut-off value was 489.52, and the evaluation sensitivity and specificity were 55.5% and 82.9%, respectively (Fig. 2). Based on the cut-off value, 771 patients were divided into L-SII group ($SII \leq 489.52$, $n = 531$) and H-SII group ($SII > 489.52$, $n = 240$).

Correlation between SII and clinicopathological factors of GC patients before and after PSM

Before matching, SII was related to the maximum diameter of tumor, histological type, serum albumin (ALb), white blood cell (WBC), carbohydrate antigen 125 (CA125) and TNM stage ($P < 0.05$), but not related to the age, gender, smoking, drinking, carcinoembryonic antigen (CEA), CA199 and operation method ($P > 0.05$). PSM was performed in L-SII group and H-SII group, and a total of 354 patients were eligible, including 177 cases in H-SII group and 177 cases in L-SII group. The above differences were evenly distributed after matching ($P > 0.05$) (Table 1).

Table 1
Correlation between SII and clinicopathological factors of GC patients before and after PSM

Parameters	Before PSM		After PSM					
	H-SII (n = 240)	L-SII (n = 531)	χ^2	P	H-SII (n = 177)	L-SII (n = 177)	χ^2	P
Age(years)			3.415	0.065			0.011	0.915
≤ 60	129(53.75%)	323(60.83%)			100(56.50%)	99(55.93%)		
> 60	111(46.25%)	208(39.17%)			77(43.50%)	78(44.07%)		
Gender			0.097	0.756			0.000	1.000
Male	187(77.92%)	419(78.91%)			138(77.97%)	138(77.97%)		
Female	53(22.08%)	112(21.09%)			39(22.03%)	39(22.03%)		
Smoking			0.234	0.628			0.011	0.915
Yes	113(47.08%)	260(48.96%)			84(47.46%)	83(46.89%)		
No	127(52.92%)	271(51.04%)			93(52.54%)	94(53.11%)		
Drinking			1.246	0.264			1.457	0.227
Yes	91(37.92%)	224(42.18%)			72(40.68%)	61(34.46%)		
No	149(62.08%)	307(57.82%)			105(59.32%)	116(65.54%)		
Tumor size (mm)			11.357	< 0.001			0.232	0.630
≤ 50	168(70.00%)	429(80.79%)			128(72.32%)	132(74.58%)		
> 50	72(30.00%)	102(19.21%)			49(27.68%)	45(25.42%)		
Differentiation			4.188	0.041			0.000	1.000
Well and Moderately	73(30.42%)	202(38.04%)			61(34.46%)	61(34.46%)		
Poorly	167(69.58%)	329(61.96%)			116(65.54%)	116(65.54%)		
Alb (g/L)			33.695	< 0.001			0.000	1.000
≤ 39.85	183(76.25%)	288(54.24%)			121(68.36%)	121(68.36%)		
> 39.85	57(23.75%)	243(45.76%)			56(31.64%)	56(31.64%)		
WBC(W,×10⁹cells/L)			107.173	< 0.001			0.113	0.737
≤ 6.525	119(49.58%)	451(84.93%)			118(66.67%)	115(64.97%)		
> 6.525	121(50.41%)	80(15.07%)			59(33.33%)	62(35.03%)		
CEA			0.542	0.462			0.554	0.457
≤ 2.17ug/ml	128(53.33%)	268(50.47%)			89(50.28%)	82(46.33%)		
> 2.17ug/ml	112(46.67%)	263(49.53%)			88(49.72%)	95(53.67%)		
CA199			0.835	0.361			0.0137	0.711
≤ 21u/mL	179(74.58%)	412(77.59%)			135(76.27%)	132(74.58%)		
> 21u/mL	61(25.42%)	119(22.41%)			42(23.73%)	45(25.42%)		

Parameters	Before PSM				After PSM			
	H-SII (n = 240)	L-SII (n = 531)	χ^2	P	H-SII (n = 177)	L-SII (n = 177)	χ^2	P
CA125			6.671	0.010			1.532	0.216
≤ 14.6u/ml	145(60.42%)	371(69.87%)			112(63.28%)	123(69.49%)		
> 14.6u/ml	95(39.58%)	160(30.13%)			65(36.72%)	54(30.51%)		
Type of surgery			5.572	0.062			4.974	0.084
DG	80(33.33%)	217(40.87%)			57(32.21%)	39(22.04%)		
PG	78(32.50%)	172(32.39%)			59(33.33%)	63(35.59%)		
TG	82(34.17%)	142(26.74%)			61(34.46%)	75(42.37%)		
TNM stage			11.509	0.003			2.425	0.298
I	38(15.83%)	137(25.80%)			34(16.95%)	35(19.77%)		
II	43(17.92%)	105(19.78%)			31(17.51%)	40(22.60%)		
III	159(66.25%)	289(54.42%)			116(65.54%)	102(57.63%)		
H-SII: High systemic immune inflammatory group;L-SII: Low systemic immune inflammatory group; ALb: Albumin; WBC:White blood cell;DG:Distal gastrectomy;PG:Proximal gastrectomy;TG:Total gastrectomy								

Correlation between SII and OS of GC patients before and after PSM

The median survival time was 73 months (95%CI: 65.172–80.828) in L-SII group and 28 months (95%CI: 24.646–31.354) in H-SII group. The 1-, 3- and 5-year survival rates were 92.8%, 81.0% and 67.9%, respectively, in L-SII group, and 80.0%, 39.8% and 28.9%, respectively, in H-SII group. It can be seen that the survival rate in L-SII group was superior to that in H-SII group, and the OS rate had a statistically significant difference between the two groups ($P < 0.001$). After PSM, the 1-, 3- and 5-year survival rates were 89.3%, 74.3% and 60.4%, respectively, in L-SII group, and 80.8%, 45.8% and 32.9%, respectively, in H-SII group. The mean survival time was 56.771 months (95%CI: 51.613–61.929) in L-SII group. In H-SII group, the mean survival time was 38.443 months (95%CI: 34.792–42.093), and the median survival time was 32 months (95%CI: 22.446–41.554). It can be seen that the survival rate in L-SII group was significantly better than that in H-SII group, and the OS rate had a statistically significant difference between the two groups ($P < 0.001$) (Fig. 3).

Influencing Factors For Patient'S Survival Before And After Psm

Before PSM, the results of univariate analysis revealed that age, maximum diameter of tumor, histological type, SII, ALb, CEA, CA199, CA125, WBC count, operation method and TNM stage were all influencing factors for the prognosis of GC patients ($P < 0.05$). The results of multivariate analysis showed that age (HR = 1.334, 95%CI: 1.045–1.704, $P = 0.021$), histological type (HR = 0.741, 95%CI: 0.560–0.980, $P = 0.036$), SII (HR = 2.707, 95%CI: 2.074–3.533, $P < 0.001$), ALb (HR = 0.385, 95%CI: 0.283–0.525, $P < 0.001$), CEA (HR = 1.285, 95%CI: 1.007–1.640, $P = 0.044$), CA125 (HR = 1.370, 95%CI: 1.069–1.755, $P = 0.013$), WBC (HR = 1.361, 95%CI: 1.047–1.770, $P = 0.021$), operation method and TNM stage were all independent risk factors for the prognosis of GC patients (Table 2). After PSM, the results of univariate analysis manifested that age, maximum diameter of tumor, SII, ALb, CA125, operation method and TNM stage were risk factors for the prognosis of GC patients. According to multivariate analysis, the maximum diameter of tumor (HR = 1.493, 95%CI: 1.043–2.136, $P = 0.029$), SII (HR = 2.669, 95%CI: 1.881–3.788, $P < 0.001$), ALb (HR = 0.286, 95%CI: 0.183–0.447, $P < 0.001$), CA125 (HR = 1.402, 95%CI: 1.002–1.961, $P = 0.048$), operation method and TNM stage were independent risk factors for the prognosis of GC patients (Table 3). Besides, the correlation between statistically significant independent risk factors after PSM ($P < 0.05$) (except SII) and prognosis of GC patients was analyzed by Kaplan-Meier method. It was found that the larger maximum diameter of tumor, lower level of ALb, higher level of CA125 and higher TNM stage

indicated a worse prognosis. Besides, the prognosis was better among patients undergoing DG than that among patients treated with PG and TG (Fig. 4).

Table 2
Risk factors for the prognosis of GC patients before PSM

Parameters	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P	HR (95% CI)	P
Age ,y(≤ 60 VS. >60)	1.531(1.209–1.938)	< 0.001	1.334(1.045–1.704)	0.021
Gender (male VS. female)	0.792(0.584–1.072)	0.131		
Smoking (yes VS. no)	0.913(0.722–1.155)	0.448		
Drinking (yes VS. no)	0.981(0.774–1.244)	0.876		
Tumor size ,mm(≤ 5 VS. >5)	1.619(1.253–2.091)	< 0.001	1.182(0.903–1.547)	0.223
Differentiation (Poorly VS. Well and Moderately)	0.605(0.462–0.794)	< 0.001	0.741(0.560–0.980)	0.036
SII (>489.52 VS. ≤ 489.52)	3.919(3.087–4.977)	< 0.001	2.707(2.074–3.533)	< 0.001
Alb ,g/L(≤ 39.85 VS. >39.85)	0.298(0.220–0.403)	< 0.001	0.385(0.283–0.525)	< 0.001
CEA ,ug/ml(≤ 2.17 ug/ml VS. >2.17 ug/ml)	1.329(1.050–1.680)	0.018	1.285(1.007–1.640)	0.044
CA199 ,u/ml(≤ 21 u/mL VS. >21 u/mL)	1.560(1.208–2.013)	0.001	1.184(0.905–1.550)	0.218
CA125 ,u/ml(≤ 14.6 u/ml VS. >14.6 u/ml)	1.658(1.307–2.104)	< 0.001	1.370(1.069–1.755)	0.013
WBC ^{10⁹/L} (≤ 6.525 VS. >6.525)	2.158(1.698–2.742)	< 0.001	1.361(1.047–1.770)	0.021
Type of surgery		< 0.001		< 0.001
DG	1(reference)		1(reference)	
PG	1.781(1.325–2.395)		1.728(1.279–2.337)	
TG	2.331(1.736–3.129)		1.878(1.386–2.545)	
TNM stage		< 0.001		0.016
I	1(reference)		1(reference)	
II	1.574(1.004–2.466)		1.248(0.788–1.978)	
III	2.385(1.665–3.416)		1.654(1.136–2.408)	
HR: hazard ratio;CI:confidence interval;SII:Systemic immune inflammatory;ALb:Albumin;WBC:White blood cell;DG:Distal gastrectomy;PG:Proximal gastrectomy;TG:Total gastrectom				

Table 3
Risk factors for the prognosis of GC patients after PSM

Parameters	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P	HR (95% CI)	P
Age,y (≤60 VS. >60)	1.544(1.121–2.126)	0.008	1.356(0.973–1.889)	0.072
Gender(male VS. female)	0.958(0.649–1.415)	0.83		
Smoking(yes VS. no)	1.023(0.744–1.407)	0.887		
Drinking(yes VS. no)	1.098(0.790–1.527)	0.577		
Tumor size,mm (≤5 VS. >5)	1.729(1.237–2.417)	0.001	1.493(1.043–2.136)	0.029
Differentiation (Poorly VS. Well and Moderately)	0.733(0.514–1.046)	0.087		
SII (>489.52 VS. ≤489.52)	2.472(1.757–3.480)	< 0.001	2.669(1.881–3.788)	< 0.001
Alb,g/L (≤39.85 VS. >39.85)	0.306(0.198–0.474)	< 0.001	0.286(0.183–0.447)	< 0.001
CEA,ug/ml (≤2.17ug/ml VS. >2.17ug/ml)	1.380(0.999–1.906)	0.051		
CA199,u/ml (≤21u/mL VS. >21u/mL)	1.415(0.996–2.011)	0.053		
CA125,u/ml (≤14.6u/ml VS. >14.6u/ml)	1.479(1.067–2.050)	0.019	1.402(1.002–1.961)	0.048
WBC ^{10⁹/L} (≤6.525 VS. >6.525)	0.921(0.659–1.288)	0.631		
Type of surgery		0.001		0.001
DG	1(reference)		1(reference)	
PG	1.803(1.159–2.805)		2.061(1.314–3.232)	
TG	2.230(1.451–3.428)		2.324(1.490–3.623)	
TNM stage		< 0.001		0.002
I	1(reference)		1(reference)	
II	1.803(1.159–2.805)		1.120(0.579–2.167)	
III	2.230(1.451–3.428)		2.123(1.257–3.584)	
HR: hazard ratio;CI:confidence interval;SII:Systemic immune inflammatory;ALb:Albumin;WBC:White blood cell;DG:Distal gastectomy;PG:Proximal gastrectomy;TG:Total gastrectom				

Clinicopathological Characteristics Of Age-stratified Patients After Psm

After PSM, 199 non-elderly patients (56.21%) and 155 elderly patients (43.79%) were obtained for analysis. It was proved that SII was related to the TNM stage of non-elderly patients ($P < 0.05$) (Table 4).

Table 4
Age-stratified PSM

Parameters	Non-elderly patients(n = 199)				Elderly patients(n = 155)			
	H-SII(n = 100)	L-SII(n = 99)	χ^2	P	H-SII(n = 77)	L-SII(n = 78)	χ^2	P
Gender			0.106	0.744			0.102	0.749
Male	80(80.00%)	81(81.82%)			58(75.32%)	57(73.08%)		
Female	20(20.00%)	18(18.18%)			19(24.68%)	21(26.92%)		
Smoking			0.245	0.621			0.516	0.472
Yes	47(47.00%)	50(50.51%)			37(48.05%)	33(42.31%)		
No	53(53.00%)	49(49.49%)			40(51.95%)	45(57.69%)		
Drinking			0.445	0.505			1.145	0.285
Yes	42(42.00%)	37(37.37%)			30(38.96%)	24(31.17%)		
No	58(58.00%)	62(62.63%)			47(61.04%)	54(70.13%)		
Tumor size (mm)			0.834	0.361			0.102	0.749
≤ 50	70(70.00%)	75(75.76%)			58(75.32%)	57(73.08%)		
> 50	30(30.00%)	24(24.24%)			19(24.68%)	21(26.92%)		
Differentiation			0.065	0.799			0.073	0.787
Well and Moderately	33(33.00%)	68(68.69%)			49(63.64%)	48(61.54%)		
poorly	67(67.00%)	31(31.31%)			28(36.36%)	30(38.46%)		
Albumin (g/L)			0.056	0.813			0.113	0.737
≤ 39.85	61(61.00%)	62(62.63%)			60(77.92%)	59(75.64%)		
> 39.85	39(39.00%)	37(37.37%)			17(22.08%)	19(24.36%)		
WBC,10⁹/L			0.411	0.521			0.052	0.820
≤ 6.525	65(65.00%)	60(60.61%)			53(68.83%)	55(70.51%)		
> 6.525	35(35.00%)	39(39.39%)			24(31.17%)	23(29.49%)		
CEA			0.047	0.828			0.78	0.377
≤ 7ug/ml	48(48.00%)	46(46.46%)			41(53.25%)	36(46.15%)		
> 7ug/ml	52(52.00%)	53(53.54%)			36(46.75%)	42(53.85%)		
CA199			0.085	0.771			0.773	0.379
≤ 21u/mL	75(75.00%)	76(76.77%)			60(77.92%)	56(71.79%)		
> 21u/mL	25(25.00%)	23(23.23%)			17(22.08%)	22(28.21%)		
CA125			3.336	0.068			0.01	0.921
≤ 14.6u/ml	65(65.00%)	76(76.77%)			47(61.04%)	47(60.26%)		
> 14.6u/ml	35(35.00%)	23(23.23%)			30(38.96%)	31(39.74%)		
Type of surgery			2.206	0.332			3.085	0.214
DG	36(36.00%)	26(26.26%)			21(27.27%)	13(16.67%)		

Parameters	Non-elderly patients(n = 199)				Elderly patients(n = 155)			
	H-SII(n = 100)	L-SII(n = 99)	χ^2	P	H-SII(n = 77)	L-SII(n = 78)	χ^2	P
PG	32(32.00%)	36(36.37%)			27(35.07%)	27(34.62%)		
TG	32(32.00%)	37(37.37%)			29(37.66%)	38(48.71%)		
TNM stage			14.683	0.001			4.102	0.129
I	14(14.00%)	18(18.18%)			16(20.78%)	17(21.79%)		
II	9(9.00%)	28(28.28%)			22(28.57%)	12(15.38%)		
III	77(77.00%)	53(53.54%)			39(50.65%)	49(62.83%)		
H-SII: High systemic immune inflammatory group;L-SII:Low systemic immune inflammatory group;ALb:Albumin;WBC:White blood cell;DG:Distal gastrectomy;PG:Proximal gastrectomy;TG:Total gastrectomy								

Cox regression analysis on influencing factors for the prognosis of non-elderly patients

The results of univariate analysis showed that the maximum diameter of tumor, SII, ALb, CA125 and TNM stage were risk factors for the prognosis of the 199 non-elderly patients with GC. The results of multivariate analysis showed that SII (HR = 3.383, 95%CI: 1.961–5.838, $P < 0.001$) and ALb (HR = 0.400, 95%CI: 0.238–0.674, $P = 0.001$) were independent risk factors for the prognosis of non-elderly patients (Table 5).

Table 5
Univariate and multivariate analyses on OS of non-elderly patients

Parameters	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P	HR (95% CI)	P
Gender male VS. female	0.8340.459-1.516	0.552		
Smoking yes VS. no	1.0420.663-1.639	0.858		
Drinking yes VS. no	1.0900.688-1.726	0.714		
Tumor size,mm ≤5 VS. >5	1.8781.177-2.995	0.008	1.4520.893-2.359	0.133
Differentiation Poorly VS. Well and Moderately	0.9260.561-1.528	0.764		
SII >489.52 VS. ≤489.52	3.9702.334-6.751	<0.001	3.3831.961-5.838	<0.001
Albumin,g/L ≤39.85 VS. >39.85	0.4620.277-0.769	0.003	0.4000.238-0.674	0.001
CEA,ug/ml ≤2.17ug/ml VS. >2.17ug/ml	1.2080.768-1.901	0.413		
CA199,u/ml ≤21u/mL VS. >21u/mL	1.1100.660-1.867	0.693		
CA125,u/ml ≤14.6u/ml VS. >14.6u/ml	1.7831.124-2.830	0.014	1.5920.992-2.554	0.054
WBC,10 ⁹ /L ≤6.525 VS. >6.525	0.9810.616-1.561	0.934		
Type of surgery		0.104		
DG	1reference			
PG	1.8011.039-3.123			
TG	1.2960.723-2.324			
TNM stage		0.004		0.090
I	1reference		1reference	
II	1.6440.549-4.920		2.1110.686-6.493	
III	3.5391.423-8.799		2.7761.091-7.063	
HRhazard ratio; CIconfidence interval; SIISystemic immune inflammatory; ALbAlbumin; WBCWhite blood cell; DGDistal gastrectomy; PGProximal gastrectomy; TGTototal gastrectom				

Cox regression analysis on influencing factors for the prognosis of elderly patients

The results of univariate analysis showed that the histological type, SII, ALb, CEA, CA199, operation method and TNM stage were risk factors for the prognosis of the 155 elderly patients with GC. It was confirmed by multivariate analysis that SII (HR = 2.372, 95%CI: 1.444–3.896, P = 0.001), ALb (HR = 0.164, 95%CI: 0.059–0.457, P = 0.001) and operation method were independent risk factors for the prognosis of elderly patients (Table 6).

Table 6
Univariate and multivariate analyses on OS of elderly patients

Parameters	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P	HR (95% CI)	P
Gender male VS. female	0.976 0.574-1.661	0.928		
Smoking yes VS. no	0.960 0.610-1.513	0.861		
Drinking yes VS. no	1.024 0.635-1.652	0.922		
Tumor size,mm ≤5 VS. >5	1.610 0.993-2.611	0.053		
Differentiation Poorly VS. Well and Moderately	0.532 0.321-0.881	0.014	0.689 0.410-1.158	0.160
SII >489.52 VS. ≤489.52	1.612 1.012-2.567	0.044	2.372 1.444-3.896	0.001
Alb,g/L ≤39.85 VS. >39.85	0.132 0.048-0.361	<0.001	0.164 0.059-0.457	0.001
CEA,ug/ml ≤2.17ug/ml VS. >2.17ug/ml	1.704 1.069-2.715	0.025	1.530 0.930-2.517	0.094
CA199,u/ml ≤21u/mL VS. >21u/mL	1.848 1.138-3.002	0.013	1.457 0.877-2.421	1.146
CA125,u/ml ≤14.6u/ml VS. >14.6u/ml	1.142 0.718-1.818	0.574		
WBC,109/L ≤6.525 VS. >6.525	0.940 0.574-1.539	0.806		
Type of surgery		<0.001		0.002
DG	1 reference		1 reference	
PG	1.893 0.875-4.098		1.548 0.709-3.381	
TG	4.004 1.937-8.279		3.174 1.490-6.760	
TNM stage		0.020		0.090
I	1 reference		1 reference	
II	0.902 0.397-2.049		0.766 0.328-1.786	
III	1.914 1.020-3.590		1.539 0.798-2.967	
HR hazard ratio CI confidence interval SII Systemic immune inflammatory ALb Albumin WBC White blood cell DG Distal gastrectomy PG Proximal gastrectomy TG Total gastrectom				

Postoperative Survival Rate Of Non-elderly And Elderly Patients

Among the non-elderly patients, the 1-, 3- and 5-year survival rates were 92.9%, 82.8% and 62.1%, respectively, in L-SII group, and 85.0%, 47.6% and 31.9%, respectively, in H-SII group. The survival rate of GC patients in L-SII group was better than that in H-SII group, and the OS rate had a statistically significant difference between the two groups ($P < 0.001$). Among the elderly patients, the 1-, 3- and 5-year survival rates were 83.3%, 63.2% and 43.8%, respectively, in L-SII group, and 75.3%, 42.2% and 19.4%, respectively, in H-SII group. The survival rate of GC patients in L-SII group was better than that in H-SII group, and the OS rate had a statistically significant difference between the two groups ($P = 0.041$) (Fig. 5).

Discussion

In recent years, related studies have demonstrated that the inflammatory response (tissue necrosis, activation of immune response, secretion of cytokines) plays an important role in the occurrence, development and metastasis of tumors [15, 16]. In the early stage, the inflammatory response is induced and the anti-tumor immune response is inhibited, thereby promoting growth of

tumor cells. With the growth of tumor, the inflammatory response is further enhanced. To eliminate inflammation, the recruitment of immune cells to tumor tissues is reduced, resulting in immune escape of tumor cells. In the late stage, the body's immunity is almost lost, and the inflammatory response is further worsened, thus facilitating the tumor progression^[17]. SII, a new inflammatory index integrating the NE, LY and PLT counts^[18], can reflect the balance status between tumor immunity and inflammatory response more comprehensively than platelet-to-lymphocyte ratio (PLR), neutrophil-to-lymphocyte ratio (NLR) and monocyte-to-lymphocyte ratio (MLR), and better predict the prognosis of cancer patients^[18,19,20].

In the present study, the optimal cut-off value of SII was determined using the ROC curve, based on which the patients were divided into H-SII group and low L-SII group. In terms of the correlation between SII and the clinicopathological characteristics of patients, SII was related to the maximum diameter of tumor, histological type, ALb, WBC, CA125 and TNM stage ($P < 0.05$). In addition, the survival time of GC patients in H-SII group was significantly shortened, and the 5-year survival rate was significantly lower than that in L-SII group. Wang^[21] *et al.* retrospectively analyzed the clinical data of 444 GC patients treated with gastrectomy and found that the 5-year survival rate is 29.4% and 46.1%, respectively, in H-SII group and L-SII group, suggesting that the SII level is related to the prognosis of GC patients, consistent with the results of this study. Han BL^[22] also showed that SII is closely related to the prognosis of GC patients, and high SII indicates a poor prognosis of GC patients. Its mechanism can be explained by the functions of NE, PLT and LY. In the case of tumor invasion, the inflammatory response can promote the increase in NE and the secretion of NO, reactive oxygen species, arginase and other active substances, which can regulate the activity of NK cells and LY, thereby facilitating tumor growth and metastasis^[23, 24]. In addition, the PLT count rises in inflammation, and PLT promotes tumor angiogenesis *via* releasing vascular endothelial growth factor and platelet-derived growth factor^[25], creating favorable conditions for tumor growth. Moreover, PLT can form colonizing tumor thrombi with tumor cells, thus promoting the further invasion of tumor cells into the body. On the contrary, LY can effectively suppress the occurrence and development of tumors, and induce the death of tumor cells. The long-term inflammatory stimulus will lead to decline in LY subsets, thereby causing immune dysfunction, and the decrease in LY count can raise the risk of immune escape of tumor cells^[26, 27]. Therefore, SII can reflect the relation between systemic inflammatory response and body's immunity during tumor progression. The higher the SII is, the severer the inflammatory response and immunosuppression in GC patients will be.

In this study, it was confirmed by univariate analysis before and after PSM that age, maximum diameter of tumor, SII, ALb, CA125, operation method and TNM stage were risk factors for the prognosis of GC patients. Before PSM, the results of multivariate analysis showed that age, histological type, SII, ALb, CEA, CA125, WBC, operation method and TNM stage were independent risk factors for the prognosis of GC patients. After PSM, the results of multivariate analysis revealed that the maximum diameter of tumor, SII, ALb, CA125, operation method and TNM stage were independent risk factors for the prognosis of GC patients. Then the statistically significant factors after PSM (except SII) were analyzed. It was found that the larger maximum diameter of tumor, lower level of ALb, higher level of CA125 and higher TNM stage indicated a worse prognosis. The prognosis was better among patients undergoing DG than that among patients treated with PG and TG. Wang *et al.*^[28] found in the study on the correlation between SII and the prognosis of GC patients that SII and TNM stage are independent risk factors for the prognosis of GC patients, consistent with the results of this paper. As the main component of plasma protein, ALb can reflect the nutritional status of patients, and it has also been widely used as an antidote and transporter in predicting the survival status and disease progression of cancer patients^[29, 30]. CA125 is mainly present in epithelial tissues and serum of patients, and its expression level in the serum of GC patients is higher than that in normal people, which is positively correlated with the TNM stage and negatively correlated with the prognosis of GC patients, consistent with the results of this study. This indicates that CA125 can be used to assess the severity of GC^[31]. In addition, the multivariate analysis before and after PSM demonstrated that no positive result was obtained in age after PSM, but other scholars argued that there is a correlation between age and the prognosis of GC patients^[32, 33]. The reasons why age was not an independent risk factor for the prognosis of GC patients in this study are as follows: PSM was used in this study to validate the impact of SII on the prognosis on the basis of excluding other confounding factors, which stressed the effect of SII on the prognosis, thus affecting the prognostic assessment by other indexes to a certain extent. Besides, it is possibly because the sample size was small after PSM and this study itself had limitations, rather than simply considering age unrelated to the prognosis of patients. Therefore, the sample size remains to be further expanded in the future. To further explore the correlation between age and the prognosis of GC patients, age was further subjected to stratification analysis to exclude the value deviation in a certain stage caused by the small sample size or the bias of age stratification in this study. It is

reported in the literature that elderly patients are aged ≥ 60 years old [34, 35]. In this study, the median age of 354 patients was 59.0 years old after PSM. Therefore, patients aged ≥ 60 years old were defined as elderly patients in this study. The results revealed that the mean SII of elderly patients (495.73) was significantly higher than that of non-elderly patients (443.21), suggesting the correlation between SII and patient's age. In addition, the risk factors for the prognosis of GC patients were analyzed under age stratification. It was found that the high SII was correlated with the poor prognosis of elderly patients with GC, and the 5-year OS rate of elderly patients was lower than that of non-elderly patients in H-SII group (31.9% vs. 62.1%). The specific mechanism remains to be further confirmed by a large number of prospective studies.

There were certain limitations in this study. First, this was a small-sample retrospective study on GC patients in high-altitude areas. To better control the bias, PSM was used for data analysis, but all the impact of covariates on the outcome failed to be fully eliminated. Second, this was a single-center study, so the conclusion cannot fully represent the characteristics of other study centers and populations, limiting its popularization. Therefore, the preliminary results need to be validated by larger-sample randomized controlled studies in the future, so as to offer more convincing theoretical support to the existing conclusion.

In conclusion, SII, that remains simple, universal, non-invasive, cheap and reproducible, is expected to be an index for assessing the prognosis of GC patients.

List Of Abbreviations

systemic immune inflammation index	SII
gastric carcinoma	GC
receiver operating characteristic	ROC
propensity score matching	PSM
high SII	H-SII
low SII	L-SII
hazard ratio	HR
confidence interval	CI
neutrophil	NE
lymphocyte	LY
platelet	PLT
distal gastrectomy	DG
proximal gastrectomy	PG
total gastrectomy	TG
Overall survival	OS
albumin	ALb
white blood cell	WBC
carbohydrate antigen 125	CA125
carcinoembryonic antigen	CEA
platelet-to-lymphocyte ratio	PLR
neutrophil-to-lymphocyte ratio	NLR
monocyte-to-lymphocyte ratio	MLR

Declarations

Author Contributions

Xu Zhaojun and Chen Xiaobin: Designed the research and wrote the paper; Chen Xiaobin, Jiang Shuyun, Liu Tao and Wang Cheng: Revised the paper; Xu Zhaojun, Cai Baojia and Chen Xiaobin: Participated in research work; Xu Zhaojun, Chen Xiaobin: Collected samples; Xu Zhaojun, Chen Xiaobin and Yuan Jiaqi : Analyzed data and constructed figures; Ma Xiaoming and An Juan : Responsible for project guidance.

Conflicts of Interest and Source of Funding:

Drs. Xu Zhaojun, Chen Xiaobin, An Juan, Yuan Jiaqi, Jiang Shuyun, Liu Tao, Cai Baojia, Wang Cheng and Ma Xiaoming have no conflicts of interest or financial ties to disclose.

This work was supported by the Key R & D and transformation program of Qinghai Province-Special project of science and technology assistance (No.2021-QY-213); the CAS (Light of the West China) Program (No.2019-33), the National Natural Science Foundation of China (No.81460429), the Open Project of State Key Laboratory of Plateau Ecology and Agriculture, Qinghai University (No.2019-ZZ-07), the Scientific Research Project of Research Team of Medical College of Qinghai University (No. 2020-KYT-2).

Data availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

Research reported in this publication was supported by grant number 81460429 from the National Natural Science Foundation of China.

Additional Information

Ethical approval

Ethical approval was not required in the treatment of the patient in this report.

Consent

Written consent has been received from the subject.

References

- 1 Freddie Bray BSc, MSc, PhD et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries[J]. CA: A Cancer Journal for Clinicians, 2018, 68(6) : 394-424.
- 2 Kun Gao and Jun Wu. National trend of gastric cancer mortality in China (2003–2015): a population-based study[J]. 2019, 39(1) : 24.
- 3 Park, Lee, HoJae, et al. Overview of gastrointestinal cancer prevention in Asia[C]// Asia-Pacific Microwave Conference. 2015.
- 4 Prete A D , Allavena P , Santoro G , et al. Molecular pathways in cancer-related inflammation[J]. Biochimica Medica, 2011, 21(3):264-275.
- 5 Cancer-Related Inflammation[J]. Journal of Clinical Immunology, 2013, 33(1):79-84.

- 6 The pretreatment neutrophil-to-lymphocyte ratio predicts therapeutic response to radiation therapy and concurrent chemoradiation therapy in uterine cervical cancer.[J]. International Journal of Clinical Oncology, 2015, 20(5):989-996.
- 7 LABELLE M, BEGUM S, HYNES RO. Direct signaling between platelets and carcinoma cells induces an epithelial-mesenchymal-like transition and promotes metastasis. 2011, 20050:576-90.
- 8 Zhu Lizhen et al. A new prognostic score based on the systemic inflammatory response in patients with inoperable non-small-cell lung cancer.[J]. OncoTargets and therapy, 2016, 9 : 4879-86.
- 9 GAO YB, GUO W, CAI SH, et al. Systemic immune-inflammation index (SII) is useful to predict survival outcomes in patients with surgically resected esophageal squamous cell carcinoma. 2019, 100140:3188-3196.
- 10 WANG P, YUE WS, LI WY, et al. Systemic immune-inflammation index and ultrasonographic classification of breast imaging-reporting and data system predict outcomes of triple-negative breast carcinoma. 2019, 11:813-819.
- 11 HU B, YANG XR, XU Y, et al. Systemic immune-inflammation index predicts prognosis of patients after curative resection for hepatocellular carcinoma. 2014, 200230:6212-22.
- 12 AZIZ MH, SIDERAS K, AZIZ NA, et al. The Systemic-immune-inflammation Index Independently Predicts Survival and Recurrence in Resectable Pancreatic carcinoma and its Prognostic Value Depends on Bilirubin Levels: A Retrospective Multicenter Cohort Study. 2019, 270010:139-146.
- 13 Shan Fei, Li Ziyu, Zhang Lianhai, Li Shuangxi, Jia Yongning, Miao Rulin, Xue Kan, Li Zhemin, Gao Xiangyu, Wang Yinkui, Yan Chao, Li Shen, Ji Jiafu. The Union for International Cancer Control (UICC) and the American Joint Committee on Cancer (AJCC) gastric cancer TNM staging system (8th edition) explanation and elaboration [J]. Chinese Journal of Practical Surgery, 2017,37(01): 15-17
- 14 Hu Xiang. Changes of Japanese "Guidelines" and New Trends of Gastric Cancer Treatment [J]. Chinese Journal of Practical Surgery, 2019,39(03): 231-236
- 15 Sergei I. Grivennikov, Florian R. Greten, Michael Karin. Immunity, Inflammation, and carcinoma. 2010, 140060:883-899.
- 16 Singh R, Mishra MK, Aggarwal H. Inflammation, Immunity, and Cancer. Mediators Inflamm. 2017;2017:6027305.
- 17 Huang B. Regulation of immune response and inflammation in tumor microenvironment[J]. Chinese Journal of Cancer Biotherapy, 2012,19(02):111-115.
- 18 Fu Hongyuan et al. Systemic Immune-Inflammation Index (SII) is Useful to Predict Survival Outcomes in Patients After Liver Transplantation for Hepatocellular Carcinoma within Hangzhou Criteria.[J]. Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology, 2018, 47(1) : 293-301.
- 19 Diem S, Schmid S, Krapf M, Flatz L, Born D, Jochum W, Templeton AJ, Früh M. Neutrophil-to-Lymphocyte ratio (NLR) and Platelet-to-Lymphocyte ratio (PLR) as prognostic markers in patients with non-small cell lung cancer (NSCLC) treated with nivolumab. Lung Cancer. 2017 Sep;111:176-181.
- 20 Trifan G, Testai FD. Systemic Immune-Inflammation (SII) index predicts poor outcome after spontaneous supratentorial intracerebral hemorrhage. J Stroke Cerebrovasc Dis. 2020 Sep;29(9):105057.
- 21 Kang Wang, Feiyu Diao, Zhijun Ye, Xinhua Zhang, Ertao Zhai, Hui Ren, Tong Li, Hui Wu, Yulong He, Shirong Cai, Jianhui Chen. Prognostic value of systemic immune-inflammation index in patients with gastric carcinoma[J]. Chinese Journal of carcinoma, 2017, 360090:420-426.
- 22 Han BL, Wang YM, Xue YW. Effect of preoperative systemic immune-inflammation index on the prognosis of patients with gastric cancer[J]. Chinese Journal of General Surgery, 2019(04):306-309.

- 23 Hu XX, He YF, Luo HQ, Chen WJ, Ke LH, Yan Y, Wu SS, Niu JY, Li HM, Xu HJ, Hu B. The relationship between peripheral blood NLR, PLR and clinical prognosis of small cell esophageal cancer[J]. Chinese Clinical Oncology, 2019,24(06):543-547.
- 24 Teramukai S , Kitano T , Kishida Y , et al. Pretreatment neutrophil count as an independent prognostic factor in advanced non-small-cell lung cancer: An analysis of Japan Multinational Trial Organisation LC00-03[J]. European Journal of Cancer, 2009, 45(11):1950-1958.
- 25 Brill A, Dashevsky O, Rivo J, Gozal Y, Varon D. Platelet-derived microparticles induce angiogenesis and stimulate post-ischemic revascularization. Cardiovasc Res. 2005 Jul 1;67(1):30-8.
- 26 Hu Hong et al. Prognostic value of preoperative NLR, dNLR, PLR and CRP in surgical renal cell carcinoma patients.[J]. World journal of urology, 2017, 35(2) : 261-270.
- 27 Ozmen S et al. Neutrophil-lymphocyte ratio (NLR) and platelet-lymphocyte ratio (PLR) may be superior to C-reactive protein (CRP) for predicting the occurrence of differentiated thyroid cancer.[J]. Endocrine regulations, 2017, 51(3) : 131-136.
- 28 Wang Qingshan and Zhu Dayong. The prognostic value of systemic immune-inflammation index (SII) in patients after radical operation for carcinoma of stomach in gastric cancer.[J]. Journal of gastrointestinal oncology, 2019, 10(5) : 965-978.
- 29 Atsushi Abe et al. Correlation between prognostic nutritional index and occlusal status in gastric cancer[J]. Oral Diseases, 2020, 26(2) : 465-472.
- 30 Maria Mantzorou, Antonios Koutelidakis, Stamatios Theocharis, et al. Clinical Value of Nutritional Status in Cancer: What is its Impact and how it Affects Disease Progression and Prognosis? 2017, 69(8):1151-1176.
- 31 Huang Chao et al. Clinical Significance of Serum CA125, CA19-9, CA72-4, and Fibrinogen-to-Lymphocyte Ratio in Gastric Cancer With Peritoneal Dissemination.[J]. Frontiers in oncology, 2019, 9 : 1159.
- 32 Zou XN, Wan X, Dai Z, Yang GH. Epidemiological characteristics of cancer in elderly chinese. ISRN Oncol. 2012;2012:381849.
- 33 Su Yan,Bin Li,Zhen-Zhong Bai,Jun-Qi Wu,Da-Wei Xie,Ying-Cai Ma,Xu-Xiang Ma,Jun-Hui Zhao,Xin-Jian Guo.Clinical epidemiology of gastric cancer in Hehuang valley of China:A 10-year epidemiological study of gastric cancer[J].World Journal of Gastroenterology,2014,20(30):10486-10494.
- 34 Yang LD, Wang XD, Li Q, Liang YH, Chen Y, Wu MJ, Xu F. The safety in more than 60-year-old gastric cancer patients treated by fast-track surgery[J]. Journal of Modern Oncology, 2015,23(19):2814-2817.
- 35 Xu PW, Zhang H. Clinical manifestations, pathological features and prognosis of elderly patients with gastric cancer[J]. Journal of Nanjing Medical University (Natural Science), 2015,35(03):398-400.

Figures

Figure 1

Study flow diagram.

Figure 2

ROC curves were plotted to determine the optimal cut-off value of SII.

Figure 3

Postoperative OS curve in the two groups before and after PSM. a: Before PSM, the 5-year OS rate in L-SII group was significantly higher than that in H-SII group (67.9% vs. 28.9%) ($P < 0.001$). b: After PSM, the 5-year OS rate in L-SII group was significantly higher than that in H-SII group (60.4% vs. 32.9%) ($P < 0.001$).

Figure 4

Survival curves under different risk factors for the prognosis after PSM. a: different tumor sizes (H-SII group vs. L-SII group). b: Different ALb levels (H-SII group vs. L-SII group). c: Different CA125 levels (H-SII group vs. L-SII group). d: Different operation methods (DG vs. PG vs. TG). e: Different TNM stages (stage I vs. stage II vs. stage III).

Figure 5

Correlation between the prognosis of non-elderly and elderly patients and SII after PSM. a: The 5-year OS rate in L-SII group was significantly higher than that in H-SII group among the non-elderly patients (62.1% vs. 31.9%) ($P < 0.001$). b: The 5-year OS rate in L-SII group was significantly higher than that in H-SII group among the elderly patients (43.8% vs. 19.4%) ($P = 0.041$).