

Structure and surgical dissection layers of the bare area of the liver

Takamichi Ishii (✉ taishii@kuhp.kyoto-u.ac.jp)

Graduate School of Medicine Kyoto University <https://orcid.org/0000-0002-7461-9653>

Satoru Seo

Graduate School of Medicine Kyoto University

Takashi Ito

Graduate School of Medicine Kyoto University

Satoshi Ogiso

Graduate School of Medicine Kyoto University

Ken Fukumitsu

Graduate School of Medicine Kyoto University

Kojiro Taura

Graduate School of Medicine Kyoto University

Toshimi Kaido

Graduate School of Medicine Kyoto University

Shinji Uemoto

Graduate School of Medicine Kyoto University

Research article

Keywords: Bare area, liver, diaphragm, cadaver, Laennec's capsule

Posted Date: October 31st, 2019

DOI: <https://doi.org/10.21203/rs.2.16652/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 31st, 2020. See the published version at <https://doi.org/10.1186/s12893-020-00830-8>.

Abstract

Background : The bare area, which lies between the right liver and diaphragm, should be dissected to mobilize the right liver. The bare area was reportedly formed by direct adhesion between the liver and diaphragm, meaning that the bare area lacked serosal components. This study aimed to analyze the structure of the bare area by an integrated study of surgical and laparoscopic images and pathological studies and describe surgical procedures focusing on the multilayered structure.

Methods : Surgical specimens of hepatectomy were analyzed histologically to evaluate the macroscopic structure of the bare area. Laparoscopic images and cadaver anatomy of the bare area were also examined, changing the dissection layers of the bare area.

Results : The bare area comprised a multilayered structure that resulted in fusions in the hepatic development among the parietal peritoneum (peritoneum of the diaphragm part) and the visceral peritoneum (liver serosa). The multilayered structure of the bare area comprised the liver, sub-serosal connective tissue, liver serosa, parietal peritoneum, retroperitoneal connective tissue, epimysium of the diaphragm, and diaphragm, in order from the liver to the diaphragm. Laparoscopic images sometimes enabled us to recognize the multilayered structure of the bare area.

Conclusions : Histopathological findings showed the bare area to be a multilayered structure. In cases where tumors are located underneath the bare area, it could be important to dissect the bare area, with careful attention to its multilayered structure. Surgical dissection of the bare area in the outer layer of the fused peritoneum could allow a sufficient safety margin.

Background

In the process of human development, a liver grows, rotates, and adheres to the right diaphragm [1, 2]. The attachment site of the liver to the right diaphragm is known as the bare area. The liver surface under the bare area is thought to lack a peritoneal covering in adults [3, 4]. The term, bare area, originates from this recognition. However, the bare area actually has a multilayered structure, namely a fused peritoneum generated from the hepatic visceral peritoneum and the diaphragm parietal peritoneum. In hepatic surgery, the bare area sometimes needs to be dissected to provide mobilization of the right liver. Especially, when the tumors are located below the bare area, it could be important to recognize the multilayer structure when surgically approaching the tumors, which might result in prevention of tumor cell dissemination.

The aim of this paper was to analyze the structure of the bare area by an integrated study of surgical laparoscopic images and histopathological studies and to describe surgical procedures focusing on the multilayered structure.

Methods

Representative patients

All patients who underwent operations in our department from January 1990 to March 2018 were candidates of this retrospective and observational study. Written informed consent was obtained from all patients in accordance with the ethical guidelines of Kyoto University Hospital. This study was approved by the ethics committee of Kyoto University (approval number: R-1721) and performed in accordance with the 1964 Helsinki Declaration.

Surgical procedure and patient care

All patients underwent multi-detector computed tomography (CT) scan before the operation. Three-dimensional images were reconstructed from the CT data using SYNAPSE VINCENT (Fujifilm, Tokyo, Japan).

The surgical procedures used in the study have been described in previously published articles [5–7]. In open hepatectomy, hepatic resection was performed by means of a Mercedes incision or an inverted L-shaped incision. The liver was mobilized, and an intraoperative ultrasound was performed to assess the lesions. An intermittent Pringle maneuver, or selective vascular clamping if necessary, was applied to occlude the blood inflow of the liver. Hepatic parenchymal resection was performed using CUSA Excel (Integra LifeSciences, Plainsboro, NJ, USA) and bipolar cautery equipped with a channel for water dripping.

In laparoscopic hepatectomy, the patient was placed in the left half-lateral decubitus position to expose the lateral and posterior aspect of the right lobe. Pneumoperitoneal pressure by carbon dioxide was maintained at 8–12 mmHg. For the assessment of the tumor location and its relationship with the vasculature, intraoperative contrast-enhanced ultrasonography was routinely performed. The transection of the liver surface was performed using an ultrasonically activated scalpel, and the parenchymal resection was performed by clamp crush methods using an intermittent Pringle maneuver. Bipolar electrocoagulation was performed for minor bleeding.

Cadaver anatomy

The study utilizing donated cadavers was approved by the ethics committee of Kyoto University (approval number: R-1785). One liver was studied in a Thiel-embalmed female cadaver in her seventies. The bare area of the liver was dissected sharply from the diaphragm using surgical scissors, and the right liver surface was exposed, changing the dissection layers.

Histological examination

Surgical materials were fixed by formalin for approximately 48 hours. After a thorough rinsing, the tissues were then embedded in paraffin and sectioned into 3 μ m-thick sections. Hematoxylin-eosin (HE) staining, Masson trichrome staining, and Sirius red staining were performed according to the standard protocol [8–10].

Results

Histological findings of the bare area of the liver

A 75-year-old man underwent a right hemi-hepatectomy for a huge hepatocellular carcinoma with partial resection of the right diaphragm. The right liver was removed without dissection of the bare area. Histological specimens were obtained from the bare area. HE staining revealed no direct tumor invasion into the diaphragm. The serial sections were also stained by the Masson trichrome and Sirius red stains to clarify the structure the connective tissues between the liver and the diaphragm. There were sparse connective tissues lying in a width of 100–200 μ m between the liver and the diaphragm in low-magnification micrographs (Fig. 1a, 1c, and 1e). The high-magnification micrographs showed that the liver parenchyma was covered with liver serosa (visceral peritoneum) and approximately 100 μ m-thick dense sub-serosal tissues. In addition, the diaphragm tendon and its epimysium were covered with parietal peritoneum (Fig. 1b, 1d, and 1f).

The multilayered structure of the bare area was composed of the liver parenchyma, sub-serosal connective tissue, liver serosa, parietal peritoneum, retroperitoneal connective tissue, epimysium of the diaphragm, and the diaphragm, in order from the liver to the diaphragm (Fig. 2). There were 200–300 μ m-thick connective tissues between the liver parenchyma and epimysium of the diaphragm of the bare area, and the liver serosa was in close proximity to the parietal peritoneum that caused peritoneal fusion. There are two layers in the dissection of the bare area in surgical procedures, an outer layer of the fused peritoneum (near the diaphragm) and an inner layer of the fused peritoneum (near the liver).

Macroscopic findings of the bare area of the liver

In order to examine the possibility of recognizing the multilayered structure of the bare area intraoperatively, a cadaver and two cases of laparoscopic hepatectomy were analyzed. In the cadaver study, when dissecting the bare area near the diaphragm, the surface of the liver was seen to be covered with dense fibrous tissues (Fig. 3a). On the other hand, when dissecting near the liver, the raw surface of the liver was exposed (Fig. 3b). In case of laparoscopic bare area dissection, surgeons could sometimes recognize the line between the sparse connective tissue on the diaphragm side and the dense fibrous tissue on the liver side, when the liver exerted traction in the counter direction of the diaphragm (Fig. 3c). When dissecting the bare area near the liver (the inner side of the fused peritoneum), the liver surface covered only by a thin connective tissue was exposed (Fig. 3d). Therefore, it was possible to recognize

the multilayered structure of the bare area and select the dissection layer under the magnified view of laparoscopy.

Case presentation of a colorectal liver metastatic tumor located directly beneath the right diaphragm

One representative case of a colorectal liver metastatic tumor is presented to demonstrate the importance of selecting the appropriate dissection layer of the bare area for good surgical therapeutic effect. A 64-year-old woman underwent laparoscopic right hemi-colectomy for colon cancer. Five months after the operation, a metastatic liver tumor, 15 mm in diameter, was found at hepatic segment 7. She received chemotherapy for six months and then underwent laparoscopic partial hepatectomy. Preoperative CT revealed a tumor 10 mm in diameter, located directly underneath the right dome of the diaphragm (Fig. 4a and 4b). Because of a possible direct invasion of the tumor into the right diaphragm, the bare area was dissected carefully near the diaphragm (the outer layer of the fused peritoneum) (Fig. 4c). There was no gross invasion observed, and the right liver was dissected from the diaphragm at the outer layer of the fused peritoneum with a dense white connective tissue (Fig. 4d).

The low-magnification micrographs revealed that the tumor was located quite close to the liver serosa, although there was no direct invasion into the diaphragm (Fig. 5a, 5c, and 5e). However, tumor cells expanded to the sub-serosal connective tissue in the high-magnification micrographs (Fig. 5b, 5d, and 5f). In this surgery, the bare area was dissected at the outer layer of the fused peritoneum, and the parietal peritoneum and the partial retroperitoneal connective tissue were removed with the liver. Therefore, surgical margins of approximately 500 μ m were reserved.

Discussion

The present study results revealed that the bare area has a multilayered structure. The bare area has been recognized as a hepatic surface area lacking the peritoneum [3, 4]. The term “bare area” is derived from this recognition. However, in the fetal hepatic developmental process, the liver enlarges, rotates, and adheres to the diaphragm, resulting in the formation of the bare area [1, 2]. This indicates that even the liver surface of the bare area has the peritoneum [11]. The histopathological examination in this study revealed that the bare area had a multilayered structure. The parietal peritoneum covering the diaphragm and the visceral peritoneum, which is the liver serosa, are in close proximity and fused to each other. There is a relatively dense connective tissue on the liver side and sparse connective tissue on the diaphragm side of the fused peritoneal peritoneum. The structure composed of dense connective tissue on the liver side and liver serosa is the Laennec’s capsule [12, 13]. Because the liver surface of the bare area is covered with the Laennec’s capsule, as shown in our study, the phrase “bare area” may not be appropriate.

Application of the histological findings in surgical procedures

Recognition of the multilayer structure of the bare area during surgery: Cadaver anatomy revealed a glossy white connective tissue attached to the liver surface when the diaphragm side was dissected. On the other hand, the dissection on the liver side exposed raw liver surface. Therefore, it is possible to dissect the bare area by changing the dissection layer. It might be difficult to recognize the fusion peritoneum itself because both the parietal peritoneum and the liver serosa are a monolayer peritoneum structure. However, under laparoscopic magnification, the borderline between the loose connective tissue of the outer layer of the fused peritoneum and the dense connective tissue of the inner layer can be recognized. Especially in patients with liver cirrhosis, this boundary might be easy to recognize because the Laennec's capsule has often thickened in such patients. Therefore, it is possible to select the dissection layer at least inside and outside the fusion peritoneum.

Selection of dissection layers of the bare area's multilayers could affect surgical control of tumors: When the tumor is located just below the diaphragm dome, as shown in the representative case presented in this study, the tumor cells could invade very close to the liver surface. Consciously dissecting the outer layer of the fused peritoneum could secure a safe surgical margin from tumors and possibly result in prevention of the dissemination.

Conclusion

The bare area is a multilayered structure. There are dissection layers on the outside and inside of the fused peritoneum, respectively. These dissection layers are visible macroscopically. If tumors are located in close proximity to the diaphragm, dissection of the bare area should be performed with attention to this multilayered structure.

Declarations

Availability of data and materials

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Abbreviations

CT: computed tomography

HE: hematoxylin-eosin

Declarations

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Research involving human participants

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee (ethics committee of Kyoto University, approval number: R-1721 and R-1785) and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Written informed consent was obtained from all patients in accordance with the ethical guidelines of Kyoto University Hospital.

Acknowledgements

Funding

This work was supported in part by grants from the Scientific Research Fund of the Ministry of Education, Culture, Sports, Science, and Technology of Japan (19K09145).

Author contributions

T Ishii: Project development, Data collection, Data analysis, Manuscript writing

S Seo: Project development

T Ito: Data collection, Data analysis

S Ogiso: Data collection

K Fukumitsu: Data collection

K Taura: Data management

T Kaido: Data management

S Uemoto: Project development, Data management

All authors have read and approved the final version of the manuscript.

References

1. Jin ZW, Cho BH, Murakami G, Fujimiya M, Kimura W, Yu HC (2010) Fetal development of the retrohepatic inferior vena cava and accessory hepatic veins: Re-evaluation of the Alexander Barry's hypothesis. *Clin Anat* 23:297–303. <https://doi.org/10.1002/ca.20930>
2. Severn CB (1972) A morphological study of the development of the human liver. II. Establishment of liver parenchyma, extrahepatic ductus and associated venous channels. *Am J Anat* 133:85–107. <https://doi.org/10.1002/aja.1001330106>
3. Lowe MC, D'Angelica MI (2017) Anatomy of hepatic resectional surgery. *Surg Clin North Am* 96:183–195. <https://doi.org/10.1016/j.suc.2015.11.003>
4. Abdel-Misih SR, Bloomston M (2010) Liver anatomy. *Surg Clin North Am* 90:643–653. <https://doi.org/10.1016/j.suc.2010.04.017>
5. Yamamoto Y, Ikai I, Kume M, Sakai Y, Yamauchi A, Shinohara H, Morimoto T, Shimahara Y, Yamamoto M, Yamaoka Y (1999) New simple technique for hepatic parenchymal resection using a Cavitron Ultrasonic Surgical Aspirator and bipolar cautery equipped with a channel for water dripping. *World J Surg* 23:1032–1037. <https://doi.org/10.1007/s002689900619>
6. Ishii T, Hatano E, Yasuchika K, Taura K, Seo S, Uemoto S (2014) High risk of lung metastasis after resection of hepatocellular carcinoma more than 7 cm in diameter. *Surg Today* 44:1900–1905. <https://doi.org/10.1007/s00595-013-0792-1>
7. Fuji H, Seo S, Toda R, Toh T, Ikeno Y, Fukumitsu K, Ishii T, Taura K, Hatano E, Kaido T, Uemoto S (2018) Optimal introduction of laparoscopic liver resection for Child-Pugh B. *Asian J Endosc Surg* 21. <https://doi.org/10.1111/ases.12634>
8. Ishii T, Yasuchika K, Suemori H, Nakatsuji N, Ikai I, Uemoto S (2010) Alpha-fetoprotein producing cells act as cancer progenitor cells in human cholangiocarcinoma. *Cancer Lett* 294:25–34. <https://doi.org/10.1016/j.canlet.2010.01.019>
9. Iezzoni JC (2018) Diagnostic histochemistry in hepatic pathology. *Semin Diagn Pathol* 35:381–389. <https://doi.org/10.1053/j.semdp.2018.10.003>
10. Tanabe K, Taura K, Koyama Y, Yamamoto G, Nishio T, Okuda Y, Nakamura K, Toriguchi K, Yamanaka K, Iwaisako K, Seo S, Asagiri M, Hatano E, Uemoto S (2015) Migration of splenic lymphocytes promotes liver fibrosis through modification of T helper cytokine balance in mice. *J Gastroenterol* 50:1054–1068. <https://doi.org/10.1007/s00535-015-1054-3>
11. Macchi V, Porzionato A, Bardini R, Picardi EE, De Caro R (2013) Surgical anatomy of the posterior liver surface: the retrohepatic lamina as the basis for mobilisation of the right liver. *J Gastrointest Surg* 17:1766–1773. <https://doi.org/10.1007/s11605-013-2299-x>
12. Sugioka A, Kato Y, Tanahashi Y (2017) Systematic extrahepatic Glissonean pedicle isolation for anatomic liver resection based on Laennec's capsule. proposal of a novel comprehensive surgical anatomy of the liver. *J Hepatobiliary Pancreat Sci* 24:17–23. <https://doi.org/10.1002/jhbp.410>

13. Hayashi S, Murakami G, Ohtsuka A, Itoh M, Nakano T, Fukazawa (2008) Connective tissue configuration in the human liver hilar region with special reference to the liver capsule and vascular sheath. *J Hepatobiliary Pancreat Surg* 15:640–647. <https://doi.org/10.1007/s00534-008-1336-8>

Figures




Figure 1

Histopathological images of the bare area including the liver and the diaphragm Fig. 1a and b Hematoxylin-eosin staining Fig. 1c and d Masson trichrome staining Fig. 1e and f Sirius red staining L; liver, SS; sub-serosal connective tissue, VP; visceral peritoneum, PP; parietal peritoneum, E; epimysium, D; diaphragm. Bars in a, c, and e indicate 500 μm , and bars in b, d, and f indicate 100 μm .




Figure 2

Schema of the multilayered structure of the bare area. The visceral peritoneum fuses with the parietal peritoneum. *: inner layer of the fused peritoneum, **: outer layer of the fused peritoneum.




Figure 3

Fig. 3a The liver surface (*) is covered with glossy white connective tissues when dissected in the outer layer of the fused peritoneum (cadaver anatomy). Fig. 3b When dissected in the inner layer of the fused peritoneum, the liver surface (*) is naked compared to the adjacent surface with the liver serosa (cadaver anatomy). Fig. 3c Intraoperative image of a laparoscopic partial hepatectomy for hepatocellular carcinoma in a 69-year-old man: The line between the outer layer and the inner layer of the fused peritoneum is shown by yellow arrowheads. Fig. 3d Intraoperative image of a laparoscopic partial hepatectomy for colorectal liver metastasis in a 71-year-old man: When the inner layer of the fused peritoneum is dissected, the liver surface (*) covered only by thin connective tissue is exposed.




Figure 4

Preoperative and intraoperative images of the 64-year old woman with a colorectal liver metastasis. Fig. 4a Preoperative computed tomography image. The tumor is indicated by a yellow arrowhead. Fig. 4b Three-dimensional image reconstructed by SYNAPSE VINCENT. The tumor is indicated in yellow. Fig. 4c Laparoscopic partial hepatectomy was performed. The bare area dissected is the outer layer of the fused peritoneum. Fig. 4d The right liver upon dissection revealed a dense white connective tissue.




Figure 5

Histopathological images of the liver surface of the resected specimen Fig. 5a and b Hematoxylin-eosin staining Fig. 5c and d Masson trichrome staining Fig. 5e and f Sirius red staining T; tumor, L; liver, SS; sub-serosal connective tissue, VP; visceral peritoneum, PP; parietal peritoneum. Bars in a, c, and e indicate 1000 μm , and bars in b, d, and f indicate 200 μm .