Deciphering the Influence of Multiple Anthropogenic Inputs on Taxonomic and Functional Profiles of the Microbial Communities in Yitong River, Northeast China

Ying Zhang

Northeast Normal University

Yang Huo (✉ huo0814@outlook.com)

Northeast Normal University

Zhiruo Zhang

Jilin Jianzhu University

Suiyi Zhu

Northeast Normal University

Wei Fan

Northeast Normal University

Xianze Wang

Northeast Normal University

Mingxin Huo

Northeast Normal University

Research Article

Keywords: Anthropogenic inputs, Wastewater treatment plant effluent, River ecosystem, Bacterioplankton communities, 16S rRNA gene sequencing, Functional profiles, Fecal contamination indicator

Posted Date: August 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-729563/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Environmental Science and Pollution Research on February 3rd, 2022. See the published version at <https://doi.org/10.1007/s11356-021-18386-2>.

Abstract

We conducted physicochemical parameters analysis, 16S rRNA amplicon sequencing and real-time quantitative polymerase chain reaction to explore the impact of human inputs on the bacterioplankton communities within a tributary of the largest river flowing through a megacity in northeast China. Agriculture largely accounted for the alteration of diversity and functions of the microbial communities. Furthermore, nutrients were significantly declined at the reservoir outlet, and WWTP effluent discharge caused changes in the river microbial community. $\text{NH}_3\text{-N}$ and $\text{NO}_3^-\text{-N}$ were the main environmental factors that drive the shift of the bacteria community, and rare taxa played a more important role in the response to environmental changes compared with the abundant ones. The occurrence of the human-specific fecal indicator was mostly derived from agriculture, and its increase in relative abundance was observed in the effluent. Thus, our study provides guidance for ecological assessment and management of rivers by revealing the response pattern of river bacterioplankton to multiple types of anthropogenic stressors.

1 Introduction

The river ecosystem is vital for the persistence of human civilization (Clark et al., 2018). Microorganisms are essential components for the freshwater environment, playing an important role in material and energy flow in aquatic ecosystems (Falkowski et al., 2008; Zeglin, 2015; Yang and Zhang, 2020). Stream microbial communities drive organic chemical degradation, carbon flow, nitrification or denitrification, and phosphate uptake in rivers, which have been suggested as sensitive bioindicators for water quality (Guecker et al., 2006; Valett et al., 2008; Tank et al., 2010).

As a typical lotic ecosystem, rivers are influenced not only by climate factors but also by the disturbance of anthropogenic effects, such as hydraulic condition, land-use alteration, and wastewater discharge by wastewater treatment plants (WWTPs) (Garcia-Armisen et al., 2014; Liao et al., 2018; Price et al., 2018; Burdon et al., 2020; Yang and Zhang, 2020). The construction of a reservoir may significantly change hydrological regime and biological activity, and consequently impact the water quality and denitrification effect downstream of the river (Li et al., 2015; Schmadel et al., 2018). Land-use alteration is proved to induce the change of the existence form of nitrogen, affecting the efficiency of biological absorption and denitrification, and also contaminants from agricultural land discharge into river by irrigation runoff (Mulholland et al., 2008; Eggen et al., 2014). Aquaculture and human activities are important sources of fecal pollution, causing potential health risks (Vadde et al., 2019).

Rivers in arid and semi-arid regions receive a major portion of their base flow from WWTP effluents (Yu et al., 2020). China discharges more than 60 billion m^3 municipal wastewater per year, of which more than 90% is treated through WWTPs (Tong et al., 2020). Treated effluent discharge into the rivers is considered to alter the availability of nitrogen and phosphorus, and also the relative content of various forms of nitrogen in the receiving water (Ekka et al., 2006; Lewis and Grimm, 2007). The decrease of ammonia nitrogen in the downstream is usually accompanied by an increase of nitrate, which results from nitrification by nitrifying bacteria to a great extent (Sonthiphand et al., 2013).

Until now, many researches have focused on the spatial and temporal taxonomic profile variation of microbes exposed to inputs. However, the functional diversity of river microbial communities and their direct relevance to various anthropogenic stresses have yet to be comprehensively understood (Wang et al., 2017; Mansfeldt et al., 2020). In this study, we selected a river (Yitong River) in northeast China that experiences different human disturbances, including land-use partition, hydraulic changes, and WWTP effluent discharge. High-throughput amplicon sequencing and quantitative polymerase chain reaction (qPCR) were performed to depict the taxonomic and functional diversity of bacterioplankton communities alongside the highly disturbed watershed in different sampling times. The aim of this study is to investigate the relationship between the diversity pattern of river bacterioplankton and multiple types of anthropogenic disturbances for guidance in evaluation and management of the river ecosystem.

2 Materials And Methods

2.1 Study area and sample collection

Yitong River is a secondary tributary of Songhua River in Jilin Province, northeast China, with a total length of 343 km, covering 8,440 km² of the watershed area. The mean annual precipitation of the watershed is 622 mm, with 78.1% accounting for rainfall events from June to September. The river experiences a four-month freezing period from mid-November to late March the following year. We selected a stretch of 80.04 km of the river (43°10' N to 43°57' N, 125°16' E to 125°29' E), which successively flows through an agricultural area and a constructed reservoir (Xinlicheng Reservoir), and then enters a highly urbanized area (Changchun City) that serves a population of 7.538 million. In southern Changchun, the river receives effluent discharged from a domestic WWTP (Dongnan WWTP), with daily wastewater treatment capacity of 150,000 m³ with the oxidation ditch treatment process.

Sampling campaigns were conducted along the above mentioned stretch on October 11, 2019, and August 22, 2020, which represented normal and wet periods, respectively. Eight sampling sites were selected, including agriculture sites (Y0–Y3), of which Y0 contained aquaculture activity and Y3 was the reservoir outlet, and urban sites (Y4–Y7), upstream and downstream of the WWTP effluent discharge (Fig. 1). The latitude and longitude information of each sampling site could be found in the Table S1. For each site, a total volume of 5.5 L surface water was collected into sterile containers (4.5 L in a polypropylene bucket and 1 L in a brown glass bottle). All the containers were carefully rinsed using the river water prior to collection. After collection, the samples were immediately transported to a laboratory and preserved at 4 °C.

2.2 Physicochemical analysis

For each site, water temperature (T), pH, electric conductivity (COND), oxidation–reduction potential (ORP), and total dissolved solid (TDS) were measured in the field using a multiparameter water quality analyzer (ULTRAMETER II™ 6PFC^E, MYRON Co., USA). Ammonia nitrogen (NH₃-N), total phosphorus (TP), and soluble reactive phosphorus (SRP) were measured using an ultraviolet spectrophotometer (Thermo

Scientific Co., USA). Nitrite (NO_2^- -N) and nitrate (NO_3^- -N) were measured by using ion chromatography (Metrohm Co., Switzerland). Fe, Mn, K, Mg, Ca, and Si were measured using inductively coupled plasma-optical emission spectroscopy (ICP-OES, EWA Inc., Beijing), and total organic carbon (TOC) was measured by a total organic carbon analyzer (Shimadzu Co., Japan).

2.3 DNA extraction

Each water sample (500 mL to 1100 mL) was vacuum pump-filtered using a Millipore 0.45 μm filter membrane. DNA was extracted from the membranes using a DNeasy PowerWater Kit (Qiagen Co., USA) following the manufacturer's instructions. Extracted DNA concentration was measured using a NanoPhotometer N50 (IMPLEN Co., Germany). Prepared DNA samples were stored at -80°C until further analysis.

2.4 Amplicon sequencing and bioinformatic analysis

16S rRNA gene V4 region amplicon sequencing was performed on an Illumina MiSeq PE300 platform (Illumina, USA) at Shanghai Majorbio Bio-pharm Technology Co., Ltd. Raw read files were uploaded to the National Center for Biotechnology Information Sequencing Read Archive database (SRA; <https://trace.ncbi.nlm.nih.gov/Traces/sra/>) with the BioProject Accession ID PRJNA683092. A list of the sample names and their corresponding BioSample Accession IDs are shown in Table S2.

The quality of reads and the effect of merge were filtered for quality control to remove barcode primers and low-quality reads by using a Galaxy pipeline in Denglab (Feng et al., 2017). A total of 770,634 high-quality bacterial sequences from 16 samples were obtained after quality control. The clean reads were clustered into OTUs with the UPARSE method at the similarity threshold of 97% (Edgar, 2013). The taxonomy annotation for each OTU in bacteria was assigned against the SILVA database (version 132) (Pruesse et al., 2012; Quast et al., 2013; Yilmaz et al., 2014).

2.5 Diversity analysis

Diversity analysis was performed using R software (version 4.0.2) (R Core Team, 2020). Alpha diversity indices, including Shannon, Inv_Simpson, and Richness, were calculated for each sample in *vegan 2.5-7* (Oksanen et al., 2020). For beta diversity, PCoA was performed based on Bray–Curtis dissimilarity to identify potential patterns and relationships within the bacteria community and the sites where the sample was collected.

Distance-based redundancy analysis was used to test the relationships between the bacterial communities and the water chemistry parameters by using *vegan 2.5-7* (Oksanen. et al., 2020; Xu et al., 2020). All environmental variables apart from pH were $\log_{10}(x + 1)$ transformed and normalized (Li et al., 2018). Forward selection was performed to select which variables were statistically significant in shaping the bacteria community composition, and the statistical significance was determined through ANOVA-like permutation tests (with 999 permutations) using the *anova.cca* function in the package (Blanchet. et al., 2008).

2.6 Co-occurrence network and functional analysis

The top 10 phyla network was constructed by calculating the Spearman's correlations with $r > 0.6$ and p value < 0.05 in *Hmisc* 4.5-0 (An et al., 2018; Wang et al., 2020; Harrell, 2021). *Cytoscape* software (version 3.8.0) was applied for network visualization (Shannon et al., 2003).

Piphillin was used to predict functional diversity aligned to the KEGG database (version Oct2018) based on microbial 16S rRNA genes at the clustering threshold of 97% similarity (Iwai et al., 2016; Narayan et al., 2020). The visualization of predicted function diversity was plotted by R package *pheatmap* 1.0.12 (Raivo, 2019).

2.6 Fecal indicator quantification by qPCR

qPCR assays were performed to investigate the quantitative distributions of total 16S rRNA gene and human-related fecal indicator bacteria. *Bacteroides dorei* plays a significant role in contributing to normal intestinal functioning, and its specie assay is highly specific to the gastrointestinal tract of humans (Bakir et al., 2006; Shanks et al., 2007; Price et al., 2018). A universal *Bacteroides* spp. protocol was used to determine the overall fecal pollution (Price et al., 2018). Standard plasmids were constructed using a pClone007 Vector Kit (TsingKe Biotech Co.,Ltd, Beijing). Each qPCR was performed in a 20 μ L reaction mixture containing 7.2 μ L of ddH₂O, 10 μ L of fast-start essential DNA green master (Roche Ltd., Germany), 0.4 μ L of each primer (10 μ M), and 2 μ L of DNA template. Each sample was conducted in triplicate. The primer pairs and thermal programs are detailed in Table S3.

3 Results And Discussion

3.1 Anthropogenic stress on spatiotemporal variation of physicochemical characteristics of surface water in Yitong River

The environmental factors of Yitong River are summarized in Table S4. Figure 2 shows that the contents of TOC, NH₃-N, and TP were at a high level across the agriculture area, which may be caused by irrigation runoff and aquaculture wastewater. As a drinking water protection area, the river exhibited a decline in the concentration of N and P at the reservoir outlet (Fig. 2a, b). The maximum concentrations of NO₂⁻-N and NO₃⁻-N were found in the WWTP effluent in both periods (10.18 mg/L and 12.67 mg/L). The variation of ammonia nitrogen in the wet period before and after the discharge of WWTP effluent fluctuated more than that in the normal period (Fig. 2a). The concentration of TP and SRP was higher in the wet period than in the normal period at most sampling points. TOC concentrations showed a significant difference between wet and normal periods (Fig. 2c).

3.2 Anthropogenic stress on bacterioplankton diversity variation and community dissimilarity among the samples

High-throughput sequencing was performed to profile the spatial and temporal distribution of the bacterioplankton community in Yitong River. After quality filtering, 770,634 high-quality reads were obtained from the eight sites. Those reads were clustered into 5,674 OTUs at the 97% sequence identity. The Shannon index values varied from 2.6 to 6.0, the Inv_Simpson index values ranged from 5.5 to 101.8, and the Richness index values varied from 783 to 2,315 (Table S5). The richness at the reservoir outlet decreased, thereby indicating lower alpha diversity (Fig.2c). Although both the maximum and minimum values of the alpha diversity indices appeared in the wet period, the normal period exhibited larger differences in terms of Shannon indices (Fig.3a). Precipitation in the wet period might accelerate the flow velocity of the river and promote the recovery of microbial diversity in the surface water, which resulted in similar richness to the upstream sites of the WWTP effluent in the wet period (Fig.3c). In the normal period, the bacterial diversity of the downstream site, being adjacent to the WWTP effluent (Y6A), increased (Fig.3b, c). This result suggested that the WWTP effluent accounted for the increasing rate of the total river flow during the normal period, and the effluent contributed a large number of microorganisms from the WWTP into the receiving water. This finding is consistent with that of previous studies (Price et al., 2018).

Figure 3 Bacteria community alpha-diversity: Shannon index of all samples, (a) grouped by period, (b) grouped by location. (c) Observed richness of bacteria community for all the samples. (d) PCoA ordination plot describing bacterial community composition (beta diversity) based on the Bray–Curtis similarity.

The PCoA plot suggested an obvious separation between the normal and wet periods in bacterial community composition with a visible dissimilarity between groups. From the agricultural area to the urban area, a clear separation was also observed (Fig. 3d). Despite the different temporal composition patterns, a remarkable shift of the bacteria community did not occur from the upstream to the downstream despite the discharge of WWTP effluent for receiving water. This finding indicates that land-use variation mostly changed the bacterial community composition (Garcia-Armisen et al., 2014; Price et al., 2018).

3.3 Anthropogenic stress on bacterioplankton taxonomic compositions and co-occurrence relationship of the phyla

The 5,674 clustered OTUs belonged to 47 phyla and 575 families with only 49 shared OTUs in all samples. At the phylum level, the top 4 phyla were *Proteobacteria* (45.7%), *Bacteroidetes* (16.9%), *Actinobacteria* (16.2%), and *Cyanobacteria* (9.0%), which accounted for more than 80% of the total sequences (Fig. 4c). *Actinobacteria* and *Cyanobacteria* decreased in the WWTP effluent, similar to a previous study (Romero et al., 2019). In terms of family level (Fig. 4d), the relative abundance of *Sporichthyaceae* decreased apparently from the agricultural area to the urban area. More interestingly, in

the normal period, we found an abrupt increase in the relative abundance of *Acidobacteria* and one of its family *Ilumatobacteraceae* at the reservoir outlet.

The predominance of *Proteobacteria* in freshwater ecosystems was well documented in many previous studies (Xia et al., 2013; Ligi et al., 2014; Dai et al., 2016). *Bacteroidetes*, which widely inhabit marine and freshwater ecosystems (Zeglin, 2015), is specialized in organic chemical degradation (Traving et al., 2017), which explains its high relative abundance in the agricultural area (Fig. 4c). High abundance of freshwater *Actinobacteria* is ubiquitous in lakes with eutrophication (Wu et al., 2007; Debroas et al., 2009; Humbert et al., 2009; Newton and McMahon, 2011), but their abundance often decreases with decreasing oxygen concentrations (Allgaier and Grossart, 2006). *Actinobacteria* has been viewed as a water quality indicator on account of its sensitivity to the condition that causes blooms (Pascual-Benito et al., 2020). On account of agriculture, substances that contained N or P flowed upstream and were stored by the reservoir with low liquidity; this situation might have caused the increase in *Actinobacteria* and *Ilumatobacteraceae* at the outlet. *Cyanobacteria* plays a key part in nutrient cycling in freshwater. Therefore, its variation might indicate the change in the content of different forms of nitrogen (Newton et al., 2011).

A co-occurrence network was constructed to better understand the interaction of microbial members in the effluent-river ecosystem (Fig. 4a, b). Both networks consisted of 11 nodes (phyla) and 54 edges (correlations) (Table S8). The positive interactions between phyla increased from the normal period to the wet period, whereas the correlations decreased. In the normal period, the positive interactions were mainly observed in *Bacteroidetes*, *Acidobacteria*, and *Chloroflexi*, which were all the main phyla in the upstream (Fig. 4c, a), and the negative correlations were mainly observed in *Actinobacteria*, *Verrucomicrobia*, and *Planctomycetes*. However, during the wet period, the positive interactions were mainly observed among *Planctomycetes* and other phyla, and the interactions enhanced obviously in the effluent (Fig. 4c, b), *Chloroflexi*, *Cyanobacteria*, and the negative correlations were mainly observed in *Bacteroidetes* and *Actinobacteria*.

3.4 Anthropogenic stress on bacterioplankton community predicted functions especially nutrients metabolism

Piphillin was applied to predict functional diversity based on the microbial community of 16S rRNA genes. The KEGG annotation results (Iwai et al., 2016; Narayan et al., 2020) show that the main 50 level-KO3 functions belonged to cellular processes, environmental information processing, genetic information processing, human diseases, metabolism, and organismal system level-KO1 categories (Fig. 5). Carbon fixation can be regularly found in *Proteobacteria*, which are dominant in Y0A, Y1A, Y4A, and Y5A. Notably, the decrease in *Proteobacteria* in Y4, Y5, Y6, and Y7 was in accordance with the decline of carbon fixation (Figs. 4c and 5). The core nitrogen cycle involves nitrogen fixation and denitrification. As mentioned above, the highest total concentration of NO_2^- -N and NO_3^- -N in Y5A and Y5 corresponded to the strong correlation between WWTP effluent discharge and the function of the nitrogen cycle. Conversely, the low

N concentration in Y3A and Y3 corresponded to the decreased function of the nitrogen cycle (Figs. 1a and 5).

The predicted functions varied considerably in both periods, and the functions in the normal period exhibited more significant changes than the composition along the river. Studies demonstrated that variations of functional genes often result in the differences of community composition to maintain the ecosystem function. Thus, the community composition may not respond comprehensively to environmental changes (Rousk et al., 2009; Wang et al., 2020).

3.5 Main environmental factors shaping community composition and comparison of the response between abundant and rare taxa

Abundant and rare taxa exhibit unequal life histories, and rare taxa sometimes display unique biogeography, which differs fundamentally from that of abundant taxa. Thus, the rare taxa are essential for understanding community changes (Lynch and Neufeld, 2015). In this study, we defined relative abundance thresholds as 0.01% for abundant and rare OTUs (Chen et al., 2019; Xu et al., 2020). The abundant and rare OTUs are described in Table S6 and Figure S1.

To explore the relationship between bacteria community composition and environmental variables, and whether both the abundant and rare taxa show the same or different response to environmental variables, 17 environmental variables were normalized to determine the evident variables that shape the community composition, as analyzed by RDA for all OTUs, abundant OTUs, and rare OTUs (Fig. 6). Calculating the variance inflation factor determines that TDS, $\text{NH}_3\text{-N}$, $\text{NO}_2^- \text{-N}$, $\text{NO}_3^- \text{-N}$, TP, SRP, and Ca were strongly related to OTU. Slight variances were discovered between all OTUs and rare OTUs (Figs. 6a and 6c). These physicochemical characters explained the comparatively low proportion of the difference of the OTUs to the abundant OTUs, which was consistent with the view that environmental variables mainly shape the community of rare OTUs rather than the abundant OTUs (Xu et al., 2020). In addition, $\text{NH}_3\text{-N}$ is a possible factor that leads to the obvious partitioning between two periods, and $\text{NO}_3^- \text{-N}$ was the main cause for the differences from upstream to downstream of the WWTP effluent, generally belonging to agricultural and urban areas, respectively.

3.6 Identification and quantification of anthropogenic microbial indicators derived from agriculture and WWTP effluent

Two *bacterioides* indicating anthropogenic fecal pollution (*Bacteroides* spp. and *Bacteroides dorei*) were quantified by using real-time PCR. An increasing trend of 16S rRNA and *Bacteroides* spp. from upstream to downstream could be seen, but a high concentration was observed in Y0, which might be caused by agriculture. As for *B. dorei*, the human specialized indicator, the qPCR results indicated few distinctions in the concentrations (Fig. 7a). Significantly, the relative abundance of *Bacteroides* spp. demonstrated that

B. dorei appeared at Y5A and Y5 sites and higher relative abundance was observed during the wet period (Fig. 7b).

Although human-specific fecal pollution was observed in the WWTP effluent, the effluent did not significantly increase the downstream abundance of fecal indicators such as *B. dorei*. On the one hand, the discrepancy observed between the qPCR and NGS results may be due to primer bias either during qPCR or amplicon library preparation or limitations of the taxonomic database used for assignment (Price et al., 2018). On the other hand, the actual absolute concentrations of some sites were lower than the limit of detection and limit of quantification, thereby leading to imprecise real-time PCR assay results in freshwater (Seurinck et al., 2005).

4 Conclusion

In this study, we investigated the response of the bacterioplankton community composition and function to multiple types of anthropogenic inputs, including agriculture, reservoir construction, and WWTP effluent discharge. Our results indicate decreased concentration of nutrients concomitant with an increase in the actors of carbon and nitrogen metabolism at the reservoir outlet. Moreover, temporal and spatial changes were observed in bacterial community composition and function, and these changes were shaped by land-use change and WWTP effluent. The change in precipitation was the main factor that facilitated the changes. All these changes were more significant in the normal period. The RDA analysis results suggest that $\text{NH}_3\text{-N}$ and $\text{NO}_3^-\text{-N}$ were the main environmental variables that drove the change of bacterial community composition. The absolute concentration acquired from qPCR showed evidence of pollution with agriculture as the source. The relative abundance of fecal indicators was significantly higher in the effluent site. Our study on Yitong River provides a reference for evaluating anthropogenic stressors on river ecology by revealing the response pattern of river bacterioplankton.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: All data generated or analyzed during this study are included in this published article and its supplementary information files.

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by the National Natural Science Foundation of China (Nos. 51908109, 52000085 and 52070038), and National Key Research and Development Program of China (2019YFE0117900).

Authors' contributions: Yang Huo: conceptualization, validation and funding acquisition; Ying Zhang: investigation, writing—original draft and software; Suiyi Zhu and Wei Fan: resources; Zhiruo Zhang and Xianze Wang: validation; Mingxin Huo: supervision. All authors have read and approved the final manuscript.

References

- Allgaier M, Grossart H-P (2006) Diversity and seasonal dynamics of Actinobacteria populations in four lakes in northeastern Germany. *Appl Environ Microbiol* 72(5): 3489-3497
- An X L, Su J Q, Li B, Ouyang W Y, Zhao Y, Chen Q L, Cui L, Chen H, Gillings M R, Zhang T, Zhu Y G (2018) Tracking antibiotic resistome during wastewater treatment using high throughput quantitative PCR. *Environ Int* 117: 146-153
- Bakir M A, Sakamoto M, Kitahara M, Matsumoto M, Benno Y (2006) *Bacteroides dorei* sp. nov., isolated from human faeces. *Int J Syst Evol Microbiol* 56(64257): 1639-1643
- Blanchet. F G, Legendre. P, Borcard D (2008) Forward selection of explanatory variables. *Ecology* 89(9): 2623-2632
- Burdon F J, Bai Y H, Reyes M, Tamminen M, Staudacher P, Mangold S, Singer H, Rasanen K, Joss A, Tiegs S D, Jokela J, Eggen R I L, Stamm C (2020) Stream microbial communities and ecosystem functioning show complex responses to multiple stressors in wastewater. *Glob Change Biol* 26(11): 6363-6382
- Chen W, Ren K, Isabwe A, Chen H, Liu M, Yang J (2019) Stochastic processes shape microeukaryotic community assembly in a subtropical river across wet and dry seasons. *Microbiome* 7(1): 138
- Clark D R, Ferguson R M W, Harris D N, Matthews Nicholass K J, Prentice H J, Randall K C, Randell L, Warren S L, Dumbrell A J (2018) Streams of data from drops of water: 21st century molecular microbial ecology. *WIREs Water* 5(4): e1280
- Dai Y, Yang Y Y, Wu Z, Feng Q Y, Xie S G, Liu Y (2016) Spatiotemporal variation of planktonic and sediment bacterial assemblages in two plateau freshwater lakes at different trophic status. *Appl Microbiol Biotechnol* 100(9): 4161-4175
- Debroas D, Humbert J-F, Enault F, Bronner G, Faubladier M, Cornillot E (2009) Metagenomic approach studying the taxonomic and functional diversity of the bacterial community in a mesotrophic lake (Lac du Bourget - France). *Environ Microbiol* 11(9): 2412-2424
- Edgar R C (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 10(10): 996-1000
- Eggen R I, Hollender J, Joss A, Scharer M, Stamm C (2014) Reducing the discharge of micropollutants in the aquatic environment: the benefits of upgrading wastewater treatment plants. *Environ Sci Technol*

48(14): 7683-7689

Ekka S A, Hagard B E, Matlock M D, Chaubey I (2006) Dissolved phosphorus concentrations and sediment interactions in effluent-dominated Ozark streams. *Ecol Eng* 26(4): 375-391

Falkowski P G, Fenchel T, Delong E F (2008) The microbial engines that drive Earth's biogeochemical cycles. *Science* 320(5879): 1034-1039

Feng K, Zhang Z, Cai W, Liu W, Xu M, Yin H, Wang A, He Z, Deng Y (2017) Biodiversity and species competition regulate the resilience of microbial biofilm community. *Mol Ecol* 26(21): 6170-6182

Harrell Jr. F E (2021) Hmisc: Harrell Miscellaneous. R package version 4.5-0. <https://CRAN.R-project.org/package=Hmisc>

Garcia-Armisen T, Inceoglu O, Ouattara N K, Anzil A, Verbanck M A, Brion N, Servais P (2014) Seasonal Variations and Resilience of Bacterial Communities in a Sewage Polluted Urban River. *PLoS One* 9(3): e92579

Guecker B, Brauns M, Pusch M T (2006) Effects of wastewater treatment plant discharge on ecosystem structure and function of lowland streams. *J N Am Benthol Soc* 25(2): 313-329

Humbert J-F, Dorigo U, Cecchi P, Le Berre B, Debroas D, Bouvy M (2009) Comparison of the structure and composition of bacterial communities from temperate and tropical freshwater ecosystems. *Environ Microbiology* 11(9): 2339-2350

Iwai S, Weinmaier T, Schmidt B L, Albertson D G, Poloso N J, Dabbagh K, Desantis T Z (2016) Piphillin: Improved Prediction of Metagenomic Content by Direct Inference from Human Microbiomes. *PLoS One* 11(11): e0166104

Lewis D B, Grimm N B (2007) Hierarchical regulation of nitrogen export from urban catchments: Interactions of storms and landscapes. *Ecol Appl* 17(8): 2347-2364

Li F, Peng Y, Fang W, Altermatt F, Xie Y, Yang J, Zhang X (2018) Application of Environmental DNA Metabarcoding for Predicting Anthropogenic Pollution in Rivers. *Environ Sci Technol* 52(20): 11708-11719

Li R, Chen Q, Tonina D, Cai D (2015) Effects of upstream reservoir regulation on the hydrological regime and fish habitats of the Lijiang River, China. *Ecol Eng* 76: 75-83

Liao K, Bai Y, Huo Y, Jian Z, Hu W, Zhao C, Qu J (2018) Integrating microbial biomass, composition and function to discern the level of anthropogenic activity in a river ecosystem. *Environ Int* 116: 147-155

Ligi T, Oopkaup K, Truu M, Preem J K, Nolvak H, Mitsch W J, Mander U, Truu J (2014) Characterization of bacterial communities in soil and sediment of a created riverine wetland complex using high-throughput

16S rRNA amplicon sequencing. *Ecol Eng* 72: 56-66

Lynch M D J, Neufeld J D (2015) Ecology and exploration of the rare biosphere. *Nat Rev Microbiol* 13(4): 217-229

Mansfeldt C, Deiner K, Machler E, Fenner K, Eggen R I L, Stamm C, Schonenberger U, Walser J C, Altermatt F (2020) Microbial community shifts in streams receiving treated wastewater effluent. *Sci Total Environ* 709: 135727

Mulholland P J, Helton A M, Poole G C, Hall R O, Hamilton S K, Peterson B J, Tank J L, Ashkenas L R, Cooper L W, Dahm C N, Dodds W K, Findlay S E, Gregory S V, Grimm N B, Johnson S L, Mcdowell W H, Meyer J L, Valett H M, Webster J R, Arango C P, Beaulieu J J, Bernot M J, Burgin A J, Crenshaw C L, Johnson L T, Niederlehner B R, O'brien J M, Potter J D, Sheibley R W, Sobota D J, Thomas S M (2008) Stream denitrification across biomes and its response to anthropogenic nitrate loading. *Nature* 452(7184): 202-205

Narayan N R, Weinmaier T, Laserna-Mendieta E J, Claesson M J, Shanahan F, Dabbagh K, Iwai S, Desantis T Z (2020) Piphillin predicts metagenomic composition and dynamics from DADA2-corrected 16S rDNA sequences. *BMC Genomics* 21(1): 56

Newton R J, Jones S E, Eiler A, McMahon K D, Bertilsson S (2011) A guide to the natural history of freshwater lake bacteria. *Microbiol Mol Biol R* 75(1): 14-49

Newton R J, McMahon K D (2011) Seasonal differences in bacterial community composition following nutrient additions in a eutrophic lake. *Environ Microbiol* 13(4): 887-899

Oksanen J, Blanchet. F G, Friendly. M, Kindt. R, Legendre. P, Mcglinn. D, Minchin. P R, O'hara. R B, Simpson. G L, Solymos. P, Stevens. M H H, Szoecs. E, Wagner. H (2020) *vegan: Community Ecology Package*. R package version 2.5-7. <https://CRAN.R-project.org/package=vegan>

Pascual-Benito M, Balleste E, Monleon-Getino T, Urmeneta J, Blanch A R, Garcia-Aljaro C, Lucena F (2020) Impact of treated sewage effluent on the bacterial community composition in an intermittent mediterranean stream. *Environ Pollut* 266: 115254

Price J R, Ledford S H, Ryan M O, Toran L, Sales C M (2018) Wastewater treatment plant effluent introduces recoverable shifts in microbial community composition in receiving streams. *Sci Total Environ* 613: 1104-1116

Pruesse E, Peplies J, Glockner F O (2012) SINA: accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics* 28(14): 1823-1829

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner F O (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 41(D1): D590-D596

R Core Team (2020) R: A Language and Environment for Statistical Computing. <https://www.R-project.org/>

Raivo K (2019) pheatmap: Pretty Heatmaps. R package version 1.0.12. <https://CRAN.R-project.org/package=pheatmap>

Romero F, Sabater S, Font C, Balcazar J L, Acuna V (2019) Desiccation events change the microbial response to gradients of wastewater effluent pollution. *Water Res* 151: 371-380

Rousk J, Brookes P C, Baath E (2009) Contrasting soil pH effects on fungal and bacterial growth suggest functional redundancy in carbon mineralization. *Appl Environ Microbiol* 75(6): 1589-1596

Schmadel N M, Harvey J W, Alexander R B, Schwarz G E, Moore R B, Eng K, Gomez-Velez J D, Boyer E W, Scott D (2018) Thresholds of lake and reservoir connectivity in river networks control nitrogen removal. *Nat Commun* 9(1): 2779

Seurinck S, Defoirdt T, Verstraete W, Siciliano S D (2005) Detection and quantification of the human-specific HF183 *Bacteroides* 16S rRNA genetic marker with real-time PCR for assessment of human faecal pollution in freshwater. *Environ Microbiol* 7(2): 249-259

Shanks O C, Domingo J W, Lu J, Kelty C A, Graham J E (2007) Identification of bacterial DNA markers for the detection of human fecal pollution in water. *Appl Environ Microbiol* 73(8): 2416-2422

Shannon P, Markiel A, Ozier O, Baliga N S, Wang J T, Ramage D, Amin N, Schwikowski B, Ideker T (2003) Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13(11): 2498-2504

Sonthiphand P, Cejudo E, Schiff S L, Neufeld J D (2013) Wastewater effluent impacts ammonia-oxidizing prokaryotes of the Grand River, Canada. *Appl Environ Microbiol* 79(23): 7454-7465

Tank J L, Rosi-Marshall E J, Griffiths N A, Entekin S A, Stephen M L (2010) A review of allochthonous organic matter dynamics and metabolism in streams. *J N Am Benthol Soc* 29(1): 118-146

Tong Y, Wang M, Penuelas J, Liu X, Paerl H W, Elser J J, Sardans J, Couture R M, Larssen T, Hu H, Dong X, He W, Zhang W, Wang X, Zhang Y, Liu Y, Zeng S, Kong X, Janssen A B G, Lin Y (2020) Improvement in municipal wastewater treatment alters lake nitrogen to phosphorus ratios in populated regions. *Proc Natl Acad Sci U S A* 117(21): 11566-11572

Traving S J, Rowe O, Jakobsen N M, Sorensen H, Dinasquet J, Stedmon C A, Andersson A, Riemann L (2017) The Effect of Increased Loads of Dissolved Organic Matter on Estuarine Microbial Community Composition and Function. *Front Microbiol* 8: 351

Vadde K K, Feng Q, Wang J, McCarthy A J, Sekar R (2019) Next-generation sequencing reveals fecal contamination and potentially pathogenic bacteria in a major inflow river of Taihu Lake. *Environ Pollut*

Valett H M, Thomas S A, Mulholland P J, Webster J R, Dahm C N, Fellows C S, Crenshaw C L, Peterson C G (2008) Endogenous and exogenous control of ecosystem function: N cycling in headwater streams. *Ecology* 89(12): 3515-3527

Wang P, Wang X, Wang C, Miao L, Hou J, Yuan Q (2017) Shift in bacterioplankton diversity and structure: Influence of anthropogenic disturbances along the Yarlung Tsangpo River on the Tibetan Plateau, China. *Sci Rep* 7(1): 12529

Wang X, Wang P F, Wang C, Chen J, Hou J, Miao L Z, Feng T, Yuan Q S (2020) Taxonomic and Functional Responses of Sediment Bacterial Community to Anthropogenic Disturbances in the Yarlung Tsangpo River on the Tibetan Plateau. *J Environ Inform* 35(1): 23-33

Wu X, Xi W, Ye W, Yang H (2007) Bacterial community composition of a shallow hypertrophic freshwater lake in China, revealed by 16S rRNA gene sequences. *FEMS Microbiol Ecol* 61(1): 85-96

Xia N, Xia X H, Zhu B T, Zheng S K, Zhuang J (2013) Bacterial diversity and community structure in the sediment of the middle and lower reaches of the Yellow River, the largest turbid river in the world. *Aquat Microb Ecol* 71(1): 43-U168

Xu H, Zhang S, Ma G, Zhang Y, Li Y, Pei H (2020) 18S rRNA gene sequencing reveals significant influence of anthropogenic effects on microeukaryote diversity and composition along a river-to-estuary gradient ecosystem. *Sci Total Environ* 705: 135910

Yang J, Zhang X (2020) eDNA metabarcoding in zooplankton improves the ecological status assessment of aquatic ecosystems. *Environment International* 134: 105230

Yilmaz P, Parfrey L W, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W, Glockner F O (2014) The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. *Nucleic Acids Res* 42(D1): D643-D648

Yu M, Liu S, Li G, Zhang H, Xi B, Tian Z, Zhang Y, He X (2020) Municipal wastewater effluent influences dissolved organic matter quality and microbial community composition in an urbanized stream. *Sci Total Environ* 705: 135952

Zeglin L H (2015) Stream microbial diversity in response to environmental changes: review and synthesis of existing research. *Front Microbiol* 6: 454

Figures

Figure 1

Sampling map of Yitong River. The distances from Y4 to Y5, Y6, and Y7 are 0.4, 0.97, and 4.44 km, respectively.

Figure 2

Main physicochemical characteristics of the sampling sites.

Figure 3

Bacteria community alpha-diversity: Shannon index of all samples, (a) grouped by period, (b) grouped by location. (c) Observed richness of bacteria community for all the samples. (d) PCoA ordination plot describing bacterial community composition (beta diversity) based on the Bray–Curtis similarity.

Figure 4

Bacteria community interaction and relative abundance. The top 10 phyla interaction in the normal period (a) and the wet period (b). The sizes of each shape represent the relative abundance of the phyla. Red line: negative correlations; Green line: positive corrections (Spearman's $r > 0.6$, $p < 0.05$). Relative abundance at phylum level (c) and family level (d).

Figure 6

Relationships between bacteria communities and environmental variables. Redundancy analysis of (a) all, (b) abundant, and (c) rare bacteria community compositions at the OTU level from the eight surface water sites with respect to the 17 measurable variables. Arrows indicate the direction and magnitude of variables.

Figure 7

(a) Copy numbers of *B. dorei*, *Bacteroides* spp., and 16S rRNA gene as determined by qPCR. (b) Relative abundance of *Bacteroides* spp. and *B. dorei* determined by qPCR. (c) Relative abundance of *Bacteroides* spp. obtained by sequencing at species level.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryinformation.docx](#)