

Is Procalcitonin (PCT) a reliable biomarker for preoperative diagnosing of low grade Periprosthetic Joint Infection? A Prospective Study

André Busch (✉ andre.busch@tissueflow.com)

University of Duisburg-Essen

Marcus Jäger

Department of Orthopaedics and Trauma Surgery University of Duisburg-Essen

Harald Engler

Institute of Medical Psychology and Behavioral Immunobiology, University of Duisburg-Essen

Marcel Haversath

Department of Orthopaedics and Trauma Surgery University of Duisburg-Essen

Charlotte Bielefeld

Department of Orthopaedics and Trauma Surgery University of Duisburg-Essen

Stefan Landgraeber

Department of Orthopedics University of Saarland

Alexander Wegner

Department of Orthopaedics and Trauma Surgery University of Duisburg-Essen

Research article

Keywords: Periprosthetic infection, total joint arthroplasty, synovial fluid procalcitonin, complication

Posted Date: March 11th, 2020

DOI: <https://doi.org/10.21203/rs.2.16660/v4>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Musculoskeletal Disorders on April 20th, 2020. See the published version at <https://doi.org/10.1186/s12891-020-03266-6>.

Abstract

Background: Since a „gold-standard“ is missing, diagnosing periprosthetic joint infection (PJI) remains a challenge in orthopedic surgery. The purpose of this study was to evaluate the accuracy of serum and synovial fluid Procalcitonin (S-PCT and SF-PCT) as a diagnostic parameter and to compare it to the biomarkers recommended in the 2018 Definition of periprosthetic hip and knee infection.

Methods: Between August 2018 and July 2019, a prospective cohort study was conducted in 70 patients with painful hip, shoulder and knee arthroplasty. Besides medical history, clinical and laboratory data was gathered. PJI was diagnosed based on the 2018 Definition of periprosthetic hip and knee infection. Preoperative blood and synovial joint fluid were taken for PCT measurement. S-PCT and SF-PCT levels were measured using standard quantitative PCT enzyme immunoassays.

Results: Twenty three patients (33%) were classified as the PJI group and forty seven patient (67%) as the aseptic group. The mean levels of S-PCT were significantly ($p < 0.001$) higher in the PJI group than those in the aseptic group (PJI 0.05 ± 0.21 ng/mL (0.0-1.03) vs. aseptic 0.02 ± 0.03 ng/mL (0.0-0.18)). In synovial fluid, the mean PCT values in the aseptic group were significantly higher ($p < 0.001$) than those of PJI group (PJI 2.7 ± 1.4 ng/mL (0.53-9.7) vs. aseptic 8.7 ± 2.5 ng/mL (0.25-87.9)). S- PCT, with a cut-off level of 0.5 ng/mL, had a sensitivity of 13.0% and a specificity of 91.0%.

SF-PCT, with a cut-off level of 5.0 ng/mL, had a sensitivity of 13.0% and a specificity of 52.0%.

Conclusion: S-PCT and SF-PCT appeared to be no reliable biomarkers in the differential diagnosis of PJI from aseptic loosening in total joint arthroplasty.

Background

Periprosthetic joint infection (PJI) is a severe complication after total joint arthroplasty. It is one of most common reasons for revision surgery in arthroplasty[i]. The 5-year incidence exceeds one percent after primary arthroplasty[ii]. The differentiation between aseptic and septic failure is crucial for surgical planning[iii]. According to the current Consensus Definition for PJI, a minimum of two positive cultures of periprosthetic tissue or the presence of a sinus tract with evidence of communication to the joint or visualization of the implant are major criteria in diagnosis[iv]. However, microbiological diagnostic is occasionally false negative or positive[v]. Conventional serum biomarkers such as white cell count (WCC) and C-reactive protein (CRP) have limited diagnostic accuracy[vi][vii]. Other serum biomarkers such as Interleukin-6 (Il-6) which are often used in inflammation diagnostics reveal also shortcomings in sensitivity and specificity[viii]. A wide spectrum of synovial fluid biomarkers (SF-alpha-1-Defensin, SF-CRP, SF-Il-6) have been utilized with the goal to diagnose PJI [ix]. Yet, there is no “gold standard” for definite diagnosis of PJI [x].

Procalcitonin (PCT) has been utilized as a serum marker in detecting bacterial infection for several years. [xi][xii][xiii]. Besides CRP serum PCT (S-PCT) seems to be the most promising biomarker to differentiate

between aseptic and septic processes [xiv]. PCT, the precursor of calcitonin, is a 116-amino-acid protein produced by the neuroendocrine and the parafollicular cells of the thyroid[xv]. In healthy patients, S-PCT level is in general very low[xvi], but markedly increased in severe bacterial and fungal infections[xvii]. It has been demonstrated that the injection of bacterial endotoxin in normal subjects induces the release of PCT systemically[xviii][xix]. Previous studies about S-PCT as a biomarker in diagnostic of PJI did not reveal consistent results^{7,13,39,40}. On the hand Glehr et al (2013) described S-PCT as a sensitive, but less specific biomarker for detection of PJI, on the other hand Randau et al (2014) and Bottner et al (2007) found S-PCT to be a very specific, but a less sensitive biomarker for diagnosis of PJI^{13,40,50}. Until now, only one study evaluated the effectiveness of synovial fluid (SF-PCT) for diagnosis of PJI in 32 patients. Ngasoongsong et al (2019) assessed SF-PCT as a specific, but less sensitive marker¹⁵.

The purpose of our study was to investigate the diagnostic value of S-PCT and SF-PCT in Periprosthetic Infection in comparison to the currently recommended parameters. As the second study, we evaluate the effectiveness of SF-PCT for diagnosis of PJI. For the first time, SF-PCT is compared to the current most frequent used biomarker (SF-CRP and SF-AD-1). The hypothesis to be tested was: Due to its properties as a reliable biomarker in bacterial infection, S-PCT and SF-PCT are significantly increased in patients with PJI.

Methods

Study design:

After approval of the institutional review board (18-8042-BO), a prospective study was performed of data gathered from Department of Orthopedics and Trauma Surgery from University of Duisburg-Essen, Germany, in patients with persisting pain[i] after hip, knee and shoulder arthroplasty.

All patients signed informed consent forms prior to being enrolled. The study was conducted in accordance with the declaration of Helsinki.

Medical history, clinical examinations, laboratory values including C-reactive protein (CRP) and joint aspiration fluid were gathered preoperatively as routine diagnostic procedures. Based on the findings of the preoperative diagnostic tests, the patients were considered as aseptic or septic according to the 2018 Definition of periprosthetic hip and knee infection[ii].

The study focused on the differentiation between low-grade infects and aseptic cases. According to WAIOT definition patients without two or more signs or symptoms of local inflammation (pain, swelling, redness, warmth, function laesa) were classified as low-grade infects [iii]. In order to determine the impact of renal dysfunction on serum and synovial values of PCT, serum creatinine concentrations were gathered at the time of joint puncture.

Inclusion criteria were a sufficient amount of synovial fluid for all determinations, and full clinical and laboratory data to allow for diagnosis of periprosthetic infection (PJI). Patients were further excluded, if

they showed signs of early postoperative PJI (8 weeks) due to lack of reliability of synovial and serologic markers shortly after surgery [iv][v]. Metallosis, other inflammatory comorbidities (HIV, rheumatic diseases), and previous or concomitant antibiotic therapy were considered as exclusion criteria.

Sample Preparation

All patients gave their written informed consent that surplus material of their blood and synovial samples which is not needed for standard diagnostics is used for research studies.

Blood was taken from the cubital vein the day before surgery. Synovial aspiration was executed avoiding an admixture of blood with an 18-gauge needle. Synovial fluid was aseptically aliquoted into sterile tubes and centrifuged for 8 minutes at 4°C with 2000 g. The synovial fluid samples were put on ice and transported within 60 minutes to Laboratory of Institute of Medical Psychology and Behavior Science University of Duisburg-Essen and frozen at -80° C.

Determination of the Levels of Serum and Synovial Fluid biomarkers:

S-PCT levels were quantified under the use of immunoassay (Centaur, Siemens, Germany) with lower limit of detection of 0.02 ng/mL (normal < 0.5 ng/mL). Serum CRP was analyzed by immune turbidimetry (Centaur, Siemens, Germany) (normal < 0.5 mg/dl). Synovial leukocyte level and percentage of polymorphic neutrophils was measured by flow cytometry with EDTA plasma (normal range, <3000/ μ l and <80%). SF-PCT levels were measured using a standard quantitative PCT enzyme immunoassay kit, according to the manufacturer's instructions (Anti-Procalcitonin antibody ab166963, ABCAM, Cambridge, UK). Synovial alpha-1-Defensin was analyzed using a standard quantitative enzyme immunoassay kit (Human α -Defensin 1 Antibody, R&D Systems Bio-Techne, Minneapolis, USA)(cut-off level 4800 ng/mL). The results were given as standardized signal relative to a tolerance limit value (interpretation values: < 0.9 aseptic, 0.9–0.99 unspecific, \geq 1.0 septic). Synovial CRP was analyzed under use of a quantitative enzyme-linked immunoassay (CRP ELISA (EU59131), IBL International GmbH, Hamburg, Germany)(cut-off level (> 6,9 mg / l)).

Statistical Analysis:

The data were processed with the statistical software package SPSS. Basic descriptive statistics were used to analyze clinical and laboratory values. Normally distributed continuous data were shown as mean \pm standard deviation (SD) and compared using student's t-test. Non-normally distributed continuous data were shown as mean and compared using the Mann–Whitney U test. A p value < 0.05 was considered statistically significant. Sensitivity, specificity, area under the curve (AUC) and their 95% confidence interval (CI) for any cut-off level were calculated via receiver operating characteristic (ROC) analysis.

Results

Patients:

From July 2018 to June 2019, 78 patients introduced themselves with persisting pain after hip, knee and shoulder arthroplasty in the consultation hour. 70 patients could be included in the study. Three patients were excluded due to insufficient amount of synovial fluid via preoperative puncture. Two patients were excluded due to inflammatory diseases, another two due to early postoperative PJI. One patient was excluded because antibiotic therapy was already started prior to the puncture. All 78 patients who introduced themselves in consultation hour were operatively treated. In all 78 cases histological specimens were taken according to the current recommendations in the 2018 Definition of periprosthetic hip and knee infection ²¹.

From the 70 included patients, 47 patients were identified as having an aseptic joint effusion according to the Definition of Parvizi et al. (2018) were included into the study. The group included 27 women and 20 men with a mean age of 66 ± 12.5 (38-88) years. There were 18 knees, 27 hips and 2 shoulders. The group consisted of 45 patients with polyethylene wear debris induced osteolysis and 2 hips with corrosion of modular head-neck junction. The mean BMI (Body Mass Index) was 26.7 ± 3.1 (22-37).

In the same period, 23 patients were classified as having a PJI according to the Definition of Parvizi et al. (2018). No patient was classified as having a high-grade infect according to the WAIOT definition. The group consisted of 15 women and 8 men with a mean age of 72 ± 11.3 (47-89) years. There were 3 knees, 17 hips and 3 shoulders. The mean BMI was 27.1 ± 7.3 (19-45). In 16 aspirations joint fluid was tested positive in microbiological culture. Bacteria were identified in 16 (70%) of 23 patients of the infection group. Staphylococci were found in 11 (69%), Propioni bacteria and Enterococci in each two (13%) and Serratia marcescens were found in one (6%). In 7 patients (29%) in the infection group with positive histologic specimens for infection, no bacteria could be isolated after 14 days incubation. The patients who were identified as having PJI were operatively revised via two-stage revision with implantation of an intermittent antimicrobial-impregnated spacer.

There were no significant differences in age ($p=0.32$) sex ($p=0.53$) and age at time of surgery ($p=0.70$) between the two groups. The distribution of site of joint arthroplasty was significantly different between the two groups ($p=0.01$) with higher rates of hip arthroplasties in the PJI group.

The S-PCT measurement was positive in 1 joint and negative in 69 (see figure 1). The mean S-PCT level in the PJI and aseptic groups was 0.05 ng/ml (0.00 to 1.03) and 0.02 ng/ml (0.00 to 0.18), respectively ($p < 0.001$). With a cut-off value of 0.5 ng/ml, S-PCT showed a specificity of 91% and a sensitivity of 13%. Comparing these data with the diagnosis criteria of PJI according to the Definition of Parvizi et al (2018), it was found that the PCT-assay was false-positive in 0 and false-negative in 22 cases. The mean SF-PCT in the PJI and aseptic groups was 2.7 ng/ml (0.53 to 9.7) and 8.7 ng/ml (0.25 to 87.9), respectively ($p < 0.001$) (See figure 2). SF-PCT showed with a cut-off level of 5.0 ng/ml a sensitivity of 13% and a specificity of 52%. The mean serum CRP values in the PJI and aseptic groups was 2.3 mg/dl (0.0 to 8.6) and 0.35 mg/dl (0.0-1.9) respectively ($p < 0.001$) (See figure 3). The mean SF-CRP in the PJI and aseptic groups was 19.6 $\mu\text{g/ml}$ (0.6 to 339) and 1.4 $\mu\text{g/ml}$ (0.4 to 5.3), respectively ($p < 0.001$) (See figure 4). The mean synovial fluid alpha-1-defensin levels were significantly higher ($p=0.006$) in PJI group with 3.6

µg/ml (0.2-5.7) than in aseptic group with 2.0 µg/ml (0.2-5.7). Synovial alpha-1-defensin showed a sensitivity of 52% and a specificity of 88% with a cut-off of 4,8 µg/ml. The data of statistical analysis are presented in table 2.

There were no significant differences ($p=0.98$) in the creatinine values between the aseptic and the PJI group. Furthermore, there was no significant correlation between PCT and creatinine values ($p=0.68$).

Discussion

A periprosthetic joint infection (PJI) is a serious complication after total joint replacement. Despite the existence of an international consensus for the definition of PJI, there is no “gold standard” for definite diagnosis of PJI⁹¹⁰. The differentiation between aseptic and septic failure remains a key challenge in orthopedic surgery as the treatment of aseptic failure is completely different to the treatment of PJI [i].

In recent years, several studies reported on the determination of synovial and serum biomarkers for diagnosing periprosthetic infection[ii][iii][iv][v][vi].

CRP is a protein that is synthesized in the liver in response to acute inflammation when there are increased macrophages[vii]. Several studies have endorsed the role of synovial CRP in diagnosing patients with PJI. Most studies reported that synovial CRP is a parameter with high sensitivity and specificity in diagnosing chronic periprosthetic hip infection and favorable to serum CRP [viii][ix][x]. In contrast, Tetreault et al (2014) found no advantage to the use of synovial-fluid CRP over serum CRP in the diagnosis of PJI[xi]. In our study, as expected the additional determination of synovial CRP increases the specificity, but not the sensitivity.

Alpha-Defensins are microbicidal peptides that are active against many Gram-negative and Gram-positive bacteria, fungi, and enveloped viruses[xii]. Bingham et al (2014) concluded that the sensitivity and specificity of the synovial fluid α -defensin assay is superior to other currently available clinical tests [xiii][xiv]. However, there are also reports about low sensitivity values (64%) of synovial alpha-1-defensin [xv]. In our study, synovial alpha-1-defensin was presented as very specific, but less sensitive biomarker to distinguish between aseptic and septic loosening.

The reports about PCT as a diagnostic biomarker for periprosthetic infection are inconsistent (see table 3). Randau et al (2014) and Bottner et al (2007) demonstrated that S-PCT is a very specific, but a less sensitive biomarker for diagnosis of PJI¹³[xvi][xvii]. In contrast, Glehr et al (2013) classified S-PCT as a sensitive, but not specific biomarker for PJI detection⁷. In a systematic review and meta-analysis, Yoon et al (2018) concluded that S-PCT is not recommended for use as a rule-out diagnostic tool for PJI²⁷. Sa-Ngasoongsong et al (2019) reported on SF-PCT as a reliable test for PJI diagnostic with high specificity and sensitivity¹⁵. The results of the current study show that S-PCT is a specific, but less sensitive marker for PJI diagnostic. Interestingly, the aseptic group presented significantly higher SF-PCT values than septic patients.

We could not confirm the initially described hypothesis. We believe that there is no solid evidence to recommend a single determination of S-PCT to rule out PJI. Also, the use of SF-PCT as a parameter in PJI diagnostic's does not appear to be expedient. The lower PCT values in synovial fluid in PJI in comparison with aseptic patients may base on different reasons. Firstly, since not all patients suffering from PJI show bacteremia[xviii], there is no trigger for release of PCT into the blood. It is conceivable that low grade infections such as the majority of PJI do not have the virulence to trigger PCT release. Secondly, in healthy patients transient bacteremia, even after tooth brushing, is a frequent phenomenon that may induce low-grade PCT release [xix][xx][xxi]. Thirdly, it is well known, that in patients with chronic kidney disease PCT levels can be increased due to reduced renal elimination [xxii][xxiii]. Thus, the retention of PCT in patients with kidney diseases could result in false higher PCT values. In our cohort, we could not find any correlation between PCT and creatinine values. Finally, the penetration of PCT into the joint fluid has been infrequently studied. The penetration of PCT into synovial fluid is maybe different in each patient.

Another reason which may have an influence on the results of this study is the possible high rate of false-negatives despite the measurement of serum and synovial fluid biomarkers in addition to conventional microbiological diagnostics. Kheir et al (2018) pointed out that surgeons should be aware of the high rate of false-negatives associated with low-virulence organisms and culture-negative cases due to low sensitivity rates. The sensitivity of the serum and synovial biomarkers appears to be related to organism type[xxiv].

In comparison to serum and synovial CRP as well as synovial AD-1, S-PCT and SF-PCT show a lack of specificity (synovial) and sensitivity (serum) and thus cannot be counted as reliable biomarkers for the differentiation between aseptic processes and PJI (see table 1 and figure 6).

Our study has some limitations. There have been used many different wear couples. It is well known that first generation polyethylene inlays in total joint arthroplasty show higher rates of wear debris induced periprosthetic osteolysis than modern polyethylene inlays[xxv]. Thus, the inflammatory response with release of inflammatory biomarkers is depended on the used material. The duration from initial assessment or symptom onset to fluid collection could not be exactly assessed in all patients. Therefore, the results may be influenced by the incubation period of the germ. Furthermore, unknown factors for elevated infection parameters could have affected the outcome.

One major strength of our study is the design as a prospective trial. To our knowledge, this is one of the first reports about PCT determination in synovial fluid. In addition, patients with chronic diseases (HIV, rheumatic diseases) which could affect the laboratory values were excluded.

Conclusion

S-PCT and SF-PCT appeared to be no reliable alternative biomarker in the differential diagnosis of PJI from aseptic loosening in total joint arthroplasty. The frequently described high sensitivity of alpha-1-Defensin in PJI diagnosis could not be confirmed with our data.

In future studies, the detection of direct parameters for a periprosthetic infection should probably play a more prominent role .

Declarations

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Availability of data and materials: All patient-related data were collected by file research from the archives of the participating centers

Consent to publish: All patients consented to publish personal data in an anonymized form.

Ethics approval and consent to participate: The study was approved by the ethics committee University of Duisburg-Essen (18-8042-BO).

Competing Interests: None of the authors had competing interests.

Acknowledgements: A special thank goes to Alexandra Kornowski who performed the immunoassays.

Authors contribution: All authors ensured that they had furnished a substantial contribution to the article and that they are in agreement with form and contents of the manuscript.

MJ analyzed and interpreted the patient data regarding the scientific relevance and supervised the study as chairman of the department.

AB and AW conceived the study and wrote the article.

CB and HE were responsible for sample analysis.

MH and SL as senior surgeons were responsible for sample collection.

Abbreviations

PJI= Periprosthetic joint infection

WCC= white cell count

CRP= C-reactive protein

IL-6= Interleukin-6

SF=synovial fluid

PCT=Procalcitonin

BMI= body mass index

SD= standard deviation

CI= confidence interval

HIV= Human immunodeficiency virus

References

1. Otto-Lambertz C, Yagdiran A, Wallscheid F, Eysel P, Jung N. Dtsch Arztebl Int. 2017 May 26;114(20):347-353. doi: 10.3238/arztebl.2017.0347. Periprosthetic Infection in Joint Replacement.
2. Gundtoft PH, Overgaard S, Schonheyder HC, Moller JK, Kjaersgaard-Andersen P, Pedersen AB. The "true" incidence of surgically treated deep prosthetic joint infection after 32,896 primary total hip arthroplasties. Acta Orthop 2015. January 30:1–9.
3. Parvizi J, Jacovides C, Antoci V, Ghanem E. Diagnosis of periprosthetic joint infection: the utility of a simple yet unappreciated enzyme. J Bone Joint Surg Am. 2011;93:2242–2248. doi: 10.2106/JBJS.J.01413.
4. Parvizi J, Tan TL, Goswami K, Higuera C, Della Valle C, Chen AF, et al. The 2018 Definition of periprosthetic hip and knee infection: an evidence-based and validated criteria. J Arthroplast 2018. doi:10.1016/j.arth.2018.02.078..
5. Bjerke-Kroll B.T., Christ A.B., McLawhorn A.S., Sculco P.K., Jules-Elysée K.M., Sculco T.P. Periprosthetic joint infections treated with two-stage revision over 14 years: an evolving microbiology profile. J. Arthroplasty. 2014;29(5):877–882. doi: 10.1016/j.arth.2013.09.053.
6. Yuan K, Chen HL, Cui ZM. Diagnostic accuracy of C-reactive protein for periprosthetic joint infection: a meta-analysis. Surg Infect (Larchmt) 2014;15:548–559. doi: 10.1089/sur.2013.066.
7. Kheir MM, Tan TL, Shohat N, Foltz C, Parvizi J. Routine Diagnostic Tests for Periprosthetic Joint Infection Demonstrate a High False-Negative Rate and Are Influenced by the Infecting Organism. J Bone Joint Surg Am. 2018 Dec 5;100(23):2057-2065. doi: 10.2106/JBJS.17.01429.
8. Glehr M, Friesenbichler J, Hofmann G, Bernhardt GA, Zacherl M, Avian A, Windhager R, Leithner A. Novel biomarkers to detect infection in revision hip and knee arthroplasties. Clin Orthop Relat Res. 2013 Aug;471(8):2621-8. doi: 10.1007/s11999-013-2998-3. Epub 2013 Apr 23
9. Deirmengian C, Kardos K, Kilmartin P, Cameron A, Schiller K, Parvizi J. Diagnosing periprosthetic joint infection: has the era of the biomarker arrived? Clin Orthop Relat Res. 2014 Nov;472(11):3254-62. doi: 10.1007/s11999-014-3543-8
10. Patel R, Alijanipour P, Parvizi J., Advancements in Diagnosing Periprosthetic Joint Infections after Total Hip and Knee Arthroplasty., Open Orthop J. 2016 Nov 30;10:654-661. doi: 10.2174/1874325001610010654. eCollection 2016.

11. Simon L, Gauvin F, Amre DK, Saint-Louis P, Lacroix J. Serum procalcitonin and C-reactive protein levels as markers of bacterial infection: a systematic review and meta-analysis. *Clin Infect Dis*. 2004;39(2):206–217.
12. Whicher J, Bienvenu J, Monneret G. Procalcitonin as an acute phase marker. *Ann Clin Biochem*. 2001;38(Pt 5):483–493.
13. Arkader R, Troster EJ, Lopes MR, Junior RR, Carcillo JA, Leone C, Okay TS. Procalcitonin does discriminate between sepsis and systemic inflammatory response syndrome. *Arch Dis Child*. 2006;91(2):117–120.
14. Pontrelli G, De Crescenzo F, Buzzetti R, Jenkner A, Balduzzi S, Calò Carducci F, Amodio D, De Luca M, Chiurchiù S, Davies EH, Copponi G, Simonetti A, Ferretti E, Di Franco V, Rasi V, Della Corte M, Gramatica L, Ciabattini M, Livadiotti S, Rossi P. Accuracy of serum procalcitonin for the diagnosis of sepsis in neonates and children with systemic inflammatory syndrome: a meta-analysis. *BMC Infect Dis*. 2017 Apr 24;17(1):302. doi: 10.1186/s12879-017-2396-7.
15. Sa-Ngasoongsong P, Wongsak S, Jarungvittayakon C, Limsamutpetch K, Channoom T, Kawinwonggowit V. Comparison of Synovial Fluid and Serum Procalcitonin for Diagnosis of Periprosthetic Joint Infection: A Pilot Study in 32 Patients. *Biomed Res Int*. 2018 Oct 1;2018:8351308. doi: 10.1155/2018/8351308. eCollection 2018.
16. Morgenthaler N. G., Struck J., Fischer-Schulz C., Seidel-Mueller E., Beier W., Bergmann A. Detection of procalcitonin (PCT) in healthy controls and patients with local infection by a sensitive ILMA. *Clinical Laboratory*. 2002;48(5-6):263–270.
17. Ferriere F. Procalcitonin, a new marker for bacterial infections. *Annales de Biologie Clinique*. 2000;58(1):49–59.
18. Dandona P, Nix D, Wilson MF, Aljada A, Love J, Assicot M, Bohuon C. Procalcitonin increase after endotoxin injection in normal subjects. *J Clin Endocrinol Metab*. 1994;79(6):1605–1608.
19. Wegner, A., S. Elsenbruch, J. Maluck, J.S. Grigoleit, H. Engler, M. Jager, I. Spreitzer, M. Schedlowski Benson S. Inflammation-induced hyperalgesia: effects of timing, dosage, and negative affect on somatic pain sensitivity in human experimental endotoxemia. *Brain Behav Immun*, 2014. 41: p. 46-54.
20. Cats-Baril W, Gehrke T, Huff K, Kendoff D, Maltenfort M, Parvizi J. International consensus on periprosthetic joint infection: description of the consensus process. *Clin Orthop Relat Res*. 2013;471:4065–4075. doi: 10.1007/s11999-013-3329-4.
21. Parvizi J, Tan TL, Goswami K, Higuera C, Della Valle C, Chen AF, et al. The 2018 Definition of periprosthetic hip and knee infection: an evidence-based and validated criteria. *J Arthroplast* 2018. doi:10.1016/j.arth.2018.02.078.
22. Romanò CL, et al. The W.A.I.O.T. Definition of High-Grade and Low-Grade Peri-Prosthetic Joint Infection. *J Clin Med* 2019. PMID 31083439 Free PMC article.
23. Bilgen O, Atici T, Durak K. Karaeminoğullari, Bilgen MS. C-reactive protein values and erythrocyte sedimentation rates after total hip and total knee arthroplasty. *J Int Med Res*. 2001;29:7–12. doi:

10.1177/147323000102900102.

24. Larsson S, Thelander U, Friberg S. C-reactive protein (CRP) levels after elective orthopedic surgery. *Clin Orthop Relat Res.* 1992;275:237–242.
25. Ellenrieder M, Lenz R, Haenle M, Bader R, Mittelmeier W. Two-stage revision of implant-associated infections after total hip and knee arthroplasty. *GMS Krankenhhyg Interdiszip.* 2011;6(1):Doc17. doi: 10.3205/dgkh000174. Epub 2011 Dec 15.
26. Lee Y. S., Koo K., Kim H. J., et al. Synovial Fluid Biomarkers for the Diagnosis of Periprosthetic Joint Infection. *The Journal of Bone & Joint Surgery.* 2017;99(24):2077–2084. doi: 10.2106/JBJS.17.00123.
27. Vicenti G, Bizzoca D, Nappi V, Pesce V, Solarino G, Carrozzo M, Moretti F, Dicuonzo F, Moretti B. Serum biomarkers in the diagnosis of periprosthetic joint infection: consolidated evidence and recent developments. *Eur Rev Med Pharmacol Sci.* 2019 Apr;23(2 Suppl):43-50. doi: 10.26355/eurrev_201904_17473.
28. Xie K., Dai K., Qu X., Yan M. Serum and Synovial Fluid Interleukin-6 for the Diagnosis of Periprosthetic Joint Infection. *Scientific Reports.* 2017;7(1) doi: 10.1038/s41598-017-01713-4.
29. Yuan K, Li WD, Qiang Y, Cui ZM. Comparison of procalcitonin and C-reactive protein for the diagnosis of periprosthetic joint infection before revision total hip arthroplasty. *Surg Infect (Larchmt).* 2015 Apr;16(2):146-50. doi: 10.1089/sur.2014.034. Epub 2015 Feb 6.
30. Yoon JR, Yang SH, Shin YS. Diagnostic accuracy of interleukin-6 and procalcitonin in patients with periprosthetic joint infection: a systematic review and meta-analysis. *Int Orthop.* 2018 Jun;42(6):1213-1226. doi: 10.1007/s00264-017-3744-3. Epub 2018 Jan 2.
31. Parvizi J, Jacovides C, Adeli B, Jung KA, Hozack WJ Mark B. Coventry Award: synovial C-reactive protein: a prospective evaluation of a molecular marker for periprosthetic knee joint infection. *Clin Orthop Relat Res.* 2012 Jan; 470(1):54-60.
32. Ghanem E, Antoci V Jr, Pulido L, Joshi A, Hozack W, Parvizi J The use of receiver operating characteristics analysis in determining erythrocyte sedimentation rate and C-reactive protein levels in diagnosing periprosthetic infection prior to revision total hip arthroplasty. *Int J Infect Dis.* 2009 Nov; 13(6):e444-9.
33. Greidanus NV, Masri BA, Garbuz DS, Wilson SD, McAlinden MG, Xu M, Duncan CP. Use of erythrocyte sedimentation rate and C-reactive protein level to diagnose infection before revision total knee arthroplasty: a prospective evaluation. *J Bone Joint Surg Am.* 2007;89:1409–1416. doi: 10.2106/JBJS.D.02602.
34. Omar M, Ettinger M, Reichling M, Petri M, Guenther D, Gehrke T, Krettek C, Mommsen P. Synovial C-reactive protein as a marker for chronic periprosthetic infection in total hip arthroplasty. *Bone Joint J.* 2015 Feb;97-B(2):173-6. doi: 10.1302/0301-620X.97B2.34550.
35. Tetreault MW, Wetters NG, Moric M, Gross CE, Della Valle CJ. Is synovial C-reactive protein a useful marker for periprosthetic joint infection? *Clin Orthop Relat Res.* 2014 Dec;472(12):3997-4003. doi: 10.1007/s11999-014-3828-y. Epub 2014 Jul 29.

36. White SH, Wimley WC, Selsted ME Structure, function, and membrane integration of defensins. *Curr Opin Struct Biol.* 1995 Aug; 5(4):521-7.
37. Bingham J, Clarke H, Spangehl M, Schwartz A, Beauchamp C, Goldberg B The alpha defensin-1 biomarker assay can be used to evaluate the potentially infected total joint arthroplasty. *Clin Orthop Relat Res.* 2014 Dec; 472(12):4006-9.
38. Shahi A, Parvizi J. The role of biomarkers in the diagnosis of periprosthetic joint infection. *EFORT Open Rev.* 2017 Mar 13;1(7):275-278. doi: 10.1302/2058-5241.1.160019. eCollection 2016 Jul.
39. Frangiamore SJ, Saleh A, Grosso MJ, Kovac MF, Higuera CA, Iannotti JP, Ricchetti ET. α -Defensin as a predictor of periprosthetic shoulder infection. *J Shoulder Elbow Surg.* 2015 Jul;24(7):1021-7. doi: 10.1016/j.jse.2014.12.021. Epub 2015 Feb 8.
40. Randau TM, Friedrich MJ, Wimmer MD, Reichert B, Kuberra D, Stoffel-Wagner B, Limmer A, Wirtz DC, Gravius S. Interleukin-6 in serum and in synovial fluid enhances the differentiation between periprosthetic joint infection and aseptic loosening. *PLoS One.* 2014 Feb 21;9(2):e89045. doi: 10.1371/journal.pone.0089045. eCollection 2014.
41. Bottner F, Wegner A, Winkelmann W, Becker K, Erren M, Götze C. Interleukin-6, procalcitonin and TNF-alpha: markers of peri-prosthetic infection following total joint replacement. *J Bone Joint Surg Br.* 2007 Jan;89(1):94-9.
42. Klement MR, Siddiqi A, Rock JM, Chen AF, Bolognesi MP, Seyler TM. Positive Blood Cultures in Periprosthetic Joint Infection Decrease Rate of Treatment Success. *J Arthroplasty.* 2018 Jan;33(1):200-204.e1. doi: 10.1016/j.arth.2017.08.034. Epub 2017 Sep 4.
43. Lucas VS, Gafan G, Dewhurst S, Roberts GJ. Prevalence, intensity and nature of bacteraemia after toothbrushing. *J Dent.* 2008;36(7):481–487.
44. Lockhart PB, Brennan MT, Sasser HC, Fox PC, Paster BJ, Bahrani-Mougeot FK. Bacteremia associated with toothbrushing and dental extraction. *Circulation.* 2008;117(24):3118–3125
45. Schlein RA, Kudlick EM, Reindorf CA, Gregory J, Royal GC. Toothbrushing and transient bacteremia in patients undergoing orthodontic treatment. *Am J Orthod Dentofacial Orthop.* 1991;99(5):466–472.
46. Yunus I, Fasih A, Wang Y. The use of procalcitonin in the determination of severity of sepsis, patient outcomes and infection characteristics. *PLoS One.* 2018 Nov 14;13(11):e0206527. doi: 10.1371/journal.pone.0206527. eCollection 2018.
47. Herget-Rosenthal S, Klein T, Marggraf G, Hirsch T, Jakob HG, Philipp T, Kribben A. Modulation and source of procalcitonin in reduced renal function and renal replacement therapy. *Scand J Immunol.* 2005 Feb;61(2):180-6.
48. Kheir MM, Tan TL, Shohat N, Foltz C, Parvizi J. Routine Diagnostic Tests for Periprosthetic Joint Infection Demonstrate a High False-Negative Rate and Are Influenced by the Infecting Organism. *J Bone Joint Surg Am.* 2018 Dec 5;100(23):2057-2065. doi: 10.2106/JBJS.17.01429.
49. Scemama C., Anract P., Dumaine V., Babinet A., Courpied J.P., Hamadouche M. Does vitamin E-blended polyethylene reduce wear in primary total hip arthroplasty: a blinded randomised clinical trial. *Int Orthop.* 2017 Jun;41(6):1113-1118. doi: 10.1007/s00264-016-3320-2. Epub 2016 Nov 4.

Tables

2018 Definition of periprosthetic hip and knee infection

Major criteria (at least one of the following)	Decision
Two positive cultures of the same organism	Infected
Sinus tract with evidence of the communication to the joint or visualization of the prosthesis	

Minor criteria	Score	Decision
Elevated serum CRP or D-Dimere	2	≥6 Infected
Elevated serum ESR	1	
Elevated synovial WBC or LE (++)	3	2-5 Possibly Infected
Positive Alpha-Defensin	3	
Elevated synovial PMN	2	0-1 Not Infected
Elevated synovial CRP	1	

Inconclusive pre-op Score or dry tap	Score	Decision
Preoperative Score	-	≥6 Infected
Positive Histology	3	4-5 Inconclusive
Positive Purulence	3	
Positive Single Culture	2	≤3 Not Infected

Table 1: (CRP (C-reactive protein), ESR (Erythrocyte sedimentation rate), LE (leucocyte esterase), PMN (polymorphonuclear leukocyte), AD-1 (alpha-1-defensin)).

Diagnostic Accuracy

Parameter	PJI (n=23)	Aseptic (n=47)	Cut-Off	Sensitivity (%)	Specificity (%)	AUC	p-value
Serum CRP (mg/dl)	2.3 (0.0-8.6)	0.35 (0.0-1.9)	0.5	57	81	0.70 (0.59-0.87)	<0.001
Synovial CRP (mg/ml)	19.6 (0.6-339)	1.4 (0.4-5.3)	6.9	26	100	0.79 (0.65-0.91)	<0.001
Serum PCT (ng/ml)	0.05 (0.0-1.03)	0.02 (0.0-0.18)	0.1 0.3 0.5	26 17 13	81 84 91	0.53 (0.39-0.68)	<0.001
Synovial PCT (ng/ml)	2.7 (0.53-9.7)	8.7 (0.25-87.9)	1.0 5.0	87 13	0 52	0.26 (0.13-0.38)	<0.001
Synovial AD-1 (µg/ml)	3.6 (0.2-5.7)	2.0 (0.2-5.7)	4.8	52	88	0.67 (0.52-0.82)	0.006

Table 2: Diagnostic accuracy of PJI (Periprosthetic Joint Infection) diagnosis using serum or synovial fluid biomarkers (CRP (C-reactive protein), AUC (Area under the curve), PCT (Procalcitonin) AD-1 (alpha-1-defensin))

Literature overview

Author	Parameter	Cut-Off	Sensitivity	Specificity	p-value
Current study	Serum PCT	0.5 ng/ml	13	91	<0.001
		1.0 ng/mL	87	0	
	Synovial fluid PCT	5.0 ng/mL	13	52	
Glenn et al (2013)	Serum PCT	0.055 ng/mL	81	54	0.038
		0.36 ng/mL	90	33	
Randau et al (2014)	Serum PCT	46 ng/mL	13	100	
Sa-Ngasoong-song P et al (2019)	Serum PCT	0.1 ng/mL	65	92	< 0.001
		0.3 ng/mL	50	100	
		0.5 ng/mL	40	100	
	Synovial fluid PCT	0.08 ng/mL	90	83	< 0.001
		0.12 ng/mL	80	92	
		0.16 ng/mL	55	91	
Bottner et al (2007)	Serum PCT	0.3 ng/mL	33	98	n.a.

Table 3: Procalcitonin: Overview of sensitivity and specificity values in different studies (PCT (Procalcitonin))

Figures




Figure 1

Serum PCT (S-PCT): Log-scale dot plots demonstrate the diagnostic separation of study groups




Figure 2

Synovial PCT (SF-PCT): Log-scale dot plots demonstrate the diagnostic separation of study groups




Figure 3

Serum CRP (S-CRP): Log-scale dot plots demonstrate the diagnostic separation of study groups




Figure 4

Synovial CRP (SF-CRP): Log-scale dot plots demonstrate the diagnostic separation of study groups




Figure 5

Synovial alpha-1-Defensin (SF-AD-1): Log-scale dot plots demonstrate the diagnostic separation of study groups




Figure 6

ROC analysis of all parameters

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [StrobePCTArticle.doc](#)