TRIB3 Regulates FSHR Expression in Human Granulosa Cells Under High Levels of Free Fatty Acids

Nan Wang

Department of Gynecology and Obstetrics, Ruijin Hospital Affiliated with the Medical School of Shanghai Jiao Tong University

Chenchen Si

Department of Gynecology and Obstetrics, Ruijin Hospital Affiliated with the Medical School of Shanghai Jiaotong University

Lan Xia

Department of Gynecology and Obstetrics, Ruijin Hospital Affiliated with the Medical School of Shanghai Jiao Tong University

Xian Wu

Department of Gynecology and Obstetrics, Ruijin Hospital Affiliated with the Shanghai Jiao Tong University

Sheng Zhao

Department of Gynecology and Obstetrics, Ruijin Hospital Affiliated with Shanghai Jiao Tong University Medical School

Huihui Xu

Department of Gynecology and Obstetrics, Ruijin Hospital Affiliated with the Medical School of Shanghai Jiao Tong University

Zhide Ding

Department of Histology, Embryology, Genetics and Development Biology, Shanghai Key Laboratory for Reproductive Medicine, Shanghai Jiao Tong University School of Medicine

Zhihong Niu (✉ kangniu@sina.com)

Department of Obstetrics and Gynecology, Ruijin Hospital Affiliated with the Medical School of Shanghai Jiao tong University <https://orcid.org/0000-0003-2840-0981>

Research Article

Keywords: tribbles pseudokinase 3 (TRIB3), follicle stimulating hormone receptor (FSHR), Akt/GSK3 β pathway, palmitic acid, human granulosa cells

Posted Date: July 21st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-730466/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Granulosa cells (GCs) in cumulus oophorus highly express follicle stimulating hormone receptor (FSHR), which is the most important mediator of both estradiol synthesis and oocyte maturation. Obese women have elevated free fatty acids (FFAs) levels in their follicular fluids and decreased FSHR expression in GCs, which is related to an altered protein kinase B/glycogen synthase kinase 3 β (Akt/GSK3 β) signaling pathway. Such FFA increases accompany 3-fold rises in pseudokinase 3 (TRIB3) expression and reduce the Akt phosphorylation status in both the human liver and in insulinoma cell lines. Therefore, in a high FFA environment, we determined if TRIB3 mediates regulation of FSHR via the Akt/GSK3 β signaling pathway in human GCs.

Methods: GCs from women undergoing in vitro fertilization were collected and designated as high and low FFAs cohorts based on their follicular fluid FFA content. GCs with low FFA levels and a human granulosa-like tumor (KGN) cell line were exposed to palmitic acid (PA), which is a dominant FFA follicular fluid constituent. The effects were assessed of this substitution on the Akt/GSK3 β signaling pathway activity as well as the expressions of TRIB3 and FSHR at both the gene and protein levels by qPCR, Western blot and immunofluorescence staining analyses. Meanwhile, the individual effects of *TRIB3* knockdown in KGN cells and p-AKT inhibitors were compared to determine the mechanisms of FFA-induced FSHR downregulation.

Results: The average FSH dose consuming per oocyte (FSH dose/oocyte) was elevated and Top embryo quality ratio was decreased in women with high levels of FFAs in their follicular fluid. In these women, the GC TRIB3 and ATF4 protein expression levels were upregulated which was accompanied by FSHR downregulation. Such upregulation was confirmed based on corresponding increases in their gene expression levels. On the other hand, the levels of p-Akt decreased while p-GSK3 β increased in the GCs. Moreover, *TRIB3* knockdown reversed declines in FSHR expression and estradiol (E2) production in KGN cells treated with PA, which also resulted in increased p-Akt levels and declines in the p-GSK3 β level. In contrast, treatment of *TRIB3*-knockdown cells with an inhibitor of p-Akt (Ser473) resulted in rises in the levels of both p-GSK3 β as well as FSHR expression whereas E2 synthesis fell.

Conclusions: During exposure to a high FFA content, TRIB3 can reduce FSHR expression through stimulation of the Akt/GSK3 β pathway in human GCs. This response may contribute to inducing oocyte maturation.

Background

Cumulus oophorus bear mature follicles that are composed of a secondary oocyte and surrounded by granulosa cells (GCs, derived from follicular cells). Their formation is one of the necessary conditions for natural fertilization to occur. Normal follicular growth depends on both timely endocrine stimulation and their nutrient and energy statuses [1]. GCs can foster oocyte support localized in the center of the follicle by providing the nutrients via the zona pellucida until oocyte maturation or ovulation. In clinics,

folliculogenesis is severely impaired in some patients afflicted with a metabolic disease such as obesity [2]. As a characteristic marker of nutritional imbalances, both nutritional deprivation and excess were found to have the detrimental effects on non-esterified fatty acids (NEFAs) or free fatty acids (FFAs) and in turn on folliculogenesis in both human and animal models [3]. During in vitro fertilization (IVF), obese women have high FFAs in their follicular fluid [4] and lower estradiol (E2) levels in their serum [5], and usually require more gonadotrophin to induce ovarian stimulation [6–8]. On the other hand, both in vitro and in vivo animal experiments demonstrated that ovarian follicles exposed to high levels of FFAs had an abnormal ovarian steroidogenesis, reduced ovulation rates, and decreased oocyte competence for fertilization [9, 10]. However, the mechanism underlying FFA-induced cytotoxicity in GCs even resulting in oocyte maturation failure remains not fully understood.

Recently, some results demonstrated that the Akt/GSK3 β signal pathway modulates the down-regulation of follicle stimulating hormone receptor (FSHR) expression in the GCs of overweight/obese women [11], but the mechanism controlling this pathway is still unclear. Yan reported that palmitic acid (PA), an important FFA constituent in GCs can induce endoplasmic reticulum stress (ERs) in a human hepatic cell line, accompanied by a significant induction of tribbles of pseudokinase 3 (TRIB3) expression, which was associated with decreased levels of phosphorylated Akt (p-Akt) [12]. Moreover, some other reports showed that TRIB3 induces abnormal Akt phosphorylation under high PA conditions in various cells [13–15]. On the other hand, TRIB3 was involved in cell proliferation and fatty acid oxidation signaling in bovine cumulus cells, which are also derived from follicular cells as granulosa cells and played a key role in oocyte meiotic resumption regulation [16]. Therefore, we presumed that TRIB3 may participate in regulating FSHR expression in GCs. In this study, human GCs were obtained from follicular fluids in clinical infertile subjects and the TRIB3 effects were investigated on FSHR expression in GCs and relevant signal pathways. Meanwhile, some possible targets which can reduce the impact of high FFA levels on GCs function were also evaluated.

Methods

Subject enrollment

A total of 150 women undergoing IVF treatment due to tubal pathology were recruited from the Center for Reproductive Medicine, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine. Patients who were more than 35 years of age or diagnosed with polycystic ovarian syndrome (PCOS, according to Rotterdam criteria [17]), endometriosis and other medical abnormality that could affect folliculogenesis were excluded from this research project. All participants gave their written informed consent regarding the use of clinical data, blood, and follicular fluid samples, and this study was approved by the ethics committee of Ruijin Hospital (2020104a). All patients underwent the standard GnRH antagonist protocol treatment with recombinant human FSH (rFSH; follitropin alfa; Merck, Geneva, Switzerland), which was started on the 2nd day of the menstrual cycle.

Collection of human granulosa cells and follicular fluid

Human follicular fluid was collected during oocyte retrieval. Only follicular fluid from follicles with diameters of 16–20 mm and free of blood upon macroscopic analysis were collected for further analyses. Each sample was centrifuged (250× *g*, 10 min,) and the supernatant was collected and stored at – 80°C until analyzed. The pellets were resuspended with phosphate-buffered saline (PBS), and the suspension was layered over 40 % Percoll (Sigma-Aldrich; Merck KGaA) and centrifuged at 450 × *g*, 4°C for 20 min. GCs were collected from the interphase between follicular fluid and the percoll layer and washed with PBS thrice, then incubated with trypsin (ThermoFisher Scientific, Waltham, MA, USA) at 37°C for 2 min. Next, cells were incubated with red blood cell lysis buffer for 5 min at 4°C to lyse the surplus red blood cells. Finally, the GCs were cultured overnight in DMEM/F12 (Hyclone, Logan, UT, USA) medium supplemented with 10% fetal bovine serum (FBS, Gibco, Waltham, MA, USA) and 1% penicillin-streptomycin (Hyclone).

Cell transfection and varied treatments

GCs and human granulosa-like tumor (KGN) cell line (Feiya Biotechnology Co., Ltd. Jiangsu, China) were used for in vitro study. TRIB3-shRNA and negative control (NC) plasmids were constructed by GeneChem Biotechnology Co., Ltd (Shanghai, China). KGN cells were transfected with the indicated lentiviruses (multiplicity of infection = 50) for 24 h. Then the transfected cells were selected using 1 µg/mL puromycin (Beyotime, Jiangsu, China) to generate stable cell lines. Western blotting analysis confirmed *TRIB3* knockdown in target cells.

All cell lines were cultured in DMEM/F12 supplemented with 10% FBS and 1% penicillin-streptomycin, and incubated at 37°C with 5% CO₂. In this study, palmitic acid (PA) (Sigma-Aldrich, St. Louis, MO, USA) was employed to investigate the FFAs effect on the GCs and KGN cells. PA was diluted with 5% FA-free BSA(Sigma) at 70°C (50 mM stock solutions) and p-Akt(Ser473) inhibitor Palomid 529 (P529) (Selleck, Houston, TX, USA) was dissolved with DMSO (100 mM stock solutions). Before each experiment, 50 mM PA was diluted in cell culture medium and used at a final concentration of 200 µM, and P529 was used at a final concentration of 60 µM in cell culture medium. Recombinant FSH (rFSH) was added to the culture dishes at a final concentration of 10 IU/mL to test the FSH-stimulated protein expression of genes in the FSHR signal pathway. To test the level of E2, 100 nM testosterone (Sigma) was added to the culture medium of shRNA-NC cells and shRNA-TRIB3 cells as a substrate for the synthesis of estrogen in vitro. Subsequently, collected human GCs with low FFAs level (FFA ≤ 0.41mM/L) were divided into four groups and treated with the reagents as follows for 24 h, respectively: (a) Blank control (no intervention); (b) rFSH; (c) 200 µM PA + rFSH; (d) 300 µM PA + rFSH. shRNA-TRIB3 cells were divided into six groups and stimulated with reagents as follows for 24 h: (a) Blank control (no intervention); (b) rFSH; (c) 200 µM PA + rFSH; (d) 300 µM PA + rFSH; (e) 200 µM PA + rFSH + P529; (f) 300 µM PA + rFSH + P529. Meanwhile, KGN cells transfected with shRNA- NC were designated as the control.

Laboratory analysis

Serum and follicular FFA concentrations were quantified as described previously [18]. Serum anti-Mullerian hormone (AMH) levels were measured with the Human AMH ELISA kit (Biotra, Guangzhou,

China), and the levels of FSH, LH, E2, progesterone (P), and total testosterone were measured using a chemiluminescence immunoassay (ECLIA) kit (Beckman Coulter, Brea, CA, USA) according to the manufacturer's instructions, respectively.

RNA extraction and qRT-PCR

Real-time PCR assay was used to determine the mRNA expression levels of *ATF4*, *TRIB3*, *CYP19A1*, *CYP17A1*, and *FSHR*. Total RNA was isolated using a Takara RNA Extraction Kit (Takara, Dalian, China). cDNA was synthesized by reverse transcription using an RT reaction kit (Takara), according to the manufacturer's instructions. The SYBR Green qPCR Mix (Takara) was used to perform real-time quantitative PCR (qPCR). The cycling conditions included 30 min incubation at 95°C, followed by 40 cycles at 95°C for 5 s, 60°C for 34 s and 95°C for 15 s (Applied Biosystems 7500, Fisher Scientific, USA). Meanwhile, *GAPDH* was used as an internal control. The $2^{-\Delta\Delta C_t}$ method was used to calculate relative expression levels (defined as fold-change). Each cell sample in every group was measured thrice and a *P*-value < 0.05 was considered statistically significant. All primer sequences were as follows (5' → 3'):

FSHR forward primer: GGCCATGCTCATCTTCACTG,

FSHR reverse primer: ATAGAGGAAGGGGTGGCAC;

TRIB3 forward primer: GTCCGAGTGAAAAAGGCGTA,

TRIB3 reverse primer: TGCCCTACAGGCACTGAGTA;

ATF4 forward primer: ATGACCGAAATGAGCTTCCTG,

ATF4 reverse primer: GCTGGAGAACCCATGAGGT;

CYP17A1 forward primer: GCTGCTTACCCTAGCTTATTTG,

CYP17A1 reverse primer: ACCGAATAGATGGGGCCATATTT;

CYP19A1 forward primer: CGAAAGTGCTATCGTGGTT,

CYP19A1 reverse primer: TGTGGAAATCCTGCGTCT;

GAPDH forward primer: CACATCGCTCAGACACCATG,

GAPDH reverse primer: TGACGGTGCCATGGAATTTG

Immunofluorescence staining

Immunofluorescence staining was performed according to a previously described protocol [19]. Briefly, collected GCs were incubated for 24 h and then fixed in 4% paraformaldehyde for 30 min. After fixation, the cells were blocked with 3% bovine serum albumin (BSA, Servicebio, Wuhan, China) for 30 min. Then, the cells were incubated with rabbit anti-TRIB3 (cat. 3868, 1:200; Proteintech, Rosemont, IL, USA), rabbit

anti-FSHR (cat. 22665-1-AP 1:200; Proteintech), rabbit anti-ATF4 (cat. GB111137, 1:200; Servicebio) antibody at 4°C overnight, followed by incubation with secondary antibodies AlexaFluor488 (cat. GB25303, 1:200; Servicebio) for 1 h in the dark. Finally, coverslips were mounted on slides with antifade mounting medium containing DAPI (Servicebio) after the slides were washed with PBS thrice. Three independent experiments were performed for each condition. Images were captured using an panoramic desk (P250, 3D Histech, Hungary).

Western blotting

Total protein from the GCs and KGN cells was obtained with radio-immunoprecipitation assay (RIPA, Beyotime, Jiangsu, China) lysis buffer containing 1% protease inhibitor cocktail (Roche, Basel, Switzerland), and protein concentrations were measured using the BCA protein assay kit (Beyotime) [20]. Then, 30 µg protein samples were loaded per well in 4%-12% Bis-Tris polyacrylamide gels (Tanon) and separated by electrophoresis for 1 h at 120 V. Proteins were transferred onto polyvinylidene difluoride (PVDF) membranes (Millipore, Billerica, MA, USA) with transmembrane equipment (Tanon) for 50 min at 400 mA, 4°C. PVDF membranes containing proteins were blocked with protein free rapid blocking buffer (Epizyme, Cambridge, MA, USA) for 20 min and then incubated with specific primary antibodies overnight at 4°C. Primary antibodies included TRIB3 (1:5000, Abcam, Cambridge, UK), FSHR (1:1000, Proteintech), p-GSK3β (1:3000, Abcam), GSK3β (1:5000, Abcam), phospho-Akt (1:2000, CST), Akt (1:1000, Cell Signaling Technology [CST] Danvers, MA, USA), and tubulin (1:5000, Abcam). After washing with tris-buffered saline containing 0.1% of Tween-20 (TBST) for 10 min thrice, the PVDF membranes were incubated with horseradish peroxidase-conjugated anti-rabbit secondary antibody (1:5000, CST) for 1 h and target proteins were detected using the Western Chemiluminescent HRP Substrate Kit (Millipore), according to the manufacturer's instructions. The results of Western blots were quantified by densitometry using ImageJ software (National Institutes of Health, Bethesda, MD, USA) and data were normalized as compared to the control treatment.

Statistical analyses

Quantified data are expressed as mean ± standard error of the mean (SEM). All statistical analysis was performed with SPSS version 24.0 (SPSS Inc., Chicago, IL, USA). After inspection for normal distribution of the data, student's *t*-test was employed to compare two groups. Non-parametric Kruskal-Wallis (KW) test was employed to analyze differences between more than two groups. *P* < 0.05 was considered statistically significant.

Results

Basic characteristics of infertile subjects and IVF cycle outcomes

This study totally enrolled 150 women aged 24–35 years who were infertile due to tubal pathology. All the subjects were divided into two groups (highest quartile vs. other quartiles) based on their FFA levels in

the follicular fluid samples: Group I, patients below the third quartile (FFA $\leq$ 0.41 mM/L, n = 113); Group II, patients above the third quartile (FFA > 0.41 mM/L, n = 37) according to the method of group division [21]. As shown in Table 1, we found that the average FSH dose consumption per oocyte (FSH dose/oocyte) in Group II was higher than that in Group I. Moreover, the number of top quality embryos in Group II was much less than that in Group I. However, there was no statistical significance when other parameters such as age, BMI, or ovarian reserve, were compared between the two groups.

Table 1
Comparison of basic characteristics of study subjects

Parameter	Group I (n = 113)	Group II (n = 37)	P Value
Age (y)	30.6 $\pm$ 2.3	30.9 $\pm$ 2.1	0.77
BMI (kg/m ²)	23.06 $\pm$ 3.75	23.09 $\pm$ 3.06	0.15
AMH (ng/mL)	2.9 $\pm$ 0.8	3.1 $\pm$ 1.0	0.41
Serum FFA (mmol/L)	0.38 $\pm$ 0.09	0.41 $\pm$ 0.11	0.29
Basic sex hormones			
FSH (IU/L)	8.9 $\pm$ 1.7	8.7 $\pm$ 2.0	0.38
LH (IU/L)	3.8 $\pm$ 2.1	4.0 $\pm$ 1.8	0.55
P (ng/mL)	0.71 $\pm$ 0.52	0.86 $\pm$ 0.41	0.89
E2 (pg/mL)	42.4 $\pm$ 11.7	45.9 $\pm$ 14.6	0.49
Total testosterone (ng/mL)	0.32 $\pm$ 0.11	0.37 $\pm$ 0.21	0.32
Ovarian stimulation cycle characteristics			
Total FSH dose (IU)	2325.5 $\pm$ 216.7	2623.0 $\pm$ 253.6	0.09
Duration of stimulation (d)	9.6 $\pm$ 0.9	10.1 $\pm$ 1.2	0.15
E2 on trigger day (pg/mL)	3418.3 $\pm$ 788.4	3077.5 $\pm$ 696.7	0.22
No. of oocytes retrieved	11.6 $\pm$ 2.1	10.8 $\pm$ 2.6	0.19
FSH dose (IU)/oocyte	200.4 $\pm$ 22.6	242.8 $\pm$ 25.4	0.04
Normal fertilization (%)	77.5 $\pm$ 8.4	72.3 $\pm$ 10.6	0.27
Viable embryo (%)	67.9 $\pm$ 12.3	65.2 $\pm$ 10.9	0.13
Top quality embryo (%)	25.9 $\pm$ 8.8	17.3 $\pm$ 10.2	0.04
Values are shown as mean $\pm$ SD. FSH dose /oocyte was calculated as FSH dose (IU) per number of retrived oocytes. Abbreviations: BMI, body mass index; AMH, anti-Mullerian hormone; FSH, follicle-stimulating hormone; P, progesterone; E2, estradiol; LH, luteinizing hormone; FFA, free fatty acid.			

Differences in mRNA and protein expression in GCs

We measured the expression levels of related genes in all of the 150 GC samples. Compared with Group I, *TRIB3* and *ATF4* (Fig. 1A,B) expression levels were upregulated whereas the expression level of *FSHR* (Fig. 1C) was downregulated in Group II, but there were no differences in the expression of the steroid synthesis-related genes *CYP19A1* and *CYP17A1* between the two groups (Fig. 1D,E). To further validate this finding, we used immunofluorescence analysis to detect the expression levels of TRIB3, ATF4 and FSHR protein in human GCs. As shown in Fig. 2A, B, the protein levels of TRIB3 and ATF4 in Group II were also significantly upregulated compared with that in Group I, while the expression of FSHR was downregulated (Fig. 2C).

PA effects on TRIB3 and FSHR expression in vitro

Based on previous work [22], we coincubated 200 μ M PA, 300 μ M PA and 10 IU/mL rFSH with the human GCs (FFA $\leq$ 0.41mM/L) and then measured their protein expression levels, respectively (Fig. 3A). Western blotting analyses showed that TRIB3 protein expression level significantly increased in the PA-treated group (Fig. 3B). In contrast, the FSHR levels exhibited an opposite trend, which means that exposure to PA had concentration dependent inhibitory effects on FSHR expression levels (Fig. 3C). As shown in Fig. 3D, E, the expression level of p-Akt significantly decreased while the expression levels of p-GSK3 β increased in the PA-treated group, while the levels of total AKT and GSK3 β protein remained unchanged.

TRIB3 Knockdown reverses dysregulated Akt/GSK3 β signaling and declines in E2 production

In human GCs, the high-FFA levels induced FSHR dysregulation and changed Akt/GSK3 β signaling. However, it was unclear whether TRIB3 interferes either with FSHR expression or this effect occurs via Akt/GSK3 β signaling. To clarify the underlying mechanism, we knocked down *TRIB3* gene and treated shRNA-TRIB3 cells with P529, an inhibitor of p-Akt (Ser473) formation. As shown in Fig. 4B, both shRNA-NC cells and shRNA-TRIB3 cells displayed significantly increased TRIB3 protein expression when treated with either 200 μ M or 300 μ M PA. *TRIB3* gene knockdown reversed the decreased expression level of FSHR in the PA-treated group (Fig. 4C), This result suggested that in the presence of P529 this inhibitory effect could be prevented.

On the other hand, *TRIB3* gene knockdown instead attenuated the downregulation of p-Akt induced by PA, and co-treatment with inhibitor P529 significantly suppressed Akt phosphorylation without altering total Akt expression (Fig. 4D). Moreover, TRIB3 gene knockdown reversed the rises in the p-GSK3 β expression levels and this decline was inhibited in the presence of P529 (Fig. 4E), but the total GSK3 β expression level remained unchanged. Meanwhile, PA treatment decreased the E2 levels in shRNA-NC cell culture supernatants (Fig. 4F). In addition, knocking down TRIB3 expression attenuated the inhibitory effects induced by palmitate on E2 expression levels, whereas inhibition of p-AKT formation with P529 addition reversed the dampened inhibitory effect of PA on E2 expression levels. Thus, these results strongly

indicate that *TRIB3* knockdown can prevent PA-induced declines in E2 expression levels, whereas P529 co-incubation reversed the blunting effect of PA on E2 expression levels.

Discussion

During oocyte maturation, the granulosa cells play an absolutely indispensable role in fulfilling this procedure. In the present study, some infertile subjects had higher levels of FFA in their follicular fluid than that in the other subjects. There are a growing number of studies in which it was reported that FFAs have detrimental effects on ovarian function. For instance, data from bovine cumulus cells showed that elevated follicular palmitic acid and stearic acid were associated with both impaired oocyte maturation, and fertilization rates as well as poor-quality embryos [23]. In IVF treatment, elevated follicular FFA levels were linked with poor cumulus-oocyte complex (COC) morphology, and they adversely influence ovarian follicular function [24]. Moreover, our previous work also found that the oocyte developmental competence had an association with FFA concentrations in follicular fluid from women undergoing IVF [18]. However, the mechanism of the FFA influence on the developmental potential of oocytes is still not fully elucidated.

The major components of FFAs in the follicular fluid from infertility women are oleic acid (31%), palmitic acid (27%), linoleic acid (25%) and stearic acid (12%) [24]. Amongst these FFAs, palmitic acid was considered as a major factor of lipotoxicity associated with endoplasmic reticulum (ER) stress formation [25]. Studies from bovine models demonstrated that in vitro exposure to pathophysiological palmitic acid concentrations can result in compromised oocyte quality and reduced oocyte developmental competence [26]. On the other hand, mouse oocytes were exposed to lipid-rich follicular fluid from patients, which can induce the dramatic decreased rate of mature oocytes in metaphase II (MII). Additionally, the expression of ER stress markers ATF4, ATF6, and GRP78 significantly increased in cumulus-oocyte complexes (COCs) which matured in lipid-rich follicular fluid [9]. The mechanism is that palmitic acid may inhibit the viability of mouse ovarian GCs and KGN cells through inducing ER stress and stimulating the cAMP/PKA/CREB signaling pathway [10]. In the present study, we found that ATF4 expression, one of the ER stress markers, significantly increased in the GCs from some women who underwent IVF treatment and had high FFA levels in their follicular fluid. Furthermore, we found decreased FSHR expression in the GCs from these women with high follicular FFA content, which may account for why the dose consumed of FSH per oocyte during ovarian stimulation was elevated. Furthermore, after IVF treatment the harvested embryos had poor quality in these women.

More importantly, we also observed that high follicular FFA levels upregulated the *TRIB3* content of GCs. *TRIB3* is a member of the pseudokinase family and several findings within the last decade indicated that it can differentially regulate many important metabolic processes, including lipid metabolism, oxidative stress, and ER stress [27]. As a novel ER stress-inducible gene, *TRIB3* is induced via the ATF4–CHOP pathway and it plays a key role in cell apoptosis during ER stress [28]. Besides, the regulation of *TRIB3* has been proven to be effective in antitumor contexts [29, 30].

TRIB3 mediates differential regulation of the degree of Akt phosphorylation in different cells. In endometrial cancer cells, TRIB3 enhanced cell apoptosis and suppressed cell proliferation and migration ability through inhibition of Akt [14]. However, TRIB3 promoted oral squamous cell carcinoma cell proliferation by increasing Akt phosphorylation [31]. Additionally, TRIB3 hyperexpression in rat muscle cells can impair phosphorylation of Akt [32]. In the current report, we show for the first time that there is a relationship between TRIB3 expression and Akt activity in human granulosa cells. We found that in both human granulosa cells and KGN cell line, PA (one main element in FFAs) stimulation increased TRIB3 expression. Such upregulation accompanied declines in the expression levels of p-Akt and FSHR. Knockdown of TRIB3 diminished the impact of PA on FSHR expression by increasing the phosphorylation of Akt (Ser473) and reducing p-GSK3 β (Tyr216). Meanwhile, decreased E2 synthesis after PA treatment was rescued by TRIB3 knockdown in KGN cell line. Therefore, these results strongly suggest that TRIB3 has a crucial role in mediating PA-induced FSHR downregulation via Akt/GSK-3 β signaling. Such control is also linked with impairment of E2 synthesis. Based on these results, we presume that TRIB3 is a functional upstream regulator of FSHR in human CGs.

The PI3K/Akt pathway is associated with the initiation and progression of a variety of cancers, including breast, colorectal, ovarian and pancreatic cancers, and endometrial carcinoma [33, 34]. As one of the first identified substrates of Akt, GSK3 β was initially identified as an inhibitor of cellular responses to insulin by deactivating glycogen synthase phosphorylation. This enzyme can also influence cell division, growth, and development as an endogenous inhibitor of canonical wingless related integration site (Wnt) signaling [35, 36]. In patient-derived lymphoma cell lines, activated AKT phosphorylates GSK3 β Ser9 which results in the inactivation of GSK-3 β , which elevated p- β -catenin (Ser-675) [37]. On the other hand, phosphorylation of Tyr216 counteracts the inhibitory effect of Ser9 phosphorylation on GSK3 β . GSK3 β activity is instead facilitated by GSK3 β / β -catenin and FSHR downregulation [11]. In this study it was shown that KGN cells under hyper-insulin treatment displayed reduced p-Akt along with increased p-GSK3 β (Tyr216) levels. As a consequence, activated GSK3 β downregulated FSHR expression through decreasing the phosphorylation status of β -catenin at sites Ser552 and Ser675.

During follicle development, FSH induces proliferation and steroidogenesis in GCs through binding to its receptor (FSHR) localized on GCs membrane. Impaired FSHR expression usually results in decreased GC responses to FSH stimulation. In this study, we found that the ovarian response was reduced in patients with high follicular FFA levels, which indicated higher consumption of gonadotrophin to obtain an oocyte. In the shRNA-NC cell culture supernatant, E2 production declined after PA-stimulation and this effect was reversed by *TRIB3* knockdown. On the other hand, treatment with P529 of TRIB3-shRNA cells decreased the E2 to levels that were below those obtained with PA stimulation. Therefore, inhibition of TRIB3 expression may be a new target to improve ovarian response under high FFA conditions.

In the current study, our results showed that TRIB3/Akt/GSK3 β represents an important point of convergence and crosstalk between the PA response and FSHR expression. However, this study still has some limitations including our use of KGN cells as a *TRIB3* gene knockdown model. Although the KGN cell line is more likely to be closer to granulosa cells in normal physiological conditions than other human

cell lines [38], additional studies are warranted because this model does not completely replicate the physiological environment of the human body. Furthermore, the TRIB3 effect on the granulosa cells or even on the folliculogenesis in conditional *TRIB3* gene knockout mouse model also needs further study.

Conclusions

In summary, the present study demonstrated that the elevated FFAs in follicular fluid were associated with increased TRIB3 expression levels and reduced FSHR expression in human GCs. Palmitic acid, one main constituent of free fatty acids, can induce decreased FSHR expression levels via the TRIB3/Akt/GSK3 β pathway in both human GCs and KGN cell line. This research is the first study to elucidate TRIB3 biological roles in GCs exposed to a high level of FFAs. Moreover, it also provides insight into a potential target for improvement of poor reproductive performance in women with metabolic disorders and female infertility treatment in clinics as well.

Abbreviations

BMI: Body mass index; TC: Total cholesterol; TG: Triglycerides; FFAs: Free fatty acids; GCs: Granulosa cells; IVF: In vitro fertilization; PCOS: Polycystic ovary syndrome; ER: Endoplasmic reticulum; Akt: protein kinase B; Wnt: Wingless-related integration site; qPCR: Quantitative PCR; rFSH: Recombinant human FSH; T: Testosterone; TBST: Tris-buffered saline–Tween 20; p: phosphorylated; PI3K: Phosphatidylinositol 3-kinase; KGN: Human granulosa-like tumor cells; GSK3: Glycogen synthase kinase 3; GAPDH: Glyceraldehyde 3-phosphate dehydrogenase; FSHR: FSH receptor; E2: Estradiol; CYP19A1: Cytochrome P450 19A1; CYP17A1: Cytochrome P450 17A1; ATF4: Activating transcription factor 4; BSA: Bovine Serum Albumin; COCs: Cumulus-oocyte complexes; TRIB3: Tribbles pseudokinase 3; NEFAs: Non-esterified fatty acids; PA: Palmitic acid; PBS: Phosphate-buffered saline; FBS: Fetal bovine serum; AMH: Anti-Mullerian hormone; MII: Metaphase II. rFSH: Recombinant FSH; P529: Palomid 529; ECLIA: chemiluminescence immunoassay.

Declarations

Acknowledgements

The authors are very grateful to Prof. Peter Reinach for his extensive and detailed support in improving the manuscript.

Authors' contributions

NZ and WN designed the study. WN and SC performed the experiments. XL, WX, ZS and XH collected human GCs samples. WN wrote the initial manuscript. NZ and DZ supervised and supported the whole project, and edited the manuscript. DZ provided final approval of the manuscript. All authors read and agreed on the final version of the manuscript.

Funding

This work received grant support from the National Natural Science Foundation of China (No.81671530, 82071694) and Science and Technology Commission of Shanghai Municipality (No. 21140904000, 201409005800).

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

All participants signed informed consent forms approved by the Shanghai Jiao Tong University Committee on the Use of Human Subjects in Medical Research. Ethical approval of follicular fluid collection was obtained from the institutional Ethics Committee of Ruijin Hospital, Shanghai Jiao Tong University School of Medicine.

Consent for publication

Not applicable.

Competing interests

The authors declared that they have no competing interests.

References

1. Warzych E, Lipinska P. Energy metabolism of follicular environment during oocyte growth and maturation. *J Reprod Dev.* 2020; 66:1-7.
2. Yang PK, Chou CH, Huang CC, Wen WF, Chen HF, Shun CT, Ho HN, Chen MJ. Obesity alters ovarian folliculogenesis through disrupted angiogenesis from increased IL-10 production. *Mol Metab.* 2021; 49:101189.
3. Baddela VS, Sharma A, Vanselow J. Non-esterified fatty acids in the ovary: friends or foes? *Reprod Biol Endocrinol.* 2020; 18:60.
4. Valckx SDM, Arias-Alvarez M, De Pauw I, Fievez V, Vlaeminck B, Fransen E, Bols PEJ, Leroy JLMR. Fatty acid composition of the follicular fluid of normal weight, overweight and obese women undergoing assisted reproductive treatment: a descriptive cross-sectional study. *Reproductive biology and endocrinology : RB&E.* 2014; 12:13-13.
5. Shah DK, Missmer SA, Berry KF, Racowsky C, Ginsburg ES. Effect of Obesity on Oocyte and Embryo Quality in Women Undergoing In Vitro Fertilization. *Obstetrics & Gynecology.* 2011; 118:63-70.

6. Mulders AGMJ, Laven JSE, Eijkemans MJC, Hughes EG, Fauser BCJM. Patient predictors for outcome of gonadotrophin ovulation induction in women with normogonadotrophic anovulatory infertility: a meta-analysis. *Human Reproduction Update*. 2003; 9:429-449.
7. Hamilton-Fairley D, Kiddy D, Watson H, Paterson C, Franks S. Association of moderate obesity with a poor pregnancy outcome in women with polycystic ovary syndrome treated with low dose gonadotrophin. *British journal of obstetrics and gynaecology*. 1992; 99:128-131.
8. Ozekinci M, Seven A, Olgan S, Sakinci M, Keskin U, Akar ME, Ceyhan ST, Ergun A. Does obesity have detrimental effects on IVF treatment outcomes? *BMC women's health*. 2015; 15:61-61.
9. Yang X, Wu LL, Chura LR, Liang X, Lane M, Norman RJ, Robker RL. Exposure to lipid-rich follicular fluid is associated with endoplasmic reticulum stress and impaired oocyte maturation in cumulus-oocyte complexes. *Fertility and Sterility*. 2012; 97:1438-1443.
10. Wan J, Liu S. Rosiglitazone ameliorates palmitic acid-induced endoplasmic reticulum stress and steroidogenic capacity in granulosa cells. *Reprod Biol*. 2020; 20:293-299.
11. Xu P, Huang B-Y, Zhan J-H, Liu M-T, Fu Y, Su Y-Q, Sun Q-Y, Wang W-H, Chen D-J, Liu J-Q. Insulin Reduces Reaction of Follicular Granulosa Cells to FSH Stimulation in Women With Obesity-Related Infertility During IVF. *The Journal of Clinical Endocrinology & Metabolism*. 2018; 104:2547-2560.
12. Yan W, Wang Y, Xiao Y, Wen J, Wu J, Du L, Cai W. Palmitate induces TRB3 expression and promotes apoptosis in human liver cells. *Cell Physiol Biochem*. 2014; 33:823-834.
13. Kwon M, Eom J, Kim D, Kim J, Heredia J, Kang SW, Song Y. Skeletal Muscle Tissue Trib3 Links Obesity with Insulin Resistance by Autophagic Degradation of AKT2. *Cellular Physiology and Biochemistry*. 2018; 48:1543-1555.
14. Qu J, Liu B, Li B, Du G, Li Y, Wang J, He L, Wan X. TRIB3 suppresses proliferation and invasion and promotes apoptosis of endometrial cancer cells by regulating the AKT signaling pathway. *OncoTargets and therapy*. 2019; 12:2235-2245.
15. Nicoletti-Carvalho JE, Nogueira TC, Gorjão R, Bromati CR, Yamanaka TS, Boschero AC, Velloso LA, Curi R, Anhê GF, Bordin S. UPR-mediated TRIB3 expression correlates with reduced AKT phosphorylation and inability of interleukin 6 to overcome palmitate-induced apoptosis in RINm5F cells. *J Endocrinol*. 2010; 206:183-193.
16. Brisard D, Chesnel F, Elis S, Desmarchais A, Sánchez-Lazo L, Chasles M, Maillard V, Uzbekova S. Tribbles expression in cumulus cells is related to oocyte maturation and fatty acid metabolism. *J Ovarian Res*. 2014; 7:44.
17. Group TREA-SPCW. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertility and Sterility*. 2004; 81:19-25.
18. Niu Z, Lin N, Gu R, Sun Y, Feng Y. Associations between insulin resistance, free fatty acids, and oocyte quality in polycystic ovary syndrome during in vitro fertilization. *The Journal of clinical endocrinology and metabolism*. 2014; 99:E2269-E2276.
19. Aimé P, Sun X, Zareen N, Rao A, Berman Z, Volpicelli-Daley L, Bernd P, Crary JF, Levy OA, Greene LA. Trib3 Is Elevated in Parkinson's Disease and Mediates Death in Parkinson's Disease Models. *J*

Neurosci. 2015; 35:10731-10749.

20. Xu B, Zhou M, Wang J, Zhang D, Guo F, Si C, Leung PCK, Zhang A. Increased AIF-1-mediated TNF- α expression during implantation phase in IVF cycles with GnRH antagonist protocol. *Human reproduction (Oxford, England)*. 2018; 33:1270-1280.
21. Ødegård SS, Sand T, Engstrøm M, Zwart J-A, Hagen K. The impact of headache and chronic musculoskeletal complaints on the risk of insomnia: longitudinal data from the Nord-Trøndelag health study. *The journal of headache and pain*. 2013; 14:24-24.
22. Mu Y-M, Yanase T, Nishi Y, Tanaka A, Saito M, Jin C-H, Mukasa C, Okabe T, Nomura M, Goto K, Nawata H. Saturated FFAs, Palmitic Acid and Stearic Acid, Induce Apoptosis in Human Granulosa Cells. *Endocrinology*. 2001; 142:3590-3597.
23. Leroy JL, Vanholder T, Mateusen B, Christophe A, Opsomer G, de Kruif A, Genicot G, Van Soom A. Non-esterified fatty acids in follicular fluid of dairy cows and their effect on developmental capacity of bovine oocytes in vitro. *Reproduction*. 2005; 130:485-495.
24. Jungheim ES, Macones GA, Odem RR, Patterson BW, Lanzendorf SE, Ratts VS, Moley KH. Associations between free fatty acids, cumulus oocyte complex morphology and ovarian function during in vitro fertilization. *Fertility and Sterility*. 2011; 95:1970-1974.
25. Zeng X, Zhu M, Liu X, Chen X, Yuan Y, Li L, Liu J, Lu Y, Cheng J, Chen Y. Oleic acid ameliorates palmitic acid induced hepatocellular lipotoxicity by inhibition of ER stress and pyroptosis. *Nutrition & metabolism*. 2020; 17:11-11.
26. Van Hoeck V, Sturmey RG, Bermejo-Alvarez P, Rizos D, Gutierrez-Adan A, Leese HJ, Bols PE, Leroy JL. Elevated non-esterified fatty acid concentrations during bovine oocyte maturation compromise early embryo physiology. *PLoS One*. 2011; 6:e23183.
27. Mondal D, Mathur A, Chandra PK. Tripping on TRIB3 at the junction of health, metabolic dysfunction and cancer. *Biochimie*. 2016; 124:34-52.
28. Ohoka N, Yoshii S, Hattori T, Onozaki K, Hayashi H. TRB3, a novel ER stress-inducible gene, is induced via ATF4-CHOP pathway and is involved in cell death. *The EMBO journal*. 2005; 24:1243-1255.
29. Li K, Zhang TT, Hua F, Hu ZW. Metformin reduces TRIB3 expression and restores autophagy flux: an alternative antitumor action. *Autophagy*. 2018; 14:1278-1279.
30. Muñoz-Guardiola P, Casas J, Megías-Roda E, Solé S, Perez-Montoyo H, Yeste-Velasco M, Erazo T, Diéguez-Martínez N, Espinosa-Gil S, Muñoz-Pinedo C, et al. The anti-cancer drug ABTL0812 induces ER stress-mediated cytotoxic autophagy by increasing dihydroceramide levels in cancer cells. *Autophagy*. 2020:1-18.
31. Shen P, Zhang T-Y, Wang S-Y. TRIB3 promotes oral squamous cell carcinoma cell proliferation by activating the AKT signaling pathway. *Experimental and therapeutic medicine*. 2021; 21:313-313.
32. Liu J, Wu X, Franklin JL, Messina JL, Hill HS, Moellering DR, Walton RG, Martin M, Garvey WT. Mammalian Tribbles homolog 3 impairs insulin action in skeletal muscle: role in glucose-induced insulin resistance. *American journal of physiology Endocrinology and metabolism*. 2010; 298:E565-E576.

33. Pandurangan AK. Potential targets for prevention of colorectal cancer: a focus on PI3K/Akt/mTOR and Wnt pathways. *Asian Pac J Cancer Prev*. 2013; 14:2201-2205.
34. Grunt TW, Mariani GL. Novel approaches for molecular targeted therapy of breast cancer: interfering with PI3K/AKT/mTOR signaling. *Curr Cancer Drug Targets*. 2013; 13:188-204.
35. Rotte A, Pasham V, Eichenmüller M, Yang W, Qadri SM, Bhandaru M, Lang F. Regulation of basal gastric acid secretion by the glycogen synthase kinase GSK3. *J Gastroenterol*. 2010; 45:1022-1032.
36. Roe ND, Ren J. Akt2 knockout mitigates chronic iNOS inhibition-induced cardiomyocyte atrophy and contractile dysfunction despite persistent insulin resistance. *Toxicol Lett*. 2011; 207:222-231.
37. Chung J, Karkhanis V, Baiocchi RA, Sif S. Protein arginine methyltransferase 5 (PRMT5) promotes survival of lymphoma cells via activation of WNT/ β -catenin and AKT/GSK3 β proliferative signaling. *The Journal of biological chemistry*. 2019; 294:7692-7710.
38. Heng D, Wang Q, Ma X, Tian Y, Xu K, Weng X, Hu X, Liu W, Zhang C. Role of OCT4 in the Regulation of FSH-Induced Granulosa Cells Growth in Female Mice. *Frontiers in endocrinology*. 2020; 10:915-915.

Figures

Figure 1

mRNA expression in GCs from 150 infertility women (A) Expression of TRIB3 was increased in patients of Group II. (B) Expression level of ATF4 was increased in patients of Group II. (C) Expression of FSHR was decreased in patients of Group II. (D,E) No obvious difference in mRNA expression levels of CYP19A1 and CYP17A1 were observed. Data are presented as mean $\pm$ SEM of three independent experiments. * $P < 0.05$ compared with of Group I.

Figure 2

Intracellular localization and expression of TRIB3 ATF4 and FSHR in GCs was detected by immunofluorescence. (A) Expression of TRIB3 protein was upregulated in patients of Group II. (B) Expression of ATF4 protein was upregulated in patients of Group II. (C) Expression of FSHR protein was downregulated in patients of Group II. AlexaFluoro488 (green) was detected at the cell membrane in GC

and cell nuclei were counterstained with DAPI (blue). All the images were acquired at 630× magnification. Scale bar: 20 μ m.

Figure 3

hGCs treated with PA in vitro. (A) Western blot analysis with tubulin used as the loading control. (B) PA increased TRIB3 levels in a concentration-dependent manner. (C) PA decreased FSHR levels in a

concentration-dependent manner. (D,E) Reduced p-Akt and increased p-GSK3 β protein levels indicating the Akt/GSK3 β signaling was aberrantly activated in hGCs treated with PA. *P < 0.05.

Figure 4

Western blot analysis of the Akt/GSK3 β pathway following TRIB3 knockdown in KGN cells. (A) Western blot analysis was performed in shRNA-control and shRNA-TRIB3 cells treated with PA; tubulin was used as the loading control. (B) PA increased TRIB3 levels in a concentration-dependent manner, TRIB3

knockdown reduced this effect. (C) TRIB3 knockdown prevented PA-induced FSHR decrease. (D) TRIB3 knockdown blocked PA-induced decrease of p-AKT. (E) TRIB3 knockdown blocked PA-induced increase in p-GSK3 β . (b–e) P529, together with rFSH and PA, enhanced the reduction of FSHR, p-Akt and the increase of p-GSK3 β protein expression. (F) PA treatment inhibited E2 production, TRIB3 knockdown reversed this effect; P529 enhanced it.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.pptx](#)