Prediction of COVID-19 Cases in Afghanistan Using ARIMA model

Mujtaba Haidari (✉ mujtabahaidari2016@gmail.com)

Lovely Professional University

Research Article

Keywords: COVID-19, ARIMA model, positive cases, Afghanistan, Prediction.

Posted Date: July 21st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-732435/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

In Afghanistan, the novel coronavirus disease 2019 (COVID-19) is spreading rapidly. Currently, we are in third wave of pandemic, in Afghanistan. And recently government of Afghanistan recorded the highest confirmed cases since the start of pandemic. In order to prepare ourselves and make right decision, we need to predict the future. Past information has been used to predict future, but we need to confess that no prediction can be as real future. The data have been used to predict is from twenty first March 2021 to fifteenth July 2021. The forecasting period is from sixteenth July 2021 for 40 days till twenty fourth August 2021. To examine stationarity of data, we used Augmented Dickey-Fuller unit-root. In this study autoregressive integrated moving average ARIMA model has been used to predict future COVID-19 cases. In order to find the best ARIMA parameters, we used Akaike's Information Criterion (AIC) and Bayesian Information Criteria (BIC). The data which we have used in this study are taken from Ministry of Public Health. We found out that ARIMA (0, 2, 1) is best fit model. By using ARIMA (0, 2, 1), we predicted COVID-19 cases in 80 and 95 per cent level of confidence. In 95 (high) and 80 (high) per cent level of confidence COVID-19 positive cases reach 216,159 and 203,979 cases respectively. On the other hand, in 95 (low) and 80 (low) per cent level of confidence after a period of 40-day positive cases will reach 145,780 and 157,961 cases respectively. This study is the first study which predict COVID-19 cases in Afghanistan, so government of Afghanistan, non-government organizations (NGOs) and scholars can use it to plan and prepare themselves to confront uncertain future and protect people against this virus.

Introduction

First case of COVID-19 virus found in China on December 2019, and on March of this year WHO (World Health Organization) declared COVID-19 pandemic (prevalent over a whole country or the world) (The New York Times, 2021). Globally, since the first positive case of COVID-19, 188,928,123 cases have been detected till fifteenth July 2021, and 4,066,605 people are died (John Hopkins Coronavirus Resource Centre, 2021). In Afghanistan, 137853 COVID-19 cases have been confirmed and 5983 people have been died, since 24 February 2020 (WHO, 2021). Afghanistan government announced that till 29 June 2021, 1,024,168 does vaccines have been administered and 253,939 people fully vaccinated which makes 1.3% of all population (Reuters, 2021). On 26 June 2021, ministry of public health announced that among 19 samples they took 11 of them were Delta variant of COVID-19 virus and remaining were Alpha variant (Tolo News, 2021). According to the data, Delta variant of COVID-19 is 40 to 60 per cent more transmissible than the Alpha variant which was 50 percent more transmissible than the initial variant first detected in Wuhan, China (Scientific American, 2021). By rising COVID-19 positive cases, Afghanistan government announced extension of lockdown in capital city Kabul for three weeks (Garda world, 2020) which included a ban on inter-city travel and a reduction of all non-essential services (occurred on 2 May 2020).

Since start of COVID-19 pandemic many researchers have been tried to predict the pandemic using various statistical and non-statistical models (Li L., 2020; Hamizah F.A.B., 2020; Zareie B., 2020; Wynants L., 2021; Alazab M., 2020; Samuel Sanchez-C., 2020; Santosh K. C., 2020; Sree K. S., 2020; Pinter G.,

2020). Number of scholars have been used SIR (Susceptible–Infected–Removed) model and ARIMA (Autoregressive Integrated Moving Average) model to predict COVID-19 cases (Nesteruk, 2021, Singh R. K., 2020).

In this study, I used a time series model which is ARIMA model. This model is a statistical model which is used to predict time series data. By using this model, I am trying to forecast COVID-19 cases in Afghanistan.

The remaining parts of the study is structured as follow: the second section reviews existing literature on application of varied forecasting models for estimation of the pandemic. The third section the objectives of the present study have been elaborated. The fourth section gives information on description of data, research methodology and selection of best ARIMA model using Akaike's Information Criterion (AIC) and Bayesian Information Criteria (BIC). The sixth section dissentions and results. The seventh section summarises the results and implications.

Literature Review

Abolmaali S. (2021), Used Susceptible-Infected-Recovered (SIR) Model, Linear Regression, Logistic Function and ARIMA Model to forecast COVID-19 cases in India, Russia, United States of America and Brazil. This study compares these four models in term of accuracy magnitude of error and tries to forecast COVID-19 cases.

Banik A. (2020). Analysed the cause of death rate during COVID-19 period in 29 different developed and developing countries. This study examines the effect of various factors which lead to an impact in fatality rate.

Benvenuto D. (2020), used Augmented Dickey-Fuller (ADF) unit root to find weather the data is stationary or not. ARIMA model has been used to predict the prevalence and incidence of COVID-19 on eleventh and twelfth of Feb 2020.

Claris S. (2020), used ARIMA model to forecasting daily COVID-19 cases in south Africa. This model predicted positive cases for 20-days. According to this prediction model, COVID-19 cases will rise during this span of time and reach 1744 cases.

Dahesh T. (2020), considered 41-day past data to make ARIMA model and for showing the stationarity of the data Augmented Dickey-Fuller (ADF) unit root used. In this study, ARIMA model forecasted covid-19 cases for 17 days in five different countries.

Earnest A. (2005), suggested ARIMA model to predict bed occupied in Singapore during SARS outbreak. Researchers found out that ARIMA (1,0,3) was suitable for this study and training MAPE was 5.7%. it is advised that this model could be used for bed-capacity during infection diseases.

Elsheikh H. A. (2020) forecasted COVID-19 cases in Saudi Arabia by using long short-term memory (LSTM) network and compared with nonlinear autoregressive artificial neural networks (NARANN) and ARIMA model. In order to assess the accuracy of the model, the researcher used root mean square error (RMSE), coefficient of determination, mean absolute error (MAE), efficiency coefficient (EC), overall index (OI), coefficient of variation (COV), and coefficient of residual mass (CRM). Furthermore, authors used the LSTM model to predict confirmed and death cases in five more countries. Coefficient of determination of total cases and for total death are 0.976 and 0.944, respectively.

Fanelli D. (2020), forecasted and analysed COVID-19 dynamic in China, Italy and France. The Authors apply the susceptible-infected-recovered-deaths (SIRD) model, and it shows recovery, infection and death rate in all three countries.

Gupta R. (2020), analysed COVID-19 cases in India and other south Asia countries. Author proposed time-series model to predict COVID-19 cases in India. According to this study cases will reach a million in 30 days.

In the work of Kufel T. (2020), 6 ARIMA models for each 32 European countries used to predict dynamic cases of COVID-19. ARIMA (1,2,0) predicted cases for 7 days. Finding the usefulness of ARIMA model in prediction of COVID-19 dynamic cases was the aim of this study.

Katoch R. used the ARIMA model to predict COVID-19 cases in four different states and also all cases of India. In order to check stationarity Dickey–Fuller (ADF) had been used. Test Authors used different ARIMA parameters for instance for India ARIMA (4,2,7) and for Tamil Nadu ARIMA (0,2,1).

Khaliq R. (2020), proposed ARIMA model to forecast COVID-19 cases in Jammu & Kashmir. ARIMA (1, 2, 3), ARIMA (0, 2, 2) and ARIMA (0, 2, 2) are applied to forecast confirmed cases, recovered cases and deceased cases, respectively. The time span used in this study is from 9th March 2020 to 30th September 2020. An ARIMA model predicted COVID-19 cases for one month (October 2020). The study reveals cases will rise during the forecasted period.

Kim Y. (2018) studied the transiting dynamic of MERS-CoV in South Korea. Authors used an agent-based model to predict. This model is used for diseases which have rapid separation.

Malki Z. (2020) has developed a SARIMA model in order to predict COVID-19 end and existing cycle of virus. SARIMA (9,0,8) *(0,0,0,3) had been considered fit parameters. This model proposed that if prevention guidelines and precautions are not followed, virus second rebound will happen.

Moein S. (2021) suggested that the SIR model is inefficient to predict the actual spread COVID-19 in the long run. This study analysed and forecasted COVID-19 cases in Isfahan (a province in Iran).

Namasudra S. (2021) considered Nonlinear Autoregressive Neural Network Time Series (NAR-NNTS) which trained by three different algorithms which are Scaled Conjugate Gradient (SCG), Levenberg Marquardt (LM) and Bayesian and compared with model with Root Mean Square Error (RMSE), Mean

Square Error (MSE) and correlation coefficient, in case of India. After comparison, the researcher found out that a better model is the NAR-NNTS which trained by Levenberg Marquardt algorithm.

Perone G. (2020), fitted ARIMA model to forecast COVID-19 final size and separation of this virus in Italy. Emilia Romagna (0, 2, 1), Italy (4, 2, 2) and Lombardy (1, 2, 1) and two different ARIMA parameters are used by the author. This study indicates that cases will reach zero in the 50 days. The final size of the COVID-19 cases will reach between 254,000 and 272,000 and the death toll will reach between 31,318 to 33,538.

Petropoulos F. (2020) proposed sample time series method. The object of this study is to analyse and forecast the death rate, confirmed cases and recoveries of COVID-19 virus globally in a specific span of time. This study forecasted COVID-19 cases in five different rounds which start from second Feb 2020 to twenty first Mar 2020.

Roy S. (2020) used weighted overlay analysis and ARIMA model to forecast COVID-19 cases in different states of India. In order to examine the models, mean absolute error (MAE) and Root mean square error (RMSE) had been used. Study shows two parts of India are more vulnerable which are south and west.

Saba T. (2020) used six different forecast methods (Random forests, K-nearest neighbors, Support vector regression, ARIMA model, SARIMA model, Decision trees, Holt winter model, Polynomial regression and Gradients boosting regressor) to predict COVID-19 cases under different lockdown strategies. The study examines three different lockdown strategies which are partial, herd and complete, and suggests herd strategy as the best among all these three strategies.

Sujath R. (2020) proposed machine learning models (Linear regression (LR), Multilayer perceptron (MLP), Vector autoregression (VAR)) to forecast COVID-19 cases in India. The data which was used in this study is from Kaggle data. The study suggests MLP is a good method for predicting the cases more than LR and VAR models.

Objectives and rational of the study

The main objective of this study is to predict COVID-19 positive cases. The time period which we have used is from sixteenth July 2021 for a period of 40 days. Currently Afghanistan is witnessing the third wave of the COVID-19 pandemic. Presently, COVID-19 hospitals are full patients and don't have any space for hospitalization, so people hospitalize patients in hospitals' yard. So there is a great need to estimate and forecast the prevalence of this virus. The Afghanistan government tries to break the transmission chain of the virus. In order to do this, the government has imposed lockdown in capital city Kabul and other big cities. In this unprecedented situation, forecasting the future of pandemic is really critical for the government and all non-government organizations who work in health sector.

Data description

For this study, I have used daily confirmed cases of COVID-19 in Afghanistan from twenty first March 2021 to fifteenth July 2021. Data was collected from the Ministry of Public health (MoPH), Afghanistan. In this study R studio and Excel have been used to make the model and predict the confirmed cases.

Methodology

In order to predict COVID-19 confirmed cases, 119-day (twenty first March 2021 to sixteenth July 2021) data were collected to develop a time series model. ARIMA model is a time series model, if we review literature of time series forecasting, we can see number of scholars have used this model to predict the future. ARIMA (p, d, q) model consists of Auto Regressive (AR) model, Integrated (I) and Moving Average (MA) model. In this model, p and q are showing lag in AR and MA components, respectively, and d is the number of non-seasonal differences needed for stationarity. ARIMA model is constructed as shown below:

$$Y_t = \alpha + \beta_1 Y_{t-1} + \beta_2 Y_{t-2} + \dots + \beta_p Y_{t-p} + \varepsilon_t + \phi_1 \varepsilon_{t-1} + \phi_2 \varepsilon_{t-2} + \dots + \phi_q \varepsilon_{t-q}$$

Akaike's Information Criterion (AIC) and Bayesian Information Criteria (BIC) have been used to determine the order of ARIMA models.

$$AIC = -2\log(L) + 2(p+q+k+1)$$

L likelihood of the data

$$k = 1 \text{ if } c \neq 0$$

$$k = 0 \text{ if } c = 0$$

$$BIC = AIC + [\log(T) - 2] (p+q+k+1)$$

To forecast COVID-19 positive cases, we try to find the most suitable ARIMA model.

Results And Discussion

For determining the ARIMA parameters, AIC and BIC have been used. In this part of study, we discuss all steps which lead us to create ARIMA (p, d, q) and the result of the model. In this study we used data from twenty first March 2021 to fifteenth July 2021 (119 days).

In order to create time series model, we need stationary data, so Augmented Dickey-Fuller (ADF) unit-root test has been used to check stationarity of the data. Observing the time series plot of data also show whether the data is stationary or not.

Figure 1 illustrates dramatic increase in Covid-19 cases in Afghanistan. It is also clear from this figure that the data is not stationary. In Augmented Dickey-Fuller unit-root we consider null hypothesis (Time series data is non-stationary) and alternative hypothesis (Time series data is stationary). After calculation, we found out that Augmented Dickey-Fuller unit-root is 0.77, Lag order = 4 and p-value is 0.96. So it confirms the data is not stationary.

Table 1 ARIMA Model parameters for ARIMA (p, d, q)

Coefficients	
Model Parameters	ar1
Estimates	-0.5536
Standard E.	0.0889
sigma^2 estimated as 67386: log likelihood=-802.11	
AIC=1608.22 AICc=1608.33 BIC=1613.71	

Source: The author

Table 1 depicts ARIMA parameters. AR (p) is 0, I (d) is 2 and MA (q) is 1. Auto regression equal to 0 means we need 0 lags in the time period. As we mentioned previously, the data is not stationary so we need to take 2 order differences. Moving Average in time series model is past error, so in our model it is 1.

Figure 2 illustrates the COVID-19 positive cases for two different periods. First period starts from 21 March 2021 to 15 July 2021 which is before forecast period, and second period starts from 16 July 2021 to 24 August 2021 which is the forecasted period (for 40 days). Blue line indicates forecast value and the darker shaded area shows 80% Confidence Level and the lighter shaded area shows 95% confidence level. It is clear from the graph that COVID-19 cases started rising before 30 May 2021 and it continues till the end of the forecast period. On 16 July 2021, total cases were 138,930, and in 40 days it reached 180,970. If we see the outcome of the ARIMA model, there is no hint of reduction in confirmed cases. According to data from 2017 for every thousand patients in Afghanistan there are 0.39 beds (WHO, 2021). These data really show the severity of the pandemic situation in Afghanistan.

Conclusions

Afghanistan is a developing country, from August of 2021, we are in the third wave of COVID-19 pandemic. Like other countries, Afghanistan also was not prepared to confront a pandemic.

In this study, we used the ARIMA (0, 2, 1) model to forecast COVID-19 cases in Afghanistan for 40 days from 16 July 2021 to 24 August 2021. COVID-19 is not a big challenge just for Afghanistan but it is a

global challenge. Result of this study shows COVID-19 confirmed cases rise dramatically. If we consider 95% confidence level, COVID-19 cases soar from 139,437 confirmed cases to 216,159 confirmed cases.

For those scholars who would like to work on this topic, it is recommended to investigate the effect of lockdown and other protection measures on the outbreak prevalence.

Significance Statements

Before this study, there was no such study to predict the future of COVID-19 cases in Afghanistan. So there is great need for such study. This study helps the government and policy makers and non-government organizations to plan ahead in order to combat the pandemic.

Declarations

Declarations

Source

The author

Funding

The authors received no financial support for the research, authorship and/or publication of this article.

References

1. Abolmaali, S. (2021). *A comparative study of SIR Model, Linear Regression, Logistic Function and ARIMA Model for forecasting COVID-19 cases*. medRxiv. 1-9.
<https://doi.org/10.1101/2021.05.24.21257594>.
2. Alazab, M., Awajan, A., Mesleh, A., Abraham, A., Jatana, V., & Alhyarl, S. (2020). *COVID-19 Prediction and Detection Using Deep Learning*. International Journal of Computer Information Systems and Industrial Management, 12. 168-181.
3. Banik, A., Nag, T., Chowdhury, S. R., & Chatterjee, R. (2020). *Why Do COVID-19 Fatality Rates Differ Across Countries? An Explorative Cross-country Study Based on Select Indicators*. Global Business Review, 21(3) 607–625.
4. Benvenuto, D., Giovanetti, M., Vassallo, L., Angeletti, S., & Ciccozzi, M. (2020). *Application of the ARIMA model on the COVID-2019 epidemic dataset*. Data in Brief, <https://doi.org/10.1016/j.dib.2020.105340>.
5. Claris, S., & Delson, C. (2020). *Predicting South Africa's daily COVID-19 cases using ARIMA forecasting model: 6 March to 6 July 2020*. Research Square. <https://doi.org/10.21203/rs.3.rs->

6. Deshesh, T., Mardani-Fard, H. A., & Dehesh, P. (2020). *Forecasting of COVID-19 Confirmed Cases in Different Countries with ARIMA Models*. medRxiv. 1-12.
<https://doi.org/10.1101/2020.03.13.20035345>.
7. Earnest, A., Chen, M. I., Ng, D., & Sin, L. Y. (2005). *Using autoregressive integrated moving average (ARIMA) models to predict and monitor the number of beds occupied during a SARS outbreak in a tertiary hospital in Singapore*. BMC Health Services Research, 5(36), 1–8.
8. Elsheikh, A. H., Saba, A. I., Elaziz, A. M., Lu, S., Shanumgan, S., Muthuramalingm, T., Kumar, R., Mosleh, A. O., Essa, F. A., & Shehabeldeenn, T., A. (2020). *Deep learning-based forecasting model for COVID-19 outbreak in Saudi Arabia*. Process Safety and Environmental Protection, 149 (2021), 223–233.
9. Fanelli, D., & Piazza, F. (2020). *Analysis and forecast of COVID-19 spreading in China, Italy and France*. Chaos, Solitons & Fractals, 134, 109761.
10. GardaWorld. (2020, May 5). *Afghanistan: Authorities extend COVID-19 lockdown until May 24*.
<https://www.garda.com/crisis24/news-alerts/339176/afghanistan-authorities-extend-covid-19-lockdown-until-may-24-update-5>.
11. Gupta, R., & Pal, S. K. (2020). *Trend Analysis and Forecasting of COVID-19 outbreak in India*. medRxiv. 1-19. <https://doi.org/10.1101/2020.03.26.20044511>.
12. Hamizah F. A. B., Lau, C. H., Nazri, H., Ligot, D. V., Lee, G., Tan, C. L., Shaib, M. K. B. M., Zaidon, U. H. B., Abdullah, A. B., Chung, M. H., Ong, C. H., Chew, P. Y., & Salunga, R. E. (2020). *Corona Tracker: World-wide COVID-19 Outbreak Data Analysis and Prediction. [Preprint]*. Bull World Health Organ.
<http://dx.doi.org/10.2471/BLT.20.255695>.
13. Johns Hopkins Coronavirus Resource Centre. (2021, July 12). *Covid-19 dashboard*. Center for systems science and engineering. Coronavirus Resource Centre, Johns Hopkins University.
<https://coronavirus.jhu.edu/>.
14. Katock, R., & Sidhu, A. (2021). *An Application of ARIMA Model to Forecast the Dynamics of COVID-19 Epidemic in India*. Global Business Review, 1–15. <https://doi.org/10.1177/0972150920988653>.
15. Khaliq, R., Iqbal, P., Sheergjrl, A. R., Ilyas, A. M., & Wani, A. Y. (2020). *COVID-19 Forecasting using ARIMA model: A special reference to Jammu and Kashmir*. Wesleyan Journal of Research, 3 (4), 364–374.
16. Kim, Y., Ryu, H., & Lee, S. (2018). *Agent-Based Modeling for Super-Spreading Events: A Case Study of MERS-CoV Transmission Dynamics in the Republic of Korea*. Int. J. Environ. Res. Public Health, 15, 2369. doi:10.3390/ijerph15112369.
17. Kufel, T. (2020). *ARIMA-based forecasting of the dynamics of confirmed Covid-19 cases for selected European countries*. Equilibrium. Quarterly Journal of Economics and Economic Policy, 15(2), 181–204. doi:10.24136/eq.2020.009.
18. Li, L., Yang, Z., Dang, Z., Meng, C., Huang, J., Meng, H., Wang, D., Chen, G., Zhang, J., Peng, H., & Shao, Y. (2020). *Propagation analysis and prediction of the COVID-19*. Infectious Disease Modelling, 5, 282–

19. Malki, Z., Atlam, E., Ewis, A., Dagnew, G., Alzighaibi, A. R., Elmarhomy, G., Elhosseini, M. A., Hassanien, E. A., & Gad I. (2020). *ARIMA models for predicting the end of COVID-19 pandemic and the risk of second rebound*. Neural Computing and Applications, 33:2929–2948.
<https://doi.org/10.1007/s00521-020-05434-0>.
20. Moein, S., Nickaeen, N., Roointan, A., Borhani, N., Heidary, Z., Javanmard, S. H., Ghaisari, J., & Geisari, J. (2021). *Inefficiency of SIR models in forecasting COVID-19 epidemic: a case study of Isfahan*. Scientific Reports, 11:4725. <https://doi.org/10.1038/s41598-021-84055-6>.
21. Namasudra, S., Dhamodharavadhani s., & Rathipriya, R. (2021). *Nonlinear Neural Network Based Forecasting Model for Predicting COVID-19 Cases*. Neural Processing Letters.
<https://doi.org/10.1007/s11063-021-10495-w>.
22. Nesteruk, I., & Benlagha N. (2021). *Predictions of Covid-19 Pandemic Dynamics in Ukraine And Qatar Based on Generalized Sir Model*. Innov Biosyst Bioeng, 5(1), 37–46.
<https://10.20535/ibb.2021.5.1.228605>.
23. Perone, G. (2020). *An ARIMA model to forecast the spread and the final size of COVID-2019 epidemic in Italy*. medRxiv. <https://doi.org/10.1101/2020.04.27.20081539>.
24. Petropoulos, F., & Makridakis, S. (2020). *Forecasting the novel coronavirus COVID-19*. PLOS ONE, 15(3), e0231236.
25. Pinter, G., Felde, I., Mosavi, A., Ghamisi, P., & Gloaguen, R. (2020). *COVID-19 Pandemic Prediction for Hungary; A Hybrid Machine Learning Approach*. Mathematics, 8, 890.
<https://doi:10.3390/math8060890>.
26. Reuters. (2021, July 7). *COVID-19 Tracker*. <https://graphics.reuters.com/world-coronavirus-tracker-and-maps/countries-and-territories/afghanistan/>.
27. Roy, S., Bhunia, G. S., & Shit, P. K. (2021). *Spatial prediction of COVID-19 epidemic using ARIMA techniques in India*. Modeling Earth Systems and Environment. 7:1385–1391.
<https://doi.org/10.1007/s40808-020-00890-y>.
28. Saba, T., Abunadi, I., Shazad, M. N., & Khan, A. R. (2020). *Machine learning techniques to detect and forecast the daily total COVID-19 infected and deaths cases under different lockdown types*. Microsc Res Tech, 1–13. <https://doi.org/10.1002/jemt.23702>.
29. Sanchez-Caballero, S., Selles, M. A., Peydro, M. A., & Perez-Bernabeu, E. (2020). *An Efficient COVID-19 Prediction Model Validated with the Cases of China, Italy and Spain: Total or Partial Lockdowns*. Journal of Clinical Medicine, 9, 1547. <https://doi:10.3390/jcm9051547>.
30. Santosh, K. C. (2020). *COVID-19 Prediction Models and Unexploited Data*. Journal of Medical Systems, 44: 170. <https://doi.org/10.1007/s10916020-01645-z>.
31. Scientific American. (2021, June 29). *How Dangerous Is the Delta Variant, and Will It Cause a COVID Surge in the U.S*. <https://www.scientificamerican.com/article/how-dangerous-is-the-delta-variant-and-will-it-cause-a-covid-surge-in-the-u-s/>

32. Singh R, Rani M, Bhagavathula A, Sah R, Rodriguez-Morales A, Kalita H, Nanda C, Sharma S, Sharma Y, Rabaan A, Rahmani J, Kumar P. (2020). *Prediction of the COVID-19 Pandemic for the Top 15 Affected Countries: Advanced Autoregressive Integrated Moving Average (ARIMA) Model*. JMIR Public Health Surveill, 6(2): e19115. <https://doi.org/10.2196/19115>.
33. Sree, P. K., & Nedunuri, S. S. S. N. U. D. (2020). *A novel cellular automata classifier for COVID-19 trend prediction*. Journal of Health Sciences. 10(1):34-38. <https://doi.org/10.17532/jhsci.2020.907>.
34. Sujath, R., Chatterjee, J., M., & Hassanien, A., E. (2020). *A machine learning forecasting model for COVID-19 pandemic in India*. Stochastic Environmental Research and Risk Assessment, 34:959–972.
35. The New York Times. (2021, March 17). *A Timeline of the Coronavirus Pandemic*. <https://www.nytimes.com/article/coronavirus-timeline.html>.
36. TOLONews. (2021, June 26). *Delta Variant Makes 50% of COVID-19 Infections in Afghanistan*. <https://tolonews.com/health-173095>.
37. World Health Organisation (WHO). (2021, July 16). *Hospital beds (per 1,000 people) – Afghanistan*. <https://data.worldbank.org/indicator/SH.MED.BEDS.ZS?locations=AF>.
38. World Health Organisation (WHO). (2021, July 7). *Afghanistan Situation*. <https://covid19.who.int/region/emro/country/af>.
39. Wynants L. et al. (2020). *Prediction models for diagnosis and prognosis of covid-19: systematic review and critical appraisal*. bmj 369. <https://doi.org/10.1136/bmj.m1328>.
40. Zareie, B., Roshani, A., Mansournia M. A., Rasouli, M. A., & Moradi, G. (2020). *A model for COVID-19 prediction in Iran based on China parameters*. medRxiv. 1-20. <https://doi.org/10.1101/2020.03.19.20038950>.

Appendix A

Table A1. Prediction of COVID-19 confirmed cases in Afghanistan for 40 days by ARIMA model

Date	Forecast	lo 80	Hi 80	Lo 95	Hi 95
16-Jul-21	138930	138599	139262	138424	139437
17-Jul-21	140008	139426	140591	139118	140899
18-Jul-21	141086	140231	141942	139778	142395
19-Jul-21	142164	141010	143318	140399	143929
20-Jul-21	143242	141765	144719	140983	145501
21-Jul-21	144320	142496	146144	141530	147110
22-Jul-21	145398	143205	147591	142044	148752
23-Jul-21	146476	143892	149060	142525	150427
24-Jul-21	147554	144560	150548	142974	152133
25-Jul-21	148632	145207	152056	143395	153869
26-Jul-21	149710	145837	153583	143787	155633
27-Jul-21	150788	146449	155127	144152	157424
28-Jul-21	151866	147043	156688	144490	159241
29-Jul-21	152944	147621	158266	144803	161084
30-Jul-21	154022	148183	159860	145092	162951
31-Jul-21	155099	148729	161470	145356	164843
01-Aug-21	156177	149260	163095	145598	166757
02-Aug-21	157255	149776	164735	145817	168694
03-Aug-21	158333	150278	166389	146014	170653
04-Aug-21	159411	150766	168057	146189	172633
05-Aug-21	160489	151240	169738	146343	174635
06-Aug-21	161567	151701	171434	146477	176657
07-Aug-21	162645	152148	173142	146591	178698
08-Aug-21	163723	152583	174863	146686	180760
09-Aug-21	164801	153005	176597	146761	182841
10-Aug-21	165879	153415	178343	146817	184941
11-Aug-21	166957	153812	180101	146854	187059
12-Aug-21	168035	154198	181871	146873	189196

13-Aug-21	169113	154572	183653	146875	191350
14-Aug-21	170190	154934	185446	146858	193522
15-Aug-21	171268	155286	187251	146825	195712
16-Aug-21	172346	155625	189067	146774	197919
17-Aug-21	173424	155954	190894	146706	200142
18-Aug-21	174502	156272	192732	146622	202382
19-Aug-21	175580	156580	194581	146521	204639
20-Aug-21	176658	156876	196440	146405	206911
21-Aug-21	177736	157163	198309	146272	209200
22-Aug-21	178814	157439	200189	146124	211504
23-Aug-21	179892	157705	202079	145960	213824
24-Aug-21	180970	157961	203979	145780	216159

Source: The author

Figures

Figure 1

Daily COVID-19 cases in Afghanistan from twenty first March to fifteenth July, 2021 Source: The Author

Figure 2

Forecasting plot of COVID-19 confirmed cases in Afghanistan Source: The Author