Prevalence and Risk Factors of Untreated Thyroid Dysfunctions in the Older Caucasian Adults: Results of Polsenior 2 Survey

Piotr Kocelak (✉ pkocelak@sum.edu.pl)

Medical University of Silesia: Śląski Uniwersytet Medyczny w Katowicach <https://orcid.org/0000-0001-8135-8482>

Malgorzata Mossakowska

International Institute of Molecular and Cell Biology in Warsaw: Międzynarodowy Instytut Biologii Molekularnej i Komórkowej w Warszawie

Monika Puzianowska-Kuznicka

Mossakowski Medical Research Centre Polish Academy of Sciences: Instytut Medycyny Doświadczalnej i Klinicznej im M Mossakowskiego Polskiej Akademii Nauk

Krzysztof Sworczak

Medical University of Gdansk: Gdański Uniwersytet Medyczny

Adam Wyszomirski

Medical University of Gdansk: Gdański Uniwersytet Medyczny

Gabriela Handzlik

Medical University of Silesia: Śląski Uniwersytet Medyczny w Katowicach

Adrian Stefanski

Gdański Uniwersytet Medyczny

Tomasz Zdrojewski

Gdański Uniwersytet Medyczny

Jerzy Chudek

Medical University of Silesia: Śląski Uniwersytet Medyczny w Katowicach

Research

Keywords: elderly population, thyroid dysfunction, hypothyroidism, hyperthyroidism, thyroid hormones, gender differences

Posted Date: July 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-735681/v1>

Abstract

Background: To determine the prevalence of treated and untreated thyroid dysfunction and to identify factors associated with increased risk of undiagnosed thyroid dysfunction in older adults.

Methods The population of 5987 community-dwelling Polish Caucasian seniors aged 60 years and above who participated in the PolSenior 2 study (2018 – 2019). Population-based cross-sectional multidisciplinary study in design. Data from structured questionnaires, geriatric tests and scales were obtained from all study participants who underwent anthropometric and blood pressure measurements during three home visits. Assessment of thyroid function was based on TSH serum measurements.

Results

The prevalence of thyroid dysfunction in the Polish population aged 60 years or above was estimated at 15.5% (21.5% in women and 7.2% in men), with 3.2% of undiagnosed individuals among them. The prevalence of hypothyroidism and hyperthyroidism in the studied group was 13.9% (19.4% in women and 6.3% in men) and 1.6% (2.1% in women and 0.9% in men) respectively. In multiple regression analysis independent risk factors for thyroid disorders being untreated were older age (> 75 years), male sex, a low education level (primary or lower) and low utilization of medical services.

Conclusions

One-fifth of Polish Caucasian seniors with hypothyroidism and one-third with hyperthyroidism is untreated. Older, poorly educated and rarely utilizing medical services seniors, especially men, are more frequently untreated for thyroid dysfunction and some of them do not benefit from contemporary achievements of medicine.

Background

Thyroid dysfunction, mostly hypothyroidism, is the leading endocrinopathy in older adults [1], frequently with clinical presentation diverse from that in younger ones [2]. Current clinical guidelines recommend the same cut-off points for thyroid hormones in the diagnosis of thyroid dysfunction, irrespective of age, sex or race and population iodine supply [3, 4], which may explain vast differences in the prevalence of thyroid dysfunction reported in different populations [5]. One of the major points is the regional variability of iodine supply resulting in the differences in populational TSH distribution. Low iodine intake is associated with a shift of TSH towards lower values, and an increase in the prevalence of hyperthyroidism [6], while higher intake, related to population iodine supply, with a shift towards higher TSH values [7]. Nowadays, hypothyroidism associated with severe iodine deficiency is rarely seen in Europe. The policy of iodine fortification with the adoption of iodine-fortified salt, which currently covers about 71% of the world's population [8] may force change in the references ranges of TSH [9].

The prevalence of thyroid dysfunction was analyzed in the general European population in the years 1975–2012. Meta-analysis of 7 population-based and cohort studies [10] covering Western European countries (excluding individuals with already diagnosed thyroid diseases) revealed that the overall prevalence of thyroid dysfunction was estimated at 6.7% (95%CI: 6.5–6.9), including hypothyroidism – 4.9% (95%CI: 4.8–5.1) and hyperthyroidism – 1.7% (1.7–1.9). Of those 80.1% of disturbances were subclinical. No sub-analysis for older adults was done. In summarized observational studies, the incidence rate of thyroid dysfunction was estimated at 420 / 100,000 / year in women and 85 / 100,000 / year in men [10].

Thyroid dysfunction is highly prevalent in older adults. Based on the PolSenior1 study, the frequency of thyroid dysfunction in the Polish population aged 55 yrs. or over was almost 11%, including hypothyroidism in 8%, and hyperthyroidism in 3% of study subjects [11].

Notwithstanding, much higher numbers were recently shown in a cohort of 5,392 Americans aged 65 yrs. or over, participants of the Atherosclerosis Risk in Communities (ARIC) study performed in years 2011–2013 [12]. Thyroid dysfunction was diagnosed in 24.9% of the study population, including 16.9% and 0.1% already treated for hypothyroidism and hyperthyroidism, respectively. Among untreated subjects, subclinical hypothyroidism (6.1%) was about seven times more prevalent than overt hypothyroidism (0.8%), while subclinical hyperthyroidism (0.8%) was about three times than overt hyperthyroidism (0.3%).

The data from Central and Eastern Europe are sparse. A cross-sectional case-control study carried out in a single tertiary referral centre in Bulgaria [13] included in the meta-analysis of the prevalence of hypothyroidism, suggests a higher prevalence of hypothyroidism in the general population in Eastern than in Western Europe. A similar observation was presented in another meta-analysis [14], which covered the prevalence of only undiagnosed hypothyroidism in the general population in Europe. It has included mostly population-based and cohort studies from old European Union countries published from 2008 to 2017 [14]. The prevalence of undiagnosed hypothyroidism in an older population (≥ 65 yrs.) was greater than in the younger cohort, and estimated at 6.6% (95% CI 2.5–12.4), including subclinical 5.1% (95%CI: 3.3–7.3) and overt disease 1.1% (95%CI: 0–3.4). The published analysis of thyroid function in the PolSenior 1 project has omitted this aspect [11] that is important for public health policies and programs, as health consequences related to untreated thyroid disturbances can be serious and, in some cases, life-threatening. The most important are cardiovascular diseases, diabetes and rarely myxedema coma or thyroid storm [15, 16, 17].

The aim of this study was to analyze the frequency of treated and untreated thyroid dysfunction and to identify factors associated with increased risk of undiagnosed thyroid dysfunction in older adults, based on a representative national cohort.

Material And Methods

The PolSenior 2 study enrolled 5987 subjects aged 60 years and above, a representative for the community-dwelling Polish population. The study was carried out in 2018 and 2019 by nurses

deliberately trained for study purpose and specificity. The study procedures included structured questionnaires, geriatric tests and scales, as well as anthropometric and blood pressure measurements performed during three home visits. Participants were asked to show all medicines (including products containing levothyroxine or antithyroid drugs) taken during the last two weeks. Details of applied procedures are described elsewhere [18].

Blood samples were donated by 5825 participants (97.3%). Blood samples were not obtained from 162 participants due to refusal or difficulty in venepuncture. Thyroid-stimulating hormone (TSH) was assessed in 5775 (99.1%) serum samples in a single central laboratory (Brus Laboratory in Gdynia) using Atellica Solution Immunoassay & Clinical Chemistry Analyzers, and reagents from Siemens Healthineers. The reference range for TSH was 0.55–4.78 mIU/L. In the remaining 50 cases, TSH was not evaluated due to insufficient serum sample volume.

The analysed group (N = 5800; 93.5% of the PolSenior2 cohort) included all subjects with measured TSH concentration and 25 participants treated with levothyroxine or antithyroid drugs without measured TSH level.

All the subjects included in the study gave written informed consent for participation in the study. The study was approved by the ethics committee of the Medical University of Gdansk. The study was conducted according to the Declaration of Helsinki requirements.

Data analysis

Thyroid disorders were classified as 1) hypothyroidism if levothyroxine-containing medicine was used or when elevated (> 4.78 mIU/L) TSH level was present in participants not using antithyroid drugs, 2) clinically significant hyperthyroidism if the antithyroid drug was taken or decreased (< 0.1 mIU/L) TSH level, was found in people not using levothyroxine (Fig. 1).

The subgroup of untreated hypothyroidism (not using levothyroxine preparations) was divided into subclinical ($TSH \leq 10$ mIU/L) and overt hypothyroidism ($TSH > 10$ mIU/L). Among subjects treated with levothyroxine preparations and assessed TSH concentration, the percentage of subjects with elevated TSH levels (> 4.78 mIU/L) was calculated.

In the subgroup with clinically significant hyperthyroidism, the percentage of people using and not using antithyroid drugs was calculated, and among those treated with antithyroid drugs, the percentage of subjects with persistent hyperthyroidism despite the treatment ($TSH < 0.1$ mIU/L) was also evaluated.

Treated subjects both with levothyroxine or antithyroid drugs with missing TSH assessments were excluded from the analysis of the treatment efficacy.

The analysis included: sex, place of residence, marital and living status, educational level, years of education, type of work currently or in the past (grouped into 'blue collar' or farmer, 'white collar', and other), years of work, current professional activity, use of the Internet, frequency of visits to the general

practitioner, visits to specialists and hospitalization during last 5-year period, economic status, the Lawton Instrumental Activities of Daily Living (iADL) [19], the Katz Activities of Daily Living (ADL) [20], Mini-Mental State Examination (MMSE), corrected by age and education according to Mungas [21] and Geriatric Depression Scale (GDS).

Statistical analysis

Statistical analysis and data visualization were conducted using the R statistical package (R Core Team, version 3.6.3.). Sampling weights were included in calculations of proportions (prevalence rates) and 95% confidence intervals to account for the complex survey design. The post-stratification procedure was used to match the age-sex sample distribution to the national population. We applied unweighted logistic regression to assess the relationship between untreated thyroid dysfunctions and a set of risk factors. A multivariable model was built based on a step-wise backward selection procedure. Regression coefficients were expressed as odds ratios with a 95% confidence interval. The between-group differences were tested using the unpaired t-test for quantitative variables and the chi-square test in the case of categorical data. The 2-tailed tests were carried out at a significance level of $p < 0.05$.

Results

The prevalence of thyroid dysfunction

Treated and untreated thyroid dysfunction was found in 849 participants (625 women and 224 men). Based on this data the prevalence of thyroid dysfunctions was estimated at 15.5% (95%CI: 14.1–16.9) in the whole Polish population aged 60 yrs. or above, 21.5% (95%CI: 19.2–23.8) in women and 7.2% (95%CI: 6.0–8.4) in men ($p < 0.001$). The frequency of thyroid dysfunction across the analyzed age cohorts was similar (Fig. 2).

Hypothyroidism

Hypothyroidism was found in 732 participants (544 women and 188 men). Based on this data, the prevalence of hypothyroidism was estimated at 13.9% (95%CI: 12.5–15.4) in the whole Polish population aged 60 yrs. or above, 19.4% (95%CI: 17.1–21.7) in women and 6.3% (96%CI: 5.1–7.5) in men ($p < 0.001$). The disorder was untreated in 21.9% cases ($N = 160$), including 100 women (18.4%) and 60 men (31.9%), ($p < 0.001$). The majority ($N = 125$; 78.1%) of subjects with untreated hypothyroidism presented subclinical dysfunction ($TSH \leq 10$ mIU/L), with no difference between women and men ($N = 83$ –85% and $N = 42$ –70%, respectively; $p = 0.055$). The subgroup with TSH concentration > 10 mIU/L consisted of 35 participants, 17 women and 18 men (4.8%, 3.1% and 9.6% of subjects with untreated hypothyroidism, respectively; $p < 0.001$).

Among 572 individuals treated with levothyroxine, elevated TSH levels were found in 59 patients – 9.0% (95%CI: 6.4–11.6), similarly often among women – 8.5% (95%CI: 5.7–11.3) and men – 11.6% (95%CI: 4.6–18.5), ($p = 0.37$).

Hyperthyroidism

Clinically significant hyperthyroidism was found in 120 participants (83 women and 37 men). Its prevalence was estimated at 1.6% (95%CI: 1.3–1.9) in the studied population, 2.1% (95%CI: 1.6–2.6) in women and 0.9% (95%CI: 0.5–1.3) in men ($p = 0.003$). The disorder was untreated in 41 (34.2%) cases with $TSH < 0.1$ mIU/L, including 29 women (34.9%) and 12 men (32.4%), ($p = 0.79$).

Hyperthyroidism was treated with antithyroid drugs in 79 subjects, but 10 of the – 12.7% had decreased TSH levels, similarly often women and men ($p = 0.11$).

Untreated thyroid dysfunction

There were 201 participants (129 women and 72 men) with untreated thyroid dysfunction: 160 with hypothyroidism (21.9% of all with hypothyroidism) and 41 with hyperthyroidism (34.2% of all with hyperthyroidism). Therefore, the prevalence of untreated dysfunctions was estimated at 3.2% (95%CI: 2.5–3.8) in the whole Polish population aged 60 yrs. or above, 4.0% (95%CI: 3.0–5.1) in women and 1.9% (95%CI: 1.3–2.5) in men ($p = 0.001$).

There was a marked increase in the frequency of untreated thyroid dysfunction in the two oldest analyzed cohorts: 50.6% (95%CI: 37.5–63.7) in ≥ 85 yrs. old vs 17.4% (95%CI: 14.2–20.7) in 60–84 yrs. old cohorts, $p < 0.001$ (Fig. 3).

Participants with untreated thyroid disorders were significantly older and more frequently dependent on activities of daily living (Table 1). There was a higher percentage of men among untreated ones.

Table 1
Comparison of subjects with treated and untreated thyroid dysfunctions.

	Treated thyroid dysfunction N = 648	Untreated thyroid dysfunction N = 201	P value
Age [yrs]	73.9 (8.6);	79.1 (10.2);	< 0.001
Mean (SD); Median (IQR)	73 (67, 80)	79 (71, 87)	
Age > 75 yrs [N; %]	260 (40.1)	119 (59.2)	< 0.001
Women [N; %]	496 (76.5)	129 (64.2)	< 0.001
Place of residence [N; %]			
Village	164 (25.3)	68 (33.8)	0.05
Town < 50,000 citizens	183 (28.2)	57 (28.4)	
City 50,000-200,000 citizens	145 (22.4)	42 (20.9)	
City > 200,000 citizens	156 (24.1)	34 (16.9)	
Marital status [N; %]			
Never married	19 (3.0)	4 (2.0)	0.21
Married	339 (53.6)	100 (50.0)	
Widowed	245 (38.7)	91 (45.5)	
Divorced or separated	30 (4.7)	5 (2.5)	
Living status [N; %]			
Alone	175 (28.7)	45 (23.6)	0.10
With spouse	232 (38.1)	67 (35.1)	
Living with others	202 (33.2)	79 (41.4)	
Educational level [N; %]			

SD: standard deviation, IQR: interquartile range.

IADL – and Instrumental Activities of Daily Living (Lawton and Brody, 1969).

ADL – Activities of Daily Living (Katz et al., 1970)

MMSE - Mini-Mental State Examination (Folstein et al., 1975), corrected by age and education/
corrected according to Mungas ((Mungas et al., 1996)

	Treated thyroid dysfunction	Untreated thyroid dysfunction	P value
	N = 648	N = 201	
Primary or lower	163 (25.5)	80 (39.8)	< 0.001
Vocational	137 (21.4)	41 (20.4)	
Secondary	239 (37.4)	59 (29.4)	
Higher	100 (15.6)	21 (10.4)	
Years of education [years]	11.3 (3.8)	9.9 (4.3)	< 0.001
Mean (SD); Median (IQR)	11 (8,13)	10 (7,12)	
Type o work (current or in the past) [N; %]			
‘Blue collar’ or farmer	288 (48.7)	107 (61.1)	< 0.001
‘White collar’	247 (41.8)	51 (29.1)	
Other	56 (9.5)	17 (9.7)	
Years of work	33.8 (11.0)	35.0 (12.1)	0.24
Median (IQR)	35 (28, 40)	35 (29, 42)	
Professionally active [N; %]	41 (6.4)	5 (2.5)	0.033
Using the Internet [N; %]	197 (31.5)	36 (18.9)	< 0.001
Frequency of visits to the general practitioner [N; %]			
More than once a year	601 (96.9)	165 (88.7)	< 0.001
Once a year or less	19 (3.1)	21 (11.3)	
Utilization of visits to specialists during last 5-year period [N; %]			
Yes	523 (82.9)	143 (72.6)	0.001

SD: standard deviation, IQR: interquartile range.

IADL –and Instrumental Activities of Daily Living (Lawton and Brody, 1969).

ADL – Activities of Daily Living (Katz et al., 1970)

MMSE - Mini-Mental State Examination (Folstein et al., 1975), corrected by age and education/
corrected according to Mungas ((Mungas et al., 1996)

	Treated thyroid dysfunction N = 648	Untreated thyroid dysfunction N = 201	P value
No	108 (17.1)	54 (27.4)	
Hospitalization during last 5-year period [N; %]			
Yes	332 (52.6)	93 (47.0)	0.17
No	299 (47.4)	105 (53.0)	
Amount of money [N; %]			
Enough money to meet the needs	589 (93.6)	180 (90.9)	0.42
Enough money for food and clothes	27 (4.3)	12 (6.1)	
Not enough money for food and clothes	13 (2.1)	6 (3.0)	
No enough money for medicines [N; %]	75 (11.8)	21 (10.4)	0.69
iADL [N; %]			
8–18 pts – dependent	82 (12.7)	62 (30.8)	< 0.001
19–23 pts – partially dependent	113 (17.4)	31 (15.4)	
24 pts – independent	453 (69.9)	108 (53.7)	
ADL [N; %]			
0–2 pts – dependent	16 (2.5)	16 (8.0)	< 0.001
3–4 pts – partially dependent	18 (2.8)	13 (6.5)	
5–6 pts – independent	614 (94.8)	171 (85.5)	
MMSE (corrected) [N; %]			
27–30 pts – no dementia	407 (63.2)	112 (56.3)	0.07
24–26 pts – impaired cognition	119 (18.5)	36 (18.1)	
< 24 pts – dementia	118 (18.3)	51 (25.6)	

SD: standard deviation, IQR: interquartile range.

IADL –and Instrumental Activities of Daily Living (Lawton and Brody, 1969).

ADL – Activities of Daily Living (Katz et al., 1970)

MMSE - Mini-Mental State Examination (Folstein et al., 1975), corrected by age and education/
corrected according to Mungas ((Mungas et al., 1996)

	Treated thyroid dysfunction N = 648	Untreated thyroid dysfunction N = 201	P value
GDS [N; %]			
6–15 pts – depression	164 (28.0)	41 (25.6)	0.55
0–5 pts – no depression	421 (72.0)	119 (74.4)	
SD: standard deviation, IQR: interquartile range.			
IADL –and Instrumental Activities of Daily Living (Lawton and Brody, 1969).			
ADL – Activities of Daily Living (Katz et al., 1970)			
MMSE - Mini-Mental State Examination (Folstein et al., 1975), corrected by age and education/ corrected according to Mungas ((Mungas et al., 1996)			

Untreated were more frequently living in villages, had primary or lower education corresponding to lower years of education and a higher percentage of 'blue collars' and farmers. Of note untreated less frequently were utilizing medical services for outpatients, both primary and specialist.

Multiple regression analysis taking into account age, sex, iADL, place of residence, education level, type of work (in the past), professional activity, cognitive function, utilization of visits to the general practitioner and specialists and using the Internet, revealed that older age (> 75 yrs.), male sex, low education level (primary or lower) and low utilization of medical services (once a year or less) were independent risk factors for thyroid disorders being untreated (Table 2).

Table 2

Factors explaining the occurrence of untreated thyroid dysfunctions. There were 805 participants in the final multivariable model.

		Univariate model	Multivariable model
Age	> 75 yrs	2.17 (1.57–2.99) ^{***}	1.75 (1.22–2.50) ^{**}
	60–75 yrs	Reference	Reference
Sex	Men	1.82 (1.30–2.56) ^{***}	2.04 (1.41–2.94) ^{***}
	Women	Reference	Reference
Place of residence	Village	1.51 (1.07–2.12) [*]	-
	City	Reference	-
Educational level	Primary or lower	1.92 (1.39–2.70) ^{***}	1.59 (1.10–2.33) [*]
	other	Reference	Reference
Type o work (current or in the past)	‘Blue collar’ or farmer	1.80 (1.24–2.62) ^{**}	-
	Other	1.47 (0.79–2.74)	-
	‘White collar’	Reference	-
Professionally active	Yes	0.37 (0.15–0.96) [*]	0.37 (0.13–1.10) [^]
	No	Reference	Reference
Using the Internet	Yes	0.49 (0.33–0.73) ^{***}	-
	No	Reference	-
Frequency of visits to the general practitioner	Once a year or less	4.03 (2.11–7.67) ^{***}	4.20 (2.16–8.18) ^{***}
	More than once a year	Reference	Reference
Utilization of visits to specialists during last 5-year period	Yes	0.55 (0.38–0.80) ^{**}	-
	No	Reference	-

Statistical significance: ^ p < 0.1; * p < 0.05; ** p < 0.01; *** p < 0.001

		Univariate model	Multivariable model
iADL	Dependent (8–23 pts)	2.00 (1.45–2.77)***	-
	Independent (24 pts)	Reference	-
ADL	Dependent (0–4 pts)	3.06 (1.81–5.17)***	-
	Independent (5–6 pts)	Reference	-
MMSE (corrected)	Dementia (	1.57 (1.06–2.32)	-
	Impaired cognition	1.10 (0.72–1.69)	-
	No dementia	Reference	-
Statistical significance: ^ p < 0.1; * p < 0.05; ** p < 0.01; *** p < 0.001			

Discussion

Results of this study allowed us to estimate the prevalence of thyroid dysfunction in the Polish senior population at 15.5%. In this age group, thyroid dysfunction is about 3-fold greater among women (21.5%), than in men (7.2%), including hypothyroidism in 13.9% (19.4% in women and 6.3% in men) and hyperthyroidism in 1.6% (2.1% in women and 0.9% in men). As expected, these figures are higher than those for the whole European population [10], most possibly due to a different age range. Previous studies reported overt and subclinical hypothyroidism in up to 5.7% and 12.5%, respectively, in older adults in various geographical areas [22, 23]. Noteworthy, recent North American data showed a 24.9% prevalence of total thyroid dysfunction in people aged 65 and over [12].

In our study, 21.9% of subjects with hypothyroidism (18.4% of women and 31.9% of men with this pathology) and 34.2% with hyperthyroidism (24.2% women and 32.4% men respectively) were untreated, which corresponds to 3.2% of all Polish seniors – 4.0% of women and 1.9% of men aged 60 yrs. or above. Our data show a lower prevalence of untreated thyroid dysfunction than 6.7% [10] (with 4.9% of hypothyroidism and 1.7% of hyperthyroidism) reported for the general European population. This discrepancy may result from increasingly frequent use of medical services by older adults, commonly suffering from chronic illnesses. Notwithstanding, the prevalence of unrecognized thyroid disease in the Australian population aged 49 years or older was 3.6% (including 3.0% of those with hypothyroidism) [24], whereas in The Colorado Thyroid Disease Prevalence Study 9.9% of the population has a functional abnormality of the thyroid gland that has been unrecognized [25]. In the latter study, however, no separate analysis for older adults was performed.

The present study shows the spectrum of factors associated with the probability of occurrence of untreated thyroid dysfunction. The strongest factor explaining the lack of treatment was the low utilization of primary health care. This proves that family doctors can effectively diagnose thyroid dysfunction and initiate treatment or referral to an endocrinologist. It was shown that a lack of insurance by affecting utilization of outpatient clinics health care may also influence the diagnosis and treatment of thyroid disorders [26].

A second strongest predictor for untreated thyroid dysfunction was male sex. Typical symptoms of thyroid dysfunction are characteristic for young patients and their occurrence predict more accurate thyroid dysfunction in men, but clinical symptoms are less obvious and illusory in older aged patients [27]. In older subjects, the clinical manifestation of hypothyroidism might be similar to signs and symptoms associated with natural ageing. The symptoms may be limited to weight gain, fatigue, muscle cramps, constipation and skin changes [2]. The multimorbidity and adverse drug reactions can also mask or mimic the signs and symptoms of hypothyroidism. A higher percentage of undiagnosed hypothyroidism in men may suggest poor awareness of thyroid disease in men, strongly affected by up to eight to nine times lower prevalence of hypothyroidism than in women [28].

Difficulties also occur in the diagnosis of hyperthyroidism, as in the older population the most common symptom of thyroid hormones excess are tachycardia or atrial fibrillation or exacerbation of cardiovascular disease [29]. Other symptoms of this pathology are depression, apathy, anorexia, weight loss and fatigue. In our study, the frequency of untreated hyperthyroidism may influence the general health of the patients since untreated thyroid hormones excess may lead to the development of cardiac complications and the loss of bone mass [30] and is, therefore, more dangerous than untreated hypothyroidism, was 34.2%. Moreover, the awareness of atypical symptoms of thyroid dysfunction and their clinical presentation in older patients seem to be insufficient by family doctors. Moreover, men utilize less often visits to their primary care physician. The awareness may be effectively increased by educational programs for patients [31].

The next factor associated with undiagnosed thyroid dysfunction is the age of 75 yrs. and above that is also strongly associated with dependence in activities of daily living which, in turn, can limit access to medical services. Moreover, multimorbidity and atypical clinical manifestation of thyroid disease are also associated with age [2]. Of note, dependence in ADL and dementia were less important than utilization of medical services.

Finally, low educational status increased the risk of undiagnosed thyroid dysfunction and improper treatment [26]. In our study group, poor education was more important than economic status. However, this may be different in other societies in which low economic status corresponds to poor insurance status and, therefore, limits the accessibility to health care systems and laboratory tests.

The results also showed that 9.0% of individuals treated for hypothyroidism and 11.6% treated for hyperthyroidism are not properly managed. The previous studies also showed that almost one-third of the patients with diagnosed hypothyroidism did not have proper hormonal substitution [32]. The clinical

consequences of undiagnosed, untreated or undertreated hypothyroidism seem to be especially important in patients with co-morbidities like diabetes mellitus [33] and cardiovascular disease [5]. In addition, hypothyroidism is associated with decreased quality of life [34] and increased mortality [15]. Untreated hyperthyroidism may lead to atrial fibrillation, the development of cardiovascular disease (including cardiomyopathy and heart failure), bone loss and thyroid storm [35]. Notwithstanding, clinical consequences of subclinical hypothyroidism are controversial, especially in the oldest. Even though the recent data negate the benefits from levothyroxine supplementation [36].

Conclusions

Physicians should be aware of possible thyroid dysfunction in their older patients of both sexes. The unanswered question remains on the purposefulness of screening thyroid function in the whole senior population. The results of our study are important in the view of a high frequency of thyroid dysfunction in the elderly population. Moreover, the percentage of untreated thyroid dysfunction is quite high suggesting that a huge number of patients still are left without treatment which may have a detrimental effect on their health.

Abbreviations

TSH – thyroid-stimulating hormone

ARIC study - the Atherosclerosis Risk in Communities study

iADL - the Lawton Instrumental Activities of Daily Living

ADL - the Katz Activities of Daily Living

MMSE - Mini-Mental State Examination

GDS - Geriatric Depression Scale

Declarations

Ethics approval and consent to participate

The study was approved by the ethics committee of the Medical University of Gdansk. The study was conducted according to the Declaration of Helsinki requirements.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

None of the authors declares a conflict of interest and competing interest.

Funding

This paper was implemented under contract No. 6/5 / 4.2 / NPZ / 2017/1203/1257 for the implementation of the task in the field of public health of the Operational Objective No. 5 points 4.2. of the National Health Program for years 2016-2020, entitled "Health Status and Its Socio-economic Covariates of the Older Population in Poland - the Nationwide PolSenior2 Survey" (PolSenior2)

Authors' contributions

Conceptualization: P.K., J.C.; Methodology: P.K., J.C., T.Z.; software: A.W. A.S.; validation: A.S., J.C.; formal analysis: P.K. J.C.; investigation: K.S., A.S., T.Z.; resources: T.Z.; data curation: A.W., A.S., J.C.; writing—original draft preparation: P.K, M.M., J.C.; writing—review and editing: M.P-K. G.H.; visualization: A.W., J.C.; supervision: K.S., J.C.; project administration: A.S., T.Z.; funding acquisition: T.Z.

All authors have read and agreed to the published version of the manuscript.

Acknowledgements

Not applicable

References

1. Wilson S, Parle J, Roberts LM, Roalfe A, Hobbs R, Clark P, et al. Prevalence of subclinical thyroid dysfunction and its relation to socioeconomic deprivation in the elderly: A community-based cross-sectional survey. *J Clin Endocrinol Metab.* 2006;91(12):4809-4816.
2. Boelaert K, Torlinska B, Holder RL, Franklyn JA. Older subjects with hyperthyroidism present with a paucity of symptoms and signs: A large cross-sectional study. *J Clin Endocrinol Metab.* 2010;95(6):2715-2726.
3. Garber JR, Cobin RH, Gharib H, Hennessey J, Klein I, Mechanick J, et al. Clinical practice guidelines for hypothyroidism in adults: Cosponsored by the american association of clinical endocrinologists and the American thyroid association. *Endocr Pract.* 2012;18(6):988-1028.
4. Bollerslev J, Rejnmark L, Marcocci C, Shoback D, Sitges-Serra A, van Biesen W, et al. European Society of Endocrinology clinical guideline: Treatment of chronic hypoparathyroidism in adults. *Eur J Endocrinol.* 2015;173(2): G1-G20.
5. Chaker L, Bianco AC, Jonklaas J, Peeters RP. Hypothyroidism. *Lancet.* 2017;390(10101):1550-1562.

6. Laurberg P, Nøhr SB, Pedersen KM, Hreidarsson A, Andersen S, Pedersen S, et al. Thyroid disorders in mild iodine deficiency. *Thyroid*. 2000;10(11):951-963.
7. Ittermann T, Khattak RM, Nauck M, Cordova CMM, Völzke H. Shift of the TSH reference range with improved iodine supply in Northeast Germany. *Eur J Endocrinol*. 2015;172(3):261-267.
8. Nations U, Unicef F. *The State of the World's Children, 2011. Adolescence. An Age of Opportunity*. Vol 17.; 2011.
9. Cai J, Fang Y, Jing D, Xu S, Ming J, Gao B, et al. Reference intervals of thyroid hormones in a previously iodine-deficient but presently more than an adequate area of Western China: A population-based survey. *Endocr J*. 2016;63(4):381-388.
10. Madariaga AG, Santos Palacios S, Guillén-Grima F, Galofré JC. The incidence and prevalence of thyroid dysfunction in Europe: A meta-analysis. *J Clin Endocrinol Metab*. 2014;99(3):923-931.
11. Bar-Andziak E, Milewicz A, Jędrzejuk D, Arkowska A, Mieszczanowicz U, Krzyanowska-Świniarska B. Thyroid dysfunction and thyroid autoimmunity in a large unselected population of elderly subjects in Poland - The "PolSenior" multicentre crossover study. *Endokrynol Pol*. 2012;63(5):346-355.
12. Diab N, Daya NR, Juraschek SP, Martin S, McEvoy J, Schultheiss U, et al. Prevalence and Risk Factors of Thyroid Dysfunction in Older Adults in the Community. *Sci Rep*. 2019;9(1):1-8.
13. Elenkova A, Atanasova I, Kirilov G, Natchev E, Ivanova R, Kovatcheva R, et al. Autoimmune hypothyroidism is three times more frequent in female prolactinoma patients compared to healthy women: data from a cross-sectional case-control study. *Endocrine*. 2017;57(3):486-493.
14. Mendes D, Alves C, Silverio N, Marques FB. Prevalence of Undiagnosed Hypothyroidism in Europe: A Systematic Review and Meta-Analysis. *Eur Thyroid J*. 2019;8(3):130-143.
15. Rodondi N, Den Elzen WPJ, Bauer DC, Cappola A, Rayvi S, Walsh J, et al. Subclinical hypothyroidism and the risk of coronary heart disease and mortality. *JAMA - J Am Med Assoc*. 2010;304(12):1365-1374.
16. Gencer B, Collet TH, Virgini V, Bauer D, Gussekloo J, Cappola A, et al. Subclinical thyroid dysfunction and the risk of heart failure events an individual participant data analysis from 6 prospective cohorts. *Circulation*. 2012;126(9):1040-1049.
17. Gronich N, Deftereos SN, Lavi I, Persidis AS, Abernethy DR, Rennert G. Hypothyroidism Is a Risk Factor for New-Onset Diabetes: A Cohort Study. *Diabetes Care*. 2015;38(9):1657-1664.
18. Wierucki Ł, Kujawska-Danecka H, Mossakowska M, Grodycki T, Bledowski P, Chudek J, et al. Health status and its socio-economic covariates in the older population in Poland – the assumptions and methods of the nationwide, cross-sectional PolSenior2 survey. *Arch Med Sci*. Published online 2020; DOI: <https://doi.org/10.5114/aoms.2020.100898>
19. Lawton M, Brody E. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969;9:179–86. *Gerontologist*. 1969;9:1979-1986.
20. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of Illness in the Aged The Index of ADL: A Standardized Measure of Biological and Psychosocial Function Table 1.—Index of Independence in Activities of Daily Living. *JAMA*. 1963;185(12):94-99.

21. Mungas D, Marshall SC, Weldon M, Haan M, Reed BR. Age and education correction of mini-mental state examination for English- and Spanish-speaking elderly. *Neurology*. 1996;46(3):700-706.
22. Sawin CT, Castelli WP, Hershman JM, McNamara P, Bacharach P. The Aging Thyroid: Thyroid Deficiency in the Framingham Study. *Arch Intern Med*. 1985;145(8):1386-1388.
23. Razvi S, Ingoe L, Ryan V, Pearce SHS, Wilkes S. Study of Optimal Replacement of Thyroxine in the Elderly (SORTED) - results from the feasibility randomised controlled trial. *Thyroid Res*. 2016;9(1):1-9.
24. Empson M, Flood V, Ma G, Eastman CJ, Mitchell P. Prevalence of thyroid disease in an older Australian population. *Intern Med J*. 2007;37(7):448-455.
25. Canaris GJ, Manowitz NR, Maoy G, Ridgway C. The Colorado Thyroid Disease Prevalence Study. 2000;160:526-534.
26. Olmos RD, De Figueiredo RC, Aquino EM, Lotufo PA, Bensenor IM. Gender, race and socioeconomic influence on diagnosis and treatment of thyroid disorders in the Brazilian longitudinal study of adult health (ELSA-Brasil). *Brazilian J Med Biol Res*. 2015;48(8):751-758.
27. Carlé A, Bülow Pedersen I, Knudsen N, Perrild H, Ovesen L, Laurberg P. Gender differences in symptoms of hypothyroidism: A population-based DanThyr study. *Clin Endocrinol (Oxf)*. 2015;83(5):717-725.
28. Aoki Y, Belin RM, Clickner R, Jeffries R, Phillips L, Mahaffey KR. Serum TSH and total T4 in the United States population and their association with participant characteristics: National Health and Nutrition Examination Survey (NHANES 1999-2002). *Thyroid*. 2007;17(12):1211-1223.
29. Klein I, Ojamaa K. Thyroid hormone and the cardiovascular system. *N Engl J Med*. 2001;344(7):501-509.
30. Faber J, Jensen IW, Petersen L, Nygaard B, Hegedüs L, Siersbæk-Nielsen K. Normalization of serum thyrotrophin by means of radioiodine treatment in subclinical hyperthyroidism: effect on bone loss in postmenopausal women. *Clin Endocrinol (Oxf)*. 1998;48(3):285-290.
31. Canaris GJ, Tape TG, Wigton RS. Thyroid disease awareness is associated with high rates of identifying subjects with previously undiagnosed thyroid dysfunction. *BMC Public Health*. 2013;13(1):351.
32. Dew R, Okosieme O, Dayan C, Eligar V, Khan I, Razvi S, et al. Clinical, behavioural and pharmacogenomic factors influencing the response to levothyroxine therapy in patients with primary hypothyroidism – protocol for a systematic review. Published online 2017:1-7.
33. Raval AD, Sambamoorthi U. Incremental healthcare expenditures associated with thyroid disorders among individuals with diabetes. *J Thyroid Res*. 2012;2012.
34. Vigário PDS, Vaisman F, Coeli CM, Ward L, Graf H, Carvalho G, et al. Inadequate levothyroxine replacement for primary hypothyroidism is associated with poor health-related quality of life-a a Brazilian multicentre study. *Endocrine*. 2013;44(2):434-440.
35. Yamamoto H, Monno S, Ohta-Ogo K, Ishibashi-Ueda H, Hashimoto T. Delayed diagnosis of dilated thyrotoxic cardiomyopathy with coexistent multifocal atrial tachycardia: a case report. *BMC Cardiovasc Disord*. 2021;21(1):1-10.

36. Stott D, Rodondi N, Kearney P, Ford I, Westendorp R, Rudi G. Thyroid Hormone Therapy for Older Adults with Subclinical Hypothyroidism. N Engl J Med. 2017;376:2534-2544.

Figures

Figure 1

Chart flow of the study

Figure 2

The frequency of thyroid dysfunction across age cohorts

Figure 3

The frequency of untreated thyroid dysfunction in age cohorts