

Investigating the Effectiveness of Self-management Program based on 5A Model on Fatigue and Dyspnea in Patients with Heart Failure

Maryam Hajmohamadi

Kerman University of Medical Sciences

Sakineh Sabzvari

Kerman University of Medical Sciences

Yones Jahani

Kerman University of Medical Sciences

Zahra Imani- Goghary (✉ imanigoghary@yahoo.ca)

Sirjan school of medical sciences

Research Article

Keywords: Dyspnea, Fatigue ,Heart failure, 5A self- management model, self-management

Posted Date: September 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-738117/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Some symptoms such as fatigue and dyspnea decrease the quality of life in patients with heart failure. The effectiveness of self-management programs on management chronic conditions was discussed. So, this study aimed to investigate the effectiveness of self-management program based on 5A model on fatigue and dyspnea in patients with heart failure.

Materials and Methods: In this clinical trial study, 60 patients with heart failure were included. The intervention group underwent self-management program based on 5A model and the control group received routine care. All the enrolled patients were evaluated once at baseline and once after 3 months using fatigue severity and Borg dyspnea scales. The obtained data were analyzed using SPSS software version 16 by descriptive statistics and independent t-test, covariance test, and paired sample t-test. The significance level was set at 0.05.

Results: There was no significant difference in the mean scores of fatigue and dyspnea at the beginning of the study between control and intervention groups ($p>0.05$), but 3 months after intervention a significant difference was found in mean scores of fatigue and dyspnea between two groups, ($p<0.05$). The difference between pre- and post-intervention scores in terms of the dyspnea and fatigue variables was significant based on the result of paired sample t-test ($p<0.05$).

Conclusion: according to the results of this study, self-management program based on 5A Model can be used to reduce the severity of fatigue and dyspnea as well as improve the quality of life in patients with heart failure.

Trial registration number: IRCT20141109019862N8 **Trial registration date:** 11/Aug/2019

First participant enrollment: 30 /Sep/ 2019:

Background

Heart failure (HF) is one of the most common cardiovascular disorders, which is chronic, progressive, and debilitating (1). Therefore, in this disease, following the weakness of the heart muscle due to myocardial infarction, cardiomyopathy or valvular heart diseases, the heart is unable to meet all the body's metabolic needs (2). USA statistics showed an increasing trend in the number of patients with HF, as about 5 million people are currently living with the disease and more than 150000 new cases are diagnosed each year (3). Statistics in Iran also showed a high rate of the incidence of HF, as the number of patients were estimated to be about 3500 per 100000 (4).

The chronic, progressive and debilitating nature of this disease causes chronic symptoms such as: dyspnea, fatigue, confusion, weakness, pulmonary and limbs edema, chest pain and palpitation (5). Dyspnea as one of the most common symptoms in these patients, could reduce their quality of life (6).

Accordingly, dyspnea is a mental sensation of hard breathing (7), which occurs during activities (exertional dyspnea) in patients under primary heart conditions and as the disease is progressing, dyspnea can be seen with milder activities and may eventually occur even at the time of rest (orthopnea) (8). Another common symptom in patients with HF is fatigue, which is described as a mental sensation of exhaustion and lack of energy. In addition, it is a multidimensional sense influenced by various physical, psychological, and social factors (9).

Physicians prescribe some medications to control these symptoms, but interventions such as self-management programs besides pharmacological treatments are mostly used to control the symptoms and improve the quality of life of patients with HF (10). Self-management programs provide strategies for treatment of chronic health conditions, in which the patients play a pivotal role in promoting their health, preventing the disease, and successful control of the disease (11). In a self-management program, care and treatment activities are performed by focusing on patients and with the aim of achieving the maximum independence, self-determination, health promotion based on abilities and lifestyle, and increasing the quality of life (12). Meanwhile, 5A self-management model, as a habit change counseling is an evidence-based approach, appropriate for habit modification and health care, which includes the following five stages: Assess, Advise, Agree, Assist, and Arrange, that runs in three months (13). This model was firstly developed by Glasgow and it has been used by health care providers on a number of occasions, including habits change and smoking cessation counseling (14).

Mulder et al (2015) in their study used 5A self-management model for counseling and habits changing in patients with type two diabetes and their results showed that this model can improve physical activity and eating habits of the patients (15). Heidari et al (2015) in their clinical study evaluated the effectiveness of 5A model on 50 patients with COPD, and the results showed the reduction in dyspnea and fatigue (16). In another study, some positive effects of using 5A model on providing some advice for obesity patients in primary care level were reported (17). A positive effect of this model was also reported on behaviors like smoking (18).

According to the emphasis of the above-mentioned studies on the positive effects of this model on self-management, symptom control and changing unhealthy behaviors under chronic conditions, considering the high prevalence of HF and due to the limited number of studies performed on controlling symptoms of chronic disease like HF, further studies on the use of 5A self-management program are necessary.

Objectives of the study

The primary goal of the present study was to evaluate the effectiveness of self-management program based on 5A model on common disabling symptoms in patients with HF. As the fatigue and dyspnea are two common symptoms in these patients, so the Secondary objectives of this study were: investigating the effectiveness of self-management program based on 5A model, on fatigue in patients with HF as well as investigating the effectiveness of this model, on dyspnea in mentioned patients.

We anticipate that the self-management program based on 5A model will reduce the chronic symptoms of patients with HF and consequently increase their quality of life. Furthermore, we expect that this intervention will be cost-effective with reducing hospital readmission rates.

Method

Study design

The study was a randomized, single-blinded clinical trial with control and intervention groups. The aim of this trial was to investigate the effectiveness of self-management program based on 5A model on fatigue and dyspnea in patients with HF.

After obtaining the study approval (97000789) from the research council of Kerman University of medical science in 4 Feb 2019, obtaining ethics code IR.KMU.REC.1397.479 of ethics committee of Kerman University of medical science, registering clinical trial in 11/Aug/2019, with Trial registration number of: IRCT20141109019862N8 as well as receiving a written informed consent from all participants, the researcher registered the patients admitted in CCU of sirjan's (a city in Iran) hospitals in the study.

Sample size determination

According to a similar article (16) and using the sample size formula (considering the probable drop out of the participants), 60 HF patients who met the inclusion criteria were finally included in the study using convenience sampling.

$$n = \frac{(z_{1-\frac{\alpha}{2}} + z_{1-B})^2 (d_1^2 + d_2^2)}{(\mu_1 - \mu_2)^2} = \frac{(1/96 + 1/28)^2 (7/16^2 + 6/57^2)}{(34/4 - 41)^2} = 23$$

Study population and eligibility

The study population consisted of the patients with HF who were admitted in the cardiac units of Sirjan's hospitals. The inclusion criteria were as follows: age above 18 years old, being conscious, no acute respiratory infection or other respiratory diseases such as chronic obstructive pulmonary disease (COPD), and desire to participate in this study. The exclusion criteria were as follows: exacerbation of patient's condition and not attending the training sessions.

Randomization

After explaining the purpose of the research to the participants, if they wanted to participate, were added to the list of participants. After registration of all participants, they were randomly assigned to the intervention and control groups through the simple random sampling method, (30 patients in each group). Sampling was done from 30 September 2019 to 1 January 2020.

The patients in the control group received routine care, but the intervention group received a self-management program based on 5A model in addition to their routine care. (See flow diagram, Fig. 1)

Routine care

patients in the control group received standard care only. Standard care consists of medication, monitoring and general education about diet, rest and activity and next visits, without any follow-up program.

Intervention

After stabilizing the acute condition of patient, the intervention was performed at the 5 following stages during 3 months from 10 January 2020 until the 11th of April 2020. The first stage (assess): assessing patients' knowledge, life-styles, and health beliefs, as well as reviewing patients' medical history, assessing the symptoms related to their heart condition such as; dyspnea, fatigue, drugs side effects and required pharmacological knowledge, sleep status, nutrition, activity tolerance, and beliefs related to the disease and treatment. The second stage (advice): providing some advices about health risk factors and benefits of changing unhealthy life-styles, at this stage, according to the assessment results, health risks were identified for each patient and nursing diagnosis were written and reported to the patient. The third stage (agree): a written agreement was reached between the patient and the researcher on the patient's performance (agreed with the patient upon realistic goal setting). Thus, according to each nursing diagnosis, appropriate and agreed behavioral objectives were identified and an action plan was designed for each objective. The first three steps were performed in a session which lasted an hour and a half for each participant. The fourth stage (assist): assistance in developing a practical plan, three 2-hour training sessions were held in small groups consisting of five participants (considering common problems of the individuals) during 6 weeks to inform the patients about their disease. Training lectures, face to face consulting and group discussions according to the nursing diagnoses and objectives were held, also some educational brochures along with motivational programs such as walking exercise were presented. Some of the educational contents were as follows: training about diet, how to take drugs and manage their side effects and choose proper activity and exercises due to the patients' needs and how to manage it according to heart disease. The fifth stage (arrange a follow up plan): at this stage, telephone calls were made in the first two weeks daily and then weekly, up to three months for following the patient's performance.

Data collection

The following instruments were used for data collecting: 1- a questionnaire on personal information, including age, sex, marital status, educational level, income, employment status, insurance status, and place of residence; in addition information related to the participant medical history such as: reasons for referring to the hospital, suffering from other chronic disease, duration of the HF, history of taking a particular medication, family history, history of smoking and regular opium use gained through interviewing patients.

2- Fatigue severity scale (FSS): this is a valid self-report scale designed by Krupp et al. (1989) to measure fatigue in patients with multiple sclerosis and Lupus (19). In Iran, Azimiyani et al. (2010) determined reliability of the instrument through test-retest ($r = 0/93$) and Cronbach's alpha as 0.96 (20). This scale

includes 9 items, each one of which is scored in a Likert spectrum of zero to 7. Accordingly, zero score means strongly disagree and score 7 means strongly agree. The total score was obtained from the sum of scores, which would be between 0 and 63, and if a patient achieved a score equal or higher than 36, it would show that person had fatigue, so the higher the score, the more severe the fatigue.

3- Borg dyspnea scale: this is a standard numerical scale designed by Pfeiffer et al. in 2002 to examine patients' dyspnea at the times of resting and doing activities (21). Correspondingly, its reliability has been calculated to be 0.78 based on Cronbach's alpha (21). Scores would be between zero (no dyspnea) and ten (maximum dyspnea), during rest and physical activity separately.

At first personal information questionnaire was completed through interview and physical examination, the degree of HF checked out by reviewing the patients' medical documents, intensity of dyspnea and fatigue of all of participants were assessed using fatigue severity scale (FSS) and Borg's dyspnea scale, when the patients were not at the acute condition of disease. Then after performing the self- management program, the patients in both control and intervention groups also were assessed using FSS and Borg dyspnea scale.

Statistical analyses

Data were analyzed using SPSS software version 16. Demographic variables were analyzed by descriptive statistics (mean \pm Sd, frequency and percent), The compression of medical history of patients in control and intervention groups were conducted using Chi-square.

Paired t-test, independent t-test, and covariance were used to assess the intervention effect on the mean score of fatigue and dyspnea severity. The significant level was set to 0.05.

Results

In this study, 60 patients (30 in the intervention group and 30 in the control group) with mean age of 64.63 \pm 2.19 years old in the control group and 65.26 \pm 1.93 years old in the intervention group participated. The two groups had equal demographic features, but they were significantly different in opium use, so that 63 % of control group and 30% of intervention group had opium use ($p < 0.05$). Participants' medical history in both groups were also compared, which are presented in Table 1.

Table 1
 comparing medical history of patients in control and intervention groups

variable		control		intervention		test statistics	p.value
		frequency	percent	frequency	percent		
Reasons for referring to the hospital	dyspnea and chest pain	6	20	5	16.66	0.29	0.89
	dyspnea, chest pain, high blood pressure, and fatigue	12	40	11	36.67		
	dyspnea, chest pain, high blood pressure, fatigue and edema	12	40	14	46.67		
History of other chronic disease	musculoskeletal disorders, diabetes, hypertension and visual disturbances	20	66.67	15	50	1.71	0.19
	Musculoskeletal disorders, hypertension, visual disturbances and kidney disease	10	33.33	15	50		
History of taking a particular medication for...	cardiac, diabetes, hyperlipidemia and neurological medication	5	16.67	10	33.33	2.22	0.14
	cardiac, diabetes and hyperlipidemia medication	25	83.33	20	66.67		
Family history	Hypertension	12	40	13	43.34	0.11	0.95
	diabetes	8	26.67	7	23.33		
	Coronary artery disease	10	33.33	10	33.33		
History of smoking	yes	12	40	7	23.33	1.92	0.16
	No	18	60	23	76.67		
Regular	yes	19	63.33	9	30	6.69	0.01*

opium use	No	11	26.67	21	70		
Duration of the heart failure	less than three years	4	13.33	11	36.67	4.42	0.11
	3–5 years	12	40	8	26.66		
	more than 5 years	14	46.67	11	36.67		
Heart failure stage	grade 2	22	73.33	27	90	1.92	0.09
	grade 3	8	26.66	3	10		

In this study, all the participants were at the second or third stages of HF and suffered from dyspnea and fatigue. All of them experienced exertional dyspnea and dyspnea which occur during milder activities and even at rest.

In order to determine and compare the mean score of fatigue and dyspnea severity (during rest and activity) between control and intervention groups before the intervention, independent t-test was used (Table 2).

Table 2
determining and comparing mean scores of resting and exertional dyspnea and fatigue severity between two groups before intervention

Variable	groups	Mean ± SD	Test statistics	P value
Fatigue severity	Control	49.43 ± 0.89	-0.87	0.39
	Intervention	50.63 ± 1.04		
Resting Dyspnea	Control	1.66 ± 0.26	2.01	0.49
	Intervention	1.35 ± 0.15		
Exertional dyspnea	Control	4.66 ± 0.34	-1.62	0.11
	Intervention	5.46 ± 0.35		

As the results show, there weren't any significant difference between the mean scores of variables between control and intervention groups before intervention ($p > 0.05$).

However, after performing the intervention, covariance test was used with controlling the effect of opium and pre-test on results, to determine and compare the mean score of resting and exertional dyspnea and fatigue severity between control and intervention groups, which results are presented in Table3.

Table 3
 comparison of mean scores of resting and exertional dyspnea and fatigue severity between two groups after intervention by controlling confounders

Variable	DF	Mean of squares	Test statistics	P.value
Addiction (opium)	1	0.07	0.35	0.55
Resting dyspnea (before intervention)	1	50.29	251.44	0.001
Group(intervention/control)	1	7.48	37.40	0.001*
error	56	0.2		
Addiction (opium)	1	0.85	1.43	0.24
Exertional dyspnea (before intervention)	1	81.48	137.05	0.001
Group (control/intervention)	1	94.61	159.13	0.001*
error	56	0.59		
Addiction(opium)	1	31.44	2.96	0.091
Fatigue before intervention	1	266.41	25.11	0.001
Group(control/intervention)	1	6570.12	619.36	0.001*
Error	56	10.6		

According to the results there were significant differences between the mean scores of variables between control and intervention groups after intervention ($p \leq 0.001$)

Additionally, paired t-test was used to evaluate changes in scores of resting and exertional dyspnea and fatigue severity, before and after the intervention, in both groups separately, the results are presented in Table 4.

Table 4

determining and comparing mean scores of resting and exertional dyspnea and fatigue severity before and after intervention in each group separately

Group	Variable	Time	Mean \pm SD	Test statistics	P value
Control group	Fatigue severity	Before intervention	49.43 \pm 0.89	- 0.89	0.39
		After intervention	49.60 \pm 0.81		
	Resting dyspnea	Before intervention	1.66 \pm 0.26	1	0.33
		After intervention	1.63 \pm 0.26		
	Exertional dyspnea	Before intervention	4.66 \pm 0.34	-0/63	0.54
		After intervention	4.73 \pm 0.33		
intervention group	Fatigue severity	Before intervention	50.63 \pm 1.04	19.14	0.001*
		After intervention	28.26 \pm 0.59		
	Resting dyspnea	Before intervention	1.05 \pm 0.15	5.20	0.001*
		After intervention	0.40 \pm 0.05		
	Exertional dyspnea	Before intervention	5.46 \pm 0.35	12.22	0.001*
		After intervention	2.41 \pm 0.14		

As the results show there weren't any significant difference between mean scores of main variables (resting and exertional dyspnea and fatigue severity) before and after intervention in control group ($p > 0.05$), but these differences were significant in intervention group before and after intervention ($p \leq 0.001$).

Discussion

In this study, we examined the impact of using 5A self-management model on fatigue and dyspnea in patients with chronic HF. According to the results, 5a self-management model was found to have a positive effect on reducing fatigue and dyspnea in patients with HF. The results of Heidari et al.'s study (2014) have also shown that applying 5A self-management model has a positive effect on fatigue severity and dyspnea in patients with chronic obstructive pulmonary disease (16). Accordingly, fatigue and dyspnea are the two most common complaints in patients under chronic conditions such as HF, that effect their quality of life (22). The results of the present study showed that this model is not only effective on reducing exertional dyspnea, but it can also be effective on decreasing resting dyspnea. Numerous

studies have examined the effects of various non-pharmacological treatments on decreasing dyspnea. In this regard, Beniaminovitz et al. (2002) found that strengthening legs muscles reduce fatigue and dyspnea in HF disease (23). The impact of muscles training was also reported on increasing functional status of chronic HF patients (24). Therefore, the positive effect of non-pharmacological treatments can be acknowledged, so based on the results of this study and those of some other studies, it can be said that by applying self-management techniques, patients can overcome disease complications (16). Wongpiriyayothar et al. (2011) mentioned the positive effect of training patients and follow up by phone call on reducing dyspnea and improving physical function of patients with HF (25). As mentioned previously, the fifth step of 5A model is following up patients, that it's effectiveness was mentioned in Wongpiriyayother's study too.

Moreover, this study also found that using 5A self-management model has a positive effect on fatigue of chronic HF patients. Fatigue is a common symptom under chronic condition such as HF, and it is mostly related to the reduced blood and oxygen uptakes by the tissues (8, 26), which consequently affects patients' independence and quality of life. Various studies based on self-management models have been done to control fatigue symptoms in patients. Roozitalab et al. in their study (2020) stated a significant positive effect of 5A self-management model on fatigue in cancer patients undergoing chemotherapy immediately and 3 month after the end of the intervention (27). Accordingly, this result was in line with the findings of the present study, confirming the impact of 5A self-management model on fatigue. Although few studies have examined the effects of this care model, but similar self-management models showed their positive effects in this context. Wang et al. in 2015 stated positive effects of supportive nursing care on fatigue and quality of life of HF patients (28). In addition, some other studies stated positive effects of applying respiration muscles training on fatigue and quality of life of patients with HF (29–31). Furthermore, Masoudi et al. (2018) in a study pointed to a significant inverse relationship between fatigue and general self-care abilities and also between fatigue and self-care capacity (32). In most of the above-mentioned studies, the effect of a single intervention on controlling symptoms has been investigated, whereas in the comprehensive 5A self-management model, after assessment the patient a specific plan is designed for each patient with involving the patient and his family, then some trainings and consultation are provided according to patients' needs, their personal characteristics, beliefs, and abilities, afterward, this participatory program would be followed by a nurse, which will be more successful (33).

Conclusion

In chronic diseases like HF, some existing complications such as fatigue and dyspnea affect patients' performance. Self-management programs enable patients to rely on their abilities, gain sufficient knowledge about their disease and control the related symptoms and problems. Therefore, self-management and self-care models, especially 5A model, play effective roles in better controlling these symptoms, reducing the related complications, enhancing nursing care and improving patients' quality of life. Another purpose of using self-management model is active participation of patients for managing their disease, rather than complete dependency on care providers.

Abbreviations

HF: heart failure

Declarations

Ethics approval and consent to participate:

This study was approved by the ethics committee of of Kerman University of medical science (IR.KMU.REC.1397.479). The aim of the study, confidentiality of the information and the right to withdraw were explained to the participants and a written consent was obtained. The trial was performed in accordance with the declaration of Helsinki, Questionnaires were anonymous.

Consent for publication:

Not applicable

Availability of data and materials:

The datasets used and/or analyzed during the current study are available from the corresponding authors on reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

Self-funded

Authors' contributions:

All authors (M H, S S, Y J and Z I) conceptualized the study and all were major contributors in writing the manuscript. M H as a MSC students did the intervention with guidance of S S and Z I. Y J as bio-statistician supervised the data analysis. All authors approved the final manuscript.

Acknowledgements:

This study is a part of a thesis entitled “examining the effect of self-management program based on the 5A model on fatigue severity and dyspnea in HF patients” approved by Kerman medical university. We would like to thank all the participants cooperating in the implementation of this study as well as their families who helped the research team.

References

1. Ponikowski P, Anker SD, AlHabib KF, Cowie MR, Force TL, Hu S, et al. Heart failure: preventing disease and death worldwide. *ESC heart failure*. 2014;1(1):4–25.
2. Vedin O, Hagström E, Gallup D, Neely M, Stewart R, Koenig W, et al. Tooth loss is highly prevalent and associated with cardiovascular risk factors in patients with chronic coronary heart disease in the global stability trial. *Journal of the American College of Cardiology*. 2013;61(10S):E1368-E.
3. Karnik AA, Gopal DM, Ko D, Benjamin EJ, Helm RH. Epidemiology of atrial fibrillation and heart failure: a growing and important problem. *Cardiology clinics*. 2019;37(2):119–29.
4. Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM, et al. Heart disease and stroke statistics—2011 update: a report from the American Heart Association. *Circulation*. 2011;123(4):e18-e209.
5. Aghamohammadi T, Maddah SSB, Mohammadi SF, Dalvandi A, Khaleghpour M. The impact of self-management program on self-efficacy of elderly patients with heart failure. *J Urmia Nurs Midwifery Fac*. 2017;14(12):1013–23.
6. Retrum JH, Boggs J, Hersh A, Wright L, Main DS, Magid DJ, et al. Patient-identified factors related to heart failure readmissions. *Circulation: Cardiovascular Quality and Outcomes*. 2013;6(2):171–7.
7. Nava S, Larovere MT, Fanfulla F, Navalesi P, Delmastro M, Mortara A. Orthopnea and inspiratory effort in chronic heart failure patients. *Respiratory medicine*. 2003;97(6):647–53.
8. Brunner LS, Smeltzer S, Suddarth D. *Brunner & Suddarth's textbook of medical-surgical nursing; Vol. 1. Language*. 2010;27:1114–2240p.
9. Rejeh N, HEARAVI KM, Bahrami T, Raeesi M, Tadrissi D. The assessment of factors affecting fatigue in older people with hemodialysis. 2015.
10. Moattari M, Ghobadi A, Beigi P, Pishdad G. Impact of self management on metabolic control indicators of diabetes patients. *Journal of Diabetes & Metabolic Disorders*. 2012;11(1):1–6.
11. Fang X, Wang X, Bai C. COPD in China: the burden and importance of proper management. *Chest*. 2011;139(4):920–9.
12. Ghaderpanah N, Mohaddesi H, Vahabzadeh D, Khalkhali H. The effect of 5A model on behavior change of physical activity in overweight pregnant women. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. 2017;20(9):101–14.
13. Moradi M, Nasiri M, Hajjahmadi M, Jahanshah M. The effect of self-management program based on 5A model on coping strategies in the elderly men patients with hypertension. 2018.

14. Eliaszadeh P, Yarmohammadi H, Nawaz H, Boukhalil J, Katz DL. Congestive heart failure case management: a fiscal analysis. *Disease Management*. 2001;4(1):25–32.
15. Mulder BC, van Belzen M, Lokhorst AM, van Woerkum CM. Quality assessment of practice nurse communication with type 2 diabetes patients. *Patient education and counseling*. 2015;98(2):156–61.
16. Heidari M, Fayazi S, Borsi H, Moradbeigi K, Akbari Nassaji N. Effect of a self-management program based on 5A model on dyspnea and fatigue severity among patients with chronic obstructive pulmonary disease: a randomized clinical trial. *Hayat Journal*. 2015;20(4):89–99.
17. Vallis M, Piccinini–Vallis H, Sharma AM, Freedhoff Y. Modified 5 As: Minimal intervention for obesity counseling in primary care. *Canadian Family Physician*. 2013;59(1):27–31.
18. Flocke SA, Antognoli E. Assessing the accuracy of patient report of the 5As (ask, assess, advise, assist, and arrange) for smoking cessation counseling. *Family Medicine and Community Health*. 2017;5(3):164–9.
19. Krupp LB, LaRocca NG, Muir-Nash J, Steinberg AD. The fatigue severity scale: application to patients with multiple sclerosis and systemic lupus erythematosus. *Archives of neurology*. 1989;46(10):1121–3.
20. A'zimiyan M, Fallah-Pour M, Karimlou M. Fatigue severity scale (fss): evaluation of reliability of the Persian version among persons with multiple sclerosis. *Archives of Rehabilitation*. 2010;10(4):0-.
21. Pfeiffer KA, Pivarnik JM, Womack CJ, Reeves MJ, Malina RM. Reliability and validity of the Borg and OMNI rating of perceived exertion scales in adolescent girls. *Medicine and science in sports and exercise*. 2002;34(12):2057–61.
22. Backé E-M, Seidler A, Latza U, Rosnagel K, Schumann B. The role of psychosocial stress at work for the development of cardiovascular diseases: a systematic review. *International archives of occupational and environmental health*. 2012;85(1):67–79.
23. Beniaminovitz A, Lang CC, LaManca J, Mancini DM. Selective low-level leg muscle training alleviates dyspnea in patients with heart failure. *Journal of the American College of Cardiology*. 2002;40(9):1602–8.
24. Laoutaris I, Dritsas A, Brown MD, Manginas A, Alivizatos PA, Cokkinos DV. Inspiratory muscle training using an incremental endurance test alleviates dyspnea and improves functional status in patients with chronic heart failure. *European Journal of Cardiovascular Prevention & Rehabilitation*. 2004;11(6):489–96.
25. Wongpiriyayothar A, Piamjariyakul U, Williams PD. Effects of patient teaching, educational materials, and coaching using telephone on dyspnea and physical functioning among persons with heart failure. *Applied Nursing Research*. 2011;24(4):e59-e66.
26. Najafi Ghezelejh T, Salehzadeh H, Rafii F. Comparing the effects of back massage and music on fatigue in patients with chronic heart failure. *Nursing And Midwifery Journal*. 2016;14(6):516–25.
27. Roozitalab M, Hosseini N, Vossoughi M, Zoladl M. Effect of Intervention Based on 5A Self-Management Model on the Fatigue Severity of Patients with Cancer under Chemotherapy; A Randomized Clinical Trial. *Journal of Clinical Care and Skills*. 2020;1(2):63–9.

28. Wang T-C, Huang J-L, Ho W-C, Chiou A-F. Effects of a supportive educational nursing care programme on fatigue and quality of life in patients with heart failure: a randomised controlled trial. *European Journal of Cardiovascular Nursing*. 2016;15(2):157–67.
29. Bosnak-Guclu M, Arikan H, Savci S, Inal-Ince D, Tulumen E, Aytemir K, et al. Effects of inspiratory muscle training in patients with heart failure. *Respiratory medicine*. 2011;105(11):1671–81.
30. Chen D-M, Yu W-C, Hung H-F, Tsai J-C, Wu H-Y, Chiou A-F. The effects of Baduanjin exercise on fatigue and quality of life in patients with heart failure: a randomized controlled trial. *European Journal of Cardiovascular Nursing*. 2018;17(5):456–66.
31. Pozehl B, Duncan K, Hertzog M. The effects of exercise training on fatigue and dyspnea in heart failure. *European journal of cardiovascular nursing*. 2008;7(2):127–32.
32. Masoudi R, Reiszadeh I, Abolhassani S, Heidari Beni F. Investigating the Relationship between Self-Care Ability Based on Orem's Self-Care Model and Fatigue in COPD Patients. *Journal of Research in Medical and Dental Science*. 2018;6(3).
33. Javanvash Z, Mojdekanloo M, Rastaghi S, Rad M. The effect model-based self-management program 5A on quality of life of elderly patients with acute coronary syndrome Bojnourd Year 1395. *Journal of Sabzevar University of Medical Sciences*. 2018;25(1):75–82.

Figures




Figure 1

The flow diagram of the study