

# enChIP systems with MSCV-based retroviral plasmids expressing 3xFLAG-Sp-dCas9

Miyuki Yuno

Osaka Daigaku

Toshitsugu Fujita

Hirosaki Daigaku Igakubu Daigakuin Igaku Kenkyuka

Hodaka Fujii (✉ [hodaka@hirosaki-u.ac.jp](mailto:hodaka@hirosaki-u.ac.jp))

Hirosaki University <https://orcid.org/0000-0003-1296-4256>

---

## Research note

**Keywords:** enChIP, dCas9, ChIP, chromatin immunoprecipitation, CRISPR, MSCV

**Posted Date:** July 23rd, 2020

**DOI:** <https://doi.org/10.21203/rs.2.16753/v2>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Biology Methods and Protocols on July 9th, 2021. See the published version at <https://doi.org/10.1093/biomethods/bpab013>.

# Abstract

**Objective:** Engineered DNA-binding molecule–mediated chromatin immunoprecipitation (enChIP) is a technology for purifying specific genomic regions to facilitate identification of their associated molecules, including proteins, RNAs, and other genomic regions. In enChIP, the target genomic region is tagged with engineered DNA-binding molecules, e.g., a variant of the clustered regularly interspaced short palindromic repeats (CRISPR) system consisting of a guide RNA (gRNA) and a catalytically inactive form of Cas9 (dCas9). In this study, to increase the flexibility of enChIP and expand the range of target cells, we generated murine stem cell virus (MSCV)-based retroviral plasmids for expressing dCas9.

**Results:** We constructed MSCV-based retroviral plasmids expressing *Streptococcus pyogenes* dCas9 fused to a 3xFLAG-tag (3xFLAG-Sp-dCas9) and various drug resistance genes. We showed that it is feasible to purify target genomic regions with high yields using these plasmids. These systems might give enChIP users greater flexibility in choosing optimal systems for drug selection of transduced cells. In addition, they could be used to analyze different types of target cells.

## Method Details

To understand the molecular mechanisms underlying regulation of genome functions such as epigenetic regulation and transcription, it is necessary to identify the regulatory molecules binding to a genomic region of interest. We recently developed engineered DNA-binding molecule–mediated chromatin immunoprecipitation (enChIP) technology for isolation of genomic regions of interest to facilitate identification of their associated molecules [1, 2]. The engineered DNA-binding molecules that can be used to tag the target locus include transcription activator-like (TAL) proteins [3] and the clustered regularly interspaced short palindromic repeats (CRISPR) system [4, 5] consisting of a guide RNA (gRNA) and a catalytically inactive form of Cas9 (dCas9) (see our recent review [6] for the CRISPR-based systems). Locus-tagging can be achieved by expression of engineered DNA-binding molecules (in-cell enChIP) [1, 2, 7-10]. Alternatively, it can be achieved *in vitro* using recombinant or synthetic engineered DNA-binding molecules (*in vitro* enChIP) [11, 12]. After a target locus is tagged using an engineered DNA-binding molecule, it is isolated by affinity purification. Subsequently, associated proteins are identified by mass spectrometry (MS) [1, 2, 7], and associated nucleic acids such as RNAs and other genomic regions are identified by next-generation sequencing (NGS) [8, 10, 12].

In this study we generated murine stem cell virus (MSCV)-based retroviral plasmids expressing 3xFLAG-Sp-dCas9 and various drug resistance genes. Using these plasmids, we were able to purify target genomic regions with high yields. These systems might give enChIP users greater flexibility in choosing optimal systems for drug selection of transduced cells. In addition, they could be used to analyze different types of target cells.

# Materials

## Plasmids

To construct MSCV expression plasmids expressing 3xFLAG-Sp-dCas9, the pMSCVneo, pMSCVhyg, and pMSCVpuro vectors (631461, Takara Bio, Kusatsu, Japan) were digested with *Bgl*II (1021A, Takara Bio) and *Xho*I (1094A, Takara Bio). After a blunting reaction, the plasmids were treated with bacterial alkaline phosphatase (*E. coli* C75) (2120A, Takara Bio). The cleaved vectors were purified by agarose gel electrophoresis and ligated with the coding sequence of 3xFLAG-Sp-dCas9, which was isolated from 3xFLAG-dCas9/pMXs-puro (Addgene (Watertown, MA, USA) #51240) [13] by digestion with *Pac*I (R0547, New England Biolabs, Ipswich, MA, USA) and *Not*I (1166A, Takara Bio).

To construct gRNA-hIRF-1 #12/pSIR-hCD2 (Addgene #135392), pSIR-hCD2 (Addgene #51143) [13] was digested with *Eco*RI (1040A, Takara Bio) and treated with bacterial alkaline phosphatase. The cleaved vector was purified by agarose gel electrophoresis and ligated with the gBlock targeting the *interferon (IFN) regulatory factor (IRF)-1* gene promoter, which was isolated from gRNA-hIRF-1 #12 (Addgene #61079) [1] by digestion with *Eco*RI.

## Cell lines

The 293T cell line was derived by transformation of human embryonic kidney (HEK) 293 cells with the SV40 large T antigen [14]. HT1080 is a human fibrosarcoma cell line purchased from ATCC (Manassas, VA, USA) (CCL-121) [15]. 293T, HT1080, and HT1080-derived cells were cultured in DMEM (FUJIFILM Wako Pure Chemical, Osaka, Japan) supplemented with 10% fetal calf serum (FCS).

# Procedures

## Transduction of retroviral plasmids

To increase the flexibility of enChIP and expand the range of target cells, we constructed retroviral plasmids expressing *S. pyogenes* dCas9 fused with the 3xFLAG-tag (3xFLAG-Sp-dCas9), using the MSCV system along with various drug selection markers.

To confirm the performance of the system, we first transduced the retroviral plasmids expressing 3xFLAG-Sp-dCas9 into HT1080, a human fibrosarcoma cell line. 5.5  $\mu$ g of each plasmid was transfected into  $1 \times 10^6$  of 293T cells along with 5.5  $\mu$ g of pPAM3 [16] using Lipofectamine 3000 (Thermo Fisher Scientific, Waltham, MA, USA). Two days after transfection, viral supernatant was harvested and used for infection

of HT1080 cells. Infected cells were selected in culture media containing G418 (Nacalai Tesque, Kyoto, Japan) (0.8 mg/ml), hygromycin (Nacalai Tesque) (75 µg/ml), or puromycin (Nacalai Tesque) (0.75 µg/ml). After drug selection, expression of 3xFLAG-Sp-dCas9 was confirmed by immunoblot analysis with antibody against the FLAG-tag (Fig. 1a). For immunoblot analysis, cytoplasmic extracts were prepared using the NE-PER Nuclear and Cytoplasmic Extraction Reagents (Thermo Fisher Scientific). Ten micrograms of cytoplasmic extract was subjected to immunoblot analysis with anti-FLAG M2 antibody (Ab) (F1804, Sigma-Aldrich, Saint Louis, MO, USA) as described previously [1].

Subsequently, gRNA-hIRF-1 #12/pSIR-hCD2, a self-inactivating retroviral vector expressing a sgRNA targeting human *IRF-1* gene promoter and the human CD2 selection marker protein, was transduced into the HT1080 cells expressing 3xFLAG-Sp-dCas9. 5.5 µg of gRNA-hIRF-1 #12/pSIR-hCD2 was transfected into  $1 \times 10^6$  of 293T cells along with 5.5 µg of pPAM3 using Lipofectamine 3000. Two days after transfection, viral supernatant was harvested and used for infection of HT1080 cells expressing 3xFLAG-dCas9. Two days after transduction, cells expressing the sgRNA were selected by MACS sorting. MACS sorting of human CD2 (hCD2) (+) cells was performed using CD2 Microbeads (130-091-114, Miltenyi Biotec B.V. & Co. KG, Bergisch Gladbach, Germany). To monitor the results of MACS sorting, cells were stained with phycoerythrin (PE)-conjugated anti-hCD2 Ab (347597, BD Life Sciences, San Jose, CA, USA) for 30 min at 4°C. Flow cytometry was performed on a FACSCalibur (BD Life Sciences), and data were analyzed using the FlowJo software (BD Life Sciences).

## enChIP analysis

After expansion,  $2 \times 10^6$  cells were subjected to enChIP analysis. Briefly, the cells were crosslinked with formaldehyde, and the crosslinked chromatin was fragmented by sonication. Subsequently, fragmented chromatin tagged with the CRISPR complex was purified using anti-FLAG antibody. enChIP-real-time PCR to calculate the yields of the target genomic region was performed as previously described [13]. The primers used in this study were reported previously [1]. As shown in Fig. 2, the yields of enChIP (7 - 9% of input) were high and comparable to those obtained by other systems (e.g., ca. 10% of input for the pMXs system [13]). These results revealed that the enChIP system using the MSCV retroviral vectors can be used for downstream applications such as identification of molecules associated with the target genomic regions.

## Declarations

**Acknowledgements:** We thank Dr. George M. Church for providing a plasmid (Addgene #43802). This work was supported by the Takeda Science Foundation (T.F.), a Grant-in-Aid for Scientific Research (C) (#15K06895, #18K06176) (T.F.), and a Grant-in-Aid for Scientific Research (B) (#15H04329) (T.F., H.F.),

'Transcription Cycle' (#15H01354) (H.F.), from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

### Declaration of interests:

☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

T.F. and H.F. have patents on enChIP (Patent name: Method for isolating specific genomic region using molecule binding specifically to endogenous DNA sequence; Patent number: Japan 5,954,808; Patent application number: WO2014/125668). T.F. and H.F. are founders of Epigeneron, Inc. H.F. is a member of Advisory Board of Addgene.


**Supplementary material *and/or* Additional information:** The preprint of this manuscript has been placed on the preprint server *Research Square* (DOI: 10.21203/rs.2.16753/v2).

## References

1. Fujita T, Fujii H: **Efficient isolation of specific genomic regions and identification of associated proteins by engineered DNA-binding molecule-mediated chromatin immunoprecipitation (enChIP) using CRISPR.** *Biochem Biophys Res Commun* 2013, **439**:132-136.
2. Fujita T, Asano Y, Ohtsuka J, Takada Y, Saito K, Ohki R, Fujii H: **Identification of telomere-associated molecules by engineered DNA-binding molecule-mediated chromatin immunoprecipitation (enChIP).** *Sci Rep* 2013, **3**:3171.
3. Bogdanove AJ, Voytas DF: **TAL effectors: customizable proteins for DNA targeting.** *Science* 2011, **333**(6051):1843-1846.
4. Harrison MM, Jenkins BV, O'Connor-Giles KM, Wildonger J: **A CRISPR view of development.** *Genes Dev* 2014, **28**:1859-1872.
5. Wright AV, Nuñez JK, Doudna JA: **Biology and applications of CRISPR systems: harnessing nature's toolbox for genome engineering.** *Cell* 2016, **164**:29-44.
6. Fujita T, Fujii H: **Purification of specific DNA species using the CRISPR system.** *Biology Methods & Protocols* 2019, **4**(1):bpz008.
7. Fujita T, Fujii H: **Identification of proteins interacting with genomic regions of interest in vivo using engineered DNA-binding molecule-mediated chromatin immunoprecipitation (enChIP).** *Bio-Protoc* 2014, **4**:e1124.
8. Fujita T, Yuno M, Okuzaki D, Ohki R, Fujii H: **Identification of non-coding RNAs associated with telomeres using a combination of enChIP and RNA sequencing.** *PLoS One* 2015, **10**:e0123387.


9. Fujita T, Yuno M, Fujii H: **Allele-specific locus binding and genome editing by CRISPR at the p16INK4a locus.** *Sci Rep* 2016, **6**:30485.
10. Fujita T, Yuno M, Suzuki Y, Sugano S, Fujii H: **Identification of physical interactions between genomic regions by enChIP-Seq.** *Genes Cells* 2017, **22**:506-520.
11. Fujita T, Yuno M, Fujii H: **Efficient sequence-specific isolation of DNA fragments and chromatin by in vitro enChIP technology using recombinant CRISPR ribonucleoproteins.** *Genes Cells* 2016, **21**(4):370-377.
12. Fujita T, Kitaura F, Yuno M, Suzuki Y, Sugano S, Fujii H: **Locus-specific ChIP combined with NGS analysis reveals genomic regulatory regions that physically interact with the Pax5 promoter in a chicken B cell line.** *DNA Res* 2017, **24**:537-548.
13. Fujita T, Fujii H: **Identification of proteins associated with an IFN $\gamma$ -responsive promoter by a retroviral expression system for enChIP using CRISPR.** *PLoS One* 2014, **9**(7):e103084.
14. DuBridg e RB, Tang P, HC H, Leong PM, Miller JH, Calos MP: **Analysis of mutation in human cells by using an Epstein-Barr virus shuttle system.** *Mol Cell Biol* 1987, **7**:379-387.
15. Rasheed S, Nelson-Rees WA, Toth EM, Arnstein P, Gardner MB: **Characterization of a newly derived human sarcoma cell line (HT-1080).** *Cancer* 1974, **33**:1027-1033.
16. Miller AD, Buttimore C: **Redesign of retrovirus packaging cell lines to avoid recombination leading to helper virus production.** *Mol Cell Biol* 1986, **6**:2895-2902.

## Figures



**Figure 1**

enChIP systems using MSCV-based retroviral expression vectors. (a) Schematics of MSCV-based retroviral expression constructs of 3xFLAG-Sp-dCas9. (b) Expression of 3xFLAG-Sp-dCas9. Immunoblot analysis was performed with anti-FLAG Ab. Coomassie Brilliant Blue (CBB) staining is shown as a protein loading control.



**Figure 2**

Isolation of the IRF-1 locus by enChIP systems using MSCV-based retroviral expression vectors. Real-time PCR analysis was performed on chromatin complexes isolated by enChIP. An irrelevant locus (SOX2) was analyzed as a negative control.