

Gait, cognition and falls over 5 years, and Motoric Cognitive Risk in New Zealand octogenarians: Te Puawaitanga o Nga Tapuwae Kia Ora Tonu, LiLACS

Sue Lord (✉ sue.lord@aut.ac.nz)

<https://orcid.org/0000-0002-6670-6191>

Simon Moyes

University of Auckland

Ruth Teh

University of Auckland

Waiora Port

University of Auckland

Marama Muru-Lanning

University of Auckland

Catherine Bacon

University of Auckland

Tim Wilkinson

University of Otago Christchurch

Ngaire Kerse

University of Auckland

Research article

Keywords: Gait, cognition, falls, Motoric-Cognitive Risk, oldest old

Posted Date: November 5th, 2019

DOI: <https://doi.org/10.21203/rs.2.16800/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 5th, 2020. See the published version at <https://doi.org/10.1186/s12877-020-1420-8>.

Abstract

Introduction Understanding falls risk in advanced age is critical with people over 80 a rapidly growing demographic. Slow gait and cognitive complaint are established risk factors and together comprise the Motoric Cognitive Risk Syndrome (MCR). **Objectives** To examine trajectories of gait and cognition, and associations with falls over 5 years, and to document MCR in Māori and non-Māori of advanced age living in New Zealand.

Method Falls frequency was ascertained retrospectively at annual assessments. Gait speed (3m), cognition (3MS) and MCR were reported. Gait and cognition trajectories were modelled and clusters identified from Latent Class Analysis. GEE examined association between changes in gait, cognition, MCR and falls.

Results At baseline, 138 of 408 Māori (34%) and 205 of 512 non-Māori (40%) had fallen. Mean (SD) gait speed (m/s) for Māori was 0.66 (0.29) and 0.82 (0.26) for non-Māori. Respective 3MS scores were 86.2 (15.6) and 91.6 (10.4). Ten (4.3%) Maori participants met MCR criteria, compared with 7 (1.9%) non-Māori participants. Māori men were more likely to fall (OR 1.56; 95% CI 1.0 – 2.43 (P = 0.04) whilst for non-Māori slow gait increased falls risk (OR 0.40; 95% CI 0.24 – 0.68(P <0.001). Non-Māori with MCR were more than twice as likely to fall than those without MCR (OR 2.45; 95% CI 1.06 – 5.68 (P = 0.03).

Conclusions Māori and non-Māori of advanced age show a mostly stable pattern of gait and cognition over time. Risk factors for falls differ for Māori, and do not include gait and cognition.

Introduction

Falls in older people are burdensome and their impact far-reaching. The effect of even a sporadic fall can be significant, especially for adults of advanced age. Falls are the third most common cause of accidental injury-related death globally, and the dominant cause of injury-related disability-adjusted life years (DALY) in older people[1]. Compounding this issue is the rapid demographic change with the number of adults aged 80 years and over projected to triple by 2050, rising globally from 137 million in 2017 to 425 million [2]. This age group is most vulnerable to the effect of falls. In New Zealand alone people aged 85 years and over experience 8 times more hospital attendances than those aged 65–74 years, with 8.5% admitted as a result of a fall [3].

Falls are heterogenous and complex. They occur ultimately because of a failure to respond rapidly and effectively to perturbation during walking or when transitioning from one position to another Discerning the cause of failure is difficult because it is multifactorial, and numerous risk factors for falls have been identified. Gait and balance (postural control) disorders are consistently implicated as the strongest risk factors [4 5] and restoration of both through exercise are integrated into falls prevention programmes[5]. Cognition also plays a key role, both as an independent risk factor and through its modifying effect on gait and postural control [6]. Cognitive retraining is therefore often integrated into falls prevention programmes, as is use of cholinesterase inhibitors to improve cognitive function [7]. Understanding the

impact of gait and cognition on falls is important to improve early recognition of the potential for falls and inform effective prevention, which is challenging to implement and subject to failure [5].

The interaction between gait and cognition is also characterised as the Motoric Cognitive Risk (MCR) syndrome which is a pre-dementia state comprising slow gait and subjective cognitive complaint [8]. MCR syndrome is prognostic of both dementia [9] and falls. Callisaya and colleagues recently reported an increased risk of falling in adults classified as having MCR syndrome in over 6,000 older adults from 5 longitudinal cohorts, likely due to the multiplicative effect of gait and cognitive complaints [10].

The contribution of gait and cognition to falls in adults over 80 years has not been studied extensively and is the focus of this research, embedded in a 5-year longitudinal study which commenced in New Zealand in 2010: Te Puāwaitanga o Ngā Tapuwae Kia Ora Tonu/Life and Living in Advanced Age: A Cohort Study in New Zealand (LiLACS NZ). The overarching study aim was to determine predictors of successful ageing and to understand trajectories of health and wellbeing for Māori and non-Māori of advanced age, in line with the global drive to study ageing [11]. We extended this work by examining the trajectory of gait speed and cognition over 6 years and the relationship of both to falls prevalence. We hypothesised that: 1) gait and cognition would decline over 5 years; 2) gait and cognition and their decline would be independently associated with falls risk; and 3) MCR prevalence rates would be comparable to international reports.

Methods

Recruitment and Sampling

Study methodology has been described in full previously [11 12]. Briefly, this study was part of a population based longitudinal study involving Māori (indigenous New Zealanders) and non-Māori octogenarians. Potential participants were identified through the New Zealand electoral roll, Primary Health Care data bases, Hauora (Māori Health Services), and community outreach. Exhaustive sampling was attempted where an attempt to contact and enroll every eligible person was made. LiLACS NZ recruited 937 participants from the Bay of Plenty and Rotorua regions of New Zealand in 2010—421 Māori born between 1920–1930 (aged 80–90 years, 56 % of those eligible) and 516 non-Māori born in 1925 (aged 85 years, 59 % of those eligible) [13]. All participants gave informed consent and The Northern X Regional Ethics Committee of New Zealand granted ethical approval for the longitudinal study in December 2009 (NTX/09/09/088).

Measures

Participants were assessed at baseline and annually to 5 years follow up by trained interviewers using standardised techniques. A comprehensive assessment battery was administered via a home-based interview, followed by a physical and general health assessment [12]. *Falls frequency* was measured retrospectively via self-report. Participants were asked: How many times have you fallen in the past 12

months? A fall was defined as 'unintentionally coming to rest on the ground or other lower surface without being exposed to overwhelming external force or a major internal event'[14] *Gait speed* was measured over 3 metres as part of the Short Performance Physical Battery. Participants were asked to walk at a comfortable pace for two trials, and the fastest speed was recorded [15]. *Cognition* was measured using the Modified Mini Mental Status Examination (3MS) which includes four extra items and is more nuanced in terms of memory items compared with the original MMSE [16 17]. Depression was measured using the 15 item Geriatric Depression Scale (GDS) [18]. Participants met the criteria for *Motoric-Cognitive Risk (MCR) Syndrome* if they presented with a gait speed slower than 1 SD below the cohort mean as well as subjective cognitive impairment, indicated by a positive response to a question from the Geriatric Depression Scale 'Do you feel you have more problems with your memory than most?' Both criteria were used by Verghese and colleagues in the original MCR definition [8]. Frequency and amount of physical activity were measured using the 12-item Physical Activity Scale for the Elderly (PASE). Scores range from 0 (no activity) to a possible 400 [19]. Almost all answered a core set of questions including falls. About two-thirds of Māori and three quarters of non-Māori answered a full questionnaire including gait speed and cognition,

Data Analysis

Māori and non-Māori cohorts were analysed separately. We did not compare outcomes by gender given the limited data set for fallers especially in later years, but did include gender in final models (see below). Cognition (3MS) was converted to logit of the score due to its distribution which was extremely left skewed. Baseline demographic, clinical data and presence of MCR syndrome were reported. Trajectories of cognition and gait were modelled over 5 years and Latent Class Analysis was used to classify participants into exclusive groups or clusters based on longitudinal gait and cognition scores. Frequency of participants in each gait*cognition cluster were then described. Two models examined the association between gait, cognition and falls using Generalised Estimating Equation (GEE). For both analyses, falls were entered as the dependent variable and considered as *any prospective fall* over the duration of the study rather than restricted to a single year. For the first model (the 'gait & cognition' model) age, gait and cognition were entered as time-varying variables and gender as a fixed variable. The second model (the 'MCR' model) tested main effects and interaction effects for MCR, defined as the combination of subjective cognitive complaint and slow gait according to the thresholds described above. Independent variables included gender (male as the reference), age, gait speed, and cognition (logit of 3MS scores) and two gait*cognition groups: subjective cognitive complaint and normal gait; and no subjective cognitive complaint and slow gait. This enabled us to discern any differential effects of the gait*cognition interaction on falls. GEE confers advantages over other methods for analysing longitudinal data in that it does not penalise for missing data and it accommodates data that is not normally distributed. A criterion for using GEE is that a within-subject correlation structure must be selected prior to analysis [20]. We used an auto-regressive correlation structure, which denotes that observations on the same individual one year apart are higher than those on the same individual two years apart. Parameter

estimates and confidence intervals were computed and odds ratios (OR) for falls risk were calculated from exponentiated estimates, with 95% CIs reported for significant findings only. Underlying statistical assumptions were tested and verified. All statistical analyses were carried out using SAS Version 9.4.

Results

Four hundred and twenty-one Māori over 82 years and 516 non-Māori over 84 years were assessed at baseline, and 104 Māori (24.7%) and 198 non-Māori (38.3%) completed 5 years follow-up. Just over half (54%) of all LiLAC survivors were enrolled in the study at Year 5 (Table 1, Suppl data). Falls data were recorded at baseline for 408 Māori and 512 non-Māori which attenuated to 102 Māori and 197 non-Māori by Year 5. At baseline, 138 (33.8%) Māori and 205 (40%) non-Māori had fallen in the previous 12-months. Falls prevalence did not change markedly over the study, with comparable proportion for both cohorts other than for Years 3 and 5 when significantly more non-Māori fell. Māori were significantly younger than non-Māori by study design, because of the broader eligibility for age, and also presented with a higher body mass index (BMI) and higher scores for physical activity. MCR criteria were met by 10 Māori and 7 non-Māori (Table 1).

Gait speed was assessed for 228 Māori and 374 non-Māori, and cognition (3MS) completed by 266 Māori and 393 non-Māori. Non-Māori men walked significantly faster at baseline than Māori men and had higher scores for cognition (Table 1). Latent class analysis revealed three gait clusters for both cohorts (slow, medium, fast gait), four cognition clusters for Māori (low, low-moderate, moderate, and high) and three cognition clusters for non-Māori (low, moderate, high). Trajectories for gait and cognition differed for both cohorts (Fig 1). For Māori, gait speed declined for the fast gait group who comprised 10.5% of the cohort. For all other Māori and for non-Māori gait speed was fairly stable. For cognition, Māori show a sharp decline between years 1 and 3 followed by cessation of data for the low cognition group who comprised 5% of the Māori cohort. Cognition also declined for the low-moderate group with 3MS scores almost 90/100 at Year 1, lowering to 60/100 and below by Year 5. The wide confidence intervals for this group indicate heterogeneity. Trajectories for the two highest scoring cognitive groups (over two thirds of the total Māori cohort) were stable over time. Compared with Māori, cognitive trajectories for non-Māori were overall more stable. CIs for the highest cognitive groups for Māori and non-Māori were very narrow for all years, due to the left skew of the distribution, and are not visible in Fig 1. For example, baseline Māori 95% CI was 96.4 [95.6, 97.1]; and baseline non-Māori CI was 97.9 [97.6, 98.2].

Fig 2 shows the interaction between gait and cognition for each cluster. There were no fast walkers in the low cognitive group, for either Māori or non-Māori, and only 8 Māori (2.8%) and 28 non-Māori (6.7%) participants met criteria for the 'low cognition and slow gait speed' group.

Table 2 shows the association between prospective falls and gender, age, gait and cognition scores. For the 'gait and cognition' model where gait and cognition were entered as separate variables, gender was the only significant predictor of falls for Māori. Being male increased the odds of falling by 56% (OR 1.56;

95% CI 1.0–2.43 ($P = 0.04$). For non-Māori a slow gait significantly increased the odds of falling by 60% (OR 0.40; 95% CI 0.24–0.68) ($P < 0.001$). Lower 3MS scores also increased falls risk although this was not statistically significant ($P = 0.07$). For the 'MCR model' where MCR and cognition*gait interactions were included as predictors, there were no significant MCR findings for Māori. Non-Māori with MCR were more than twice as likely to fall than those without MCR (OR) 2.45; 95% CI 1.06–5.68 ($P = 0.03$). The interaction effects show that gait speed was a stronger predictor than cognition within this model (OR for gait 1.87. 95% CI 1.34–2.64 ($P < 0.001$) compared with cognition (OR 1.65; 95% CI 1.07–2.55); $P = 0.02$).

Lastly, MCR prevalence for both cohorts was similar from baseline values (Table 1) until years 4 and 5 when no Māori reported MCR, with only 58 participants in Year 4 and 45 participants in Year 5 (data beyond baseline not reported in Table 1).

Discussion

This study is the first to examine the trajectory of gait and cognition over five years with respect to falls in Māori, and to our knowledge the first in non-Māori of advanced age. Main findings were that gait and cognition did not decline markedly over five years, a slower gait significantly increased falls risk for non-Māori, and MCR prevalence was low especially for non-Māori.

Of note are the high attrition rates for both cohorts, especially in the later years. Around 65% of the total sample was lost to follow-up, or death which although not surprising given the advanced age, will have undoubtedly biased results towards participants who were cognitively and motorically resilient, and who fell less. Kerse and colleagues report retention rates and reasons for drop out for the first two years of LiLACS, and note the higher drop-out for Māori for reasons other than death [11]. They also report significant differences in baseline characteristics for those retained in the study compared with those who died or were lost to follow-up. Exceptions to this stability were the two lowest cognitive scoring groups for Māori whose trajectory markedly declined, and who together comprised just over a quarter of the total Māori cohort. A further exception was the increase in the number of non-Māori fallers at Year 5. The modest decline in gait and cognition for those who remained in the study is an important finding and perhaps surprising given the multiple morbidities associated with advanced age, also evident for 93% of LiLAC NZ participants who reported at least two diagnosed health conditions at baseline [22].

Around a third of Māori and approximately 40% of non-Māori fell at any one time over the course of the study, which is broadly comparable with published data for adults over 65 years [23]. Non-Māori were slightly older which may have impacted on falls prevalence, especially by Year 5 where almost half non-Māori fell compared to under a third of Māori. Based on PASE scores at baseline, Māori were significantly more active than non-Māori, suggesting that lower falls prevalence was not related to lower amounts of activity. Falls prevalence for Māori did not increase over time, although the reasons for this are not clear.

Gait speed for non-Māori participants was comparable to norm-referenced data. Oh-Park [24] reports 0.95 m/s for men 80 to 84 years lowering to 0.88 m/s over 85 years, and for women 80 to 84 years 0.87 m/s lowering to 0.78 m/s over 85 years. Average gait speed for Māori was significantly slower, which may

reflect anthropometric differences related to BMI which was significantly higher, or culturally different gait dynamics. Importantly, trajectories for gait speed for over 97% of Māori did not vary over the course of the study, and they were more active than non-Māori. Cognitive outcomes were considerably higher than published data for community dwelling older adults, for both cohorts. Bassuk and Murphy (2003) report an average 3MS score of 75.4 for people 75–84 years, lowering to 63.3 for those over 85 years. Only the two lowest Māori cognitive clusters had scores comparable to these, with all other groups scoring higher.

MCR prevalence at baseline was low particularly for non-Māori, although estimates are comparable to younger cohorts from Australia and UK [8]. Estimates for Māori were higher, although still lower than overall multi-country prevalence of 9.7% [8]. It is unwise to draw any conclusions from these results other than to note the low estimates probably reflect the high functioning nature of both cohorts, despite their advanced age. Kerse and colleagues documented the prevalence of dementia in LiLACS NZ as defined by 3MS scores (≤ 80 for Māori; ≤ 84 non-Māori), noting that the proportion of participants with no dementia stayed steady at approximately 87% [25]. Overall these findings point to a stable cohort with mostly preserved gait and cognition, both of which are critical to independent living and overall functioning.

Results from the multivariate modelling distinguished between cohorts. For Māori, being male was the only significant predictor of falls when gait and cognition were entered separately into the model but not when MCR was included as an independent predictor. This may reflect a modifying effect of gender on gait speed. Māori men walked at a significantly slower pace than non-Māori men, but this was not the case for women. Earlier reports suggest that being female confers a higher risk of falling [26], although data are not specific to adults of advanced years. For non-Māori, gait was a significant predictor of falls, and cognition trended towards significance. The findings were comparable when MCR criteria were included in the model. These differences are interesting, although our interpretation is cautious in line with the small proportion with MCR. Results for non-Māori conform to a substantial body of research supporting the association between gait, cognition and falls in older adults [7 27]. Results for Māori suggest the need for a broader repertoire of predictors to identify associations between falls and socio-cultural features such as whānau (family), living environment and support networks. These ethnic-specific outcomes have shown that risk factors for falls are inconsistent across groups. Intervention strategies may also need rethinking.

The key strength of this study is the robust sampling which enabled reporting of novel data for a large cohort of Maori and non-Māori of advanced years with acceptable rates of attrition. The key limitation is that falls were collected retrospectively via self report, which does not conform to best practice [14]. Also, gait speed was collected over a 3 metre distance which is short, although congruent with population-based, longitudinal studies [28]. Lastly, because Māori were not involved in developing the outcomes for this study they are unlikely to reflect key aspects of Māoritanga (Māori culture, beliefs, traditions), and may therefore be culturally-biased towards non-Māori participants.

In conclusion this study contributes to our understanding of falls evolution and the role of gait and cognition to falls in community-dwelling adults of advanced age. For both Māori and non-Māori who live

on into advanced years, fall frequency is relatively stable and there is minimal decline in gait and cognition. For non-Māori, slow gait speed is a significant predictor of falls. These findings resonate widely, given current and projected demographic trends. Further work is needed to identify culturally appropriate strategies to improve falls risk for those living to advanced age.

Declarations

Acknowledgements

We acknowledge the enormous contribution of the participants and their whānau, community organisations engaged in collecting the data. We also acknowledge our funding sources: the Health Research Council of New Zealand and Ngā Pae o te Māramatanga. We thank Te RōpuKaitiaki o ngā tikanga Māori: Hone and Florence Kameta, Betty McPherson, Paea Smith, Leiana Reynolds and Waiora Port for their guidance; Paea Smith, Hone Kameta and Betty McPherson had since passed away and their wisdom and kindness will always be cherished.

Funding

This work was supported by the Health Research Council of New Zealand (HRC09/ 068B), Ngā Pae o te Māramatanga (the New Zealand National Centre for Research Excellence for Māori) and New Zealand's Ministry of Health.

Ethics approval and consent to participate

The Northern X Regional Ethics Committee of New Zealand granted ethical approval for the longitudinal study in December 2009 (NTX/09/09/088). All participants gave written, informed consent.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request. The process to apply for LiLACS NZ data access is available at <https://www.fmhs.auckland.ac.nz/en/faculty/lilacs.html>, and we encourage applications to ensure the data gets maximal use for population health benefit. Each application is discussed by a leadership group of Māori and non-Māori academics.

Authors' contributions

SL study conception, data analysis & interpretation, original draft and ongoing revision; SM statistical analysis, manuscript revision; RT study conception, manuscript revision; WP study conception; MM-L study conception; CB study conception, critical revision; TW critical revision, NK LiLACS PI, study conception, data analysis & interpretation, manuscript revision.

Consent for publication

Not applicable

Competing interests

The authors declare they have no competing interests

References

1. Haagsma JA, Graetz N, Bolliger I, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. *Inj Prev* 2016;22(1):3–18
2. United Nations, Department of Economic and Social Affairs, Division P. World Population Ageing 2015 (ST/ESA/SER.A/390). New York, 2015.
3. Zealand HQSCN. Falls Atlas domain: falls in people aged 50 and over. Secondary Falls Atlas domain: falls in people aged 50 and over 2016 <https://www.hqsc.govt.nz/our-programmes/health-quality-evaluation/projects/atlas-of-healthcare-variation/falls/> Accessed 21st Oct 2019
4. Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988;319(26)
5. Hopewell S, Adedire O, Copsey BJ, et al. Multifactorial and multiple component interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2018;7:Cd012221
6. Montero-Odasso M, Verghese J, Beauchet O, Hausdorff JM. Gait and cognition: a complementary approach to understanding brain function and the risk of falling. *J Am Geriatr Soc* 2012;60(11):2127–36
7. Montero-Odasso M, Speechley M, Chertkow H, et al. Donepezil for gait and falls in mild cognitive impairment: a randomized controlled trial. *Eur J Neurol* 2019;26(4):651–59
8. Verghese J, Annweiler C, Ayers E, et al. Motoric cognitive risk syndrome: multicountry prevalence and dementia risk. *Neurology* 2014;83(8):718–26

9. Verghese J, Wang C, Lipton RB, Holtzer R. Motoric cognitive risk syndrome and the risk of dementia. *J Gerontol A Biol Sci Med Sci* 2013;68(4):412–8.
10. Callisaya ML, Ayers E, Barzilai N, et al. Motoric Cognitive Risk Syndrome and Falls Risk: A Multi-Center Study. *J Alzheimers Dis* 2016;53(3):1043–52
11. Kerse N, Teh R, Moyes SA, et al. Cohort Profile: Te Puawaitanga o Nga Tapuwae Kia Ora Tonu, Life and Living in Advanced Age: a Cohort Study in New Zealand (LiLACS NZ). *Int J Epidemiol* 2015;44(6):1823–32
12. Hayman KJ, Kerse N, Dyllal L, et al. Life and Living in Advanced Age: A Cohort Study in New Zealand - Te Puāwaitanga o Nga Tapuwae Kia Ora Tonu, LiLACS NZ: Study protocol. *BMC Geriatrics* 2012;12(1):33
13. Dyllal L, Kepa M, Hayman K, et al. Engagement and recruitment of Maori and non-Maori people of advanced age to LiLACS NZ. *Aust N Z J Public Health* 2013;37(2):124–31
14. Lamb SE, Jorstad-Stein EC, Hauer K, Becker C. Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* 2005;53(9):1618–22
15. Guralnik JM, Ferrucci L, Pieper CF, et al. Lower extremity function and subsequent disability: consistency across studies, predictive models, and value of gait speed alone compared with the short physical performance battery. *J Gerontol A Biol Sci Med Sci* 2000;55(4):M221–31
16. Teng EL, Chui HC. The Modified Mini-Mental State (3MS) examination. *J Clin Psychiatry* 1987;48(8):314–8
17. Bassuk SS, Murphy JM. Characteristics of the Modified Mini-Mental State Exam among elderly persons. *J Clin Epidemiol* 2003;56(7):622–8
18. Yesavage JA, Brink TL, Rose TL, et al. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res* 1982;17(1):37–49
19. Washburn RA, McAuley E, Katula J, Mihalko SL, Boileau RA. The physical activity scale for the elderly (PASE): evidence for validity. *J Clin Epidemiol* 1999;52(7):643–51
20. Liu S, Dixon J, Qiu G, Tian Y, McCorkle R. Using generalized estimating equations to analyze longitudinal data in nursing research. *West J Nurs Res* 2009;31(7):948–64
21. Salmond C, Crampton P, Sutton F. NZDep91: a New Zealand index of deprivation. *Aust NZ J Public Health* 1998;22:835–7
22. Teh R, Kerse N, Kepa M, et al. Self-rated health, health-related behaviours and medical conditions of Maori and non-Maori in advanced age: LiLACS NZ. *N Z Med J* 2014;127(1397):13–29
23. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing* 2006;35 Suppl 2:ii37-ii41
24. Oh-Park M, Holtzer R, Xue X, Verghese J. Conventional and robust quantitative gait norms in community-dwelling older adults. *J Am Geriatr Soc* 2010;58(8):1512–8

25. Kerse N, Lapsley H, Moyes S, Zawaly K, K H. Dementia Supplementary Findings from LiLACS NZ for Section Five, 'Service Use and Common Health Conditions' in the report 'Health, Independence and Caregiving in Advanced Age'. Wellington, New Zealand: Ministry of Health, 2017.
26. Chang VC, Do MT. Risk factors for falls among seniors: implications of gender. *Am J Epidemiol* 2015;181(7):521–31
27. Mirelman A, Herman T, Brozgol M, et al. Executive function and falls in older adults: new findings from a five-year prospective study link fall risk to cognition. *PLoS One* 2012;7(6):e40297
28. Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *Jama* 2011;305(1):50–8

Tables

Table 1 Baseline demographic & clinical outcomes, and falls status over 5 years

	Māori Mean (SD)	non-Māori Mean (SD)	<i>P</i>
Demographic and clinical outcomes			
Age <i>M</i> = 421; <i>non-M</i> = 516	82.6 (2.7)	84.6 (.51)	<0.001
BMI <i>M</i> = 247; <i>non-M</i> = 375	29.4 (5.4)	26.8 (4.0)	<0.001
Gait speed (m/s) <i>M</i> = 243; <i>non-M</i> = 376	.66 (.29)	.82 (.26)	<0.001
Gait speed (m/s) Men <i>M</i> = 96; <i>non-M</i> = 179	.70 (.25)	.87 (.27)	<0.001
Gait speed (m/s) Women <i>M</i> = 147; <i>non-M</i> = 197	.64 (.30)	.78 (.25)	.099
3MS (maximum score 100)	86.2 (15.6)	91.6 (10.4)	<0.001
<i>M</i> = 266; <i>non-M</i> = 393			
GDS_15(range 0 - 15)	2.5 (2.2)	2.1 (1.9)	<0.05
<i>M</i> = 255; <i>non-M</i> = 392			
Total PASE score (maximum score 400)	111.0 (85.0)	98.9 (66.0)	<0.05
<i>M</i> = 262; <i>non-M</i> = 403			
MCR (gait speed & GDS Item 10)	10 (4.2%)	7 (1.9%)	<0.001
<i>M</i> = 238; <i>non-M</i> = 371 (%)			
Falls status total n; falls n (%)			
Baseline	408; 138 (33.8%)	512; 205 (40%)	.055
Year 1	262; 90 (34.4%)	391; 160 (40.9%)	.101
Year 2	192; 54 (28.1%)	334; 121 (36.2%)	.068
Year 3	157; 53 (33.8%)	276; 121 (43.8%)	<0.05
Year 4	116; 35 (30.2%)	228; 83 (36.4%)	.280
Year 5	102; 30 (29.4)	197; 92 (46.7%)	<0.01

Abbreviations : Body Mass Index (BMI); Modified Mini Mental Status Examination (3MS); Geriatric Depression Scale (GDS); Physical Activity Scale for the Elderly (PASE); Motoric Cognitive Risk Syndrome (MCR)

^(a)Student *t*-test for all comparisons other than MCR and falls status (χ^2 test)

Table 2 Association of gait, cognition, and MCR with falls over 5 years

Predictors	Estimate	95% CI	OR ^(a)	P
<i>Māori gait & cognition model</i>				
Gender (Male)	0.44	0.00 to 0.88	1.56	0.04
Age	0.04	-0.02 to 0.12	1.62	0.21
Cognition(3MS)	-0.04	-0.22 to 0.13	0.95	0.64
Gait speed (m/s)	0.18	-0.46 to 0.83	1.19	0.58
<i>Māori MCR model</i>				
Gender (Male)	0.41	-0.85 to 0.20	0.66	0.06
Age	0.04	-0.02 to 0.12	1.05	0.20
MCR (subjective cognitive complaint and slow gait)	0.24	-0.87 to 1.35	1.27	0.67
Subjective cognitive complaint *normal gait	0.10	-0.38 to 0.60	1.11	0.66
No subjective cognitive complaint * slow gait	-0.33	-0.94 to 0.27	0.71	0.27
<i>Non-Māori gait & cognition model</i>				
Gender (Male)	0.12	-0.44 to 0.19	0.88	0.44
Age	0.05	-0.02 to 0.13	1.05	0.17
Cognition (3MS)	-0.11	-0.24 to 0.00	0.89	0.07
Gait speed (m/s)	-0.91	-1.44 to -0.38	0.40	<0.001
<i>Non-Māori MCR model</i>				
Gender (Male)	0.15	-0.16 to 0.46	1.16	0.33
Age	0.05	-0.02 to 0.13	1.05	0.19
MCR (subjective cognitive complaint and slow gait)	0.89	0.05 to 1.73	2.45	0.03
Subjective cognitive complaint *normal gait	0.50	0.06 to 0.93	1.65	0.02
No subjective cognitive complaint * slow gait	0.63	0.29 to 0.97	1.87	<0.001

Abbreviation ^(a)OR calculated from exponentiated estimates

Figures




Figure 1 Clusters of Māori and non-Māori participants based on gait and cognition trajectories over 5 years. Frequency of participants in each cluster described as a percent of the total cohort.

Figure 1

Clusters of Maori




Figure 2 Frequency of participants in each gait*cognition cluster to illustrate the interaction between both these characteristics.

Figure 2

Frequency of participants

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TS.docx](#)