

# Changes in Soil Chemical Properties Under Different Land-Use Systems and Soil Depth Layers: A Case Study from the Central Highlands of Ethiopia

Yadesa Bato (✉ [yadesabato2020@gmail.com](mailto:yadesabato2020@gmail.com))

Ambo University

**Tamrat Bekele**

Addis Ababa University College of Natural Sciences

**Sebebe Demissew**

Addis Ababa University College of Natural Sciences

---

## Research

**Keywords:** land-use system, soil depth, soil chemical properties, natural forest, exotic tree plantation, Yerer

**Posted Date:** September 18th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-74895/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

## Abstract

**Background:** Soil chemical properties have changed under different land-use systems and soil depth layers either by increasing or decreasing. Hence, scientifically information on the soil chemical properties dynamics under different land-use systems and soil depths are crucial for best land management practices, and to avoiding ecological negative impacts of it for sustainable development. The study aimed to evaluate the soil chemical properties dynamics under different land-use systems and soil depths in the central highlands of Ethiopia. The land-use systems included natural forest, four exotic tree plantation species (*Eucalyptus globules*, *Cupressus lusitanica*, *Grevillea robusta*, and *Pinus patula*), grassland, grazing land, and cropland.

**Results:** The analysis of variance (ANOVA) for the majority of soil chemical properties of OC, TN, Avail. P, soil pH, EC, CEC, and exchangeable bases (Ca, Mg, K, Na) were showed that significant variations among land-use systems ( $P < 0.0001$ ). The highest mean values of OC (3.49 % DM), TN (0.31 % DM), Avail.P (31.52 mg/kg of soil), CEC (33.63 meq/100gm soil), Exch. Ca (17.13 cmol(+)/kg soil), Exch. Mg (5.37 cmol(+)/kg soil), and Exch. K (3.60 cmol(+)/kg soil) were observed under natural forest than others of land-use systems. The results also showed that the lowest mean values of OC (1.47 % DM), TN (0.13 %DM), soil pH (5.38), CEC (18.98 meq/100gm soil), Exch. Ca (9.93 cmol(+)/kg soil), Exch. K (1.20 cmol(+)/kg soil), and Exch. Na (0.22 cmol(+)/kg soil) were recorded under cropland than other land-use systems. The highest mean values of EC (3.47ds/m), and Exch. Na (0.60 cmol(+)/kg soil) were observed under *Eucalyptus globulus* plantation forest. The overall mean values of OC, TN, Avail.P, CEC, Exch. Mg, Exch. Ca, Exch. K, and Exch. Na accumulation at the topsoil layer was higher than that of the subsoil layer except for soil pH and EC.

**Conclusion:** In general, the majority of soil chemical properties under cropland and *Eucalyptus globulus* plantation forest were poorer than the soils subjected to other land-use systems which indicated that changes in land use systems were significantly affected soil chemical properties.

## Background

Land use defined as the human activities, arrangements, and inputs that apply to the use of lands such as industrial zones, residential zones, and agricultural fields, while, land cover is the biophysical conditions that cover on the ground surface, like crops, forest, and grassland (Di Gregorio and Jansen 1998, FAO 2000). Land-use change is the main component of environmental change in every region that alters biodiversity, soil properties, and water resources interims of quality and quantity, and these in combine which affects the ecosystem functions and climate (Foley et al. 2005, Newbold et al. 2015). The world's forests contribute to essential ecosystem services such as carbon uptake; maintain high biodiversity, improved soil properties, and water quality (Hansen et al. 2013; Riebeek, 2011). The loss of forest cover and biodiversity due to anthropogenic activities is a growing concern in many parts of the world (Singh et al. 1997; Hegde and Enters 2000). Even though deforestation, forest degradation, and forest fragmentation are a global phenomenon, but they are severe in the tropical region. Nearly half (½) of the tropical natural closed-canopy forest land has been cleared and transformed into other land-uses for the sake of fulfilling basic needs (Wright, 2005). Plantation forests have been a significant element of land-use change. The world-wide relatively the annual expansion rate of plantation forest area is predicted to be approximately 2% (FAO, 2005; van Dijk & Keenan, 2007). The reasons for the expansion rate of plantation forest area can result from the demand of the world's increasing population for domestic and industrial wood products (Berthrong et al., 2009).

Across sub-Saharan Africa countries, natural resources remain central to rural people's livelihoods (Roe et al. 2009). Due to the cause of anthropogenic force can exert pressure on the natural resources, thereby influencing spatial and temporal scale changes on landscape resources (Roe et al. 2009). Land-use change is rapid in developing countries due to biophysical and socio-economic reasons. Ethiopia has lost the majority of natural forest resources, especially during the 20th century. On the other hand, reforestation and afforestation programs have helped to reverse the decline of the natural forest resource cover and its results of a shortage of wood products, fuelwood, charcoal, construction materials, and soil degradation (Duguma et al. 2009). At the same time, afforestation and reforestation programs have led to increasing forest area coverage in some regions and reducing the national net forest loss by lowering human pressure on natural forests. The expansion of plantation land use type with different species has significantly contributed to an increase in total forest plantation coverage area from an estimated 190,000 ha in 1990 to approximately 972,000 ha in 2010 (Jaleta, 2016). Reforestation in primary and secondary forest lands accounts for about half of the total increased area of plantations (FAO 2005). Plantation forests have the potential to alter the biogeochemical cycles of the ecosystem as a consequence of changes in tree species composition when compared with other adjacent natural forests (Aborisode & Aweto, 1990; Wall & Hytönen, 2005; Freier et al. 2010).

Yerer Mountain forest is one of the protected dry montane forests in the central highlands of Ethiopia. Rapid population growth requires additional farmlands for food crop production by converting forests and pasture lands to cropland to address their basic needs. In the study area, on one hand, forest and pasture areas have decreased significantly, while cultivated cropland areas have increased with intensively continuous cultivation, and on the other hand, plantation forests establishment has intensively expanded in large-scale from year to year in both reforestation and afforestation programs with the fast-growing and short rotation of exotic species by governmental sector and local communities. Though, the change of land-use systems by human intervention may cause significant changes in soil chemical properties in the study area. Changes in land use systems significantly alter the soil's physical, chemical, and biological composition and qualities (Lemeniha et al. 2004, Lemma et al. 2006; Fantaw et al. 2008, Lalisa, et.al. 2010, Tellen et.al. 2018). However, different researchers were reported to contradict ideas on the impact of exotic tree species plantation on environmental change particularly on soil chemical properties change as increases (positive), decreases (negative), or negligible (neutral) effect.

To understand the effect of human activities on the soil environment by land-use system change is fundamental to understanding global change and sustainable development. However, information on the soil chemical properties dynamics under different land-use systems and soil depths in the central highlands of Ethiopia are still too scanty to recommend the optimal sustainable management and utilization of land resources for the future. With this background, the present study was taken up to evaluate the soil chemical properties change under different land-use systems and soil depths, and to provide additional current scientifically information to build literature surrounding the effect of land-use systems on soils, particularly debate on the effect of exotic tree plantation species, thereby, based on the quantified evaluation results the study to recommend the positive effects would be accentuated but the negative

effects would be avoided. The study aimed to evaluate the soil chemical properties dynamics under different land-use systems and soil depths in the central highlands of Ethiopia.

## Methods And Materials

### Description of the study area

#### Location

A field study was conducted in the central highland of Ethiopia, Oromia Regional State, Yerer forests, and surrounding area. Yerer forest is one of the remnant montane evergreen natural forest and plantation forests with different exotic tree species in the central highland of Ethiopia. The study area is located in the central-eastern part of Ethiopia at 40 km southeast far from Addis Ababa, and 17 km north of Bishoftu town. Yerer forest and surrounding areas is located between 8°52'00" – 8°55'00" N latitude and 38°59'30" – 38°95'15" E longitude, with altitudinal ranges between 2000 and 3105 m.a.s.l. (Fig. 1).

#### Climate

The 31 years meteorological data(1988–2019) showed that the mean annual rainfall of the study area is 907.33 mm which characterized by a unimodal rainfall having a long dry season (6–8 months), and the mean annual minimum and maximum temperatures were recorded 11.1 °C, and 29.5°C, respectively National Meteorology Service Agency (2019). The farmers in the central highland of Ethiopia area are practicing "mixed" farming systems (i.e. both crop and livestock production activities).

#### The vegetation of the area

Yerer Mountain natural forest is one of the remnants and secondary natural forests, which are categorized under the Dry evergreen montane forest in the central highlands of Ethiopia. Yerer Mountain natural forest is consists of natural forest and established plantation forests, and covers an area of 3254 ha of this 1793 ha is plantation forest with exotic and native tree species in monoculture and mixed forms, and 1461 ha is covered by natural forest and others. Some tree, shrub, and climber species dominate the mixed deciduous forest. Vegetation composition of the Yerer Mountain natural forest is dominated by some tree species of *Juniperus procera*, *Olea europaea* subsp. *cuspidata*, *Ficus sur*, *Rhus retinorrhoea*, *Olinia rochetiana*, *Rhus vulgaris*, and shrub species of *Carissa spiranum*, *Dodonaea angustifolia*, *Myrsine africana*, *Erica arborea*, *Hypericum revolutum*, *Maytenus arbutifolia*, *Osyris quadripartite*, *Euclea racemosa*, *Heteromorpha arborescens*, and *Pterolobium stellatum*.

Recently, plantation activities were carried out by different exotic and native tree species in, and surround the Yerer natural forest area. The major plantations of exotic tree species are *Eucalyptus globulus*, *Cupressus lusitanica*, *Eucalyptus camandulensis*, *Pinus patula*, and *Grevillea robusta*; however, the native tree species include *Juniperus procera*, *Hagenia abyssinica*, *Podocarpus falcatus*, *Olea europaea* subsp.*cuspidata*, and *Cordia Africana*.

#### Land-use systems

In the Yerer forest and its surrounding area, the major land-use systems are a natural forest, exotic tree plantation species forests, grassland, grazing land, and cropland. The land-use change started from 1991 and in the past two half decades in the central highlands of Ethiopia, pasture, and forest areas have decreased significantly, while arable cropland has increased proportionally due to the rapid population growth requires additional farmlands for food crop production. In the present study context definition: Natural forest is the forest land that is composed primarily of the indigenous (native) tree, shrubs, lianas, climbers, and other herbaceous plant species, which may include both closed forest and open forest. Exotic tree plantation species is described as the forest stands established artificially by non-native tree or shrub species in reforestation and afforestation program for industrial and non-industrial purposes. Grassland is an area where characterized as vegetation land dominated by the nearly continuous cover of grasses (Poaceae) family rather than large shrubs or trees species. Grazing land – is a land covered with grass or herbage and suitable for grazing or feeding by livestock or herbivore. Croplands is referring to land covered with temporary crops production both in single or multiple cropping systems, followed by harvest and leave a bare soil period.

#### Sampling Design

For the present study, eight (8) different land-use systems (LUSs) were selected, namely: natural forest, four exotic tree plantation species (*Eucalyptus globules*, *Cupressus lusitanica*, *Grevillea robusta*, and *Pinus patula*), grassland, grazing land, and cropland. In this study, the eight land-use systems were considered as treatments and the sample quadrats in each land-use system were considered as a replication. The systematic sampling design was used to determine the soil chemical properties with transect lines laid in each land-use systems (LUS). Four transect lines were delineated and laid parallel to each other along the altitudinal gradient in each land-use system separately for soil sample collection by using the 50-meter distance between two adjacent transect lines, and between the consecutive interval of quadrat on a transect line to avoid biased of samples selection.

#### Soil sampling procedure, data collection, and sample preparation

For soil data collection of site selection from eight (8) land-use systems (LUS) were considered the relative similarity of each LUS as criteria in their topography, altitude, aspect, slope, and areas free from erosion exposed. For soil data collection the main square sample quadrat of 20 m x 20 m (400 m<sup>2</sup>) area was established by systematic sampling techniques with twelve (12) replication quadrats in each land-use system. The soil samples were collected in five replicates of subplots with 1 m x 1 m area within each a main square quadrat by placing four (4) subplots at each corner and one (1) subplot at the center of the main quadrat. The soil samples were collected close to the center of each subplot (1 m x 1 m area) by using a soil auger in adjusting with the required soil depth layers and by carefully cleaning the equipment at each soil sampling depths, to prevent soil contamination between different soil sampling layers and quadrats. The soil samples were collected at two soil depth layers; each layer is 20 cm thick at the topsoil layer (0–20 cm) and subsoil layer (20–40 cm) at the center of each sub-plot area.

The collected five soil samples from each main quadrat were bulked /combined with respect to their similar depth layer categories and mixed well thoroughly to form a one representative composite soil sample per the main quadrat to reduce variability within a quadrat. Finally, 1 kg of the composited representative soil samples from each quadrat concerning each depth layer was packed into plastic bags and transport to soil chemical laboratory analysis. The collected representative soil samples were labeled with appropriate information and transport immediately to the soil laboratory of Hawassa University, Wondo Genet College of Forestry and Natural Resources, for sample preparation and laboratory analysis. Before the analysis, the composite soil samples were air-dried, ground, and sieved to pass through a 2 mm size sieve in preparation for laboratory analysis of most soil chemical properties. The soil samples were further sieved to pass through a 0.5 mm size sieve for the analysis of total nitrogen (Ranst et al., 1999).

## **Laboratory Analysis of soil chemical properties**

The soil chemical properties analysis parameters include organic carbon, total nitrogen, C: N ratio, available phosphorous, soil pH, soil electrical conductivity, Cation exchange capacity, exchangeable bases (Ca, Mg, K, and Na), and Percent base saturation. The soil organic carbon (OC) was determined following the Walkley Black wet digestion method as described by Ranst et al. (1999). The total nitrogen level in the soil was determined following the modified Kjeldahl's digestion, distillation, and titration method as described by Ranst et al. (1999). The C: N ratio was determined by dividing the value of organic carbon to the total nitrogen corresponding to each soil sampling depths. The soil available phosphorus content was determined by following the Olsen procedure (Olsen et al., 1954). The soil pH was measured by using a digital pH meter in a 1:2.5 soil to water supernatant suspension (Jackson, 1973, Van Reeuwijk 1992). The Electric conductivity of the soil samples was determined by using an electrical conductivity meter from 1:1 soil-distill water mixture (20 g of soil in 20 ml distilled water). The cation exchange capacity (CEC) of soils was determined by saturation with 1N sodium acetate followed by replacement of sodium on the exchange complex with 1N ammonium acetate at pH 7 following the procedure described by Chapman (1965). Sodium level was determined by atomic absorption spectrometry (AAS) and CEC expressed as meq/100 g of soils. The exchangeable bases were extracted with 1M ammonium acetate at pH 7.0. Exchangeable Ca and Mg were measured from the extract with an atomic absorption spectrophotometer; whereas exchangeable Na and K were determined from the same extraction with a flame photometer (Black et al; 1965). Percent base saturation (PBS) was determined by dividing total exchangeable bases (Ca, Mg, K, and Na) to the CEC of the soil and multiplies by 100.

## **Data Analysis**

All quantitative field soil data were analyzed by the General linear model (GLM) analysis of variance (ANOVA) procedure. Statistical difference in soil chemical properties among land-use systems and between soil depth layers was analyzed by a two- way analysis of variance (ANOVA) at  $p < 0.05$  significant levels by using Statistical Analysis System (SAS) software (Version 9.4). A least significant difference (LSD) test was employed to assess the mean difference between soil variables. Means comparisons were used to soil properties change interpretation and explanation among eight land-use systems in the study area. Additionally, Pearson's correlation coefficient was employed to evaluate the relations of various soil chemical variables.

## **Results**

### **Effect of land-use systems on soil chemical properties**

#### **Soil Organic C, Total N, C: N ratio and Available P**

##### **Soil Organic Carbon (%OC)**

The analysis of variance for organic carbon concentration of the soil revealed a significant difference horizontally as a function of land-use systems ( $p < 0001$ ). The overall mean value of the organic carbon concentration under the eight different land-use systems (natural forest, four different exotic tree species plantation forest, grassland, grazing land, and cropland) was varied between 1.47 to 3.49%. Among the land-use systems (LUS) the highest concentration of organic carbon was observed in the natural forest, whereas in contrary the lowest concentration of organic carbon was observed in cropland (Table 1, and Fig. 2).

Table 1  
Least Significant Mean values of selected soil chemical properties of different land-use systems

Land-use systems	OC (%DM)	Total N (%DM)	C:N ratio	Avail. P (mg/kg of soil)	PH-H <sub>2</sub> O (1:2.5)	EC (ds/m)	CEC (meq/100 gm)
Natural forest	3.49 <sup>a</sup>	0.31 <sup>a</sup>	11.38 <sup>a</sup>	31.52 <sup>a</sup>	6.20 <sup>ba</sup>	1.78 <sup>c</sup>	33.63 <sup>a</sup>
<i>Eucalyptus globulus</i>	2.03 <sup>c</sup>	0.17 <sup>dc</sup>	11.87 <sup>a</sup>	3.12 <sup>d</sup>	5.48 <sup>cb</sup>	3.47 <sup>a</sup>	25.78 <sup>cb</sup>
<i>Cupressus lusitanica</i>	2.29 <sup>cb</sup>	0.20 <sup>c</sup>	11.75 <sup>a</sup>	2.45 <sup>d</sup>	5.41 <sup>c</sup>	2.34 <sup>b</sup>	21.26 <sup>dc</sup>
<i>Grevillea robusta</i>	3.14 <sup>a</sup>	0.26 <sup>b</sup>	12.03 <sup>a</sup>	5.82 <sup>dc</sup>	5.70 <sup>cb</sup>	2.42 <sup>b</sup>	23.24 <sup>c</sup>
<i>Pinus patula</i>	2.50 <sup>cb</sup>	0.22 <sup>cb</sup>	12.08 <sup>a</sup>	20.32 <sup>b</sup>	5.99 <sup>b</sup>	2.35 <sup>b</sup>	29.23 <sup>b</sup>
Grassland	2.58 <sup>b</sup>	0.23 <sup>cb</sup>	11.86 <sup>a</sup>	14.88 <sup>cb</sup>	6.49 <sup>a</sup>	1.97 <sup>cb</sup>	28.97 <sup>b</sup>
Grazing land	2.16 <sup>cb</sup>	0.18 <sup>dc</sup>	11.94 <sup>a</sup>	11.89 <sup>c</sup>	6.00 <sup>ba</sup>	1.88 <sup>cb</sup>	27.46 <sup>b</sup>
Cropland	1.47 <sup>d</sup>	0.13 <sup>d</sup>	11.48 <sup>a</sup>	6.52 <sup>dc</sup>	5.38 <sup>c</sup>	3.07 <sup>a</sup>	18.98 <sup>d</sup>
LSD	0.54	0.05	1.42	8.31	0.49	0.56	3.49
CV	24.61	26.13	13.48	77.13	9.44	26.50	15.02
SIGN	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>	ns	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>

**Note:** Means for land-use systems within a column followed by the same letter are not significantly different from each other at  $p < 0.05$ . However, mean values of different letters within a column are indicate significant differences by,  $a > ab > b > c > d > e > f$

Keys: \*, \*\*, \*\*\*, and ns = significant at  $P < 0.05$ , high significant at  $P < 0.01$ , very high significant at  $P < 0.001$ , and nonsignificant at  $P > 0.05$ , respectively

## Total nitrogen (%TN)

Statistically, the total nitrogen content of the soil showed a significant variation as a function of land-use systems ( $p < 0.0001$ ). The overall mean value of the total nitrogen levels in all different land-use systems varied between 0.13 to 0.22% ranges. Among the land-use systems, the maximum and minimum mean values of total nitrogen percent were recorded in natural forest and cropland in order of 0.31% and 0.13%, respectively (Table 1, and Fig. 2).

## Carbon to Nitrogen(C: N) ratio

The C: N ratio shall determine by dividing the value of organic carbon to the total nitrogen corresponding to each soil sampling depths. The overall ANOVA analysis result revealed that there was a non-significant ( $p < 0.05$ ) influenced by any land-use systems. The overall mean value of the C: N ratio under eight different land-use systems were felled between the 11.38 to 12.08 ranges (Table 1).

## Available phosphorus

The analysis of variance for available phosphorus revealed that there were significant differences ( $P < 0.0001$ ) between all land-use systems except between *Eucalyptus globulus* and *Cupressus lusitanica*, and between *Grevillea robusta* and cropland. The overall available phosphorus concentration level under different land-use systems was observed in the range between 2.45 to 31.52 mg/kg of soil. Among the land-use systems (LUS), the maximum and minimum available phosphorus concentration were recorded in natural forest and *Cupressus lusitanica* plantation forest, respectively (Table 1, and Fig. 2).

## Soil pH, EC (ds/m), and CEC

### Soil pH /Soil reaction

Soil reaction or soil pH is one of the soil chemical properties that used to measure the acidity or alkalinity of the soil. The analysis of variance for soil pH revealed that a significant difference as a function of land-use systems ( $p < 0.0001$ ) except between *Eucalyptus globulus* and *Cupressus lusitanica*. The overall mean value of pH in the soil under the different land-use systems was observed in the range between 5.38 to 6.49. Among the land-use systems (LUS), the maximum and minimum pH in soil was recorded in grassland, and cropland, respectively (Table 1, and Fig. 2).

### Soil electrical conductivity (EC)

Soil electrical conductivity (EC) is a measure of the number of salts in soil (salinity of soil). The statistical analysis for soil electrical conductivity revealed a significant variation among land-use systems ( $P < 0.0001$ ) except between *Eucalyptus globulus* and cropland, between *Grevillea robusta* and *Pinus patula*, and between grassland and grazing land. The overall mean values of soil electrical conductivity (EC) distribution under different land-use systems were varied in the range of 1.78 to 3.47 ds/m. Among all land-use systems, the maximum and minimum mean values of soil electrical conductivity (EC) concentration were observed in *Eucalyptus globulus* plantation forest, and natural forest, respectively (Table 1, and Fig. 3).

### Cation-Exchange Capacity (CEC)

The analysis of variance (ANOVA) for cation exchange capacity (CEC) was showed a significant variation among land-use systems ( $P < 0.0001$ ) except between *Pinus patula*, grassland, and grazing land. The overall mean values of Cation-exchange capacity (CEC) distribution under different land-use systems

were varied in the range of 18.98 to 33.63 meq/ 100 gm soil. Among all land-use systems, the maximum and minimum mean values of soil Cation-exchange capacity (CEC) concentration were observed in natural forest(33.63), and cropland (18.98), respectively (Table 1, and Fig. 3).

## Soil Exchangeable bases (Na, K, Ca, Mg), and Percent Base saturation (PBS)

### Exchangeable cations (Na, K, Ca, Mg)

The common exchangeable base cations are K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, and Na<sup>+</sup> which are held in the soil by organic matter and clay. The analysis of Variance (ANOVA) made for exchangeable cations of Na, K, Ca, and Mg concentrations were revealed that significant variations, laterally among land-use systems for the concentration of Na (P < 0.0001), K (P < 0.0001), Ca (P < 0.0001), and Mg (P < 0.0001). The present study results indicate that the concentration of exchangeable cations was observed generally in the order of Ca > Mg > K > Na in all different land-use systems. The overall mean values of the distribution of exchangeable Ca in the study area were found in the orders of natural forest > grassland > *Pinus patula* > grazing land > *Grevillea robusta* > *Cupressus lusitanica* > *Eucalyptus globules* > agricultural cropland, with numerical values of 17.13, 16.03, 14.93, 13.36, 12.56, 11.59, 11.25, and 9.93, respectively. Among all land-use systems, the higher concentration of exchangeable cations of Ca, Mg, and K was observed in the natural forest. The higher concentration of exchangeable cations of Na<sup>+</sup> was recorded in the *Eucalyptus globulus* plantation forest (Table 2, and Fig. 4).

Table 2  
Least Significant Mean values of selected soil chemical properties of different land-use systems

Land-use systems	Exch. Na (cmol(+)/kg soil)	Exch. K (cmol(+)/kg soil)	Exch. Ca (cmol(+)/kg soil)	Exch. Mg (cmol(+)/kg soil)	BS (%)
Natural forest	0.39 <sup>b</sup>	3.60 <sup>a</sup>	17.13 <sup>a</sup>	5.37 <sup>a</sup>	77.94 <sup>ba</sup>
<i>Eucalyptus globulus</i>	0.60 <sup>a</sup>	1.97 <sup>cb</sup>	11.25 <sup>cb</sup>	3.58 <sup>b</sup>	70.91 <sup>b</sup>
<i>Cupressus lusitanica</i>	0.27 <sup>ed</sup>	2.46 <sup>b</sup>	11.59 <sup>cb</sup>	2.91 <sup>cb</sup>	80.83 <sup>a</sup>
<i>Grevillea robusta</i>	0.34 <sup>d</sup>	2.19 <sup>cb</sup>	12.56 <sup>b</sup>	3.51 <sup>b</sup>	79.78 <sup>ba</sup>
<i>Pinus patula</i>	0.30 <sup>ed</sup>	2.45 <sup>b</sup>	14.93 <sup>ba</sup>	4.75 <sup>ba</sup>	76.35 <sup>ba</sup>
Grassland	0.29 <sup>ed</sup>	1.99 <sup>cb</sup>	16.03 <sup>a</sup>	3.42 <sup>b</sup>	73.32 <sup>ba</sup>
Grazing land	0.33 <sup>d</sup>	2.57 <sup>b</sup>	13.36 <sup>b</sup>	3.21 <sup>cb</sup>	74.53 <sup>ba</sup>
Cropland	0.22 <sup>e</sup>	1.20 <sup>d</sup>	9.93 <sup>c</sup>	3.47 <sup>b</sup>	77.80 <sup>ba</sup>
LSD	0.09	0.65	2.53	0.98	9.92
CV	24.53	31.78	21.27	28.94	14.53
SIGN	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>	0.0001 <sup>***</sup>	ns

**Note:** Means for land-use systems within a column followed by the same letter are not significantly different from each other at p < 0.05. However, mean values of different letters within a column are indicate significant differences by, a > b > c > d > e > f

Keys: \*, \*\*, \*\*\*, and ns = significant at P < 0.05, high significant at P < 0.01, very high significant at P < 0.001, and non-significant at P > 0.05, respectively.

The distribution of exchangeable cations of Mg in *Eucalyptus globules*, *Grevillea robusta*, cropland, grassland, grazing land, and *Cupressus lusitanica* in the order of 3.58, 3.51, 3.47, 3.42, 3.21, and 2.91, respectively, without significant difference among them statistical. However, the concentration level of exchangeable Mg under natural forest (5.37), and *Pinus patula* (4.75) were significantly different from the other land-use systems.

The distributions of exchangeable cations of K in the study land-use systems were observed: grazing land > *Cupressus lusitanica* > *Pinus patula* > *Grevillea robusta* > grassland > *Eucalyptus globulus* > cropland, which numerical values of 2.57, 2.46, 2.45, 2.19, 1.99, 1.97, and 1.20, respectively, without significant difference among them statistical. However, the concentration level of exchangeable K under natural forest was significantly different from the other land-use systems.

### Percent base saturation (PBS)

Percent base saturation (PBS) is determined by dividing total exchangeable bases (Ca, Mg, K, and Na) to the CEC of the soil and multiplies by 100. The analysis of variance (ANOVA) for percent base saturation revealed a non-significant difference with horizontal as a function of land-use systems, (at p < 0.05). In the study area, the mean values of Percent base saturation (PBS) in all land-use systems (LUS) were recorded more than 70%. Among all land-use systems (LUS) the minimum and maximum Percent base saturation (PBS) were observed in *Eucalyptus globulus*(70.91%), and *Cupressus lusitanica* (80.83%) plantation forests, respectively (Appendix Table 2).

### Effect of soil depth on soil chemical properties

The analysis of variance revealed that the soil OC, Total N, Avail. P, soil pH, CEC, Exch. Ca, Exch. Mg, Exch. Na, Exch. K and BS% were significantly affected by soil depth. Statistically, the analysis of variance for organic carbon concentration, available phosphorus, Exch. Na, Exch. K, Exch. Ca, and BS% were revealed that significant variations vertically as a function of soil depths (P < 0.0001). The overall mean values of OC, Total N, Avail. P, Exch. Na, Exch. K, Exch. Ca, and

BS% concentration under all different land-use systems (natural forest, *Eucalyptus globules*, *Cupressus lusitanica*, *Grevillea robusta*, and *Pinus patula*, grassland, grazing land, and cropland ) across vertical gradients at the top layer were higher than the subsoil layer (Table 3, and Appendix Table 3).

Table 3  
Least Significant Mean values of selected soil chemical properties of different land-use systems across soil depth layers

Soil Depth	OC (%DM)	Total N (%DM)	C:N ratio	Avail. P (mg/kg of soil)	PH-H <sub>2</sub> O (1:2.5)	EC (ds/m)	CEC (meq/100 gm)	Exch. Na (cmol(+)/kg soil)	Exch. K (cmol(+)/kg soil)	Exch. Ca (cmol(+)/kg soil)	Exch. Mg (cmol(+)/kg soil)	B
Top Soil layer (0–20 cm)	2.89	0.25	11.64	16.17	5.80	2.22	27.31	0.39	2.76	14.74	4.19	8
Sub-Soil layer (20–40 cm)	2.02	0.17	11.96	7.96	5.86	2.60	24.82	0.29	1.85	11.96	3.36	7
Sign.	0.0001***	0.0001***	ns	0.0002***	ns	0.0115*	0.0059**	0.0001***	0.0001***	0.0001***	0.0012**	0.

**Keys:** \*, \*\*, \*\*\*, and ns = significant at P < 0.05, high significant at P < 0.01, very high significant at P < 0.001, and nonsignificant at P > 0.05, respectively.

The statistical analysis of variance (ANOVA) for soil electrical conductivity (EC), Cation exchange capacity (CEC), and exchangeable Mg (Exch. Mg) were revealed that significant variations between soil depths (P < 0.05). The overall mean value of soil Cation exchange capacity (CEC), and exchangeable Mg (Exch. Mg) concentration under land-use systems at the topsoil layer were higher than the subsoil layer. However, the analysis of variance for soil pH and C: N ratio was revealed that the nonsignificance variations in between vertical gradients of soil depth at (p < 0.05). The overall mean value of soil pH, electrical conductivity (EC), and C: N ratio concentrations under land-use systems at the top layer were lower than the subsoil layer. In all land-use systems, the level of soil pH, electrical conductivity (EC), and C: N ratio trend between topsoil (0–20 cm) and subsoil (0–40 cm) layers were not uniformly distributed in the area (Table 3, and Appendix Table 3).

## Discussion

### Effect of land-use systems on soil chemical properties

#### Soil Organic C, Total N, C: N ratio and Available P

Organic matter has an important influence on soil chemical properties, soil fertility status, plant nutrition, and biological activity in the soil (Brady and Weil, 2002). Among the different land-use systems, the organic carbon concentration levels were observed in the order of natural forest > *G. robusta* > grassland > *P. patula* > *C. lusitanica* > grazing land > *E. globulus* > cropland. The overall maximum mean value of the organic carbon concentration was observed beneath the natural forest (3.49% DM), and *Grevillea robusta* plantation forest (3.14% DM) statistical without significant differences between them, but significant difference from the rest six land-use systems. This probably due to the higher accumulation of soil organic matters (SOM) by adding litters to soils from different heterogeneity plant species with the high rate of biomass production, and better carbon nutrient release or mineralize to the soils through decomposition. Similar results have been reported by Michelsen et. al. (1996), Lemeniha et. al. (2004), Lemma et. al. (2006) that the higher soil organic carbon concentrations observed in the natural forest than different exotic tree plantation forests in different regions of Ethiopia.

Among the land-use systems in the study area, the minimum value of organic carbon concentration was observed in cropland (1.47% DM) and followed by *Eucalyptus globulus* plantation forest (2.03% DM). The difference could be attributed due to the addition of lower organic matter to the soil via litter inputs and fractions of litter types by losses of organic matter through harvesting of woody and crops biomass from sites, continuous cultivation that aggravates organic matter oxidation lead to losing carbon from the soil in CO<sub>2</sub> form. The results were in agreement with the findings of Negassa (2001), Dhaliwal and Singh (2003), Lemeniha et. al. (2004), Duguma et.al. (2010), Sebatleab (2014), Mengiste et. al. (2015), and Tellen et.al. (2018) who's reported less organic carbon in the cultivated land and *Eucalyptus* plantation forest soils than others land-use systems.

The total nitrogen concentration levels distribution trend horizontal as a function of land-use systems were registered in the order of natural forest > *G. robusta* > grassland > *P. patula* > *C. lusitanica* > grazing land > *E. globulus* > cropland. The highest mean value of total nitrogen content was observed in natural forest (0.31% DM) and followed by the *Grevillea robusta* plantation forest (0.26% DM), statistically with significant differences between them. This may be attributed to the long-term accumulation of above and below-ground organic matter inputs from litterfall, root turn over mineralization by actions of soil microbes, and N fixation by symbiotic in leguminous plant species diversity in natural forest and other soil microorganisms. This argument is supported by the significant strong positive correlation (r = 0.947) between the total nitrogen and organic carbon. Similar studies were also reported as the higher total nitrogen content was observed in the natural forest than other land-use systems in different areas (Michelsen et. al. 1996, Lemeniha et. al. 2004, Lemma et. al. 2006, Sebatleab 2014, Tellen et.al. 2018). On the contrary, the lowest mean value of total nitrogen content was observed in cropland (0.130% DM) and followed by *Eucalyptus globulus* plantation forest (0.173% DM), statistically without significant differences between them. The reasons for the lower total nitrogen contents may be attributed due to the loss of nitrogen either to the atmosphere by evaporation or leached down through the soil by water. Because there is no real store of available nitrates in the soil as nitrates are released from organic matter breakdown or fertilizers unless they are used by plants. Similar studies

reported that lower total nitrogen concentrations were observed under agricultural cropland, and *Eucalyptus* plantation forest than the other land-use systems in different regions of Ethiopia (Michelsen et. al. 1996, Lemeniha et. al. 2004, Duguma et.al. 2010).

The overall mean value of the C: N ratio in all land-use systems were statistically not significant differences between them. The C/N ratio was significantly narrowed from 11.38 in the natural forest soils to 12.08 in the *Pinus patula* plantation forest soils. The lowest value of the C: N ratio was observed in the natural forest. This may be attributed by the factor of increase nitrogen nutrient inputs with a high rate of nitrogen fixation by different leguminous plant species diversity having a relatively higher protein and nutrient contents in the natural forest that leads to the lower C: N ratio, due to their fast decomposition rate and release of nitrogen to the soil. Faster decomposition of leaf litter enhances the transfer of fresh carbon to mineral soil (Polglase et al., 2000). Similarly, different workers have been reported the lower C: N ratio under different native plant species diversity (Yadessa, 1997; Hailu 2000, and Yadesa, et.al. 2010). The highest value of the C: N ratio (12.08) was observed in the *Pinus patula* plantation forest. This may be attributed due to lower nitrogen content in the litters of organic matters which results in a higher C: N ratio, and slow decomposition rate. Plant residue with a low C/N ratio (high nitrogen content) decompose more quickly than plant residue with a high C/N ratio (high carbon content) and do not increase soil organic matter accumulation levels as quickly (Janssen 1996, Bengtsson et al. 2003; Springob and Kirchmann 2003).

Phosphorus (P) is an essential element classified as a macronutrient because of the relatively large amounts of P required by plants. In the study area, the concentration of available phosphorus levels was found in the following order: natural forest > *P. patula* > grassland > grazing land > cropland > *G. robusta* > *E. globulus* > *C. lusitanica* land-use systems. The available phosphorus concentration mean values ranged between 2.45 and 31.52 mg/kg of soil. The overall available phosphorus concentration in natural forest (31.52 mg/kg of soil) was found to be higher than other land-use systems. This might be attributed due to a combination of low nutrient demand by natural forest woody plants as compared with different exotic trees species plantation forests, and better phosphorus nutrient release or mineralize to the soils by different heterogeneity plant species diversity during organic matters decomposition. This argument is supported by the analyzed simple linear correlation relationships between available phosphorus was positively correlated with the soil organic carbon ( $r = 0.547$ ). This result is in agreement with the findings of Michelsen et. al. (1996), Nsabimana, et.al. (2008), Sebatleab (2014) whose observe the higher concentration of available phosphorus in the natural forest than others land-use systems. The lower mean values of available phosphorus concentration were recorded in *Cupressus lusitanica* (2.45), and *Eucalyptus globulus*(3.12) plantation forests statistical without significant differences between them. This may be attributed due to fast-growing tree plantations species were associated with a more intense uptake of nutrients from the soil than slow-growing forest species, and the loss of organic matters by rotational harvesting and took away from the site. Michelsen et al. (1993); and Lisanework & Michelsen (1994) were noted that phosphorus (P) concentration limiting plant growth and leaf litter decomposition under *C. lusitanica* and *E. globulus* plantations in Ethiopia.

## Soil pH, EC (ds/m), and CEC

The present study results showed that soil pH significantly varied with land-use systems. The overall mean value of soil pH level distribution in different land-use systems was observed in the order of grassland > natural forest > grazing land > *P. patula* > *G. robusta* > *E. globulus* > *C. lusitanica* > cropland. Among the land-use systems, the maximum mean value of pH (6.49) was observed in grassland and followed by natural forest (6.20) without significant difference between them. This might be attributable to the higher litter deposition from the aboveground and below ground which gone under decomposition by the action of micro-organisms and subsequent mineralize to releases basic cations to the soil in both land-use systems. This argument is agrees with the simple positive correlation relationships between soil pH with soil organic carbon ( $r = 0.432$ ), cation exchange capacity (CEC) ( $r = 0.649$ ), exchangeable cations of Ca( $r = 0.592$ ), Mg( $r = 0.261$ ), and K( $r = 0.325$ ) in the soils. Similar studies results were reported by Michelsen et al. (1993), Michelsen et. al. (1996), and Nsabimana et al., (2008), for soils of higher pH level under natural forest than others land-use systems. Similarly, the higher soil pH under grassland than other land-use systems was reported by Kaur and Toor (2012).

The lower mean values of soil pH were recorded in cropland, *Cupressus lusitanica* plantation, and *Eucalyptus globulus* plantation forests in the orders of 5.38, 5.41, and 5.48, respectively, statistical without significant differences between them. The main reasons for the lowest value of soil pH in the cropland probably due to poorly managed cultivation; use of chemical fertilizers including urea, DAP, and potash, inappropriate use of ammonium-based fertilizers, intensive use of herbicides such as roundup which contain high amounts of cations that helps to neutralize the negative charges i.e. a higher concentration of H<sup>+</sup> (lower pH) will neutralize the negative charge in soils, and soil erosion. Balesdent et al (2000) and Tejada and Gonzalez (2009) have found that cultivation on farmlands led to the soil acidity increase. Other researchers have also observed the lower soil pH in cultivated farmland than other land-use systems (Emiru and Gebrekidan 2013, Tellen, et.al. 2018). Similarly, the decrease in soil pH under the *Cupressus lusitanica* and *Eucalyptus globulus* plantation forests could be due to the fast-growing exotic tree plantation forests were acidify the soil in nature by accumulating basic cations in the forest biomass, increasing production of organic acids from decomposing litter and by increasing cation leaching. The fast-growing exotic tree species forest that consumes high water for biomass production may increase solute concentrations and the mineralization of organic sulfur in the soil which leads to decrease soil pH. Similar studies were also reported that a decline in soil pH under the fast-growing exotic trees plantation forests (Jobbágy and Jackson, 2003; Mishra et al., 2003, Sanchez et al., 2003, Nsabimana et al. 2008, Ashagire et. al. 2008, and Tellen, et.al., 2018).

In the present finding, the mean values of electrical conductivity (EC) of soil were found numerical between 1.78 and 3.47 deciSiemen per meter (ds/m) ranges. According to the current finding the mean value of electrical conductivity (EC) of soil levels across land-use systems was found in the order of *E. globulus*,> cropland > *G. robusta* > *P. patula* > *C. lusitanica* > grassland > grazing land > natural forest. The maximum mean value of electrical conductivity (EC) was observed in the *Eucalyptus globulus* plantation forest (3.47) and followed by cropland (3.07) without significant differences between them. For the higher value of EC under *Eucalyptus globulus* plantation forest may be attributed due to the lower soil moisture content that related with soluble salts accumulate in the upper soils rather than leached down, high evaporation rate due to open canopy, and low infiltration rate in combine resulting to dissolved salts are left behind to accumulate in the upper soils layer, salts originate from the disintegration (weathering) of minerals and rocks, soils with an accumulation of exchangeable sodium are often characterized by poor tilth and low permeability making high EC. This argument is supported by the positive correlation between the electrical conductivity (EC) and exchangeable sodium, Na ( $r = 0.202$ ), and negatively correlated with organic carbon ( $r = -0.466$ ). This result is in

agreement with the findings of Michelsen et al. (1996). However, for the high level of EC in cropland probably due to tillage intensity and land management practice, cropping system and nature, salt accumulation from commercial fertilizers, chemical contamination (from herbicide, insecticide, and fungicide use by farmers), erosion, runoff, animal manures (usually high tunnels), and compost were contributed to raising EC. Similar studies were reported that higher values of EC in cropland soils than others of land-use systems (Dhaliwal and Singh 2003, Gol 2009, Kaur and Toor, 2012).

The minimum mean value of electrical conductivity was observed in the natural forest (1.783) than the other land-use systems. The lower level of EC under natural forest probably attributed due to the higher accumulation of organic matters (litter deposition) that decomposed and release higher exchangeable cations (K, Ca, Mg) to the soils, which lead to reducing the salinity level and lowering the values of electric conductivity in the soils. This explanation is supported with the negative correlation between electrical conductivity and exchangeable cations K( $r = -0.400$ ), Ca( $r = -0.532$ ), Mg( $r = -0.173$ ), and organic carbon ( $r = -0.466$ ). This finding is in agreement with a similar study report by Michelsen et al. (1996), and Gol (2009) who had reported that the lower mean value of electrical conductivity under natural forest than other land-use systems.

The mean values of CEC in the study area were found between 18.98 and 33.63 meq/100 gm of soil ranges. The overall mean values of soil CEC level distribution in different land-use systems were recorded in the order of natural forest > *P. patula* > grassland > grazing land > *E. globulus* > *G. robusta* > *C. lusitanica* > cropland. Among the land-use systems the highest concentration of cation exchange capacity (CEC) was registered in natural forest (33.63). This is probably influenced by the high amount of organic matter accumulation, and high soil pH in the natural forest soils that lead to higher CEC. This means the CEC of soils is affected mainly by the amount and degree of decomposition of the organic matter. In general, the higher soil organic matter (SOM) is resulting the higher the CEC. Because most of the CEC is originates from the amount of SOM decomposition rate that entirely pH-dependent. This argument is supported by a significant positive correlation between the soils CEC and OC ( $r = 0.644$ ), and pH ( $r = 0.649$ ). The CEC is positively correlated with pH; therefore, acid soils have a lower potential of CEC. The present result is an agreement with the findings of Michelsen et al. (1996), Dhaliwal and Singh (2003), Lemeniha et al. (2004), Nsabimana et al. 2008, and Tellen and Yerima (2018) were observed the highest cation exchange capacity (CEC) values in the natural forest than others of land-use systems in different areas.

The lowest concentration value of cation exchange capacity (CEC) was observed in agricultural cropland (18.98) and followed by the *Cupressus lusitanica* plantation forest (21.26), *Grevillea robusta* plantation forest (23.24), and *Eucalyptus globulus* plantation forest (25.78). The occurrence variation may be attributed due to the low additions of organic matters or litters deposition in the agricultural cropland which results in low liberation of exchange cation nutrients (Ca, Mg, Na, K) to the soil by decomposition. Similar to the present findings, other researchers were reported the lower values of CEC in mechanized farming (MF) by Lemeniha et al. (2004), Selassie et al. (2015), and Molla and Yalew (2018).

## Soil Exchangeable bases (Na, K, Ca, Mg), and Base saturation percent

The main ions associated with CEC in soils are the exchangeable cations calcium ( $\text{Ca}^{2+}$ ), magnesium ( $\text{Mg}^{2+}$ ), sodium ( $\text{Na}^+$ ), and potassium ( $\text{K}^+$ ) are generally referred to as the base cations (Rayment and Higginson 1992). As the function of land-use systems, the concentration of exchangeable cations was generally in the order of  $\text{Ca} > \text{Mg} > \text{K} > \text{Na}$  in all different land-use systems. These agree with the principle stated as the energy of the adsorption sequence of:  $\text{Ca} > \text{Mg} > \text{K} > \text{Na}$ . The highest concentration of exchangeable cations of Ca, Mg, and K, were observed in the natural forest in the order of 17.13, 5.37, and 3.60 cmol(+)/kg soil, respectively. These probably attributed due to the higher accumulation of soil organic matters (SOM) by adding of woody plant litters and understory herbaceous plant residues to soils from different heterogeneity plant species with the high rate of biomass production and those undergone decompositions, thereby, liberate cations nutrients of Ca, Mg, and K to the soils. These results are in agreement with the findings of Michelsen et al. (1993), and Tellen, et al. (2018). Similarly, Nsabimana et al. (2008) were observed the higher concentration of exchangeable cations of Ca, and Mg under mixed native species (MNS) forest than other exotic tree species plantation forests in southern Rwanda.

Additional reasons for the higher concentration of exchangeable cations of Ca, Mg, and K, in the natural forest, maybe due to reduced losses of cations nutrients from the soil by leaching out withholding positively charged ions (cations) by electrostatic force, reduced runoff, and soil erosion via upper surface cover approach with litters, and low cations nutrients demand by natural forest trees species might contribute to the higher of cations nutrients under natural forest. These explanations are supported by the positive correlation between the organic carbon and exchangeable cations in the soils K ( $r = 0.576$ ), Ca ( $r = 0.677$ ) and Mg ( $r = 0.426$ ). The present findings were agreed with Molla and Yalew (2018) and Kokeb et al. (2015). In general, a similar observation was reported by Michelsen et al. (1996) concerned with the higher cations nutrients under natural forest than other exotic tree species plantation forests in the highland of Ethiopia.

The highest concentration of Na was observed in the *Eucalyptus globulus* plantation forest (0.60 cmol(+)/kg soil) than the other land-use systems. The overall mean values of the distribution of exchangeable Na in the area were found in the orders of *Eucalyptus globulus* > natural forest > *Grevillea robusta* > grazing land > *Pinus patula* > grassland > *Cupressus lusitanica* > cropland. For the highest concentration of the exchangeable Na + in *Eucalyptus globulus* plantation forest may be due to the effects of high soil compaction that result with high bulk density, and lower soil moisture content could be facilitated to the soluble salts accumulate in the upper soils rather than leached down. Additionally, the accumulation of exchangeable sodium salts is probably originated from the disintegration (weathering) of minerals and rocks. Similar observations were reported by Michelsen et al. (1993), Michelsen et al. (1996) that the higher exchangeable sodium under *Cupressus lusitanica*, *Eucalyptus globules*, *Eucalyptus grandis*, and *Eucalyptus saligna* plantation forests in the highland of Ethiopia.

The lowest concentration of exchangeable cations of Na, K, and Ca, were recorded in cropland in the order of 0.21, 1.20, and 9.93 cmol(+)/kg soil, respectively. However, the minimum mean value of exchangeable Mg was recorded in the *Cupressus lusitanica* plantation forest (2.91 cmol(+)/kg soil). This could be due to the low addition of organic matters from external factors and the removal of crop residuals from the cropland by harvesting maybe contribute to lower addition of the cations nutrients of Na, K, and Ca to soils. Another explanation for the lowest concentration of exchangeable Na, K, and Ca in cropland probably due to continuous intensive tillage and cropping systems facilitate to lower bulk density, lower CEC, higher porosity, and higher infiltration rate in soils

could lead simply these cations nutrients of Na, K, and Ca were leached out down to the soil depth by water. Similarly, Duguma et. al. (2010), and Molla and Yalew (2018) were observed that the highest concentration of exchangeable Ca in cereal farmland than other land-use systems.

Among the land-use systems, the highest mean value of percent base saturation (80.83) was observed in the *Cupressus lusitanica* plantation forest. This probably due to the amount and nature of clay particles contents and low concentration of CEC, and low pH level in soils could be contributed to the existence of higher percent base saturation (PBS) under *Cupressus lusitanica* plantation forest than others land-use systems. Because of the amount and type of clay minerals are responsible factors for CEC in that both clay and colloidal organic matters (COM). This argument is supported by a significant negative correlation between the soils PBS and CEC ( $r = -0.300$ ) and between PBS and pH ( $r = -0.050$ ). Similarly, Kebede and Charles (2009) were suggested that clay and colloidal organic matters are negatively charged and can act as anions; as a result, these two materials can absorb and hold positively charged ions (cations). These findings are in agreement with a similar study report by Nsabimana, et.al. (2008), who had reported the higher percent base saturation (93.7%) under *Cupressus lusitanica* plantation forest among others plantation forests in southern Rwanda.

The lowest mean value of percent base saturation (70.91) was recorded in the *Eucalyptus globulus* plantation forest. This probably due to the lower addition of organic matters that undergone decomposition and liberate low cations nutrients of Na, K, Ca, and Mg to soils that lead to lower percent base saturation under *Eucalyptus globulus* plantation forest than the others land-use systems. This explanation is supported by the positive correlation between percent base saturation and organic carbon ( $r = 0.262$ ), and a negative correlation between PBS and pH ( $r = -0.050$ ). The percent base saturation (%BS) levels were distributed in similar values without significant difference among the LUS of *Grevillea robusta* plantation, cropland, *Pinus patula* plantation, grazing land, grassland in the order of 79.78, 77.80, 76.35, 74.53, and 73.32, respectively. The variation of the percent base saturation (PBS) means values among different land-use systems may be due to the variation of amount, and nature of organic matters addition, and degree of decomposition to release cation nutrients to soils. Similar observations were reported by Lemeniha et. al. (2004), Nsabimana, et.al. (2008), and Duguma et.al., (2010), who reported different percent base saturation (%BS) levels under different land-use systems.

## Effect of soil depth on soil chemical properties

Concerning to vertical gradient of soil depth the higher mean values of organic carbon (2.89%DM), total nitrogen (0.25%DM), available phosphorus (16.17 mg/kg of soil) concentration were recorded in the upper part of the soil layer at 0–20 cm depth than in the lower subsoil layer at 20–40 cm depth. The organic carbon, total nitrogen, and available phosphorus concentration levels distribution trend across soil depths were decreased gradually from the topsoil layer to the subsoil layer in all land-use systems. This explanation may be due to the higher organic matter inputs to the topsoil layer from plant litters, crop residues, commercial fertilizers, and animal waste materials than the subsoil layer. This is in agreement with previous studies (Michelsen et. al. 1993, 1996, Yadessa and Itanna 2001; Hailu 2000; Lemeniha et. al., 2004, Yadessa et.al., 2010, Kaur and Toor, 2012, Sugihara 2014).

The study results indicated that the lower overall mean values of the C: N ratio (11.64), pH (5.80), and electrical conductivity (EC) (2.22) were observed at topsoil layers than that of the subsoil layers. For the lower value of C: N ratio levels at topsoil is probably due to the highly decomposed organic matter releases higher N- levels on the topsoil than the subsoil layers, thereby, the lowering C: N ratio occurred at the topsoil layer. Similar to this finding, Yadessa and Itanna (2001) and Hailu (2000) also found a lower C/N ratio at the topsoil than in the subsoil layers under different native woody plant species diversity on farmland. The lower values of EC at topsoil are probably due to the addition and accumulation of organic matters at the topsoil surface than subsoil surface that liberates exchangeable cations, thereby, reducing soil EC at the topsoil layer. On the other hand, the pH levels distribution trend in soil was increased gradually from topsoil to subsoil layers in *Eucalyptus globulus* plantation, *Cupressus lusitanica* plantation, *Pinus patula* plantation forests, grazing land, and cropland of land-use systems in the study area. The result of the higher value of soil pH at subsoil than topsoil layers probably due to the leaching of more soluble soil minerals and basic cations from topsoil to the subsoil layer. Similarly, Michelsen et al. (1996) and Zewdie & Olsson, (2008) was reported that the increment of pH values from topsoil to subsoil layers in *Eucalyptus globulus* and, *Pinus patula* plantation forest land-use systems in the Ethiopian Highlands.

With regarding a vertical gradient of soil depth the overall higher mean value of exchangeable cations of Na, K, Ca, Mg, and percent base saturation (%BS) concentrations were observed at the topsoil layers than the subsoil layer with significance difference. The distribution of exchangeable cations of Na, K, Ca, Mg, and percent base saturation (%BS) concentration were decreased vertical from the topsoil layer to the subsoil layer in all land-use systems. These probably due to the effect of higher organic matters depositions or accumulation from litters of woody, herbaceous residuals, animal manures, and crops residuals on farmlands that undergone decomposition and mineralized cations nutrients of Na, K, Ca, and Mg to soils then CEC play the roles to retain the released cations at topsoil from the decomposed organic matter rather than translocating them to the subsoil layer. Cations, such as  $K^+$ ,  $Na^+$ , and  $Ca^{2+}$ , can be adsorbed onto soil or organic colloids, making the cations available for plant uptake by preventing cation leaching from the system (Brady and Weil 2007). Similar studies results have been reported by Michelsen et. al., (1996), Lemeniha et al. (2004), Zewdie and Olsson (2008) that the majority of exchangeable cations nutrient concentrations were declined as soil depth increase, except exchangeable Na in some land-use systems in different areas.

## Conclusion

The historical changes in land-use systems, land-use intensity, and land management practices by the human intervention have altered the characteristics of soil chemical properties across the horizontal as a function of land-use systems, and vertical layers as a function of soil depth. Under similar environmental conditions, the different land-use systems have a strong significant effect on soil chemical properties either positively or negatively. The present study results were showed that different land-use systems and soil depth have different potential in their influence on soil chemical properties dynamics. Among the different land-use systems, the natural forest land-use system was shown a positive effect on most soil chemical properties. In soil chemical properties, the results revealed that the highest mean values of OC, TN, Avail. P, CEC, Exch. Ca, Exch. Mg and Exch. K was observed under natural forest than others of land-use systems in the area. On the other hand, the lowest mean value of soil EC properties was registered under natural forest than other land-use systems. The study results showed that the lowest mean values of OC, TN, soil pH, CEC, and exchangeable bases (Na, K, and Ca) were recorded under cropland than other land-use systems.

Among the five forest land-use systems the lowest mean values of soil OC, TN, Avail. P and BS% were recorded under the *Eucalyptus globulus* plantation forest. However, the highest mean values of EC and Exch. Na was observed under the *Eucalyptus globulus* plantation forest than other land-use systems. In general, the majority of soil chemical properties under the *Eucalyptus globulus* plantation forest were poorer than the soils subjected to other forest land-use systems. The majority of soil chemical properties were significantly differenced as a function of vertical gradients (soil depth) factor. As a function of soil depth factor, the highest concentration of OC, TN, Avail.P, CEC, Exchangeable bases (Na, K, Ca, and Mg), and BS% were observed at the topsoil layer than subsoil layer. However, the highest concentration/ amount of soil pH and EC were recorded at the subsoil layer in the area.

Under similar climatic conditions, among exotic tree plantation forests, *Eucalyptus globulus* plantations forest has a more negative influence on soil chemical properties, which means the majority of soil chemical properties under *Eucalyptus globulus* plantation were poorer than the soils subjected to other land-use systems. The governmental bodies and land-users should be given attention to consider the future positive soils chemical properties change when allocating the land-use to specific land-use systems.

## Abbreviations

Av.P: available phosphorous; BS%: Base Saturation percent; C: N: the ratio of carbon to nitrogen; CEC: cation exchange capacity; CV: Coefficient Variation, DM: dry matter; EC: electrical conductivity; Exch. Ca: Exchangeable Calcium; Exch. K: Exchangeable Potassium; Exch. Mg: Exchangeable Magnesium; Exch. Na: Exchangeable Sodium; FAO: Food and Agricultural Organization of the united nations; JCS: Jaccard's Similarity Coefficient; LSD: Least Significant Mean; LUS: land-use system; m.a.s.l.: Meter above sea level; OC: organic carbon; sign: significant; TN: total nitrogen.

## Declarations

### Ethics approval and consent to participate

Not applicable to this manuscript.

### Consent for publication

All data and information are generated and organized by the authors

### Availability of data and material

We declare that the data and materials presented in this manuscript can be made available as per the editorial policy of the journal.

### Competing interests

The authors declare that they have no competing interests.

### Funding

The research is supported by the Ambo University and Addis Ababa University (AAU) in collaboration for the Ph.D. studies program.

### Authors' contributions

All authors played a vital role to accomplish this manuscript. The corresponding author, YB develop the idea of the research, designed the research method, soil samples collection, soil laboratory results organized, statistical analysis, and wrote the manuscript. TB and SD were contributed significant input into the successful completion of the manuscript by supervising the study, consistent and inspiring guidance, valuable suggestions; constructive comments, and reviews on the manuscript preparation.

### Acknowledgment

We are grateful to Addis Ababa University (AAU) and Ambo University for the financial support of this research. We would like to acknowledge to Oromia Wildlife and Forest Enterprise (OWFE) office, particularly, East Shewa Zone district staff members of the Enterprise. Our special thanks also go to the members of Yerer forest local guards with their bosses for their involvement and assisted in the field data collection. We would like to express our sincere thanks to Hawassa University, Wondo Genet College of Forestry and Natural Resources (WGCF and NRs) for soil laboratory analysis.

## References


1. Aborisade, K.D. & Aweto, A.O. (1990) Effects of exotic tree plantations of teak (*Tectona grandis*) and gmelina (*Gmelina arborea*) on a forest soil in south-western Nigeria. *Soil Use and Management* 6(1): 43–45. <https://doi.org/10.1111/j.1475-2743.1990.tb00796.x>
2. Ashagire T., Tilahun H., Collick AS, Addisu T., Biruk B. et al. (2008) Soil Properties and Fertility Status Dynamics of North-Western Ethiopia as Influenced by Land Use Changes: Case of Dibanke Watershed, Ethiopia. Pp.2-13. <https://www.researchgate.net/publication/258993430>
3. Balesdent, J., Chenu C., and Balabane M. (2000) Relationship of soil organic matter dynamics to physical protection and tillage. *Soil and Till Research* 53(3): 215-230. DOI: [10.1016/S0167-1987\(99\)00107-5](https://doi.org/10.1016/S0167-1987(99)00107-5)
4. Bengtsson G, Bengtson P, Månsson KF. (2003) Gross nitrogen mineralization, immobilization-, and nitrification rates as functions of soil C/N ratio and microbial activity. *Soil Biol. Biochemistry* 35(1): 143-154. DOI: [10.1016/S0038-0717\(02\)00248-1](https://doi.org/10.1016/S0038-0717(02)00248-1)

5. Berthrong, S.T., Jobbery, E.G. & Jackson, R.B. (2009) A global meta-analysis of soil exchangeable cations, pH, carbon, and nitrogen with afforestation. *Ecological Applications* 19(8):2228-2241. <https://doi.org/10.1890/08-1730.1>
6. Black, C.A., Evans D.D., White J.L. Ensminger L.E. Clark F.E. (1965) Method of Soil Analysis. Physical and mineralogical properties, including statistics of measurement and sampling. American Society of Agronomy: Crop Science Society of America: Soil Science Society of America, Inc., Madison, Wisconsin. Series: *Agronomy*, no. 9.
7. Brady NC and Weil RR (2002) The Nature and Properties of Soils, 13th Edition. Prentice Hall, USA.
8. Brady, N.C. and Weil, R.R. (2007) Nature and Properties of Soils, the 15th Edition
9. Chapman. H.D. (1965) Cation Exchange Capacity. P. 891–901. In C.A. Black (ed). Methods of soil analysis. Part. 2. Chemical and Microbiological Properties. American Society of Agronomy and Soil Science Society of America, Inc., Madison, Wisconsin. <https://www.epa.gov/sites/production/files/documents/9081.pdf>
10. Dhaliwal S.S and Singh B. (2003) Effect of different land-use systems on soil quality in the Kandi region of Punjab. Dissertation paper on soil science. Punjab Agricultural University, Indian, Ludhiana.
11. Di Gregorio, A., Jansen, L.J.M. (1998) A new concept for a land covers a classification system. *The Land* 2 (1), 55–65. <https://www.researchgate.net/publication/257312915>
12. Duguma L., and Hager H. (2009) Forest products scarcity perception and response by tree planting in the rural landscapes: farmers' views in central highlands of Ethiopia. *Ekologia Bratislava* 28(2):158–169. DOI: [10.4149/eko\\_2009\\_02\\_158](https://doi.org/10.4149/eko_2009_02_158)
13. Duguma L., Hager H., Sieghardt M. (2010) Effects of land-use types on soil chemical properties in smallholder farmers of central highland Ethiopia. *Ekologia Bratislava* 29(1):1-14. DOI: [10.4149/eko\\_2010\\_01\\_1](https://doi.org/10.4149/eko_2010_01_1)
14. Emiru N. and Gebrekidan H. (2013) Effect of land-use changes and soil depth on Organic Matter, Total nitrogen, and Available phosphorous contents of soils in Senbat Watershed, Western Ethiopia. *American Journal of Agricultural and Biological Science* 8(3):206-212. <https://www.researchgate.net/publication/264859040>.
15. Fantaw Y, Ledin S., Abdu A (2007) Changes in soil organic carbon and total nitrogen contents in three adjacent land use types in the Bale Mountains, south-eastern highlands of Ethiopia. *Forest Ecology and Management* 242(2-3): 337-342. DOI: [10.1016/j.foreco.2007.01.087](https://doi.org/10.1016/j.foreco.2007.01.087)
16. FAO (2000) Global forest resources assessment (FRA) in 2000. Forestry paper 139, Rome, Italy.
17. FAO (Food and Agriculture Organization) (2005) Global forest resources assessment 2005. FAO Forestry paper 140 Rome, Italy.
18. Foley JA, DeFries R., Asner GP, Barford C, Bonan G, Carpenter SR, Chapin FS, Coe MT, Daily GC, Gibbs HK et. al. (2005) Global consequences of land use. *Science* 309 (3534): 570–574. Vol. 309, Issue 5734, pp. 570-574. DOI: [10.1126/science.1111772](https://doi.org/10.1126/science.1111772)
19. Freier, K.P., Glaser, B. and Zech, W (2010) Mathematical modeling of soil carbon turnover in natural Podocarpus forest and Eucalyptus plantation in Ethiopia using compound-specific d13C analysis. *Global Change Biology* 16(5): 1487–1502. <https://doi.org/10.1111/j.1365-2486.2009.02096.x>
20. Gol C. (2009) the effects of land-use change on soil properties and organic carbon at Dagdami river catchment in Turkey. *Journal of Environmental Biology* 30 (5): 825-830. <https://www.researchgate.net/publication/41418129>
21. Hailu T., Legesse N., and Olsson M. (2000) Effects of *Milletia ferruginea* on soil fertility and growth performance of maize. *Agroforestry Systems* 48(1): 9–24. DOI: [10.1023/A:1006274912762](https://doi.org/10.1023/A:1006274912762)
22. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, Thau D, Stehman SV, Goetz SJ, Loveland TR et. al. (2013) High-resolution global maps of 21st-century forest cover change. *Science* 342 (6160): 850–853. DOI: [10.1126/science.1244693](https://doi.org/10.1126/science.1244693).
23. Hegde, R. and Enters, T. 2000. Forest products and household economy: a case study from Mudumalai Wildlife Sanctuary, Southern India. *Environmental Conservation* 27 (03): 250-259. DOI: [10.1017/S037689290000028X](https://doi.org/10.1017/S037689290000028X)
24. Jackson, M. L. (1973). Soil Chemical Analysis. Prentice Hall of India, Delhi.
25. Jaleta, D., Mbilinyi, B., Mahoo, H., Lemenih, M. (2016). Eucalyptus expansion as relieving and provocative tree in Ethiopia. *J. Agric. Ecol. Res. Int.* 6 (3): 1–13. DOI: [10.9734/JAERI/2016/22841](https://doi.org/10.9734/JAERI/2016/22841)
26. Janssen BH (1996) Nitrogen mineralization in relation to C: N ratio and decomposability of organic materials. *Plant Soil* 181(1): 39–45. DOI: [10.1007/BF00011290](https://doi.org/10.1007/BF00011290)
27. Jobbágy EG, Jackson RB (2003) Patterns and mechanisms of soil acidification in the conversion of grasslands to forests. *Biogeochemistry* 64: 205-229.
28. Kaur R and Toor A.S (2012) Effect of Different Agricultural Land Uses on Carbon Sequestration in Soils of Sub-Montaneous Districts of Punjab. M.Sc thesis. Punjab Agricultural University, Ludhiana, Indian. <http://krishikosh.egranth.ac.in/handle/1/5810002276>
29. Kebede F., and Charles Y. (2009) Soil Fertility Status and NumassFertilizer recommendation of TypicHapluusterts in the Northern Highlands of Ethiopia. *World ApplSci J* 6: 1473-1480.
30. Lemeniha M., Mats O., Erik K. (2004) Comparison of soil attributes under *Cupressus lusitanica* and *Eucalyptus saligna* established on abandoned farmlands with continuously cropped farmlands and natural forests in Ethiopia. *Forest Ecology and Management* 195(1-2): 57–67. DOI: [10.1016/j.foreco.2004.02.055](https://doi.org/10.1016/j.foreco.2004.02.055)
31. Lemma B., Berggren, D., Nilsson, I. & Olsson, M. (2006) Soil carbon sequestration under different exotic tree species in the southwestern highlands of Ethiopia. *Geoderma* 136(3):886-898 DOI: [10.1016/j.geoderma.2006.06.008](https://doi.org/10.1016/j.geoderma.2006.06.008)
32. Mengiste W., Mohammed M., and Yitebarek T. (2015) Evaluation of the Effect of Land Use Types on Selected Soil Physico-Chemical Properties in Itang-Kir Area of Gambella Region, Ethiopia. *Journal of Biology, Agriculture, and Healthcare* 5 (11): 92-102. <https://www.researchgate.net/publication/328662919>


33. Michelsen A., Lisanework N., Friis I. and Holst N. (1996) Comparisons of Understorey Vegetation and Soil Fertility in Plantations and Adjacent Natural Forests in the Ethiopian Highlands. *Journal of Applied Ecology*, 33(3): 627-642. <https://www.jstor.com/stable/2404991>
34. Michelsen, A., Lisanwork N & Friis I. (1993) Impacts of tree plantations in the Ethiopian highland on soil fertility shoot and root growth, nutrient utilization, and mycorrhizal colonization. *Forest Ecology and Management* 61(3-4): 299-324. [https://doi.org/10.1016/0378-1127\(93\)90208-5](https://doi.org/10.1016/0378-1127(93)90208-5)
35. Mishra A, Sharma SD, Khan GH. (2003) Improvement in physical and chemical properties of sodic soil by 3, 6, and 9 years old plantation of *Eucalyptus tereticornis* Biorejuvenation of sodic soil. *Forest Ecology and Management* 184(1-3):115-124. DOI: 10.1016/S0378-1127(03)00213-5
36. Molla FE, and Yalew AM. (2018) The effects of land-use types and soil depth on soil properties of Agedit watershed, Northwest. Ethiopia. *Ethiopian Journal of Science and Technology* 11(1):39-56. <https://www.researchgate.net/publication/323221132>
37. Muche M, Kokeb A, Molla E (2015) Assessing the Physicochemical Properties of Soil under Different Land Use Types. *J Environ Anal Toxicol* 5(5): 309. DOI:10.4172/2161-0525.1000309
38. Negassa W. (2001) Assessment of Important Physicochemical Properties of Nitosols under Different Management Systems in Bako Area, Western Ethiopia. M.Sc thesis presented to School of graduate studies, Alemaya University, Ethiopia
39. Newbold T, Hudson LN, Hill S, Contu S (2015) Global effects of land use on local terrestrial biodiversity. *Nature* 520 (7545):45– 50. DOI: 10.1038/nature14324
40. Lisanework N. and Michelsen A. (1994) Litterfall and nutrient release by decomposition in three plantations compared with a natural forest in the Ethiopian highland. *Forest Ecology and Management* 65(2-3): 149-164. [https://doi.org/10.1016/0378-1127\(94\)90166-X](https://doi.org/10.1016/0378-1127(94)90166-X)
41. Nsabimana, D. Klemetson, L. Kaplin, B.A. Wallin G (2008) Soil carbon and nutrient accumulation under forest plantations in southern Rwanda. *African Journal of Environmental Science and Technology* 2 (6):142-149. Available online at <http://www.academicjournals.org/AJest>
42. Olsen, S.R., Cole C.V., Watanabe F.S., Dean, L.A. (1954) Estimation of available Phosphorus in soil by extraction with NaHCO<sub>3</sub>. U.S. Department of Agriculture Circular, U.S. Government Printing Office, Washington, D.C. 939pp.
43. Polglase, P.J., Paul, K.I., Khanna, P.K., Nyakuengama, J.G., O'Connell, A.M., Grove, T.S., Battaglia, M. (2000) Change in soil carbon following afforestation or reforestation: a review of experimental evidence and development of a conceptual framework. Australian Greenhouse Office National Carbon Accounting System Technical Report No.20.
44. Ranst V.E., Verloo M., Demeyer A., and Pauwels J.M. (1999) Manual for the soil chemistry and fertility laboratory: analytical methods for soils and plant equipment, and management of consumables. Belgium, pp. 243-245. <http://hdl.handle.net/1854/LU-113771>
45. Rayment GE, Higginson FR (2012) Electrical Conductivity. In 'Australian New, Comprehensive Soil Chemical Methods Laboratory Handbook. Inkata Press: Melbourne. <https://doi.org/10.1080/00103624.2012.641802>
46. Riebeek H and Simmon R (2011) NASA Earth Observatory: The carbon Cycle, (C). <http://earthobservatory.nasa.gov/Features/CarbonCycle>
47. Roe D., Nelson, F., Sandbrook, C. (eds.) (2009) Community management of natural resources in Africa: Impacts, experiences, and future directions, Natural Resource Issues No. 18, International Institute for Environment and Development, London, UK. <https://www.researchgate.net/publication/235959522>
48. Sanchez PA, Palm CA, Buol SW. (2003) Fertility capability soil classification: a tool to help assess soil quality in the tropics. *Geoderma* 114(3): 157-185. <https://www.researchgate.net/publication/222885577>
49. Sebatleab M. (2014) Impact of land use and land cover change on soil physical and chemical properties: a case study of Era-Hayelom tabias, northern Ethiopia. United Nations University Land Restoration Training Programme [final project] <http://www.unulrt.is/static/fellows/document/Mulugeta2014.pdf>
50. Selassie Y G, Anemut F and Addisu S (2015) The effects of land-use types, management practices, and slope classes on selected soil Physico-chemical properties in Zikre watershed, North-Western Ethiopia. *Environmental Systems Research* 4:3 DOI 10.1186/s40068-015-0027-0
51. Singh, S.P., Rawat, Y.S, and Garkoti, S.C. (1997) Failure of brown oak (*Quercus semicarpifolia*) to regenerate in the Central Himalaya: a case of environmental semi-surprise. *Current Science* 73(4): 371-374. <https://www.researchgate.net/publication/292371111>
52. Springob G, Kirchmann H (2003). Bulk soil C to N ratio as a simple measure of net N mineralization from stabilized soil organic matter in sandy arable soils. *Soil Biol. Biochem* 35(4): 629-632. DOI: 10.1016/S0038-0717(03)00052-X
53. Sugihara S., Shibata M., Mvondo AD, Araki ZS., and Funakawa S (2014) Effect of vegetation on soil C, N, P, and other minerals in Oxisols at the forest-savanna transition zone of central Africa, *Soil Science and Plant Nutrition*, 60:1, 45-59. <https://doi.org/10.1080/00380768.2013.866523>
54. Tejada, M., and Gonzalez J. L. (2008) Influence of two organic amendments on the soil physical properties, soil Losses, sediments, and runoff water quality. *Geoderma* 145 (3-4): 325-334. <https://doi.org/10.1016/j.geoderma.2008.03.020>
55. Tellen, V.A., Yerima, B.P.K. (2018) Effects of land-use change on soil physicochemical properties in selected areas in the North West region of Cameroon. *Environmental Systems Research* 7: 3. <https://doi.org/10.1186/s40068-018-0106-0>
56. Van Dijk A.I.J and Keenan R.J. (2007) Planted forests and water in perspective. *Forest Ecology and Management*, 251(1-2): 1–9.
57. Van Reeuwijk LP (1992) Procedures for soil analysis (3<sup>rd</sup> ed) International Soil Reference and Information Center (ISRIC), Wageningen.
58. Wall A and Hytönen J (2005) Soil fertility of afforested arable land compared to continuously. *Plant Soil* 275: 247–260. <https://doi.org/10.1007/s11104-005-1869-4>
59. Wright S. J (2005) Tropical forests in a changing environment. *Trends in Ecology and Evolution* 20(10):553-560. <https://doi.org/10.1016/j.tree.2005.07.009>
60. Yadesa B, Tesfaye A, and Zebene A (2010) Management of Scattered *Faidherbia albida* and its effects on soil physical and chemical Properties and Chickpea grain yield In the Parkland Agroforestry System of Adama District, East Shoa Zone, Oromia Regional State, Ethiopia. M.Sc thesis in agroforestry, from Hawassa University, Ethiopia.

61. Yadessa A. and Itanna F. (2001) Contribution of indigenous trees to soil properties: The Case of scattered *Cordia africana* Lam. trees in croplands of western Oromia. *Ethiopian Journal of Natural Resources* 3 (2): 245-270. <https://www.researchgate.net/publication/340255635>
62. Zewdie M. (2008) Temporal Changes of Biomass Production, Soil Properties, and Ground Flora in *Eucalyptus globulus* plantations in the Central Highlands of Ethiopia. Doctoral Thesis Swedish University of Agricultural Sciences, Uppsala.

## Figures



**Figure 1**  
 A) Map of Ethiopia, B) Map of East Shewa Zone in Oromia Regional State, C) Map of study area of Yerer forest and Surrounding (developed using ArcGIS 10.4.1 Software). Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.



**Figure 2**

The mean values of soil OC, TN, and avail. P under different land-use systems (LUS)




Figure 3

The mean values of soil pH, EC, and CEC under different land-use systems (LUS)




Figure 4

The mean values of soil exchangeable bases (Na, K, Ca, Mg) under different land-use systems

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalSupplementaryfiles.docx](#)