

Daily Physical Activity and Related Risk Factors in COPD

Ali Albarrati (✉ albarrati@ksu.edu.sa)

King Saud University

NS Gale

Cardiff University

M Munnery

Cardiff Metropolitan University

JR Cockcroft

Cardiff Metropolitan University

Dennis Shale

Cardiff University

Research article

Keywords: BODE index, Breathlessness, Daily activity, Exacerbation, Handgrip strength

Posted Date: November 7th, 2019

DOI: <https://doi.org/10.21203/rs.2.16885/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on March 5th, 2020. See the published version at <https://doi.org/10.1186/s12890-020-1097-y>.

Abstract

Background : Factors associated with reduced daily physical activity (DPA) in patients with COPD are still controversial. Physical inactivity in COPD increases risk of cardiovascular disease, frequent exacerbations, reduced health status, and increased symptoms. We hypothesised that reduced DPA in patients with COPD is independent of traditional risk factors including age and spirometry.

Methods : In this cross-sectional study, DPA (over 7 days) was assessed on 88 community stable patients with COPD and 40 controls free from cardiorespiratory disease. Spirometry, body composition, number of exacerbations, handgrip strength (HGS), modified Medical Research Council (mMRC), arterial stiffness, 6-minute walking distance (6MWD) and BODE index were also determined. Frequent exacerbation was defined as ≥ 2 and non-frequent exacerbation < 2 .

Results : Patients with COPD had reduced DPA and exercise capacity compared with controls similar in age, BMI and gender, $p < 0.001$. Frequent exacerbators had less DPA than infrequent exacerbators and both less than controls, $p < 0.001$. Patients with higher BODE index were less active than those with lower index. Time spent on moderate activity was related to cardiovascular risk factors including arterial stiffness. The DPA in patients was independent of age, gender, spirometry, body composition and HGS, $p > 0.05$. The level of breathlessness was superior to lung function in predicting the level of DPA.

Conclusion : The level of DPA in COPD was independent of traditional risk factors. Breathlessness score is a better predictor of the DPA than lung function and handgrip strength.

Background

Reduced daily physical activity (DPA) is a multifactorial deficit in chronic obstructive pulmonary disease (1). Reduced DPA is a key contributor to cardiovascular morbidity, recurrent hospital admissions related to exacerbations, disease progression and reduced health related quality of life (2). Physical inactivity is also the strongest predictor for all-cause mortality in this population (3). Nevertheless, it is modifiable and studies indicate that engaging in a regular physical activity could reduce the risk of cardiovascular disease, hospitalisation rate and death related to COPD (4, 5). Garcia-Aymerich and colleagues found that participation in physical activity equivalent to walking or cycling for at least 2 hours a week was associated with a 30–40% reduction in the risk of COPD related hospital admission and mortality (5).

A number of studies have tried to identify factors related to declining of physical activity in COPD and yielded controversial results (6-8). Most common risk factors reported in these studies are reduced lung function, musculoskeletal strength, breathlessness score and inflammation. The discrepancies in the results may be attributed to the methodological variations as the patients recruited in these studies were either participating in an inpatient or outpatient rehabilitation program which may not represent typical patients with COPD. Patients with COPD are usually referred to rehabilitation program when they report significant breathlessness and may be physically inactive and deconditioned. This may not reflect the general, less symptomatic COPD population, who may remain active and have a reasonable health status. Moreover, the control subjects recruited in these studies were relatively active and may not be representative of the older population.

Therefore, the aim of this original study was to comprehensively examine risk factors related to decreased DPA in stable community patients with COPD compared with non-COPD controls similar in age and gender.

Methods

Subjects

This is a cross sectional analysis from a large prospective longitudinal study in COPD (The ARCADE study, Clinical Trials No [NCT 01656421](#)) (9). A convenience sample of 93 patients with COPD confirmed with post bronchodilator spirometry and 42 controls, either current or ex-smokers free from respiratory disease wore a physical activity monitor for 7 days. All were

clinically stable, no antibiotics or oral corticosteroids in the previous 4 weeks. Exclusions included inflammatory disease such as rheumatoid arthritis, oral maintenance corticosteroids and long-term oxygen therapy and had not attended pulmonary rehabilitation in the last year. Participants were recruited from respiratory outpatient clinics, pulmonary rehabilitation and smoking cessation referrals, general practice databases and previous participants in respiratory research at Cardiff University. Controls were recruited from previous research database at Cardiff University, smoking cessation clinic and patients' relatives. All subjects gave written informed consent and the study had approval from the South East Wales Research Ethics Committee.

Anthropometry and body composition measurement

In all subjects height and weight was measured barefoot with subjects wearing lightweight clothing and body composition was assessed using a single frequency segmental bioelectrical impedance analyser (BC-418 MA, Tanita Corp., Tokyo, Japan). Body mass index (BMI kg/m²), the fat free mass and fat mass were also determined. Waist and hip circumference were measured with an stretch resistant tape (10).

Lung function and COPD related questionnaires

All subjects performed spirometry (Vitalograph alpha, Bucks, UK) to measure the forced expiratory volume in 1 second (FEV₁), the forced vital capacity (FVC) and FEV₁/FVC ratio. Patients completed the test 10 minutes post 400µg of salbutamol via a spacer device, having been asked to abstain from using their inhalers for at least 6 hours prior to their visit (11).

Breathlessness was scored using the modified Medical Research Council (MRC) dyspnoea scale, and patients reported, the number of exacerbations in the last year (defined as requirement for antibiotic or oral corticosteroid therapy) per year and the number of previously diagnosed comorbidities in patients and comparators were also recorded. Infrequent exacerbators were defined as <2 in the previous year, while frequent exacerbators as ≥2 (12).

Patients completed the St George's Respiratory Questionnaire (SGRQ) and the COPD assessment test, both validated measures of health status and symptoms (13, 14).

Daily Physical Activity Assessment

Participants wore a physical activity monitor (SenseWear™ BodyMedia, Inc. Pittsburgh, PA), left arm throughout the day apart from whilst showering or bathing for seven days. The monitor has been found to be reliable and valid in patients with COPD (24). To avoid seasonal changes, data were collected from the beginning of May until the end of September. The variables chosen for this analysis were the total daily number of steps (Steps); daily time spent in at least moderate physical activity as defined by any activity ≥ 3 METs; and daily mean METs level (METS). Step count was divided to above or below 5000 steps/day as a cut-off value of recommended steps per day (15).

Physical Performance

The 6MWD was carried out once in accordance with a protocol adapted from the American Thoracic Society (ATS) guideline using a 30-m level, straight indoor track (16). Maximal right and left handgrip strength (HGS) was determined twice in each hand and the mean was calculated for each one using a hand dynamometer (T.K.K. 5401 grip-D, Takei, Japan).

BODE Index

BODE index score was calculated by adding the score of each variable (BMI, post bronchodilator FEV₁, mMRC and 6MWD)

Aortic Pulse Wave Velocity

After 10 minutes rest, peripheral systolic and diastolic BP seated and supine was determined (OMRON Corporation, Kyoto, Japan). Aortic PWV and central blood pressures including mean arterial pressure (MAP) measured noninvasively (SphygmoCor; AtCor Medical, Sydney, Australia) (17).

Inflammatory biomarkers

A blood sample was obtained for determination of C - reactive protein (CRP, high sensitivity) and fibrinogen by standard assays (Biochemistry, University Hospital of Wales).

Statistical Analysis

The statistical software package 18.0 (Chicago, Illinois, U.S.A.) was used for all analysis. Data were checked for normality prior to analysis. Data were presented as mean and standard deviation or median and interquartile range (IQR). Comparisons between patients and controls were performed using analysis of variance. Categorical data was analysed using the Chi-square test. Relationships between variables were explored using Pearson's and Spearman correlation coefficients. Multivariate analysis was performed using a stepwise multiple regression model. For all analysis $p < 0.05$ was considered significant.

Results

Eighty-eight patients and forty controls had complete data. Five patients and two controls were removed from the analysis. Two patients and two controls did not wear the monitor for seven days and therefore were excluded. Three patients returned the monitor after 14 days and the batteries of the monitors were dead and we could not retrieve the data after charging the monitors. The patients and controls were similar in age, gender proportion and BMI. The patients had poorer lung function (FEV_1 , FVC and their ratio), lower resting oxygen saturations and greater tobacco exposure than the control group (all $p < 0.001$) (Table 1). The severity of airflow obstruction by GOLD stratification (Table 2). The distribution of the modified MRC breathlessness score was: mMRC 0 $n=11$, mMRC 1 $n=34$, mMRC 2 $n=17$, mMRC 3 $n=18$ and mMRC 4 $n=8$. The patients were also classified according to GOLD quadrants based on CAT score (Table 2).

Measures of daily physical activity

The DPA was lower in the patients, mean (SD), 4159 (2749) steps than controls 6734 (3491) steps, $p < 0.001$, however, there was no difference in DPA for males and females in either group, $p > 0.05$. The DPA was not related to age in either patients or controls. Using the minimum recommended level of DPA of 5000 steps/day, only 31% of the patients had reached the recommended amount of physical activity; while in contrast, 73% of the controls met the minimum amount of the DPA.

	COPD (n= 88)	Comparator (n= 40)	p
Gender Male: female	46:41	19:21	0.117
Age (years)	66 (8)	66 (7)	0.923
Smoking (pack years)	42.2 (24.8)	22.9 (20.8)	0.001
FEV ₁ (L)	1.5 (0.8)	2.6 (0.6)	0.001
FVC (L)	2.4 (0.9)	3.4 (0.9)	0.001
FEV ₁ /FVC	0.52 (0.1)	0.78 (0.05)	0.001
FEV ₁ (%)	56 (17)	106 (17)	0.001
FVC (%)	84 (19)	112 (17)	0.001
Resting O ₂ saturation (%)	96 (2)	98 (1)	0.001
BMI (kg/m ²)	27.5 (5.2)	28.6 (4.9)	0.297
Waist:Hip ratio	0.95 (0.09)	0.89 (0.06)	0.002
FFM (kg)	49.5 (10.5)	50.9 (9.1)	0.486
FM (kg)	24.8 (9.9)	27.2 (10.3)	0.243
FFM/FM	2.3 (1.1)	2.2 (1.1)	0.556
FFMI (kg/m ²)	18.2 (2.4)	18.5 (2.1)	0.589
FMI (kg/m ²)	9.3 (3.9)	10.1 (4.3)	0.3
6MWD (m)	336 (107)	515 (88)	0.001
Mean handgrip (kg)	27.2 (9.6)	29.7 (10.2)	0.209
Fibrinogen (g/l) #	3.8 (1.3)	3.2 (0.6)	0.012
HsCRP (mg/l)#	3.3 (3.1)	1.6 (3.2)	0.023
No. Steps	4096 (2720)	6577 (3350)	0.001
No. Comorbidity	1.3 (1.2)	0.8 (0.8)	0.027

Table 1: Physical and Clinical Characteristics of patients with COPD and Controls.

All data mean (SD) unless otherwise indicated *Median (range) # Geometric mean. Abbreviations: 6MWD = six minute walk distance; BMI = body mass index; FEV1 = forced expiratory volume in 1 second; FVC = forced vital capacity; FFM= fat free mass; FFMI= fat free mass index; FM= fat mass; FMI= fat mass index; HsCRP= high sensitivity C- reactive protein.

Table 2: Distribution of Patients with COPD according to GOLD classifications and Quadrants

GOLD Classification	Patients Number	GOLD Quadrants	Patients Number
GOLD I	9	GOLD A	8
GOLD II	50	GOLD B	24
GOLD III	25	GOLD C	10
GOLD IV	4	GOLD D	46

The DPA related to FEV₁% predicted ($r= 0.39, p<0.001$), oxygen saturation ($r_s= 0.29, p=0.012$) and mMRC dyspnoea score ($r=-0.45, p< 0.001$), in the patients. Only was FEV₁% predicted related to DPA in the controls, $r=0.39, p=0.017$.

Daily physical activity and Disease severity

Level of DPA differed across the GOLD categories, between GOLD 1 and GOLD 4, $p=0.004$ (Figure 1). Patients with an FEV₁>50% had greater level of DPA, lower exacerbation rate, breathlessness score and CAT score than patients with an FEV₁<50%, $p<0.001$ (Table 2). There was an association between level of DPA and BODE index, $p<0.001$. Patients with BODE score of ≥ 6 walked less significantly than patients with a lower score, $p<0.001$ (Figure 2).

Figure.1: Daily physical activity disease severity (GOLD categories).

Daily physical activity and body composition and physical function

There was no relationship between body composition parameters and DPA in the patients or control group. However, in the patients, the time spent undertaking moderate activity was related to BMI, $r=-0.39$, FM, $r=-0.37$, FFM:FM, $r=0.50$, all $p<0.001$. Similar relationships were evident in the control group, all $p<0.05$.

The 6MWD and HGS were less in patients than controls, both $p<0.001$ (Table 1). In the patients, the level of daily of daily physical activity was related to the 6MWD, $r=0.45, p=0.001$, and controls, $r=0.49, p=0.002$, but it was not related to HGS, $p>0.05$. The 6MWD and HGS were all related to each other in the patients, $p<0.01$.

Figure. 2: Daily Physical activity and BODE index<or>6

Daily physical activity and aortic stiffness

In the patients, the time spent undertaking moderate activity was related to aortic stiffness ($r_s=-0.28, p<0.01$), resting oxygen saturation ($r_s=-0.38, p<0.002$). This was not evident in controls. Aortic stiffness was greater in patients, 9.5 (2.2), than controls, 8.2 (1.4), $p<0.01$.

Patient reported outcomes

The level of DPA and the SGRQ total score were related ($r=-0.38$, $p<0.001$), as was the SGRQ activity domain ($r=-0.40$, $p<0.001$). The CAT score was also related to the level DPA ($r=-0.34$, $p=0.003$). The 6MWD also related to the total SGRQ score, $r=-0.63$, and the CAT score $r=-0.54$, (all $p<0.001$).

Of the patients, 33 reported 0-1 exacerbations/ year and 47 reported two or more exacerbations/ year. The level of DPA was related to the frequency of exacerbations $r=-0.34$, $p<0.001$, and frequent exacerbators, 3348 (2221) steps, had lower physical activity of daily living than infrequent exacerbators, 5094 (3027) steps.

Systemic inflammation

Circulating CRP and fibrinogen were greater in patients than controls ($p<0.001$) and both were related to the level of DPA, CRP, $r_s = -0.29$, $p=0.013$, and fibrinogen, $r=-0.30$, $p=0.009$, but were unrelated to the level of DPA in the control group.

Predictive factors for the daily physical activity.

In the patients, a stepwise multivariate regression analysis including DPA as a dependent variable and FEV₁%, mMRC and the number of exacerbations as the independent variables showed that mMRC and number of exacerbations explained 28% of the variability in the DPA with FEV₁% predicted excluded from the analysis, adjusted $R^2=0.28$, $p<0.001$. The mMRC and the number of exacerbations explained 34% and 25% of the reduction in the DPA, respectively.

Discussion

To our knowledge, this is the first study to comprehensively examine risk factors associated with daily physical activity in stable community patients with COPD and controls similar in age and gender. The DPA in COPD was associated with level of breathlessness and frequency of exacerbations, independent of age, lung function, sex, peripheral muscle mass strength.

The GOLD classification was not discriminative enough between patients with moderate and severe impaired daily activity. The level of breathlessness was found to be superior to lung function. This may be explained as a number of studies have found that FEV₁ is less sensitive to describe the complexity of COPD and weakly related to the level of breathlessness (19, 20). Breathlessness is the most common reason for visiting primary care and respiratory physicians regardless the severity of lung function (18). A recent study on patients attending primary care found physical inactivity was associated with breathlessness independent of lung function (21). Perception of breathlessness, which patients experience during performing daily activities, is involved in a complex mechanism beyond respiratory mechanical factors (22). Patients with severe breathlessness often limit their physical and functional activity to minimise dyspnoea and creating a vicious cycle of inactivity, muscle atrophy, deconditioning and which result in greater activity intolerance (23-27). Similar findings reported by other researchers confirming the independent relationship between disease severity and physical activity (8, 28).

Patients with poor prognosis (BODE ≥ 6) had limited physical activity which is a strong predictor of poor outcomes including mortality (29, 30). Similarly, a number of studies reported in a small cohort study in patients with COPD a direct relationship between physical inactivity and increased BODE index (23, 31). This is supported by the concept that reduced physical activity is a strong predictor of morbidity and mortality in patients with COPD (3, 5). Pitta and colleagues found inactivity for just one day doubled the BODE score, which increased the probability of premature mortality (23). BODE index is a multidimensional and appears to have the capacity to stratify patients better than a single marker of airway severity, as FEV₁ cannot reflect the whole picture of COPD.

Physical inactivity increases the risk of cardiovascular disease and its related risk factors including arterial stiffness. Arterial stiffness is a validated marker of CV morbidity and mortality. A few studies have examined the relationship between daily physical activity and arterial stiffness in COPD and showed different results (32, 33). In our study, moderate physical activity was associated with reduced arterial stiffness. Reduced daily physical activity is associated with increased risk of

cardiovascular disease, rehospitalisation and arterial stiffness in COPD (34). Increased arterial stiffness is a well established risk factor in COPD and associated with increased cardiovascular disease in this population (32, 35). Increased arterial stiffness is proposed to be a causative factor of increased cardiovascular disease in COPD, which is reduced by physical activity (36, 37). Indeed, a number of studies found a reduction in aortic stiffness after patients with COPD attended a structured programme of pulmonary rehabilitation (37, 38). A recent study found that patients engaging in any type of moderate physical activity was associated with lower mortality rate after one year compared to inactive patients (39).

Altered body composition is an accepted feature of COPD with several studies reporting reduced musculoskeletal mass and strength (40-43). However, in our study, neither muscle mass nor handgrip muscle strength was associated with reduced physical activity and exercise capacity. Van Gestel and colleagues reported a similar finding to our study and showed that HGS was not associated with DPA and could not be used to predict physical activity (44). A possible explanation is that despite the limitations of lower limb activities by breathlessness, upper limb activities and strength may be maintained (1).

In addition, the larger lower limb mass may be more affected by circulating inflammation, a key feature of COPD. Increased circulatory inflammation may profoundly alter musculoskeletal function, increase muscle oxidative stress and accelerate muscle protein degradation (18, 45). Systemic inflammation (CRP) was elevated in COPD in the present study and was associated with fat mass a potential source of inflammation and impaired physical activity. Increased inflammatory biomarkers i.e. CRP and fibrinogen, which are associated with increased risk of cardiovascular disease an insulin resistance, and their association with reduced DPA has been previously shown in individuals with COPD and community dwellers (46, 47). Increased systemic inflammation is linked to an exacerbation frequency, which is associated with breathlessness and impaired physical activity (46).

Limitations

A key limitation of this study is its cross-sectional nature, and this issue is being addressed in the ongoing longitudinal ARCADE study. We had a limited number of DPA monitors so which curtailed recruitment.

Conclusions

The increased level of breathlessness and frequency of exacerbation are key factors in limiting daily physical activity in patients with COPD independent of loss of muscle mass and strength. The association between reduced DPA and breathlessness suggests that interventions to improve perception of breathlessness may optimise patient outcomes including physical activity. There is a need for long-term behavioural interventions to increase/maintain PA in patients with COPD.

Abbreviations

6MWD = six minute walk distance; BMI = body mass index; FEV1 = forced expiratory volume in 1 second; FVC = forced vital capacity; FFM= fat free mass; FFMI= fat free mass index; FM= fat mass; FMI= fat mass index; HsCRP= high sensitivity C- reactive protein.

Declarations

Ethics approval and consent to participate

All subjects gave written informed consent and the study had approval from the South East Wales Research Ethics Committee, part of the National Research Ethics Service (UK). The ARCADE study, Clinical Trials No NCT 01656421

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due other manuscripts will be published from this data, but are available from the corresponding author on reasonable request.

Competing interests

The authors declare no conflict of interest.

Funding

The ARCADE study was funded by GSK.

Authors' contributions

NG, JC and DS designed the study. AA and MM performed the study and collected data. AA, NG analyzed the data. AA, NG, DS wrote the paper. All authors critically reviewed the paper and approved it.

Acknowledgements

The authors would like to thank primary and secondary care physicians as well as the British Lung foundation for assistance with recruitment.

References

1. Maltais F, Decramer M, Casaburi R, Barreiro E, Burelle Y, Debigaré R, et al. An Official American Thoracic Society/European Respiratory Society Statement: Update on Limb Muscle Dysfunction in Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med*. 2014;189(9):e15-e62.
2. Vestbo J, Hurd SS, Agustí AG, Jones PW, Vogelmeier C, Anzueto A, et al. Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease, GOLD Executive Summary. *Am J Respir Crit Care Med*. 2012.
3. Waschki B, Kirsten A, Holz O, Müller KC, Meyer T, Watz H, et al. Physical activity is the strongest predictor of all-cause mortality in patients with COPD: a prospective cohort study. *Chest*. 2011;140(2):331-42.
4. Blair SN, Cheng Y, Holder JS. Is physical activity or physical fitness more important in defining health benefits? *Med Sci Sports Exerc*. 2001;33(6 Suppl):S379-99; discussion S419-20.
a
5. Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM. Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax*. 2006;61(9):772-8.
6. Singh S, Morgan MD. Activity monitors can detect brisk walking in patients with chronic obstructive pulmonary disease. *J Cardiopulm Rehabil*. 2001;21(3):143-8.
7. Coronado M, Janssens JP, de Muralto B, Terrier P, Schutz Y, Fitting JW. Walking activity measured by accelerometry during respiratory rehabilitation. *J Cardiopulm Rehabil*. 2003;23(5):357-64.
8. Jehn ML, Schmidt-Trucksass A. Association between daily activity and VO₂peak. *Medicine and science in sports and exercise*. 2010;42(6):1229; author reply 30.

9. Gale NS, Albarrati AM, Munnery MM, Munnery IC, Irfan M, Bolton CE, et al. Assessment of Risk in Chronic Airways Disease Evaluation (ARCADE): Protocol and preliminary data. *Chron Respir Dis*. 2014;11(4):199-207.
10. WHO. Waist Circumference and Waist–Hip Ratio: Report of a WHO Expert Consultation 2008 [Available from: http://www.who.int/nutrition/publications/obesity/WHO_report_waistcircumference_and_waisthip_ratio/en/index.html].
11. Global Initiative for Chronic Obstructive Lung Disease G. Global Strategy for the Diagnosis, Management and Prevention of COPD, Global Initiative for Chronic Obstructive Lung Disease (GOLD). [Internet]. 2015. Available from: <http://www.goldcopd.org>.
12. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*. 2007;176(6):532-55.
13. Jones PW, Harding G, Berry P, Wiklund I, Chen WH, Kline Leidy N. Development and first validation of the COPD Assessment Test. *Eur Respir J*. 2009;34(3):648-54.
14. Jones PW, Quirk FH, Baveystock CM, Littlejohns P. A self-complete measure of health status for chronic airflow limitation. The St. George's Respiratory Questionnaire. *Am Rev Respir Dis*. 1992;145(6):1321-7.
15. Martin SB, Morrow JR, Jr., Jackson AW, Dunn AL. Variables related to meeting the CDC/ACSM physical activity guidelines. *Med Sci Sports Exerc*. 2000;32(12):2087-92.
16. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med*. 2002;166(1):111-7.
17. Laurent S, Cockcroft J, Van Bortel L, Boutouyrie P, Giannattasio C, Hayoz D, et al. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *European heart journal*. 2006;27(21):2588-605.
18. O'Donnell DE, Ora J, Webb KA, Laveneziana P, Jensen D. Mechanisms of activity-related dyspnea in pulmonary diseases. *Respiratory physiology & neurobiology*. 2009;167(1):116-32.
19. Cote CG, Celli BR. BODE index: a new tool to stage and monitor progression of chronic obstructive pulmonary disease. *Pneumonol Alergol Pol*. 2009;77(3):305-13.
20. Cote CG, Dordelly LJ, Celli BR. Impact of COPD exacerbations on patient-centered outcomes. *Chest*. 2007;131(3):696-704.
21. Kelly JL, Elkin SL, Fluxman J, Polkey MI, Soljak MA, Hopkinson NS. Breathlessness and skeletal muscle weakness in patients undergoing lung health screening in primary care. *Copd*. 2013;10(1):40-54.
22. Gifford AH, Mahler DA, Waterman LA, Ward J, Kraemer WJ, Kupchak BR, et al. Neuromodulatory effect of endogenous opioids on the intensity and unpleasantness of breathlessness during resistive load breathing in COPD. *COPD*. 2011;8(3):160-6.
23. Pitta F, Troosters T, Probst VS, Lucas S, Decramer M, Gosselink R. Potential consequences for stable chronic obstructive pulmonary disease patients who do not get the recommended minimum daily amount of physical activity. *J Bras Pneumol*. 2006;32(4):301-8.
24. Pitta F, Troosters T, Probst VS, Spruit MA, Decramer M, Gosselink R. Quantifying physical activity in daily life with questionnaires and motion sensors in COPD. *The European respiratory journal*. 2006;27(5):1040-55.
25. Seymour JM, Ward K, Sidhu PS, Puthuchery Z, Steier J, Jolley CJ, et al. Ultrasound measurement of rectus femoris cross-sectional area and the relationship with quadriceps strength in COPD. *Thorax*. 2009;64(5):418-23.
26. Singer J, Yelin EH, Katz PP, Sanchez G, Iribarren C, Eisner MD, et al. Respiratory and skeletal muscle strength in chronic obstructive pulmonary disease: impact on exercise capacity and lower extremity function. *Journal of cardiopulmonary rehabilitation and prevention*. 2011;31(2):111-9.
27. Wagner PD. Skeletal muscles in chronic obstructive pulmonary disease: deconditioning, or myopathy? *Respirology*. 2006;11(6):681-6.
28. Troosters T, Sciruba F, Battaglia S, Langer D, Valluri SR, Martino L, et al. Physical inactivity in patients with COPD, a controlled multi-center pilot-study. *Respiratory medicine*. 2010;104(7):1005-11.

29. Waschki B, Kirsten AM, Holz O, Mueller KC, Schaper M, Sack AL, et al. Disease Progression and Changes in Physical Activity in Patients with Chronic Obstructive Pulmonary Disease. *American journal of respiratory and critical care medicine*. 2015;192(3):295-306.
30. Celli BR, Cote CG, Marin JM, Casanova C, Montes de Oca M, Mendez RA, et al. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *The New England journal of medicine*. 2004;350(10):1005-12.
31. Moy ML, Matthes K, Stolzmann K, Reilly J, Garshick E. Free-living physical activity in COPD: assessment with accelerometer and activity checklist. *Journal of rehabilitation research and development*. 2009;46(2):277-86.
32. Sievi NA, Franzen D, Kohler M, Clarenbach CF. Physical inactivity and arterial stiffness in COPD. *International journal of chronic obstructive pulmonary disease*. 2015;10:1891-7.
33. Stickland MK, Vogan N, Petersen SR, Wong EY, Bhutani M. Physical activity and arterial stiffness in chronic obstructive pulmonary disease. *Respiratory physiology & neurobiology*. 2013;189(1):188-94.
34. Watz H, Waschki B, Boehme C, Claussen M, Meyer T, Magnussen H. Extrapulmonary effects of chronic obstructive pulmonary disease on physical activity: a cross-sectional study. *American journal of respiratory and critical care medicine*. 2008;177(7):743-51.
35. Sabit R, Bolton CE, Edwards PH, Pettit RJ, Evans WD, McEnery CM, et al. Arterial stiffness and osteoporosis in chronic obstructive pulmonary disease. *American journal of respiratory and critical care medicine*. 2007;175(12):1259-65.
36. Macnee W, Maclay J, McAllister D. Cardiovascular injury and repair in chronic obstructive pulmonary disease. *Proceedings of the American Thoracic Society*. 2008;5(8):824-33.
37. Gale NS, Duckers JM, Enright S, Cockcroft JR, Shale DJ, Bolton CE. Does pulmonary rehabilitation address cardiovascular risk factors in patients with COPD? *BMC pulmonary medicine*. 2011;11:20.
38. Vanfleteren LE, Spruit MA, Groenen MT, Bruijnzeel PL, Taib Z, Rutten EP, et al. Arterial stiffness in patients with COPD: the role of systemic inflammation and the effects of pulmonary rehabilitation. *The European respiratory journal*. 2014;43(5):1306-15.
39. Moy ML, Gould MK, Liu I-LA, Lee JS, Nguyen HQ. Physical activity assessed in routine care predicts mortality after a COPD hospitalisation. *ERJ Open Research*. 2016;2(1).
40. Agusti A. Physical activity and chronic obstructive pulmonary disease. *American journal of respiratory and critical care medicine*. 2008;177(7):675-6.
41. Ansari K, Keaney N, Taylor I, Burns G, Farrow M. Muscle weakness, health status and frequency of exacerbations in chronic obstructive pulmonary disease. *Postgrad Med J*. 2012;88(1041):372-6.
42. Cesari M, Pedone C, Chiurco D, Cortese L, Conte ME, Scarlata S, et al. Physical performance, sarcopenia and respiratory function in older patients with chronic obstructive pulmonary disease. *Age Ageing*. 2012;41(2):237-41.
43. Eliason G, Abdel-Halim S, Arvidsson B, Kadi F, Piehl-Aulin K. Physical performance and muscular characteristics in different stages of COPD. *Scand J Med Sci Sports*. 2009;19(6):865-70.
44. van Gestel AJ, Clarenbach CF, Stowhas AC, Rossi VA, Sievi NA, Camen G, et al. Predicting daily physical activity in patients with chronic obstructive pulmonary disease. *PloS one*. 2012;7(11):e48081.
45. Reid WD, Rurak J, Harris RL. Skeletal muscle response to inflammation—lessons for chronic obstructive pulmonary disease. *Crit Care Med*. 2009;37(10 Suppl):S372-83.
46. Watz H, Waschki B, Meyer T, Magnussen H. Physical activity in patients with COPD. *The European respiratory journal*. 2009;33(2):262-72.
47. Geffken DF, Cushman M, Burke GL, Polak JF, Sakkinen PA, Tracy RP. Association between physical activity and markers of inflammation in a healthy elderly population. *American journal of epidemiology*. 2001;153(3):242-50.

Figures




Figure.1: Daily physical activity disease severity (GOLD categories).

Figure 1

Daily physical activity disease severity (GOLD categories).




Figure 2

