

Analysis of transcriptomic changes in bovine endometrial stromal cells treated with lipopolysaccharide

Xuefen Ding

Henan Agricultural University

Haimiao Lv

Henan Agricultural University

Lixin Deng

Henan Agricultural University

Wenju Hu

Henan Radio and Television University

Zhan Peng

Henan Agricultural University

Chenbo Yan

Henan Agricultural University

Dexin Yang

Henan Agricultural University

Chao Tong

Henan Agricultural University

Xin Zhuang wang (✉ happywang169@163.com)

Henan Agricultural University <https://orcid.org/0000-0002-2410-9831>

Research article

Keywords: Whole-genome sequencing, Endometritis, Bovine endometria stromal cells, Lipopolysaccharide

Posted Date: January 22nd, 2020

DOI: <https://doi.org/10.21203/rs.2.16904/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Endometritis adversely affects the ability of cattle to reproduce, and significantly reduces milk production. Consequently, it has great influence on the economic value of dairy cows. The endometrium is mainly composed of epithelial and stromal cells and they produce the first immune response to invading pathogens. Epithelial cells are the first cellular barrier through which bacteria enter the uterine endometrium. However, most of the epithelial cells are disrupted and stromal cells are exposed to an inflammatory environment when endometritis occurs, especially postpartum. A loss of the protective epithelium allows bacteria or toxins to access the underlying stromal cells. The activation of Toll-like receptor (TLRs) on stromal cells induces increased production of cytokines and chemokines. Understanding the genome-wide characterization of the bovine endometritis will be beneficial for prevention and treatment of endometritis. In this study, whole-transcriptomic gene changes in bovine stromal cells treated with LPS were compared with those treated with PBS (control group) and were analyzed by RNA sequencing (RNA-seq).

Results: Compared with the control group, a total of 366 differentially expressed genes (DEGs) were identified in LPS-induced group (234 upregulated and 132 downregulated genes), with an adjusted P -value ≤ 0.05 by DESeq. Gene ontology (GO) enrichment analysis revealed DEGs were most enriched in lymphocyte activation, interleukin-1 receptor binding, regulation of cell activation, and lymphocyte-activated interleukin-12 production. Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis revealed DEGs were most enriched in TNF signaling pathway, Toll-like receptor signaling pathway, cytokine-cytokine receptor interaction, nucleotide-binding oligomerization domain-like (NOD-like) receptor signaling pathway, NF- κ B signaling pathway, and chemokine signaling pathway.

Conclusion: The results of this study unraveled endometrial stromal cells transcriptome profile alterations in bovine affected by LPS which may have a significant effect on the eliminating or reducing inflammation by comprehending molecular mechanisms and authenticating unique genes related to endometritis.

Background

Endometritis is caused by a highly inflammatory environment created by pathogen-associated molecular patterns (PAMPs), resulting in subclinical or clinical disease [1, 2]. Endometritis severely affects the ability of cattle to reproduce, compromises animal welfare, and significantly reduces milk production [3-5]. As a consequence, it has great influence on the economic value of dairy cows [2]. Bacterial infection is the most prevalent element of bovine endometritis, especially, *Escherichia coli*. Once endometrium is infected with *E. coli*, it promotes invasion of other bacteria into the uterus after few days of calving [6].

Lipopolysaccharide (LPS) is a vital constituent of the outer membrane of gram-negative bacteria which can mimic characteristics of an actual gram-negative bacterial infection [7]. Previous studies have reported expression changes in peroxisome proliferator-activated receptors (PPARs) after intrauterine injection of *E. coli* suspension *in vivo* and in endometrial explants with exposure to LPS *in vitro*. Their

findings indicate that LPS exposure can mimic inflammation *in vitro*, which is useful for understanding intracellular mechanisms during endometritis, and specific PPAR ligands could potentially be used to cure endometritis [8], such as PPAR α and PPAR δ [9, 10].

The endometrium is mainly composed of epithelial cells and stromal cells and they produce the first immune response to the invading pathogens [11]. Epithelial cells of the endometrium are a physical barrier between the uterine lumen and the stromal cells [12]. However, after parturition the epithelial layer is damaged and invading pathogens access the stroma [13-16]. The epithelium is disrupted by trauma during parturition, and exfoliation of the surface of the uterine caruncles about 12 to 14 days postpartum [14, 15], followed by cell proliferation and regeneration of the epithelium, which is completed between 20 and 30 days after parturition [14, 15]. However, many kinds of bacteria invade endometrium only in the first weeks after parturition. There are commonly *Escherichia coli*, *Trueperella pyogenes* and Grampositive anaerobic bacteria in diseased animals [2, 17, 18], which ahead of regeneration of the epithelial cells. *T. pyogenes* in the endometrium got to the highest concentration after second and third week postpartum [19]. Once the stroma is colonized with bacteria, *T. pyogenes* infection and endometritis appear to persist [16, 17]. Exactly, infection related tissue damage may hamper regeneration of the epithelium and result in a chronic infection. The ability of *T. pyogenes* to cause pathology in the denuded endometrium is similar to the discovery of pathogenic *E. coli* adapted to the bovine endometrium [18]. In addition, previous study showed also LPS treated bovine endometrial explants *in vitro*, which resulted in characteristic histopathologic changes, including luminal epithelial cell shedding and damage. The epithelial cells and endometrial gland layers were completely detached and disappeared. A loss of the protective epithelium allows bacteria or toxins to access the underlying stromal cells. The activation of TLRs on stromal cells induces increased production of cytokines and chemokines [20]. The stromal cells are more sensitive than epithelial cells to pyolysin-mediated cytolysis [21] On the other hand, the previous reports also pointed out that stromal cells are more accessible to the vascular system and mononuclear cells, thus, the effect of stromal cytokines and chemokines is more significant [5, 22, 23]. Moreover, stromal cells outnumber epithelial cells in the endometrium and secrete soluble growth factors as much as epithelial cells in order to increase immunological function of uterus endometrium [14, 22-24]. So stromal cells play a significant role in initiating bovine endometritis response.

There are lots of study about bovine endometrial stromal cells. Such as some study was to establish and characterize an *in vitro* model of endometrial inflammation using bovine endometrial epithelial (bEEL) and stromal (bCSC) cell lines. The researchers evaluated the effects of the LPS and pro-inflammatory mediators (IL-1 β and TNF α) on bEEL and bCSC cells by detecting cytokine gene expression (such as CXCL8/IL8, IL1A, IL1B, and IL6) and eicosanoid biosynthesis pathway gene expression and production. Result showed bEEL and bCSC lines could mimic the physiological responses in mammals with endometrial infection, making them brilliant *in vitro* models for intrauterine environment studies [25, 26]. Other study used primary isolated endometrial epithelial and stromal cells and *ex vivo* organ culture (EVOCs) as well-characterized models of endometritis to explore cellular pathways that modulate inflammatory responses within the endometrial tissue. Result showed that manipulating the mevalonate pathway modulates innate immunity in endometrial cells and tissue and using topical administration of

farnesyl-diphosphate farnesyl transferase (FDFT1) inhibitors into the uterus to limit the severity of bacterial endometritis [27]. On the other hand, in order to understand the recognition of viruses by endometrial epithelial and stromal cells and the induction of pro-inflammatory mediators and interferon (IFNs), study explored the role of TLR3 and retinoic acid-inducible gene I (RIG-I) in the innate immune response to bovine herpesvirus-4 (BoHV-4) and bovine viral diarrhea virus (BVDV), and viral pathogen-associated molecular patterns (PAMPs). Result showed that endometrial stromal cells responded to viable viruses by inducing inflammatory cytokines and chemokines, and bovine endometrial cells could check and respond to virus, and their PAMPs, through TLR3 and RIG-I [28].

Studies to date of the endometrial immune response have largely been based on a candidate gene approach. Genome-wide characterization is crucial for understanding development and pathophysiology of bovine endometritis, which will be beneficial for prevention and treatment of endometritis. Bovine microarray were used for research whole-transcriptomic gene in primary cultures of mixed bovine epithelial and stromal endometrial cells that were treated with LPS for 6 h [29]. Nonetheless, array technology has limitation of exactitude of the expression measurements [30]. High-throughput sequencing technologies provide a platform for insights into genomic characterization. Specifically, high-throughput mRNA sequencing (RNA-seq) enables concurrent transcript discovery and abundance estimation [31]. Previous studies on RNA-seq analysis in mixed bovine epithelial and stromal endometrial cells treated with LPS showed 20 differentially-expressed miRNAs and 108 DEGS, as well as enriched 118 GO and 66 KEGG pathways. These results provided a understanding which effect of miRNA in bovine endometritis [32]. In addition, a previous study on RNA-seq analysis of change in miRNA and mRNA expression in 15 cows at 7 and 21 days postpartum showed 4197 DEGs in healthy cows, while only 31 genes were differentially expressed in cows with cytological endometritis from proinflammatory to hyperplasia stages [33]. Other studies have been performed on RNA-seq analysis of effects of LPS on whole transcriptome of bovine endometrial epithelial cells with focus on genes involved in embryo-maternal interactions. They identified 2,035 and 2,073 DEGs in control cells and cells treated with 2 and 8 µg/mL LPS for 24h, respectively [34]. However, comprehensive transcriptomic analysis of response of bovine endometrial stromal cells to LPS by RNA-seq remains to be elucidated. However, comprehensive transcriptomic analysis of response of bovine endometrial stromal cells to LPS by RNA-seq remains to be elucidated.

In this study, genome-wide gene expression profiles in primary bovine endometrial stromal cells treated with LPS were examined using RNA-Seq, and the results were confirmed by real-time quantitative reverse-transcriptase polymerase chain reaction (qRT-PCR). Based on these data, the functions and pathways involved in the bovine endometritis, especially postpartum in the stromal cells were characterized. The bioinformatics analysis confirmed that some of the expressed genes in our study have been shown to play important roles in bovine endometritis, especially postpartum, thus supporting the reliability of our data. These genes and pathways may be involved in the pathological process of bovine endometritis, especially postpartum and deserve further study and discussion.

Results

Immunofluorescence analysis

Stromal cells exhibited flat, fibroblast-like morphology (Additional file 1: Figure S1). Endometrial stromal cells were estimated by immunostaining for vimentin (stromal cells marker). Stromal cells expression was vimentin-positive and cytokeratin-18 negative (Additional file 2: Figure S2). Cells morphology and the positive staining for vimentin confirmed that the cultured cells were indeed stromal cells.

Quality control of the RNA sequence

RNA quality was fine (Additional file 3: Figure S3). We established three RNA libraries for control group and three RNA libraries for LPS-induced group. After filtering the low-quality reads, the average number of clean reads was 42,381,010 (92.67%) and 44,167,270 (92.77%) for the control group and LPS-induced group, respectively (Table 1). The clean reads were used for the following analyses and most of them (> 93.99%) were mapped to the *Bos taurus* genome (Table 2).

Identification of the source of variance in the expressed transcripts by principal component analysis (PCA)

The purpose of PCA analysis is to account for the origin of variance in our data. We performed PCA analysis with two elements: PC1 and PC2. PC1 was 55% and PC2 was 42% (Additional file 4: Figure S4). PCA analysis can cluster similar samples together, the closer the distance, the higher the similarity between samples. This indicated that these data could be used further.

Differential gene expression

There were a total of 366 DEGs (composed of 234 upregulated and 132 downregulated DEGs) identified in LPS-induced group compared to control group. Some of these genes are related to inflammation, such as inflammatory cytokines including *IL1A*, *IL2RB*, and *IL6*, increased 3.16, 20.53, 15.88 fold, respectively. Chemokines including *CCL2*, *CCL5*, *CXCL8*, and *CXCL2*, increased 16.91, 10.48, 3.39, 2.14 fold respectively. Antimicrobial factor *DEFB10* increased 2.79 fold (Figure 1 and Additional file 5: Table S1)

The heatmap of DEGs

Heatmap analysis of DEGs identified genes of expression levels with high correlation among samples. Some of these genes may be involved in biological processes, metabolic processes, or signaling pathways (Figure 2).

GO enrichment of DEGs

GO enrichment was used to characterize DEGs. DEGs were enriched in immunity-related GO terms, such as leucocyte-mediated immunity, intracellular signal transduction, IL-1 receptor binding, *IL-1 β* secretion, and response to cytokines. There were 638 GO terms, including 574 biological process (BP), 21 cellular

component (CC), and 43 molecular function (MF). GO enrichment results for all DEGs are illustrated (Additional file 6, Table S2), and the top 20 GO terms with the most significant enrichment were selected (Figure 3).

KEGG enrichment analysis

The purpose of KEGG enrichment analysis is to appraise pathways that are vital in inflammation in endometrial stromal cells. The results of the analysis showed that inflammation pathways in endometrial stromal cells are mainly centered on TNF signaling pathway, complement and coagulation cascades, cytokine-cytokine receptor interaction, *IL-17* signaling pathway, chemokine signaling pathway, nucleotide-binding oligomerization domain-like (NOD-like) receptor signaling pathway, NF- κ B signaling pathway, and so on. KEGG enrichment results of DEGs are illustrated (Additional file 7, Table S3). The top 20 KEGG terms with the most significant enrichment were selected (Figure 4).

Confirmation of DEGs by quantitative reverse-transcription PCR (qRT-PCR)

We selected ten DEGs which included seven upregulated genes, *NFKBIZ*, *NFKBIA*, *CCL2*, *C3*, *CSF3*, *GRO1*, *STEAP4* and three downregulated gene, *L1CAM*, *CD8A*, *SBSPON*, and confirmed them by real time qRT-PCR (as shown in Figure 5). The expression level of the ten DEGs were found to be similar with RNA-seq results.

Discussion

In this study, we described whole transcriptomic genes changes in bovine endometrial stromal cells treated with LPS compared with control group by RNA-seq. Stromal cells are exposed to an inflammatory environment when epithelial cells are disrupted (Figure 6), inflammation on endometrial stromal cells is activated (Figure 7) [1, 35]. Results indicate that more than 97% of clean tags of RNA-seq results mapped to the genome and PCA was low, which proved our data could be used for functional analyses.

In the present study, we chose passage 6 bovine endometrial stromal cells. Cultures at the fourth and sixth passage were semiconfluent and exhibited the expected interstitial Cajal-like cells morphology and cellular characteristics [36]. Previous study using P5 bovine endometrial epithelial cells to investigate effects of LPS on whole transcriptome with a special focus on genes involved in embryo-maternal interactions [34]. And transcriptome profiling analysis of on P6 bovine vaginal epithelial cell response to an isolated lactobacillus strain [37]. Mammalian two-hybrid assay to determine whether transiently transfected androgen receptors and Hic-5/ARA55 interact in P6 beard dermal papilla cells with transfected androgen receptor in beard dermal papilla cells was confirmed with mammalian two-hybrid assays [38]. Previous study also proved characteristics of bEEL and bCSC lines could mimic the physiological responses found in mammals with endometrial infection, making them excellent in vitro models for intrauterine environment studies [26]. In addition, previous work has showed the negative effects of cell passaging on mRNA expression and the secretion rate of some pro-inflammatory factors in human gingival fibroblasts [39]. However, some study indicated that the mRNA expression of some genes

in primary bovine oviductal epithelial cells (BOEC) is different from P0 to P3 *in vitro*. Such as the transcription rate of MUC4 was influenced by the number of cell culture passages, and encoded proteins of IL8 differed between distinct cell culture passage numbers. However, there are also mRNA expression of some genes not affected by the number of cell culture passages, such as *OXCT2*, *PGE2*, *PTGS2* and *IL8* mRNA. Surprisingly, the mRNA of *HK1* was twofold more highly expressed in oviductal cells in P0 than in BOEC *in vivo* situation. But BOEC in other cell culture passages showed similar HK1 mRNA expression compared with cells *in vivo* state [40]. Likewise, BOEC incubated with *T. pyogenes* showed not only different mRNA expression of some genes, but also similar mRNA expression in P0 and P3 compared with the controls (without *T. pyogenes*). the latter including *IL1A*, *IL1B*, *TNFA*, *IL8*, *CXCL3* and *CXCL5* [41]. On the other hand, in order to explore the transcriptional changes of cell lines caused by long period *in vitro* selection, study performed a gene-expression meta-analysis and compared 60 tumor cell lines from nine tissue origins to 135 human tissue and 176 tumor tissue samples. Result showed statistically significant differential expression of approximately 30% of the about 7,000 genes investigated compared to the tissues by significance analysis of microarrays [42]. We define bovine endometrial stromal cells passage from P1 to P6. Cell of P1 to P3 were pure, but number were not enough. So we think P6 cell expression profile is not exactly the same as P1, but P6 cells still represent most of the characteristics of primary stromal cells.

LPS binds to lipoprotein binding protein (LBP), which presents it to CD14 receptor. This induces CD14 to present LPS-LBP complex to MD-2 (myeloid differentiation factor 2) [43, 44]. This, in turn, promotes dimerization of Toll-like receptor 4 (TLR4)/MD-2, activating two downstream signaling pathways including MyD88-dependent pathway (MyD88-) and TRIF-dependent pathway. The former triggers NF- κ B and mitogen-activated protein kinase (MAP kinase) signaling and induces inflammatory cytokines, and the latter causes the induction of type I interferons (IFN) through interferon regulatory factor 3 (IRF3) activation and inflammatory cytokines through NF- κ B activation [43]. Our data presents evidence for the activation of the MyD88-independent pathway as LPS upregulated inflammatory cytokines, such as *IL6*, *IL1A*, *CXCL8* and activated NF- κ B pathway (Additional file 2: Table S1 and Additional file 4: Table S3). Bovine endometrial epithelial and stromal cells express the TLR4/CD14/MD2 receptor complex. We found increased expression levels of *CD14* but constant levels of TLR4 or MyD88 in our study in which bovine stromal cells were treated for 12 h with 0.5 μ g/mL LPS (Additional file 3: Table S1), consistent with previous studies that exposed mixed bovine epithelial and stromal endometrial cells to 100 ng/ml LPS for 6 h [45]. In addition, LPS upregulated TLR4 only in bovine endometrial epithelial cells which were treated with 1 μ g/mL LPS for 6 h and 12h, and the expression was greatest at 12h [46].

After the membrane surface receptor is recognized, intracellular inflammation is recognized and intracellular signaling cascades are activated by LPS, which is mediated by innate pattern recognition receptors (PRR) including TLR, retinoic acid-inducible gene-I-like (RIG-I-like) receptors, NOD-like receptors, and C-type lectin receptors [47]. These lead to the expression of inflammatory mediators and contribute to the clearance of pathogens [48]. Our data further confirm these findings because LPS triggers NOD-like receptor signaling pathway, TLR signaling pathway, and C-type lectin receptor signaling pathway in bovine endometrial stromal cells (Additional file 5, Table S3). In addition, our RNA-seq results also

demonstrate the upregulation of expression of inflammatory and chemotactic cytokines related genes, such as *IL6*, *IL1A*, *IL2RB*, *CCL2*, *CCL5*, *CXCL5*, *CXCL8*, *CXCL2*, *CX3CL1*, and *CCL20* (Additional file 3: Table S1). This is also consistent with previous studies that mRNA of proinflammatory cytokines *IL1A*, *IL1B*, *IL6*, *TNF*, and expression of chemokines *IL8* and *CXCL5* increased in endometrial epithelial cells during the estrous cycle and subclinical or clinical endometritis [49, 50]. The expression of these cytokines was also upregulated in only epithelial cells, only stromal cells, and mixed epithelial and stromal cells treated with LPS *in vitro* [45, 51, 52]. Similarly, endometrial stromal cells responded to viable viruses by inducing inflammatory cytokines and chemokines [28]. Antiviral defense of endometrial stromal cells requires coordinated recognition of PAMPs, initially via TLR3 and later via inducible RIG-I [28]. On the other hand, activation of PRR also triggers many signal transduction pathways through one or more of the IRF family of transcription factors, leading to the expression of IFNs [53]. LPS upregulated the expression IFN-stimulated genes (ISG) including *RSAD2*, *MX2*, *OAS1Y*, *ISG15*, and *BST2* in mixed epithelial and stromal cells [45]. In the present study, our results show that LPS upregulated the expression of *ISG15* and *IRF1*, 3.71 fold and 2.36 fold, respectively (Additional file 5: Table S1).

Local immune responses are activated which result in expression of proinflammatory cytokines when the uterus is exposed to bacteria. This is followed by production of antimicrobial peptides (AMP) and acute phase proteins by epithelial and innate immune cells [35]. Many of AMP genes are expressed in bovine endometrium, including lingual antimicrobial peptide (*LAP*), tracheal antimicrobial peptide (*TAP*), and some β -defensins [35, 54, 55]. Defensins are small cationic polypeptides, which have antimicrobial and immunomodulatory properties and are secreted by many cell types [56]. Some studies have identified an increase in expression levels of TAP, LAP, *DEFB1*, and *DEFB5* in endometrium of cows with serious inflammation or following LPS treatment *in vitro* [46, 51, 57]. Other studies showed LAP, TAP, neutrophil β -defensins (*BNBD4*, *DEFB5*) were all upregulated in bovine epithelial cells but not stromal cells with LPS [35, 58]. In the present study, we found *DEFB10* was upregulated 2.79 fold in bovine stromal cells with LPS (Additional file 5: Table S1).

Cascade reaction of complement signals also plays a key role in immune defense, and complement activation lead to opsonization of pathogens and their removal by phagocytes as well as cell lysis [59]. Some studies pointed out that complement factors *C2* and *CFB* increased in primary cultures of mixed bovine epithelial and stromal endometrial cells with LPS [45]. Our results agreed with previous studies and identified that the expression of other complement genes were upregulated, such as *C3*, *C1S*, *C1R*, *CF1*, and *C4A* (Additional file 5: Table S1). LPS can change expression level of many genes involved in cell adhesion, such as *VCAM1*, *SELP*, *CADM2*, *HEPACAM*, and *SDK2* (Additional file 5: Table S1). Leucocytes attach to cell adhesion molecules (CAM) on endothelial cells and are involved in inflammation and immune function [60]. Each CAM has an inherent effect on immune response process, such as vascular cell adhesion molecule-1 (VCAM-1), which promotes firm binding of T cells and induces trans-migration [61] In addition, there were reports in the 80s that CAMs may play a role in inherited inflammatory disease, which is characterized by deficient leucocyte CAMs [62].

This study also proves that LPS can alter other gene expression levels in stromal cells, such as hyaluronan synthase 2 (*HAS2*), which showed 3.36 fold increase in expression. It belongs to the hyaluronan (HA) family, holds extracellular matrix (ECM) polysaccharides and has multiple functions, such as tissue repair and promoting the expression of inflammation-related genes including *TNF*, *IL12*, *IL1B*, and *MMPs* [63]. Six-transmembrane epithelial antigen of prostate 4 (*STEAP4*), which showed 81.57 fold increase in expression, is an anti-inflammatory protein induced by inflammatory signals such as *TNF α* and *IL6* [64-66]. Notably, mice with *STEAP4*-knockout are susceptible to insulin resistance, hyperglycemia, and inflammation [67].

In our next study, we will continue to analyze and validate immune-related proteins and KEGG signaling pathways by western blot and compare genes and proteins in stromal cells treated with LPS *in vitro* with stromal cells from bovine endometritis with subclinical or clinical disease. These can better elucidate characteristics of bovine endometritis.

Conclusions

In brief, we identified a total of 366 DEGs in bovine endometrial stromal cells in LPS-induced group as compared to the control group. We analyzed the interactions between LPS and stromal cells including interaction of LPS with surface receptors of stromal cells and intracellular signaling cascades activated by PRRs. These indicate that stromal cells play a key role in bovine endometrium reaction to foreign intruders by initiating or inhibiting biochemical and molecular signals. In addition, our results identified many important immune-related genes and signaling pathways, especially, *IL6*, *IL1A*, *IL2RB*, *CCL2*, *CCL5*, *CXCL5*, *CXCL8*, *CXCL2*, and *CX3CL1*, and *TNF* signaling pathway, *NF- κ B* signaling pathway. Our results also expound the cellular processes in the endometrium and may serve as a diagnostic norm of defining and detecting endometritis, which may assist in development of more targeted and effective vaccines and drugs for the prevention and treatment of bovine endometritis. Furthermore, our analysis paves the way for the future elucidation of molecular mechanism of microbial invasion and host cell response. These results unraveled endometrial stromal cells transcriptome profile alterations in bovine affected by LPS which may have a significant effect on the eliminating or reducing inflammation by comprehending molecular mechanisms and authenticating unique genes related to endometritis.

Methods

Bovine endometrial stromal cell isolation and primary culture

Whole stromal cells were isolated from bovine endometrium as described before with some changes [68, 69]. All experimental procedures were approved by Institutional Animal Use Committee of Henan Agricultural University (approval number. 2005-0026). Bovine uteri, stored on ice, were transported to the laboratory from a local abattoir. The endometrium was cut into strips and placed in serum-free DMEM/F12 (Gibco, Grand Island, NY, USA) containing 50 IU/mL penicillin, 50 μ g/mL streptomycin, and 2.5 μ g/mL amphotericin B. Then, endometrial strips were cut into about 1 mm³ fragments and placed in

phosphate-buffered saline (PBS). They were then digested with 30 mL sterile digestive solution, composed of 60 mg trypsin III (Roche, Lewes, UK), 60 mg collagenase II (Sigma, Poole, UK), and 12 μ L deoxyribonuclease I (DNase I, Sigma) in 120 mL PBS at 37°C for about 1 h. The cell suspension was filtered with a 40 μ m mesh and the filtrate was resuspended with 2 mL DMEM/F-12 containing 10% fetal bovine serum (FBS, Gibco, Grand Island, NY, USA). After washing the cells five times with DMEM/F-12 containing 10% FBS, the filtrate was centrifuged at 100 rpm for 10 min at room temperature (RT) [18]. The cells were resuspended in DMEM/F-12 containing 15% FBS, 100 U/mL penicillin, 100 U/mL streptomycin, and 100 U/mL amphotericin B. Cells were seeded in 25 cm² culture flask at a density of 2×10^5 cells/mL, and were incubated at 37°C with 5% CO₂. To obtain stromal cells and remove epithelial cells populations, the cell suspension was removed 18 h after plating, which allowed selective attachment of stromal cells. The medium was changed every two days. Cells were passaged with 0.25% trypsin-EDTA until the cells reached ~80% confluence. Cell morphology was recorded and photographed under an inverted microscope.

Cells treatment with LPS

Stromal cells from passage 6 (P6) were seeded in 25 cm² culture flask at a density of 2×10^5 cells/mL. The cells were incubated at 37°C with 5% CO₂. Once the cells reached 80% confluence, the medium was removed and cells were washed with PBS. We divided the cells from an individual cow into two groups of three duplicates each. DMEM/F12 containing 15% FBS and PBS (control) was added to one group and DMEM/F12 containing 15% FBS and 0.5 μ g/mL ultrapure LPS obtained from an E. coli strain (serotype 055:B5, Sigma-Aldrich, Madison, USA, L2880) was added to the other group and incubated for 12 h each.

RNA extraction, quantification, and qualification

Total RNA from stromal cells was extracted by RNAisio Plus (Takara, Dalian, Liaoning, China). Concentration and quality of RNA were estimated with NanoDrop spectrophotometer (Thermo Scientific) and integrity of RNA was verified with 1% agarose gel.

Library preparation and transcriptome sequencing

Input material of 3 μ g RNA was used for RNA sample preparation. Sequencing libraries were generated using the TruSeq RNA Library Preparation Kit v2 (Illumina, San Diego, CA, USA). In summary, poly-A containing mRNA was enriched by oligo-dT magnetic beads and fragmentation was completed by divalent cations under elevated temperature in an Illumina proprietary fragmentation buffer. Random oligonucleotides and SuperScript II were used for synthesizing first strand cDNA. Second strand cDNA was synthesized. Remaining overhangs were converted into blunt ends by exonuclease/polymerase. Before hybridization, 3' ends of the DNA fragments was adenylated and Illumina paired-end (PE) adapter oligonucleotides were ligated. The amplified library fragments were enriched using Illumina PCR primer cocktail in a 15-cycle PCR reaction, then the library size was selected at 300-400 bp fragments. AMPure XP system was used to purify PCR products and Bioanalyzer 2100 system (Agilent, Santa, Clara, CA,

USA)) was used to detect library size. Finally, library sequencing of PEs was accomplished with Next-Generation Sequencing (NGS) on Illumina HiSeq platform.

Quality control

The raw data of FASTQ was generated by the software of the sequencing platform, Illumina HiSeq. Raw data of each sample was counted separately, including sample names, percentage of ambiguous base, Q20, and Q30 sequencing data along with some connectors and low-quality reads. These sequences may cause interference to subsequent data analysis, so sequencing data need to be further filtered. The criteria for data filtering included: 1) Cutadapt is used to remove 3' end connectors, and the removed parts have at least 10 bp overlap with known connectors, allowing 20% base mismatch; 2) removal of reads with average mass fraction lower than Q20.

Mapping reads to the reference genome

The reference genome used was of *Bos taurus*. ARS-UCD1.2.dna.toplevel.fa. were downloaded from Ensembl genome browser 95. The clean reads were mapped to the reference genome with HISAT2 (<http://ccb.jhu.edu/software/hisat2/index.shtml>).

Gene expression level quantification

The reads of each gene was counted by HTSeq and normalized by fragments per kilobase of transcript sequence per millions base pairs sequenced (FPKM). FPKM was then calculated [31].

Analysis of DEGs

Genes that were differentially expressed in two groups (three biological replicates per group) were screened with DESeq. The screening conditions for differentially expressed genes were: $|\log_2\text{FoldChange}| > 1$, significance of false discovery rate (FDR) < 0.05 . FDR is the adjusted P -value.

Gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis of DEGs

We performed GO enrichment analysis for function annotation and KEGG enrichment analysis for signaling pathway annotation on DEGs [70, 71]. GO and KEGG enrichment analyses were completed by the Database for Annotation, Visualization, and Integrated Discovery (DAVID) version 6.7 with threshold of FDR < 0.05 [72].

qRT-PCR

The expression profiles of randomly selected 10 differentially expressed genes were validated using SYBR Green based qRT-PCR using sequence specific primers (Additional file 8: Table S4) designed using the online primer design tool (<http://frodo.wi.mit.edu/primer3/>). The cDNA samples were synthesized by reverse transcription of equal amount of total RNA from stromal cells treated LPS (experimental group)

and PBS (control group), three duplicates each. respectively, using cDNA synthesis kit (Takara, Dalian, Liaoning, China) according to the manufacturer's instructions. The reverse transcription was first performed in 10 µl reaction volume to remove the genomic DNA reaction containing RNA samples, 1 µL; 5×gDNA Eraser Buffer, 2µL; gDNA Eraser, 1 µL, dd H₂O, 6 µL. Reaction conditions included 42°C for 2 min and 4°C 1 min, which followed 20 µl reaction volume reaction containing 10 µL reaction volume of the previous step, 1µL Prime Script RT Enzyme Mix, 1µL Prime Mix, 4 µL 5×Prime Script Buffer, 4 µL dd H₂O. Reaction conditions included 37°C for 15 min, 85°C 5 s, 4°C 1 min. Primer specificity was tested by first performing a conventional PCR and confirmed by the presence of a single melting curve during qRT-PCR. Serial dilutions(1:10, 1:20, 1:50, 1:200) were made from a pool of cDNA and calibration curves were performed for each gene. Afterwards, The qRT-PCR was then performed in 20 µl reaction volume containing TB Green Premix ExTaq II, 10 µL (Takara, Dalian, Liaoning, China); the cDNA samples, 1µL; the specific forward and reverse primer, 1µL (respectively); and dd H₂O, 7 µL in the 0.1 mL white PCR 8-strip Tubes (NEST Biotechnology, Wuxi, China) with CFX96 real-time PCR detection system (Bio-Rad, Munich, Germany). Reaction conditions included pre-degeneration at 95°C for 30 s, followed by 40 cycles of degeneration at 95°C for 5 s, and annealing at 60°C for 30 s. At the end of each PCR reaction, the specificity the amplification was confirmed by evaluating the dissociation curve. The abundance of each transcript in each sample was determined using a comparative threshold cycle comparative Ct ($2^{-\Delta\Delta CT}$) method as described previously [73]. The data obtained from qRT-PCR was analyzed after the Ct value of the target genes was normalized with the Ct value of Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*). The qRT-PCR was performed in three biological. The Student's t-test or the least significant difference test procedure was employed to detect the mRNA expression differences between the samples. Differences with $p < 0.05$ were considered as significant. Statistical significance of the data was determined by a Student's t-test carried out with SPSS (PASW Statistics for Windows, Version 18.0, Chicago: SPSS Inc., USA).

Declarations

Ethics approval and consent to participate

All experimental procedures were approved by Institutional Animal Use Committee of Henan Agricultural University (approval number. 2005-0026) and Beijing Association for Science and Technology (approval SYXK [Beijing] 2007-0023).

Consent for publication

Not applicable.

Availability of data and materials

The sequence data of this study have been deposited into Sequence Read Archive (<http://www.ncbi.nlm.nih.gov/sra>, accession number PRJNA574911). The datasets supporting the conclusions of this article are included in this article and its supplementary information files.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Key Research and Development Program (2018YFD0501803, 2018YFD0501800), The Programmes of the National Beef Cattle and Yak Industrial Technology System (CARS-37) and Project for the Training of Young Cadre Teachers in Henan Higher Education Institutions in 2016(2016GGJS-282).

Authors' contributions

XW, CT and XD designed the study and revised manuscript. XD, HL, LD, WH, ZP, CY, and DY performed the study. XD, HL, LD, CT and XW analyzed the data. XD, HL and LD wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

Abbreviations

LPS: Lipopolysaccharide; DEGs: Differentially expressed genes; GO: Gene ontology; KEGG: Kyoto Encyclopedia of Genes and Genomes; PPARs peroxisome proliferator-activated receptors; RNA-Seq: High-throughput mRNA sequencing; PCA: Principal component analysis; LBP: lipoprotein binding protein; PRRs: Innate pattern recognition receptors; AMP: Antimicrobial peptides; *LAP*: Lingual antimicrobial peptide; *TAP*: tracheal antimicrobial peptide; MD-2: Myeloid differentiation factor 2; FDR: False discovery rate; IRF3: interferon regulatory factor 3; MAP kinase: Mitogen-activated protein kinase; ISG: IFN-stimulated genes; GAPDH: Glyceraldehyde 3-phosphate-dehydrogenase; STEAP4: Six-transmembrane epithelial antigen of prostate 4; CAM: Cell adhesion molecules; VCAM-1: Vascular cell adhesion molecule-1; HA: hyaluronan; ECM: extracellular matrix; HAS2: hyaluronan synthase 2

References

1. Kelly P, Meade KG, O'Farrelly C. Non-canonical Inflammasome-Mediated IL-1beta Production by Primary Endometrial Epithelial and Stromal Fibroblast Cells Is NLRP3 and Caspase-4 Dependent. *Front Immunol.* 2019;10:102. <https://doi.org/10.3389/fimmu.2019.00102>.
2. Sheldon IM, Cronin J, Goetze L, Donofrio G, Schuberth HJ. Defining Postpartum Uterine Disease and the Mechanisms of Infection and Immunity in the Female Reproductive Tract in Cattle. *Biol Reprod.* 2009;81(6):1025-1032. <https://doi.org/10.1095/biolreprod.109.077370>.

3. Bromfield JJ, Santos JE, Block J, Williams RS, Sheldon IM. PHYSIOLOGY AND ENDOCRINOLOGY SYMPOSIUM: Uterine infection: linking infection and innate immunity with infertility in the high-producing dairy cow. *J Anim Sci*. 2015; 93(5):2021-2033. <https://doi.org/10.2527/jas.2014-8496>.
4. Ribeiro ES, Lima FS, Greco LF, Bisinotto RS, Monteiro AP, Favoreto M, et al. Prevalence of periparturient diseases and effects on fertility of seasonally calving grazing dairy cows supplemented with concentrates. *J Dairy Sci*. 2013;96(9):5682-5697. <https://doi.org/10.3168/jds.2012-6335>.
5. Chapwanya A, Meade KG, Doherty ML, Callanan JJ, Mee JF, O'Farrelly C. Histopathological and molecular evaluation of Holstein-Friesian cows postpartum: toward an improved understanding of uterine innate immunity. *Theriogenology*. 2009;71(9):1396-1407. <https://doi.org/10.1016/j.theriogenology.2009.01.006>.
6. Dohmen MJ, Joop K, Sturk A, Bols PE, Lohuis JA. Relationship between intra-uterine bacterial contamination, endotoxin levels and the development of endometritis in postpartum cows with dystocia or retained placenta. *Theriogenology*. 2000;54(7):1019-1032. [https://doi.org/10.1016/s0093-691x\(00\)00410-6](https://doi.org/10.1016/s0093-691x(00)00410-6).
7. Beutler B, Hoebe K, Du X, Ulevitch RJ. How we detect microbes and respond to them: the Toll-like receptors and their transducers. *J Leukoc Biol*. 2003;74(4):479-485. <https://doi.org/10.1189/jlb.0203082>.
8. Socha BM, Lada P, Szczepanska AA, Lupicka M, Korzekwa AJ. The influence of experimentally induced endometritis on the PPAR expression profile in the bovine endometrium. *Theriogenology*. 2018;122:74-83. <https://doi.org/10.1016/j.theriogenology.2018.09.013>.
9. Socha BM, Lada P, Szczepanska AA, Lupicka M, Korzekwa AJ. The influence of experimentally induced endometritis on the PPAR expression profile in the bovine endometrium. *Theriogenology*. 2018;122:74-83. <https://doi.org/10.1016/j.theriogenology.2018.09.013>.
10. MacLaren LA1, Guzeloglu A, Michel F, Thatcher WW. Peroxisome proliferator-activated receptor (PPAR) expression in cultured bovine endometrial cells and response to omega-3 fatty acid, growth hormone and agonist stimulation in relation to series 2 prostaglandin production. *Domest Anim Endocrinol*. 2006;30(3):155-169. <https://doi.org/10.1016/j.domaniend.2005.07.003>
11. Darren, Davies, Kieran, Meade, Shan, Herath, David, Eckersall. Toll-like receptor and antimicrobial peptide expression in the bovine endometrium. *Reprod Biol Endocrinol*. 2008;6:53. <https://doi.org/10.1186/1477-7827-6-53>.
12. Amos MR, Healey GD, Goldstone RJ, Mahan SM, Duvel A, Schuberth HJ, et al. Differential endometrial cell sensitivity to a cholesterol-dependent cytolysin links *Trueperella pyogenes* to uterine disease in cattle. *Biol Reprod*. 2014;90(3):54. <https://doi.org/10.1095/biolreprod.113.115972>
13. Sheldon IM. Genes and environmental factors that influence disease resistance to microbes in the female reproductive tract of dairy cattle. *Reprod Fert*. 2015;27(1):72-81. <https://doi.org/10.1071/RD14305>

14. Wagner WC, Hansel W. Reproductive physiology of the post partum cow. I. Clinical and histological findings. *J Reprod Fertil.* 1969;18(3):493-500. <https://doi.org/10.1530/jrf.0.0180493>.
15. Archbald LF, Schultz RH, Fahning ML, Kurtz HJ, Zemjanis R. A sequential histological study of the post-partum bovine uterus. *J Reprod Fertil.* 1972;29(1):133-136. <https://doi.org/10.1530/jrf.0.0290133>
16. Bonnett BN, Martin SW, Gannon VP, Miller RB, Etherington WG. Endometrial biopsy in Holstein-Friesian dairy cows. III. Bacteriological analysis and correlations with histological findings. *Can J Vet Res.* 1991;55(2):168-173.
17. Sheldon I, Noakes D, Rycroft A, Pfeiffer D, Dobson H. Influence of uterine bacterial contamination after parturition on ovarian dominant follicle selection and follicle growth and function in cattle. *Reproduction.* 2002;123(6):837-845.
18. Sheldon IM1, Rycroft AN, Dogan B, Craven M, Bromfield JJ, Chandler A, et al. Specific Strains of *Escherichia coli* Are Pathogenic for the Endometrium of Cattle and Cause Pelvic Inflammatory Disease in Cattle and Mice. *PLoS One.* 2010;5(2):e9192. <https://doi.org/10.1371/journal.pone.0009192>.
19. Williams EJ1, Fischer DP, Noakes DE, England GC, Rycroft A, Dobson H, et al. The relationship between uterine pathogen growth density and ovarian function in the postpartum dairy cow. *Theriogenology.* 2007;68(4):549-559. <https://doi.org/10.1016/j.theriogenology.2007.04.056>
20. Deng Y, Liu B, Mao W, Shen Y, Fu C, Gao L, et al. Regulatory roles of PGE2 in LPS-induced tissue damage in bovine endometrial explants. *Eur J Pharmacol.* 2019;852:207-217. <https://doi.org/10.1016/j.ejphar.2019.03.044>
21. Amos MR1, Healey GD, Goldstone RJ, Mahan SM, Düvel A, Schuberth HJ, et al. Differential endometrial cell sensitivity to a cholesterol-dependent cytolysin links *Trueperella pyogenes* to uterine disease in cattle. *Biol Reprod.* 2014;90(3):54. <https://doi.org/10.1095/biolreprod.113.115972>
22. Ireland JJ, Murphee RL, Coulson PB. Accuracy of predicting stages of bovine estrous cycle by gross appearance of the corpus luteum. *J Dairy Sci.* 1980;63(1):155-60. [https://doi.org/10.3168/jds.S0022-0302\(80\)82901-8](https://doi.org/10.3168/jds.S0022-0302(80)82901-8).
23. Cronin JG, Turner ML, Goetze L, Bryant CE, Sheldon IM. Toll-like receptor 4 and MYD88-dependent signaling mechanisms of the innate immune system are essential for the response to lipopolysaccharide by epithelial and stromal cells of the bovine endometrium. *Biol Reprod.* 2012;86(2):51. <https://doi.org/10.1095/biolreprod.111.092718>.
24. de Moraes CN, Maia L, Dias MC, Dell'Aqua CP, da Mota LS2Chapwanya A, et al. Bovine endometrial cells: a source of mesenchymal stem/progenitor cells. *Cell Biol Int.* 2016;40(12):1332-1339. <https://doi.org/10.1002/cbin.10688>
25. Koh YQ, Mitchell MD, Almughlliq FB, Vaswani K, Peiris HN. Regulation of inflammatory mediator expression in bovine endometrial cells: effects of lipopolysaccharide, interleukin 1 beta, and tumor necrosis factor alpha. *Physiol Rep.* 2018;6(9):e13676. <https://doi.org/10.14814/phy2.13676>.

26. Almughlliq FB, Koh YQ, Peiris HN, Vaswani K, Arachchige BJ, Reed S et al. Eicosanoid pathway expression in bovine endometrial epithelial and stromal cells in response to lipopolysaccharide, interleukin 1 beta, and tumor necrosis factor alpha. *Reprod Biol.* 2018;18(4):390-396. <https://doi.org/10.1016/j.repbio.2018.10.001>
27. Healey GD, Collier C, Griffin S, Schuberth HJ, Sandra O, Smith DG et al. Mevalonate Biosynthesis Intermediates Are Key Regulators of Innate Immunity in Bovine Endometritis. *J Immunol.* 2016;196(2):823-31. <https://doi.org/10.4049/jimmunol.1501080>
28. Carneiro LC, Bedford C, Jacca S, Rosamilia A, de Lima VF, Donofrio G et al. Coordinated Role of Toll-Like Receptor-3 and Retinoic Acid-Inducible Gene-I in the Innate Response of Bovine Endometrial Cells to Virus. *Front Immunol.* 2017. <https://doi.org/10.3389/fimmu.2017.00996>.
29. Oguejiofor CF, Cheng Z, Abudureyimu A, Fouladi-Nashta AA, Wathes DC. Global transcriptomic profiling of bovine endometrial immune response in vitro. I. Effect of lipopolysaccharide on innate immunity. *Biol Reprod.* 2015;93(4):100. <https://doi.org/10.1095/biolreprod.115.128868>
30. Marioni JC, Mason CE, Mane SM, Stephens M, Gilad Y. RNA-seq: an assessment of technical reproducibility and comparison with gene expression arrays. *Genome Res.* 2008;18(9):1509-17. <https://doi.org/10.1101/gr.079558.108>.
31. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, et al. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol.* 2010;28(5):511-5. <https://doi.org/10.1038/nbt.1621>.
32. Wang J, Yan X, Nesengani LT, Yang L, Lu W. Integrated analysis of miRNA and mRNA expression profiling in bovine endometrial cells in response to lipopolysaccharide-stimulation. *Microb Pathog.* 2018. <https://doi.org/10.1016/j.micpath.2017.11.053>
33. Foley C, Chapwanya A, Callanan JJ, , Whiston R, Miranda-CasoLuengo R, Lu J, et al. Integrated analysis of the local and systemic changes preceding the development of post-partum cytological endometritis. *BMC Genomics.* 2015 <https://doi.org/10.1186/s12864-015-1967-5>
34. Guo Y, van Schaik T, Jhamat N, Niazi A, Chanrot M, Charpigny G, et al. Differential gene expression in bovine endometrial epithelial cells after challenge with LPS; specific implications for genes involved in embryo maternal interactions. *PLoS One.* 2019;14(9):e0222081. <https://doi.org/10.1371/journal.pone.0222081>.
35. Chapwanya A, Meade KG, Doherty ML, Callanan JJ, Mee JF, O'Farrelly C. Histopathological and molecular evaluation of Holstein-Friesian cows postpartum: toward an improved understanding of uterine innate immunity. *Theriogenology.* 2009;71(9):1396-407. <https://doi.org/10.1016/j.theriogenology.2009.01.006>.
36. Cretoiu SM, Cretoiu D, Suciuc L, Popescu LM. Interstitial Cajal-like cells of human Fallopian tube express estrogen and progesterone receptors. *J Mol Histol.* 2009;40(5-6):387-94. <https://doi.org/10.1007/s10735-009-9252-z>.
37. Niu C, Cheng C, Liu Y, Huang S, Fu Y, Li P. Transcriptome Profiling Analysis of Bovine Vaginal Epithelial Cell Response to an Isolated Lactobacillus Strain. *mSystems.* 2019;4(5).

<https://doi.org/10.1128/mSystems.00268-19>

38. Inui S, Fukuzato Y, Nakajima T, Kurata S, Itami S. Androgen receptor co-activator Hic-5/ARA55 as a molecular regulator of androgen sensitivity in dermal papilla cells of human hair follicles. *J Invest Dermatol.* 2007;127(10):2302-6. <https://doi.org/10.1038/sj.jid.5700883>.
39. Kent LW, Dyken RA, Rahemtulla F, Allison AC, Michalek SM. Effect of in vitro passage of healthy human gingival fibroblasts on cellular morphology and cytokine expression. *Arch Oral Biol.* 1996;41(3):263-70. [https://doi.org/10.1016/0003-9969\(95\)00127-1](https://doi.org/10.1016/0003-9969(95)00127-1).
40. Danesh Mesgaran S, Sharbati J, Einspanier R, Gabler C. mRNA expression pattern of selected candidate genes differs in bovine oviductal epithelial cells in vitro compared with the in vivo state and during cell culture passages. *Reprod Biol Endocrinol.* 2016;14(1):44. <https://doi.org/10.1186/s12958-016-0176-7>.
41. Danesh Mesgaran S, Gärtner MA, Wagener K, Drillich M, Ehling-Schulz M, Einspanier R, et al. Different inflammatory responses of bovine oviductal epithelial cells in vitro to bacterial species with distinct pathogenicity characteristics and passage number. *Theriogenology.* 2018. <https://doi.org/10.1016/j.theriogenology.2017.10.005>.
42. Sandberg R, Ernberg I. The molecular portrait of in vitro growth by meta-analysis of gene-expression profiles. *Genome Biol.* 2005;6(8):R65. <https://doi.org/10.1186/gb-2005-6-8-r65>
43. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol.* 2010;11(5):373-84. <https://doi.org/10.1038/ni.1863>.
44. Herath S, Lilly ST, Fischer DP, Williams EJ, Dobson H, Bryant CE, et al. Bacterial lipopolysaccharide induces an endocrine switch from prostaglandin F2alpha to prostaglandin E2 in bovine endometrium. *Endocrinology.* 2009;150(4):1912-20. <https://doi.org/10.1210/en.2008-1379>.
45. Oguejiofor CF, Cheng Z, Abudureyimu A, Fouladi-Nashta AA, Wathes DC. Global transcriptomic profiling of bovine endometrial immune response in vitro. I. Effect of lipopolysaccharide on innate immunity. *Biol Reprod.* 2015;93(4):100. <https://doi.org/10.1095/biolreprod.115.128868>.
46. Fu Y, Liu B, Feng X, Liu Z, Liang D, Li F, et al. Lipopolysaccharide increases Toll-like receptor 4 and downstream Toll-like receptor signaling molecules expression in bovine endometrial epithelial cells. *Vet Immunol Immunopathol.* 2013;151(1-2):20-7
47. Kayagaki N, Wong MT, Stowe IB, Ramani SR, Gonzalez LC, Akashi-Takamura S, et al. Noncanonical inflammasome activation by intracellular LPS independent of TLR4. *Science.* 2013;341(6151):1246-9. <https://doi.org/10.1126/science.1240248>
48. Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell.* 2010;140(6):805-20. <https://doi.org/10.1016/j.cell.2010.01.022>
49. Herath S, Lilly ST, Santos NR, et al. Expression of genes associated with immunity in the endometrium of cattle with disparate postpartum uterine disease and fertility. *Reprod Biol Endocrinol.* 2009. <https://doi.org/10.1186/1477-7827-7-55>.
50. Fischer C, Drillich M, Odau S, Heuwieser W, Einspanier R, Gabler C. Selected pro-inflammatory factor transcripts in bovine endometrial epithelial cells are regulated during the oestrous cycle and

- elevated in case of subclinical or clinical endometritis. *Reprod Fertil Dev.* 2010;22(5):818-29. <https://doi.org/10.1071/RD09120>.
51. Swangchan-Uthai T, Lavender CR, Cheng Z, Fouladi-Nashta AA, Wathes DC. Time course of defense mechanisms in bovine endometrium in response to lipopolysaccharide. *Biol Reprod.* 2012;87(6):135. <https://doi.org/10.1095/biolreprod.112.102376>.
 52. Cronin JG, Turner ML, Goetze L, Bryant CE, Sheldon IM. Toll-like receptor 4 and MYD88-dependent signaling mechanisms of the innate immune system are essential for the response to lipopolysaccharide by epithelial and stromal cells of the bovine endometrium. *Biol Reprod.* 2012;86(2):51. <https://doi.org/10.1095/biolreprod.111.092718>.
 53. Yanai H, Taniguchi T. Fine-tuning type I IFN signaling: A new chapter in the IFN saga. *Cell Res.* 2017;27(12):1407-1408. <https://doi.org/10.1038/cr.2017.118>.
 54. Cormican P, Meade KG, Cahalane S, Narciandi F, Chapwanya A, Lloyd AT, et al. Evolution, expression and effectiveness in a cluster of novel bovine beta-defensins. *Immunogenetics.* 2008;60(3-4):147-56. <https://doi.org/10.1007/s00251-007-0269-8>.
 55. Severa M, Coccia EM, Fitzgerald KA. Toll-like receptor-dependent and -independent viperin gene expression and counter-regulation by PRDI-binding factor-1/BLIMP1. *J Biol Chem.* 2006;281(36):26188-95. <https://doi.org/10.1074/jbc.M604516200>.
 56. Ganz T. Defensins: antimicrobial peptides of innate immunity. *Nat Rev Immunol.* 2003;3(9):710-20. <https://doi.org/10.1038/nri1180>.
 57. Wathes DC, Cheng Z, Chowdhury W, Fenwick MA, Fitzpatrick R, Morris DG, et al. Negative energy balance alters global gene expression and immune responses in the uterus of postpartum dairy cows. *Physiol Genomics.* 2009;39(1):1-13. <https://doi.org/10.1152/physiolgenomics.00064.2009>
 58. Davies D, Meade KG, Herath S, Eckersall PD, Gonzalez D, White JO, et al. Toll-like receptor and antimicrobial peptide expression in the bovine endometrium. *Reprod Biol Endocrinol.* 2008. <https://doi.org/10.1186/1477-7827-6-53>.
 59. Sarma JV, Ward PA. The complement system. *Cell Tissue Res.* 2011;343(1):227-35. <https://doi.org/10.1007/s00441-010-1034-0>.
 60. Kim SK, Moon WK, Park JY, Jung H. Inflammatory mimetic microfluidic chip by immobilization of cell adhesion molecules for T cell adhesion. *Analyst.* 2012;137(17):4062-8. <https://doi.org/10.1039/c2an35424a>.
 61. Ferran C, Peuchmaur M, Desruennes M, Ghoussoub JJ, Cabrol A, Brousse N, et al. Implications of de novo ELAM-1 and VCAM-1 expression in human cardiac allograft rejection. *Transplantation.* 1993;55(3):605-9. <https://doi.org/10.1097/00007890-199303000-00026>.
 62. Allen S, Moran N. Cell Adhesion Molecules: Therapeutic Targets for Inhibition of Inflammatory States. *Semin Thromb Hemost.* 2015;41(6):563-71. <https://doi.org/10.1055/s-0035-1556588>.
 63. Petrey AC, de la Motte CA. Hyaluronan, a crucial regulator of inflammation. *Front Immunol.* 2014. Mar 11;5:101. <https://doi.org/10.3389/fimmu.2014.00101>.

64. Gomes IM, Maia CJ, Santos CR. STEAP proteins: from structure to applications in cancer therapy. *Mol Cancer Res.* 2012;10(5):573-87. <https://doi.org/10.1158/1541-7786.MCR-11-0281>.
65. Korkmaz CG, Korkmaz KS, Kurys P, Elbi C, Wang L, Klock TI, et al. Molecular cloning and characterization of STAMP2, an androgen-regulated six transmembrane protein that is overexpressed in prostate cancer. *Oncogene.* 2005;24(31):4934-45. <https://doi.org/10.1038/sj.onc.1208677>.
66. Yoo SK, Cheong J, Kim HY. STAMPing into Mitochondria. *Int J Biol Sci.* 2014;10(3):321-6. <https://doi.org/10.7150/ijbs.8456>.
67. Wellen KE, Fucho R, Gregor MF, Morgan C, Lindstad T, Vaillancourt E, et al. Coordinated regulation of nutrient and inflammatory responses by STAMP2 is essential for metabolic homeostasis. *Cell.* 2007;129(3):537-48. <https://doi.org/10.1016/j.cell.2007.02.049>.
68. Fortier MA, Guilbault LA, Grasso F. Specific properties of epithelial and stromal cells from the endometrium of cows. *J Reprod Fertil.* 1988;83(1):239-48. <https://doi.org/10.1530/jrf.0.0830239>.
69. Hailemariam D, Ibrahim S, Hoelker M, Drillich M, Heuwieser W, Looft C, et al. MicroRNA-regulated molecular mechanism underlying bovine subclinical endometritis. *Reprod Fertil Dev.* 2014;26(6):898-913. <https://doi.org/10.1071/RD13027>.
70. Kanehisa M, Goto S, Kawashima S, Okuno Y, Hattori M. The KEGG resource for deciphering the genome. *Nucleic Acids Res.* 2004;32(Database issue):D277-80. <https://doi.org/10.1093/nar/gkh063>.
71. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* 2017;45(D1):D353-D361. <https://doi.org/10.1093/nar/gkw1092>.
72. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 2009;37(1):1-13. <https://doi.org/10.1093/nar/gkn923>.
73. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001;25(4):402-8. <https://doi.org/10.1006/meth.2001.1262>.

Tables

Table 1 Data filtering statistics after Illumina sequencing

Sample	Raw Reads	Clean Reads No.	Clean Data (bp)	Clean Reads %	Q20 (%)	Q30 (%)
PBS1	47469396	44046138	6650966838	92.78	96.08	91.45
PBS2	46293936	42854784	6471072384	92.57	96.49	92.24
PBS3	43441712	40242110	6076558610	92.63	96.25	91.79
LPS1	42929910	39806986	6010854886	92.72	96.19	91.59
LPS2	50813700	47094016	7111196416	92.67	96.16	91.6
LPS3	49082750	45600808	6885722008	92.9	96.42	91.9

N %: The percentage of fuzzy bases

Q20: The percentage of bases with a Phred value > 20

Q30: The percentage of bases with a Phred value > 30

Table 2 Summary of clean reads mapped to the reference genome

Sample	Clean_Reads	Total_Mapped	Multiple_Mapped	Uniquely_Mapped
PBS1	44046138	41598276 (94.44%)	2394502 (5.76%)	39203774 (94.24%)
PBS2	42854784	39556838 (92.30%)	2206194 (5.58%)	37350644 (94.42%)
PBS3	40242110	37935794 (94.27%)	2240965 (5.91%)	35694829 (94.09%)
LPS1	39806986	37586818 (94.42%)	2233726 (5.94%)	35353092 (94.06%)
LPS2	47094016	44530730 (94.56%)	2676831 (6.01%)	41853899 (93.99%)
LPS3	45600808	43007562 (94.31%)	2461139 (5.72%)	40546423 (94.28%)

Clean Reads: The total number of sequences used for alignment

Total Mapped: The total number of sequences identical to the reference genome in clean reads
percentage is total mapped / clean reads

Multiple Mapped: The total number of sequences aligned to multiple locations, percentage is multiple mapped / total mapped

Uniquely Mapped: The total number of sequences aligned to only one location, percentage is uniquely mapped / total mapped

Figures




Figure 1

Volcano map of DEGs. The two vertical dotted lines are the threshold of differential expression. The horizontal dotted line is the threshold FDR of 0.05. Upregulated and downregulated genes are shown as a red and blue dots, and gray dots represents non-significantly differentially expressed genes. (A: control group; B: LPS-induced group)




Figure 2

Heatmap analysis of DEGs. The horizontal lines represent genes, and each column is a sample. Red represents high-expression genes and green represents low-expression genes. The X-axis is the sample number and the Y-axis is the DEGs.

GO Enrichment




Figure 3

GO enrichment analysis displaying the first 20 GO terms with the most significant enrichment.

KEGG Pathway Enrichment




Figure 4

KEGG enrichment analysis displaying the first 20 KEGG terms with the most significant enrichment.




Figure 5

QRT-PCR verified features of DEGs by RNA-seq. The relative expression level of target mRNAs was calculated using the $2^{-\Delta\Delta Ct}$ method and expressed relative to the value in the control group. Results were displayed in mean \pm SEM (n = 3). Log2 fold change was the ratio of average log2 folds between groups.




Figure 6

Stromal cells were exposed to an inflammatory environment when epithelial cells were disrupted.




Figure 7

Summary of the immune response in bovine endometrial stromal cells exposed to bacterial LPS. LPS treatment for 12 h changed the mRNA expression of many genes involved inflammatory and innate immune response.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile3.rar](#)

- [Additionalfile5.rar](#)
- [Additionalfile8.rar](#)
- [Additionalfile7.rar](#)
- [Additionalfile6.rar](#)
- [Additionalfile4.rar](#)
- [Additionalfile2.rar](#)
- [Additionalfile1.rar](#)