Antioxidant Potential of *Pediococcus Pentosaceus* Strains from the Sow Milk Bacterial Collection in Weaned Piglets

Leli Wang

Hunan Normal University

Qihang Liu

Hunan Normal University

Yuwei Chen

Hunan Normal University

Xinlei Zheng

Hunan Normal University

Chuni Wang

Hunan Normal University

Yining Qi

Hunan Normal University

Yachao Dong

Hunan Normal University

Yue Xiao

Hunan normal university

Cang Chen

Hunan Normal University

Taohong Chen

Hunan Normal University

Qiuyun Huang

Hunan Normal University

Jialu Huang

Hunan Agricultural University

Zongzhao Zhai

Hunan Normal University

Cimin Long

Institute of Subtropical Agriculture Chinese Academy of Sciences

Huansheng Yang

Hunan Normal University

Jianzhong Li

Hunan Normal University

Lei Wang

Hunan Normal University

Gaihua Zhang

Hunan Normal University

Peng Liao

Institute of Subtropical Agriculture Chinese Academy of Sciences

Yongxin Liu

Institute of Genetics and Developmental Biology CAS: Institute of Genetics and Developmental Biology
Chinese Academy of Sciences

Peng Huang

Hunan Agricultural University

Qiye Wang

Hunan Normal University

Huanhuan Chu

Yantai Hi-tech Industrial Development Zone

Jia Yin (✉ jiayin@hunnu.edu.cn)

animal nutrition and human health laboratory <https://orcid.org/0000-0002-8867-3490>

Yulong Yin

Institute of Subtropical Agriculture Chinese Academy of Sciences

Research

Keywords: culturomics, sow milk, probiotic, oxidative stress, Drosophila, piglets

Posted Date: November 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-75522/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Antioxidant Potential of *Pediococcus Pentosaceus* Strains from the Sow Milk Bacterial Collection in Weaned Piglets

Leli Wang¹, Qihang Liu¹, Yuwei Chen¹, Xinlei Zheng¹, Chuni Wang¹, Yining Qi¹,
Yachao Dong¹, Yue Xiao¹, Cang Chen¹, Taohong Chen¹, Qiuyun Huang¹, Jialu Huang⁵,
Zongzhao Zhai¹, Cimin Long², Huansheng Yang¹, Jianzhong Li¹, Lei Wang¹, Gaihua
Zhang³, Peng Liao², Yongxin Liu⁴, Peng Huang⁵, Qiye Wang¹, Huanhuan Chu⁶, Jia
Yin^{1*}, Yulong Yin^{2*}

¹ Laboratory of Animal Nutrition and Human Health, Hunan Provincial Key Laboratory
of Animal Intestinal Function and Regulation, Hunan International Joint Laboratory of
Animal Intestinal Ecology and Health, College of Life Sciences, Hunan Normal
University, Changsha, China.

²Institute of Subtropical Agriculture, Chinese Academy of Sciences, Changsha, China

³The National and Local Joint Engineering Laboratory of Animal Peptide Drug
Development, College of Life Sciences, Hunan Normal University, Changsha, China

⁴State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental
Biology, Chinese Academy of Sciences, Beijing, China

⁵College of Animal Science and Technology, Hunan Agricultural University,
Changsha, Hunan, China.

⁶Shandong Yihe Feed Co.,Ltd. Yantai Hi-tech Industrial Development Zone, Yantai,
Shandong, China.

* Correspondence: jjayin@hunnu.edu.cn; yinyulong@isa.ac.cn

Abstract

Background: In modern animal husbandry, breeders pay increasing attention to improving sow nutrition during pregnancy and lactation to favor the health of neonates. Sow milk is a main food for piglets during their first three weeks of life, which is not only a rich repository of essential nutrients and a broad range of bioactive compounds, but also an indispensable source of commensal bacteria. Maternal milk microorganisms are important sources of commensal bacteria for neonatal gut. Bacteria from maternal milk may serve as an additive to confer a health benefit on the composition of the indigenous microbiota of piglets.

Methods: The sow milk microbiota was collected using the culturomics methods of Continuous Culture and Interval Sampling, following by the identification of 16S rDNA gene sequences. To screen potential probiotics, the functional evaluation was conducted to assess their antagonistic activity against pathogens *in vitro* and evaluate their resistance against oxidative stress in damaged *Drosophila* induced by paraquat. In piglet feeding trial, a total of 54 newborn suckling piglets were chosen from nine sows and randomly assigned to three treatments with different concentration of a candidate strain. Multiple approaches were carried out to verify its antioxidant function including western blotting, enzyme activity analysis, metabolomics and 16S rDNA sequencing.

Results: The 1240 isolates were screened out from the sow milk microbiota and grouped into 271 bacterial taxa based on a nonredundant set of 16S rDNA gene sequences. Among 80 *Pediococcus* isolates, a new *Pediococcus pentosaceus* strain

(SMM914) showed best performance in inhibition ability against swine pathogens and in *Drosophila* model challenged by paraquat. Pretreatment of piglets with SMM914 induces the Nrf2-Keap1 antioxidant signaling pathway and greatly affected the pathways of amino acid metabolism and lipid metabolism in plasma. In colon, *Lactobacillus* was significantly increased in the high dose of SMM914 group compared with the control group.

Conclusion: SMM914 functions as a promising probiotic conferring antioxidant capacity by activating the Nrf2-Keap1 antioxidant signaling pathway in piglets. Our study provided useful resources for deeply understanding the relationships between the maternal microbiota and offspring.

Keywords: culturomics, sow milk, probiotic, oxidative stress, *Drosophila*, piglets

Introduction

A unique microbiome is established in every individual in their first few years of life. In particular, the first year of life is a crucial stage of the development of the microbiome. Breastfeeding is the main factor affecting the development of the microbiome in this period [1]. Breastfed infants have a decreased risk of acute and chronic diseases, including gastroenteritis and sudden infant death syndrome, because breast milk provides essential nutrients and a broad range of bioactive compounds for developing neonates [2]. Commensal bacteria from breast milk act as pioneer bacteria during the critical stage of initial neonatal gut colonization [3]. Several studies at the strain level have specifically demonstrated that there are some bacteria shared between human breast milk and infant feces by isolating and identifying bacteria [4]. The vertical transfer concept that the breast milk microbiota can be transmitted from mothers to infants during breast feeding became accepted gradually [5]. The maternal microbiota contributes to the 'initial' intestinal microbiota establishment in infants and help to modulate both short- and long-term infant health outcomes [6-8].

Due to the high degree of similarity in anatomy, physiology and immunology between humans and pigs, piglets have been extensively used as an ideal model to study infants' gastrointestinal system and health [6]. Nowadays, pork is the most widely consumed meat in the world [9]. Early weaning is an abrupt event in piglets that often leads to severe oxidative stress and restricts the development of commercial swine husbandry. Breast milk is a main food for piglets during the first three weeks of their lives.

81 Although abundant probiotics have been isolated from human milk and bovine milk
82 [10-12], strategies for the preservation and isolation of the natural bacteria of sow milk
83 are limited currently. While culture-independent methods have allowed an
84 understanding of the composition and diversity of sow milk microbiota [9], culture-
85 dependent methods are still critical for the functional identification and utilization of
86 the sow milk microbiota, which might be indirectly or directly beneficial for precise
87 care of early-weaning piglets. However, no research has systematically investigated the
88 sow milk microbiota using culture-dependent methods so far.

89 In the present work, a comprehensive collection of sow milk microbiota was first
90 established by the culturomics of Continuous Culture and Interval Sampling. Among
91 the 1,240 bacterial isolates, 80 strains of *Pediococcus pentosaceus* (*P. pentosaceus*)
92 were identified. *P. pentosaceus*, a gram-positive, catalase-negative, homofermentative
93 bacteria strain that has long been used as biopreservatives and healthy commercial
94 starters in fermented foods [13-15]. We depended on antagonistic experiments against
95 pathogens *in vitro* and antioxidant tests in *Drosophila* model to select the candidate
96 probiotic species *Pediococcus pentosaceus* (*P. pentosaceus*). Interestingly, the
97 antioxidant activity of *P. pentosaceus* SMM914 in piglets was verified by elevated
98 antioxidant metabolites in plasma and the activation of the Nrf2-Keap1 pathway in
99 livers. It is suggested that SMM914 could confer health benefits to piglets and represent
100 a promising candidate for maintaining redox balance. These results of the sow milk

bacterium and its antioxidant function in piglets deserve more attention for promoting healthy livestock breeding.

Results

A comprehensive collection of lactic acid bacteria from sow milk

Breast milk is one of common sources of probiotic strains [16]. We hypothesized that these lactic acid bacteria (LAB) from sow milk could facilitate adaptive, functional changes for the optimal weaning transition of piglets, which are vulnerable to various stressors in early life. However, the diversity and function of probiotics in porcine milk remain relatively understudied [17]. To establish a sow milk bacterial collection (smBC), a large-scale cultivation and identification of the sow milk microbiota (SMM) were performed by sow milk collection, anerobic incubation and sequencing (Fig. 1, steps 1–3).

After the first three steps, we obtained 1240 isolates derived from the sow milk microbiota by the culturomics of Continuous Culture and Interval Sampling. These isolates were grouped into 271 bacterial taxa in the CD-HIT analysis based on a nonredundant set of sequences of the V1-V5 region of 16S rDNA gene with a cutoff value of 99% identity for classification. The Silva version 132 16S rRNA database, NCBI nucleotide collection (nr/nt) database and DAIRYdb reference database were used to classify these prokaryotes into two categories: suspected new species and previously identified prokaryote species (including known species from dairy products). A phylogenetic tree was built based on the calculated distances between pairs of

sequences (**Suppl. Data 1**). The results showed that 151 taxa were assigned to previously described species (**black, Fig. S1**), while the other 120 taxa could not be assigned to any known species (**blue, Fig. S1**), which were defined as suspected new species. The alignment against the DAIRYdb reference database revealed that 107 out of the 271 taxa were assigned to species found in dairy products (**red dots, Fig. S1**). Specifically, twenty-three species were assigned to previously described species found in both the DAIRYdb reference database and Silva version 132 16S rRNA sequences database or the NCBI nucleotide collection (nr/nt) database, including *A. lwoffii*, *Acinetobacter* sp. , *C. perfringens*, *Pelomonas*, *Enterococcus* sp., *E. durans*, *L. amylovorus*, *L. taiwanensis*, *L. garvieae*, *L. mesenteroides*, *Enterobacteriaceae* bacterium DHL-32, *Lactobacillales* bacterium, *Lactobacillaceae* bacterium, *Streptococcaceae* bacterium, *S. enterica* and *Salmonella* sp. Nine species were suspected new species and also could not be assigned to any known species in the DAIRYdb reference database. In addition, five species were assigned to a previously described species in the Silva version 132 16S rRNA database or NCBI nucleotide collection (nr/nt) database, but not in the DAIRYdb database, including *Acidovorax* sp. SEPRH9, *S. hyovaginalis*, *S. mitis*, *Streptococcus* sp. S2 and *S. thoraltensis* (**Fig. 2**).

A total of 922 out of 1240 isolates belong to *Lactobacillales* (**Suppl. Data 2**), and this group was dominated by *Lactococcus lactis*, which was roughly consistent with the microbiota composition of sow milk during lactation according to a recent sequencing report [17]. The genera *Staphylococci* and *Streptococci* represented 5.81% and 4.03%

of the total bacterial isolates, respectively. These results supported a view that commensal *Staphylococci* and *Streptococci* commonly occur in breast milk [18, 19], which may originate from the maternal skin [20]. The collection of 1240 isolates obtained through these efforts provides insight into the diversity of sow milk microbiota and enables more studies on their function associated with mammalian health.

Screening of candidate probiotic *P. pentosaceus* strains

To explore health-promoting bacterial genera, which are not limited to traditional probiotics of *Lactobacillus* and *Bifidobacterium*, we focused on *Pediococcus*. Antagonistic activity against pathogens to control their spread is a prerequisite for a potential probiotic [21]. Piglets commonly encounter pathogens on farms at increasing frequencies [22], including *Salmonella typhimurium* (*S. typhimurium*) [23], enterohemorrhagic *Escherichia coli* (EHEC) [24], enterotoxigenic *Escherichia coli* (ETEC) [21], *Klebsiella pneumoniae* (*K. pneumoniae*) [25], *Aeromonas punctate* (*A. punctate*) [26], *Staphylococcus aureus* (*S. aureus*) [27], *Listeria monocytogenes* (*L. monocytogenes*) [2] and *Clostridium perfringens* (*C. perfringens*) [28]. The fluctuating size of the inhibition zone against pathogens indicates the inhibitory activities of different *P. pentosaceus* isolates against both gram-negative and gram-positive pathogenic bacteria (Fig. 3a), revealing strain-specific antimicrobial activity against different bacteria [29]. The morphology of the top ten strains was observed (Fig. S2), and they were then used in subsequent tests.

For these *P. pentosaceus* strains with strong antimicrobial activity, we next used a simple animal model to rapidly screen bacteria that show potent antioxidant activity *in vivo* (**Fig. 3b**). Considering the similarities of the intestinal development with mammals and the cost of the mouse model [30, 31], *Drosophila* could be an appropriate model to evaluate the ability of more bacteria in protection of hosts from reactive oxygen species (ROS), whose accumulation cause damage to the health of both *Drosophila* and mammals [30, 32]. In this study, resistance to paraquat was used as a measure of free radical scavenging activity in the *Drosophila* system [32]. After paraquat treatment for 45 hours, the flies colonized with *P. pentosaceus* SMM914 showed a significantly elevated survival rate in response to paraquat challenge ($p < 0.05$, log-rank test) (**Fig. 3c**).

SMM914 showed a strong ability to inhibit the proliferation of *S. aureus*, while almost one-third of *P. pentosaceus* strains exhibited no antimicrobial ability toward this pathogen. Through the sodium hydroxide neutralization reaction, the antibacterial effect of SMM914 against *S. aureus* and ETEC was abolished, which proved its bacteriostatic effect was mainly due to the presence of organic acids (**Fig. S3**). The time curves for growth and pH of SMM914 were measured. The results revealed that the strain entered the stationary phase after 12 h of fermentation, while the pH was stabilized at approximately 3.9 after 36 h of fermentation (**Fig. S4 a, b**). A whole genome of SMM914 was sequenced, which generated 28 contigs and a total size of 1,924,818 bp genome assembly with G+C content 37.14%. A total of 1985 coding DNA

sequences, 62 RNA (including 54 tRNAs, 8 rRNAs), and 2 CRISPRs were identified in NCBI prokaryotic genome annotation pipeline. By analyzing the genome (Fig. S5), we obtained some clues about its probiotic properties. Genes related to organic acid, thioredoxin, and exopolysaccharide, which have potential of disease-resistance [33] or oxidative stress resistance activities [34, 35], were annotated in the genome (Suppl. Data 3).

Growth performance and serum biochemical parameters in the pig-feeding trial

Because SMM914 showed bacteriostatic activity and the robust resistance to paraquat-induced stress in *Drosophila*, it was selected to feed the piglets in low-dose (LD) or high-dose (HD) groups prior to early weaning (Fig. 4a). Regarding visceral indices, a higher heart coefficient, which is the relative weight of the heart [36], was observed in piglets treated with SMM914 ($p < 0.05$, $n = 7$) (Fig. S6). The heart coefficient has been reported to be negatively associated with oxidative stress via changes in angiotensin II-aldosterone-brain natriuretic peptide [36]. We speculated that the increased heart coefficient observed in this study could be an indicator of alleviated stress.

When constantly challenged by various endogenous or exogenous free radicals, the liver in vertebrates is susceptible to damages [23, 37]. Weaning is frequently associated with liver injury [38]. Results of blood biochemical parameters revealed that total protein (TP) and albumin (ALB) levels decreased in low dose of SMM914 (Fig. S7). Moreover, treatment with SMM914 no matter in low concentration or high concentration reduced alanine aminotransferase (ALT) and lactate dehydrogenase

(LDH) levels (**Fig. 4b**). The elevation of ALT and LDH activities in serum can be used as biomarkers of hepatic disorders under oxidative stress [39, 40]. Our findings demonstrated that SMM914 administration led to an amelioration on liver injury during weaning.

Pretreatment of piglets with SMM914 induces the Nrf2-Keap1 antioxidant signaling pathway

To discover the molecular mechanisms that might underlie the antioxidative effect, we investigated the alteration of the nuclear factor (erythroid-derived)-like 2 (Nrf2) signaling pathway, a conserved signaling pathway across metazoans [41], in the liver of piglets by western blotting analysis and enzyme activity assays. Kelch-like ECH-associated protein 1 (Keap1) is a specific repressor of Nrf2 via tight binding. Antioxidant metabolites can contribute to the dissociation of Keap1 and Nrf2 complex, promoting Nrf2 movement into the nucleus. Nrf2 transfers from the cytosol to the nucleus, resulting in the coordinated transcriptional upregulation of a battery of antioxidant enzymes and detoxifying proteins [42]. As expected, in the western blotting analysis of this study, the protein level of Keap1 was remarkably suppressed in piglets receiving SMM914 ($p < 0.05$). We found that SMM914 not only markedly increased the intranuclear protein expression level of Nrf2 but also led to elevated protein levels of NADPH quinineoxidoreductase-1 (NQO-1), catalase (CAT), hemeoxygenase-1 (HO-1) and superoxide dismutase (SOD) in a concentration-dependent manner (**Fig.**

4c-f & Fig. S8). CAT and SOD are widely recognized as important endogenous antioxidant enzymes that scavenge hydroxyl and superoxide anion radicals [43, 44]. In the enzyme activity assays, the HD group simultaneously increased glutathione peroxidase (GSH-Px) activity, CAT activity and SOD activity ($p < 0.05$) in livers (**Fig. 4g**). Additionally, the HD group showed a significant decrease in malondialdehyde (MDA), a product of lipid peroxidation, in livers compared with the control group. The western blotting data combined with enzyme activity tests suggested that SMM914 functions as a promising probiotic conferring antioxidant capacity through activation of the Nrf2-Keap1 antioxidant signaling pathway in piglets.

The altered pathways of amino acid metabolism and lipid metabolism in plasma

To provide a better understanding of the antioxidant effect of SMM914, we further examined the metabolic profiles of blood plasma from the three groups ($n = 7$). PLS-DA plots showed separated clusters with an optimal goodness of fit ($R^2 = 0.996$, $Q^2 = 0.681$ (**Fig. 5a**); $R^2 = 0.994$, $Q^2 = 0.479$ (**Fig. 5b**)), indicating that the models were suitable and reliable for prediction. The KEGG enrichment of differential metabolites revealed that the pathways of lipid metabolism and amino acid metabolism were the main perturbed metabolic pathways between the HD group and the control group. The possible protective effect of SMM914 on the weaned piglets is depicted in **Fig. 5c**.

In the lipid metabolism, decreased levels of cortisol and corticosterone (**Fig. 5d**) were observed in plasma, which are classic stress hormones and related to steroid hormone biosynthesis. Excessive stress can cause the development of neurological disorders [45].

Under psychological and emotional stress conditions, corticosterone and cortisol induce the oxidative load in the brain, with a significant increase in pro-oxidant markers in constantly changing environments [46]. After maternal separation, separation anxiety in human infants is an inevitable phenomenon at weaning that may raise cortisol levels and even alter the gut microbiota composition through the gut-brain axis [47, 48]. Meanwhile, the intensities of cholic acid, taurochenodeoxycholate and glycochenodeoxycholate (**Fig. 5e**) were also decreased in the LD group to 0.43-fold ($p < 0.05$), 0.58-fold ($p = 0.08$) and 0.47-fold ($p < 0.05$), respectively, compared to the levels in the control.

Conversely, in amino acid metabolism, several critical antioxidant metabolites (cysteine-S-sulfate, DL-methionine sulfoxide, L-methionine) closely related to cysteine and methionine metabolism were significantly increased to 1.41-2.03-fold in the HD group compared with the control group ($p < 0.05$) (**Fig. 5f**). D-proline, L-proline and L-glutamate, which are involved in arginine and proline metabolism, were increased in the LD group compared with the control group to 1.43-fold ($p < 0.05$), 1.30-fold ($p < 0.05$) and 1.34-fold ($p = 0.056$), respectively (**Fig. 5g**). In the glycine, serine and threonine pathway, choline was also significantly increased to 1.18-fold in the LD group ($p < 0.05$) compared with the control group (**Fig. 5h**). These three amino acid pathways are closely overlapped through several metabolites (**Fig. 5c**).

The reshaped colon microbiota in piglets by SMM914

The changeover from milk to solid feed strongly influences the development of the gut microbiota [49]. Disorders in the composition of the microbiota can induce oxidative

stress and chronic metabolic diseases through a liver-gut axis [50, 51]. In this study, colonic microbiota was further investigated by using 16S rDNA gene amplicon sequencing. No differences were observed among the control and treated groups in terms of α -diversity (**Fig. S9 a**). All samples from weaned piglets approached the saturation plateau based on Shannon-Wiener rarefaction curves (**Fig. S9 b**), suggesting that the sampling was sufficient for nearly all bacterial species. The shared and specific genera are shown in a Venn diagram (**Fig. 6a**). The bacterial community of the three groups shared 89 genera. There were 20 unique genera in the HD group, including *Akkermansia*. Eleven genera, including *Dorea* and *Lachnospiraceae_AC2044_group*, were detected in the LD and HD groups but not in the control group. To intuitively measure the extent of the similarity of the overall microbiota, the results of PCA based on distance revealed a separate clustering of samples between the HD group and the control group ($p = 0.001$, ANOSIM, comparison between the HD and control samples), but the colonic specimens of the LD group were not separated from the control group ($p = 0.274$, ANOSIM, comparison between the LD and control samples). These results indicated that the high dose of SMM914 reshaped the microbiota structure of the colon (**Fig. 6b**).

The HD group showed distinct bacterial communities compared to the control group both at family and genus level (**Fig. 6c** and **Fig. S11**). At the family level, the relative abundance of the family *Lactobacillaceae* ($p < 0.05$) and the genus *Lactobacillus* ($p < 0.05$) were both increased in HD group compared with the control group (**Fig. 7a**). For

the family *Christensenellaceae* and *Ruminococcaceae*, which are beneficial bacteria reported before [52, 53], the means of relative abundance in the LD group and HD group were increased by 89.57%, 110.27% and 804.9%, 21.59%, respectively, compared with the control group (Fig. S10). The genus *Christensenellaceae_R_7_group* ($p < 0.05$) and the genus *Ruminococcaceae* UCG-005 ($p < 0.01$) showed enrichment in the HD group (Fig. S10). Conversely, at the family level, the relative abundances of *Bacteroidaceae* in the HD group were decreased by 86.7% (Fig. S11). Specifically, the genus *Bacteroides* was observed to decrease in the HD group compared with the control group ($p < 0.05$).

Discussion

In recent years, breeders have attached great importance to improving sow nutrition during pregnancy and lactation from the source, rather than merely purchasing expensive creep feed for piglets at a later stage to effectively increase production and economic benefits. Here, we put forth the concept of Sow and Piglet Integration (SPI), which is an integral nutritional regulation scheme based on the physiological stages of sows and piglets on pig farms and the interrelationship of the microbiota between the two generations. Although the early weaning technique at 21 days is beneficial for sow productivity in intensive animal husbandry, this strategy leads to severe stress in piglets [54]. At this life stage, piglets experience a series of stressors, including separation from the mother, transport, the mixing of litters, diet transition and frequent exposure to potential pathogens, with a direct impact on breeders' profitability [55]. To face challenges from pathogens and calls for reducing the use of antibiotics, some farmers

in American have extended the lactation period to 25 days. Another promising alternative to antibiotics in animal feed is using probiotics to improve the absorption of the nutrients [56]. During the initial development of mammalian neonates, breast milk was a nutritious food and a natural reservoir of probiotics [57, 58] that adapted to satisfy neonatal needs in evolution [59]. In the present work, we focused on sow milk microbiota by using the culturomics of Continuous Culture and Interval Sampling. This identification work is an important step that could reveal the LAB repertoire of sow milk to analyze the relationships between the maternal microbiota and that of the offspring and to deeply understand the concept of SPI.

Nowadays, metagenomic approaches have generated countless sequences that have not been assigned to living purified bacteria, while the culturomics strategy has been reported to double the species number of microorganisms isolated from the human gut [60]. Although the culturomics approach is time and labor consuming, it is a more challenging task to evaluate the functional properties of such enormous isolates. To screen probiotics from sow milk quickly, we first evaluated their antagonistic activity against pathogens *in vitro*. Next, we selected several strains and further explored their antioxidant capacities *in vivo*. A rapid *Drosophila* screening model combined with a subsequent piglet experiment was used to screen out suitable probiotics from a collection of sow milk microbiota. In piglets treated with *P. pentosaceus* SMM914, we observed decreased ALT and TP levels. The shift observed in our study was in accordance with another report that the administration of *P. pentosaceus* LI05 significantly prevented increases in TP and ALT in the context of acute liver failure [61]. Besides, activation of the Nrf2-Keap1 signaling pathway was observed by western blotting analysis and enzyme activity assays in the liver of piglets. Nrf2 and its

endogenous inhibitor, Keap1, are conserved defense mechanism for regulating antioxidative activities [62].

In untargeted metabolism of plasma, several metabolic pathways were perturbed, and they can be mainly classified into lipid metabolism and amino acid metabolism. In the cysteine and methionine metabolism, L-methionine is a limiting amino acid in early lactation [63] associated with various key physiologic events. The increased availability of L-methionine in early lactation was reported to have positive effects on plasma lipid metabolism and overall antioxidant status [64]. Methionine sulfoxide is also biologically available as a methionine source. High methionine bioavailability is likely to increase the entry of L-methionine into the one-carbon metabolism cycle, where S-Adenosyl-L-methionine is then used to generate S-Methyl-5'-thioadenosine and 1-Aminocyclopropane-1-carboxylic acid. Through the transsulfuration and transmethylation pathway, L-methionine is converted into L-cysteine which can serve as a substrate for glutathione synthesis. Glutathione is an endogenous sulfur-containing antioxidant [65, 66]. Another substrate for glutathione synthesis is glutamate. In mammals, L-glutamate is an abundant amino acid in milk that is required for the synthesis of glutathione and alleviates oxidative stress by increasing antioxidant enzyme activities [45, 67]. Glutathione is decomposed into L-gamma-glutamyl amino acid, and L-gamma-glutamyl amino acid is further converted to pyroglutamic acid. A high level of pyroglutamic acid reflected glutathione deficiency and could be an indicator of the oxidative state [68]. However, in our study, the concentration of pyroglutamic acid was significantly downregulated. Collectively, the altered intensities of these metabolites might be conducive to the accumulation of glutathione. The cysteine and methionine metabolism is overlapped with choline metabolism tightly

because choline can serve as the substrate for L-methionine synthesis. Choline is an essential vitamin for humans and other mammals to regulate amino acid metabolism [69], particularly when L-methionine levels are not sufficient around parturition [70]. It has been established that choline deficiency induces the generation of ROS [71] and oxidative damage in rats [72], ruminants [73] and fishes [74]. Moreover, dietary supplementation with choline enhances the antioxidative capacity in pigs with intrauterine growth restriction [75]. New evidence has shown that choline deficiency-induced oxidative damage was associated with changes in the transcription of antioxidant enzymes and Nrf2 signaling in the liver [74].

In the primary bile acid biosynthesis and the bile secretion pathways, there are variable derivatives of cholic acid and deoxycholic acid which function as signaling molecules for the induction of oxidative stress [76, 77]. Deoxycholic acid can combine with taurine or glycine to form taurochenodeoxycholate or glycochenodeoxycholate, respectively that stimulate the phosphorylation of NADPH oxidase and the formation of ROS [78]. In addition, classic stress hormones including cortisol and corticosterone were also found to be decreased. Hypersecretion of serum cortisol concentrations was an indicator of stress in the pig [79] and contributed directly to the pathology of anxiety [80]. Kulak *et al* reported that administration of corticosterone decreased the concentration of GSH antioxidant defense [81].

We observed that high dose of SMM914 shaped the piglet microbiota. The microbiota of HD piglets was enriched with the *Lactobacillus*, *Christensenellaceae_R_7_group* and *Ruminococcaceae* UCG-005 genera, while those of piglets in control group exhibit higher relative abundances of genera from *Bacteroidaceae* and *Prevotellaceae* families.

The depletion of the *Lactobacillus* in the gut environment has been associated to oxidative damage, and members of this genus are commonly used as probiotics in animal husbandry by upregulating the expression of glutathione reductase and glutathione S-transferase during the suckling period [82, 83]. Moreover, methionine was reported to attenuate oxidative stress in rats through high abundances of *Lactobacillus* and *Lachnospiraceae* [84]. The *Ruminococcaceae* family is always negatively related to disease severity [53]. The *Christensenellaceae_R_7_group* plays a positive role in intestinal immunomodulation [85]. On the contrary, *Bacteroidaceae* and *Prevotellaceae* have been proved to be associated with several diseases. For example, chitosan oligosaccharides increase the antioxidant capacity by inhibiting the abundance of *Bacteroides* and *Prevotella* [83]. The dysbiotic microbiota of inflammatory bowel disease (IBD) is mostly characterized by an increase in *Prevotellaceae* and a reduction in *Ruminococcaceae* [86]. Improving the cellular antioxidant potential is a promising approach for prevention of IBD. In this work, it appeared that SMM914 administration in piglets positively regulate intestinal microbiota during the transition of early-weaned stress.

Conclusion

Here, we focused on bacteria profile from sow milk by using the culturomics of Continuous Culture and Interval Sampling and established a set of methods for rapid screening the beneficial microorganisms due to the importance of swine for biomedical

research and agriculture. It was indicated that SMM914 could activate the Nrf2-Keap1 antioxidant signaling pathway in piglets and relieve oxidative stress by the elevated level of amino acid metabolites in plasma and the increased beneficial microbiota in colon, although we cannot provide comprehensive mechanistic linkages that might participate in the process. The study has shed light on the effects of sow milk microbiota for deeply understanding the relationships between the sow and offspring. However, researchers are needed to further strengthen the research on safety and stability assessments of SMM914 to pave the way for the application of maternal probiotics in animal husbandry.

Methods

Culture media and bacterial strain isolation.

A total of nine healthy second-parity sows with similar breeding dates raised on a pig breeding farm (Changsha, China) were employed in this study. The sows received no antibiotics within the 4 weeks prior to breast milk sampling. The areolar skin around the teats was successively swabbed with alcohol tampons and warm saline-lubricated sterile swabs. Using sterile tubes, fresh ordinary milk was collected from six sows during lactation (**Fig. S1, step 1**).

Considering that bacterial populations can survive through cell death and the recycling of dead cells [87], the continuous culture and intermittent sampling was performed for 30 days in an anaerobic incubator ($N_2 = 90\%$, $CO_2 = 5\%$, and $H_2 = 5\%$) at $37^\circ C$ (**Fig.**

S1, step 2). Using sow milk as an inoculum, MRS (Oxoid, Code# CM0359, UK) [88], M17 (Oxoid, Code# CM0817, UK) [88], TPY (Hopebio, Code# HB0397, China) [89] and GYP (Hopebio, Code# HB8539, China) [90] media were utilized to cover as much LAB diversity as possible. The bacterial cells were harvested every other day by centrifugation at 4000 g for 10 min, and the cell pellets were resuspended in sterile normal saline. Then, the cells were spread on agar plates and anaerobically cultured in a DG250 Anaerobic workstation (DWS, UK) at 37°C for 24 h-72 h. The identification of the isolates was first carried out by using morphological method. Colonies were streaked on agar plates, and all isolates were stocked in 25% (v/v) glycerol broth at -80°C at the College of Life Science, Hunan Normal University, China (**Fig. S1, step 4**). SMM914 was deposited at the China General Microbiological Culture Collection Center (CGMCC20160).

Characterization and classification of the isolated bacteria

After isolation and purification, DNA was extracted from pure cultures. The V1-V5 region of 16S rDNA genes was amplified using Takara PrimerSTAR Max DNA Polymerase with a pair of LAB-specific primers, 15f (5'-GCTCAGGAYGAACGCGYGG -3') and 687r (5'- CACCGCTACACATGRADTTC-3') for the identification of the isolates [91]. The PCR-amplified products were identified by Sanger sequencing (Sangon Biotech Ltd., China). The sequencing error-prone areas (50 bp) at both ends were removed. The partial 16S rDNA sequences were searched

against the NCBI nucleotide collection (nr/nt) database using BLASTN. The best match for each sequence was selected, and the taxonomy was assigned (**Fig. S1, step 3**).

The nonredundant set of 16S rDNA datasets was clustered by using CD-HIT with sequence identity of 0.99 [92]. The phylogenetic relationship between isolates was determined by aligning the nonredundant set of 16S rDNA gene sequences to construct a maximum-likelihood tree by using FastTree.

All 16S rDNA sequences alignment were aligned against the Silva version 132 16S rRNA database, the NCBI nucleotide collection (nr/nt) database and the DAIRYdb reference database using BLASTN with a threshold of 1e-05 e-value, 99% coverage and 99% cutoff. A suspected new species was defined below these values against the Silva version 132 16S rRNA database and NCBI nucleotide collection (nr/nt) database [92]. The sequencing read data have been deposited in the National Center for Biotechnology Information Sequence Read Archive (**Suppl. data S2**).

Antibacterial assay

The following indicator strains were used: *S. typhimurium* ATCC 14028, EHEC ATCC 43894, ETEC O149: K88, *K. pneumoniae* ATCC 13883, *A. punctata subsp. Caviae* ATCC 15468, *S. aureus* ATCC 25923, *L. monocytogenes* ATCC 19115 and *C. perfringens* ATCC 13124.

Agar well diffusion assays [93] were utilized to evaluate antimicrobial activity against these indicator pathogenic strains *in vitro*. Briefly, pathogens were grown in Luria-Bertani (LB) broth at 37°C for 8 h and then diluted at a volume ratio of 20 µL to 4 mL

LB and mixed well. Fifty microliters of the diluted liquid were spread evenly on each soft LB agar plate containing 0.8% agar. The residual liquid was evaporated on a ventilated clean bench. Next, holes were punched in each agar plate via sterile iron pipettes, with a depth of 6 mm and a diameter of 5 mm. *P. pentosaceus* strains were grown in MRS broth at 37°C for 18 h. Then, supernatants of fermentation broth were precisely added to the holes with a 30 µL volume per well. The central well of each plate was filled with 30 µL MRS broth as the negative control. After 48 h of incubation at 37°C, antibacterial activity was observed as a halo of inhibition in the bacterial lawn formed around the sample, and the diameter of the zones of inhibition was measured. The evaluation of each sample was repeated in triplicate.

Paraquat resistance assays in *Drosophila*

The *Drosophila* experiments were conducted under a 12 h light:12 h dark cycle at 25°C on cornmeal-molasses medium. Six-day-old mated female *Drosophila* *w¹¹¹⁸* were collected under CO₂ anesthesia and starved for 2 h. Each group consisted of 3 vials, and each vial contained 20 female flies. In the bacterial association assays, the colony-forming units (CFUs) of *P. pentosaceus* were enumerated using MRS agar plates following standard microbiological procedures. Groups of adult female flies were colonized with pure cultures (1×10^{10} CFUs) of the *P. pentosaceus* strains for 3 days, including SMM847, SMM853, SMM862, SMM867, SMM881, SMM906, SMM907, SMM908, SMM914 and SMM918 isolates. Distilled water without *P. pentosaceus* was used as a negative control. Then, these flies were transferred to vials containing 2 pieces

of Whatman paper soaked with 200 μ L 5% (w/v) sucrose containing 12 mmol/mL paraquat (methyl viologen dichloride, Cat# 856177, Sigma-Aldrich, USA). Each group was supplied with fresh paraquat vials every day. The 12 mmol/mL concentration was chosen because it was empirically shown that this concentration was lethal to more than 80% of female *w¹¹¹⁸* flies within 2 days.

Piglet feeding trial and tissue sampling

The Animal Care and Use Committee of the Institute of Subtropical Agriculture, Chinese Academy of Sciences, reviewed and approved the experimental procedures involving piglets. As described in Jun Hu *et al* [91], a total of 54 newborn suckling piglets (Landrace $\times$ Yorkshire) were chosen from nine second-parity sows and randomly assigned to three treatments ($n = 18$) among each litter, including a control group (physiological saline, 2.0 mL each time, Control), a low-dose SMM914 solution group (10^8 CFU/ml, 2.0 mL each time, LD group) and a high-dose SMM914 solution group (10^9 CFU/ml, 2.0 mL each time, HD group). The solution of saline or bacterial cells was infused into each piglet's mouth by a syringe without a needle. All suckling piglets were subjected to oral gavage every other day from the age of 10 to 18 days and were weaned at 21 days.

Sample collection

Piglets per group from each of the 7 litters were selected to be euthanized for sampling at 28 days. Ten milliliters of blood were collected into heparin sodium anticoagulant tubes via direct cardiac puncture immediately after death and subjected to untargeted

metabolism analyses. Another 10 mL of blood was collected in vacuum tubes and centrifuged at 3000 rpm at 4°C for 10 min. The serum samples were kept at -80°C until analysis. After the opening of the abdomen, tissues including livers, spleens, kidneys, and hearts were weighed, dissected, and then snap-frozen in liquid nitrogen. Colon contents were stored at -80°C until the extraction of bacterial DNA.

Serum concentrations of parameters reflecting lipid (cholesterol, triglyceride, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, lipase), protein (total protein, TP; albumin, ALB), carbohydrate (glucose, alpha-amylase, lactic dehydrogenase) metabolism, liver functionality including total bilirubin (TBIL), direct bilirubin (DBIL), indirect bilirubin (IBIL), the activity of aspartate transaminase (AST), alanine aminotransferase (ALT), gamma-glutamyl transpeptidase (GGT), serglobulin (GLO), alkaline phosphatase (ALP), lactic dehydrogenase (LDH) and cholinesterase (CHE), as well as kidney functionality including creatinine (CREA), blood urea nitrogen (BUN) and uric acid (UA) were determined using commercial kits according to the manufacturer's instructions (Jiancheng Bioengineering Institute, Nanjing, China) and were identified with a TBA-120FR Automatic Chemistry Biochemistry Hiometer (Hitachi Co., Tokyo, Japan).

Western blotting analysis

Western blotting was performed as previously described with some modification [82]. Liver samples ($n = 6$) were powdered under liquid nitrogen and lysed in radioimmunoprecipitation assay buffer with the protease inhibitor

phenylmethanesulfonyl fluoride (PMSF, Beyotime Biotechnology). The supernatant was obtained by centrifugation at $12,000 \times g$ for 10 min at 4°C. The denatured proteins were separated by 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and then transferred to polyvinylidene fluoride membranes at 200 mA for 1 h. The membranes were blocked with 5% nonfat milk in Tris-buffered saline mixed with 0.5% Tween-20 (TBST) at room temperature for 2 h and then incubated with antibodies against Kelch-like ECH-associated protein 1 (Keap1) (SC-19917, Proteintech, USA), nuclear factor-E2-related factor 2 (Nrf2) (SC-98974, Proteintech, USA), NADPH quinineoxidoreductase-1 (NQO-1) (11451-1-AP, Proteintech, USA), hemeoxygenase-1 (HO-1) (27282-1-AP, Proteintech, USA), catalase (CAT) (66765-1-Ig, Proteintech, USA), Cu/Zn-superoxide dismutase (SOD1) (10269-1-AP, Proteintech, USA), PCNA (60097-1-Ig, Proteintech, USA) or β -actin (SC-47778, Proteintech, USA), which were diluted with 5% nonfat milk in TBST. The membranes were washed 3 times in TBST and then incubated with a secondary antibody. Finally, the membranes were washed with TBST solution and visualized with a chemiluminescence instrument.

Enzyme activity analysis

Liver tissue samples were homogenized in saline, followed by centrifugation ($2500 \times g$, 4°C, 10 min) to obtain supernatants ($n = 7$). Malondialdehyde (MDA), glutathione peroxidase (GSH-Px) and superoxide dismutase (SOD) in livers were determined with commercially available colorimetric diagnostic kits (Nanjing Jiancheng Bioengineering

Institute, China) following the manufacturer's instructions. The procedures were carried out in duplicate with three parallel samples.

Untargeted metabolomics

The plasma samples (100 μ L) were thawed at 4°C and homogenized in 400 μ L of precooled methanol/acetonitrile (1:1, v/v) for 60 s. The untargeted metabolic profiling analysis was conducted by using an ultra-performance liquid chromatography (UPLC) system (1290 Infinity LC, Agilent Technologies, Santa Clara, California, USA) coupled to a quadrupole time-of-flight (TOF) mass spectrometer (Triple TOF 5600, AB SCIEX) with electrospray ionization in positive and negative ionization modes. For chromatographic separation, 2 μ L of the extracted sample was injected by an autosampler system at 4°C at a delivery flow rate of 300 μ L/min into a liquid chromatography column with a column temperature of 25°C. The mobile phase consisted of A (water + 25 mM ammonium acetate + 25 mM ammonia hydroxide) and B (acetonitrile). The gradient was 95% B and 5% A for 1 min, with a linear reduction to 65% B and 35% A over 13 min, a reduction to 40% B and 60% A over 2 min, maintenance for 2 min and an increase to 95% B and 5% A over 0.1 min, with a 5 min re-equilibration period. Before injection, quality control samples were used to monitor the stability and repeatability of the data produced by the instrument. The screening criteria for differential metabolites were based on a variable importance projection score > 1 and $p < 0.05$ (Student's t-test). The metabolites were analyzed by comparing the molecular ions with compounds in the available biochemical databases, and the

pathway analysis of the identified compounds was conducted using the KEGG website (<http://www.genome.jp/kegg>).

16S rDNA gene amplicon sequencing

Intestinal digesta samples were collected after sacrifice. Bacterial DNA was extracted using a QIAamp DNA Stool Kit (Qiagen, Gaithersburg, MD, USA) according to the provided protocols ($n = 7$). The V3-V4 region of the bacterial 16S rDNA gene was amplified by PCR in triplicate in a 20 μ L mixture (2 min at 95°C, followed by 25 cycles of 30 s at 95°C, 30 s at 55°C, 30 s at 72°C and 5 min at 72°C). The primers 314-F (5'-CCTAYGGGRBGCASCAG-3') and 806-R (5'-GGACTACNNGGGTATCTAAT-3') were used to target the hypervariable region. Using the AxyPrep DNA Gel Extraction Kit (Axygen, Union City, CA, USA), the PCR products were purified and then quantified by QuantiFluorTM-ST (Promega, Madison, WI, USA). Library quality was assessed, and the amplicon library was sequenced on the Illumina MiSeq platform. The sequences were merged with FLASH (v1.2.7) [94] and quality filtered with fastp (0.19.6) [95]. Then the high-quality sequences were de-noised using DADA2 [96] plugin in the QIIME2 [97] (version 2020.2) pipeline with recommended parameters to obtain amplicon sequence variants (ASVs). To minimize the effects of sequencing depth, the number of reads from each sample was rarefied to 23834 for comparing all samples at the same sequencing level. The taxonomy of each 16S rDNA sequence was aligned against the bacteria database of SILVA [98]. QIIME2 was also used for the analysis of alpha diversity and beta diversity.

Statistical analysis and visualization

Cladograms of the Sow Milk Bacterial Collection were visualized with GraPhlAn v.0.9.7 [99], and the scripts were reused from a previously published paper [100]. SPSS software (version 19.0; IBM Corp., Chicago, IL, USA) was used to evaluate piglet experiment results with one-way analysis of variance and Duncan's multiple comparison test to determine the statistical significance of the differences among treatment groups. Different letters in a same graph indicate significant statistical differences ($p < 0.05$).

Accession codes

The numbers and GenBank IDs of 16S rRNA genes of all smBC isolates are listed in Suppl. Data s2. The Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession JAEMVT000000000.

Supplementary information

Additional file 1: Figure S1. The maximum-likelihood tree of the bacterial taxa. The 271 bacterial taxa were clustered based on 16S rDNA gene identity with a cut-off value of 99% for different taxa in the CD-HIT analysis. The closely related neighboring species of each taxon are listed next to the taxon numbers of the sow milk microbiota (SMM). The suspected new species are indicated in blue, and the species in the dairy products are indicated in red dots. **Supplementary Data S1.** The 16S rRNA sequences and classification of all isolates in smBC

Additional file 2: Figure S2. Light microscopy images obtained from *P. pentosaceus* strains using a 63x oil immersion objective. **Supplementary Data S2.** The alignment of 1240 16S rRNA sequences against the Silva version132 16S rRNA database, NCBI nucleotide collection (nr/nt) database and DAIRYdb database using BLASTN

Additional file 3: Figure S3. The antimicrobial activity of SMM914. (a) The pH of SMM914 products were determined via test papers. (b) Inhibitory effects of SMM914 against *S. aureus* and ETEC. The circular wells were filled with different products of SMM914, including MRS medium (1&2) and the cell-free supernatant without any treatment (3&4), with 15 min heat inactivation (5&6), with pH adjustment using NaOH (7&8) or with 1 mg/mL protease K treatment (9&10). The substances (1, 3, 5, 7 and 9) were obtained from the fermentation broth after cultivation under anaerobic condition, while others (2, 4, 6, 8 and 10) were obtained from culture under aerobic condition.

Figure S4. Growth curves and pH value curve of SMM914 after anaerobic and aerobic fermentation. (a) The optical density at 600 nm (OD₆₀₀) was measured from a starting OD₆₀₀ of 0.085. (b) The pH value was recorded with a pH meter. Data are the mean $\pm$ s.e.m, $n = 3$.

Additional file 4: Figure S5. The annotation and comparison of *P. pentosaceus* SMM914 genome. (a) The distribution of predicted CDSs of SMM914 in different categories of metabolic function by online software RAST. (b) A full genome comparison analysis of *P. pentosaceus* SMM914 with other *P. pentosaceus* strains, including *Pediococcus pentosaceus* SRCM100194, *Pediococcus pentosaceus*

GDIAS001, *Pediococcus pentosaceus* SL001 and *Pediococcus pentosaceus* SRCM102736, visualized by BRIG software. Colors display the percentage of sequence identity based on BLASTN. Two inner rings indicate the GC skew and the G + C content. The innermost circle shows the genome coordinates. **Supplementary Data S3.** Genes of SMM914 and predicted proteins by Pfam protein database

Additional file 5: Figure S6. Effects of SMM914 on (a) growth performance and (b) organ relative weight. Data are the mean $\pm$ s.e.m. Statistical analysis was conducted by using one-way ANOVA. Data not sharing the same letter in each point were significantly different ($p < 0.05$). **Figure S7.** Effects of SMM914 on serum biochemical parameters. Data are the mean $\pm$ s.e.m ($n = 7$). One-way ANOVA with adjustment for multiple comparisons was conducted. ^{a,b} Within a variable, values with different superscripts differ ($p < 0.05$).

Additional file 6: Figure S9. (a) Alpha diversity comparisons of the gut microbiomes including Ace, Chao, Sobs index, Shannon index, Shannoneven index, Simpson index, Simpstoneven index and PD whole tree index which were analyzed using Kruskal-Wallis H test and Tukey-Kramer post hoc test with 95% confidence level. Data are the mean $\pm$ s.d ($n = 7$). *ns*, not significant differences. (b) Rarefaction curves for Shannon indices at genus level in the colon contents. **Figure S10.** (a) Linear discriminant analysis effect size 10 was set up with 0.05 as the alpha value for Wilcoxon test and then selected with a threshold value of 4 from phylum to genus level. (b) Differences in relative abundance of the important and variable genera in colonic

microbiota among three group. Each data represented the mean and SEM of relative abundance of each genus ($n = 7$). One-way analysis of variance and Duncan's multiple comparison test to determine the statistical. Different letters in the same graph indicate significant statistical differences ($p < 0.05$). **Figure S11** (a) Bar charts of relative abundance at the family level in the control and treated groups obtained by sequencing of 16S rDNA V3-V4 region. (b) Comparison of microbial families between high dose group and control. Significantly different domains were tested by Wilcoxon rank-sum test. * $p < 0.05$; ** $p < 0.01$.

Abbreviations

SPI: Sow and Piglet Integration; Nrf2: nuclear factor-E2-related factor 2; Keap1: Kelch-like ECH-associated protein 1; NQO-1: NADPH quinone oxidoreductase-1; HO-1: heme oxygenase-1; CAT: catalase; MDA: Malondialdehyde; GSH-Px: glutathione peroxidase; SOD: superoxide dismutase; LAB: lactic acid bacteria; SMM: sow milk microbiota.

Acknowledgements

We are grateful to Jun Hu and Liyuan Cai from Huazhong Agricultural University for their constructive suggestions. We also thank Zhuomin Han for editorial assistance.

Funding

This work was supported by the National Natural Science Foundation of China (31700004), 2019 CAST Outstanding Chinese and Foreign Youth Exchange Program of China Association of Science and Technology, Hunan Province College Students

Research Learning and Innovative Experiment Project (S201910542041), Hunan Provincial Science and Technology Department (2019RS5001), Natural Science Foundation of Hunan Province (2019JJ50369) Construction of Innovative Provinces in Hunan Province (2019RS3022) and the National Students Platform for Innovation and Entrepreneurship Training Program (201810542034, 2020056).

Authors' contributions

JY and YLY contributed in conceiving the experiments. LLW designed and performed most experiments. XLZ, QYH, YX and CML collected the milk samples. THC, YNQ and YWC conducted antimicrobial activity experiment. QHL, YWC, QYW, and JLH performed paraquat resistance assays. CML, YCD, LW, PH, PL and TYY contributed to piglet-feeding trial. YXL, GHZ, JZL, HHC, PH, CC and HSY contributed the data analysis. ZZZ, JZL, HSY, CNW and QYW revised the work. JY and LLW drafted the work. All authors read and approved the manuscript.

Availability of data and materials

The raw sequence data reported in this paper have been deposited (PRJCA003003) in the Genome Sequence Archive in the BIG Data Center, Chinese Academy of Sciences, under accession code CRA002995 for bacterial 16S rRNA gene sequencing data, which are publicly accessible at <http://bigd.big.ac.cn/gsa>.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

No conflict of interest exists in the submission of this manuscript, and the manuscript has been approved by all authors for publication. The authors declare that they have no competing interests.

Author details

¹ Laboratory of Animal Nutrition and Human Health, Hunan Provincial Key Laboratory of Animal Intestinal Function and Regulation, Hunan International Joint Laboratory of Animal Intestinal Ecology and Health, College of Life Sciences, Hunan Normal University, Changsha, China.

²Institute of Subtropical Agriculture, Chinese Academy of Sciences, Changsha, China

³The National and Local Joint Engineering Laboratory of Animal Peptide Drug Development, College of Life Sciences, Hunan Normal University, Changsha, China

⁴State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China

⁵College of Animal Science and Technology, Hunan Agricultural University, Changsha, Hunan, China.

⁶Shandong Yihe Feed Co.,Ltd. Yantai Hi-tech Industrial Development Zone, Yantai, Shandong, China.

References

1. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature*. 2018;562(7728):583-8.
2. Breastfeeding and the Use of Human Milk. 1997;100(6):1035-9.
3. Li Z, Wright AG, Yang Y, Si H, Li G. Unique Bacteria Community Composition and Co-occurrence in the Milk of Different Ruminants. *Scientific reports*. 2017;7:40950.
4. Fitzstevens JL, Smith KC, Hagadorn JI, Caimano MJ, Matson AP, Brownell EA. Systematic Review of the Human Milk Microbiota. *Nutr Clin Pract*. 2017;32(3):354-64.
5. Gil-Campos M, López M, Rodríguez-Benítez MV, Romero J, Roncero I, Linares MD, et al. *Lactobacillus fermentum* CECT 5716 is safe and well tolerated in infants of 1-6 months of age: a randomized controlled trial. *Pharmacological research*. 2012;65(2):231-8.
6. Eckert R, Mucha A, Koska M, Szulc T. Relationships Between Chemical Composition of Colostrum and Milk and Rearing Performance of Piglets During a 21-Day Lactation. *Annals of Animal* 2013;13(4):771-81.
7. Meckel KR, Kiraly DD. Maternal microbes support fetal brain wiring. *Nature*. 2020;586(7828):203-5.
8. Fernandez L, Langa S, Martin V, Maldonado A, Jimenez E, Martin R, et al. The human milk microbiota: origin and potential roles in health and disease. *Pharmacological research*. 2013;69(1):1-10.
9. Chen W, Mi J, Lv N, Gao J, Cheng J, Wu R, et al. Lactation Stage-Dependency of the Sow Milk Microbiota. *Front Microbiol*. 2018;9:945.
10. Odamaki T, Bottacini F, Kato K, Mitsuyama E, Yoshida K, Horigome A, et al. Genomic diversity and distribution of *Bifidobacterium longum* subsp. *longum* across the human lifespan. *Scientific reports*. 2018;8(1):85.
11. Solieri L, Rutella GS, Tagliazucchi D. Impact of non-starter lactobacilli on release of peptides with angiotensin-converting enzyme inhibitory and antioxidant activities during bovine milk fermentation. *Food microbiology*. 2015;51:108-16.
12. Chen T, Wang L, Li Q, Long Y, Lin Y, Yin J, et al. Functional probiotics of lactic acid bacteria from Hu sheep milk. *BMC microbiology*. 2020;20(1):228.
13. Altuntas EG, Ayhan K, Peker S, Ayhan B, Demiralp DO. Purification and mass spectrometry based characterization of a pediocin produced by *Pediococcus acidilactici* 13. *Mol Biol Rep*. 2014;41(10):6879-85.
14. Anastasiadou S, Papagianni M, Filiousis G, Ambrosiadis I, Koidis P. Growth and metabolism of a meat isolated strain of *Pediococcus pentosaceus* in submerged fermentation: Purification, characterization and properties of the produced pediocin SM-1. *Enzyme and Microbial Technology*. 2008;43(6):448-54.
15. Porto MC, Kuniyoshi TM, Azevedo PO, Vitolo M, Oliveira RP. *Pediococcus* spp.: An important genus of lactic acid bacteria and pediocin producers. *Biotechnology advances*. 2017;35(3):361-74.

754 16. Wang X, Lu H, Feng Z, Cao J, Fang C, Xu X, et al. Development of Human Breast Milk
755 Microbiota-Associated Mice as a Method to Identify Breast Milk Bacteria Capable of
756 Colonizing Gut. *Frontiers in Microbiology*. 2017;8(1242).

757 17. Liu H, Hou C, Li N, Zhang X, Zhang G, Yang F, et al. Microbial and metabolic alterations
758 in gut microbiota of sows during pregnancy and lactation. *FASEB J*. 2019;33(3):4490-
759 501.

760 18. Carroll L, Osman M, Davies DP, McNeish AS. Bacteriological criteria for feeding raw
761 breast-milk to babies on neonatal units. *Lancet*. 1979;2(8145):732-3.

762 19. West PA, Hewitt JH, Murphy OM. Influence of methods of collection and storage on the
763 bacteriology of human milk. *J Appl Bacteriol*. 1979;46(2):269-77.

764 20. Cacho NT, Harrison NA, Parker LA, Padgett KA, Lemas DJ, Marcial GE, et al.
765 Personalization of the Microbiota of Donor Human Milk with Mother's Own Milk. *Front*
766 *Microbiol*. 2017;8:1470.

767 21. Dubreuil JD, Isaacson RE, Schifferli DM. Animal Enterotoxigenic *Escherichia coli*. *EcoSal*
768 *Plus*. 2016;7(1).

769 22. Barton MD. Antibiotic use in animal feed and its impact on human health. *Nutr Res Rev*.
770 2000;13(2):279-99.

771 23. Lee JW, Choi YC, Kim R, Lee SK. Multiwall Carbon Nanotube-Induced Apoptosis and
772 Antioxidant Gene Expression in the Gills, Liver, and Intestine of *Oryzias latipes*. *Biomed*
773 *Res Int*. 2015;2015:485343.

774 24. Moxley RA, Francis DH, Tamura M, Marx DB, Santiago-Mateo K, Zhao M. Efficacy of
775 Urtoxazumab (TMA-15 Humanized Monoclonal Antibody Specific for Shiga Toxin 2)
776 Against Post-Diarrheal Neurological Sequelae Caused by *Escherichia coli* O157:H7
777 Infection in the Neonatal Gnotobiotic Piglet Model. *Toxins (Basel)*. 2017;9(2).

778 25. Wei H, Shen J, Pang X, Ding D, Zhang Y, Zhang B, et al. Fatal infection in human flora-
779 associated piglets caused by the opportunistic pathogen *Klebsiella pneumoniae* from an
780 apparently healthy human donor. *J Vet Med Sci*. 2008;70(7):715-7.

781 26. Igbinosa IH, Igbinosa EO, Okoh AI. AntibioGram characterization and putative virulence
782 genes in *Aeromonas* species isolated from pig fecal samples. *Environ Sci Pollut Res Int*.
783 2016;23(12):12199-205.

784 27. Kock R, Cuny C. [Multidrug-resistant bacteria in animals and humans]. *Med Klin*
785 *Intensivmed Notfmed*. 2020;115(3):189-97.

786 28. Zheng L, Hu Y, He X, Zhao Y, Xu H. Isolation of swine-derived *Lactobacillus plantarum*
787 and its synergistic antimicrobial and health-promoting properties with ZnO
788 nanoparticles. *J Appl Microbiol*. 2020.

789 29. V R, Kumar R, R.R.B S, Kaushik J, Mann B. Comparative evaluation of selected strains of
790 lactobacilli for the development of antioxidant activity in milk. *Dairy Science and*
791 *Technology*. 2011;92.

792 30. Nie Y, Hu J, Hou Q, Zheng W, Zhang X, Yang T, et al. *Lactobacillus frumenti* improves
793 antioxidant capacity via nitric oxide synthase 1 in intestinal epithelial cells. *Faseb j*.
794 2019;33(10):10705-16.

795 31. Patel PH, Pénalva C, Kardorff M, Roca M, Pavlović B, Thiel A, et al. Damage sensing by a
796 Nox-Ask1-MKK3-p38 signaling pathway mediates regeneration in the adult *Drosophila*
797 midgut. *Nature communications*. 2019;10(1):4365.

798 32. Darby TM, Owens JA, Saeedi BJ, Luo L, Matthews JD, Robinson BS, et al. *Lactococcus*
799 *Lactis* Subsp. *cremoris* Is an Efficacious Beneficial Bacterium that Limits Tissue Injury in
800 the Intestine. *iScience*. 2019;12:356-67.

801 33. Papadopoulos GA, Poutahidis T, Tallarico N, Hardas A, Teliousis K, Arsenos G, et al.
802 Dietary supplementation of encapsulated organic acids enhances performance and
803 modulates immune regulation and morphology of jejunal mucosa in piglets. *Research in*
804 *veterinary science*. 2017;115:174-82.

805 34. Serata M, Iino T, Yasuda E, Sako T. Roles of thioredoxin and thioredoxin reductase in the
806 resistance to oxidative stress in *Lactobacillus casei*. *Microbiology*. 2012;158(Pt_4):953-62.

807 35. Zhang L, Liu C, Li D, Zhao Y, Zhang X, Zeng X, et al. Antioxidant activity of an
808 exopolysaccharide isolated from *Lactobacillus plantarum* C88. *Int J Biol Macromol*.
809 2013;54:270-5.

810 36. Zhang B, Li S, Men J, Peng C, Shao H, Zhang Z. Long-term exposure to crotonaldehyde
811 causes heart and kidney dysfunction through induction of inflammatory and oxidative
812 damage in male Wistar rats. *Toxicology Mechanisms and Methods*. 2019;29(4):263-75.

813 37. Lushchak VI, Bagnyukova TV, Lushchak OV, Storey JM, Storey KB. Hypoxia and recovery
814 perturb free radical processes and antioxidant potential in common carp (*Cyprinus*
815 *carpio*) tissues. *The International Journal of Biochemistry & Cell Biology*.
816 2005;37(6):1319-30.

817 38. Mutanen A, Lohi J, Sorsa T, Jalanko H, Pakarinen MP. Features of liver tissue remodeling
818 in intestinal failure during and after weaning off parenteral nutrition. *Surgery*.
819 2016;160(3):632-42.

820 39. Lev-Cohain N, Sapir G, Harris T, Azar A, Gamliel A, Nardi-Schreiber A, et al. Real-time
821 ALT and LDH activities determined in viable precision-cut mouse liver slices using
822 hyperpolarized [1-(13) C]pyruvate-Implications for studies on biopsied liver tissues.
823 *NMR Biomed*. 2019;32(2):e4043.

824 40. Elhelaly AE, AlBasher G, Alfarraj S, Almeer R, Bahbah EI, Fouda MMA, et al. Protective
825 effects of hesperidin and diosmin against acrylamide-induced liver, kidney, and brain
826 oxidative damage in rats. *Environ Sci Pollut Res Int*. 2019;26(34):35151-62.

827 41. Deng H, Kerppola TK. Regulation of *Drosophila* metamorphosis by xenobiotic response
828 regulators. *PLoS Genet*. 2013;9(2):e1003263.

829 42. Chen-Roetling J, Regan RF. Targeting the Nrf2-Heme Oxygenase-1 Axis after
830 Intracerebral Hemorrhage. *Curr Pharm Des*. 2017;23(15):2226-37.

831 43. Chen J, Xie H, Chen D, Yu B, Mao X, Zheng P, et al. Chlorogenic Acid Improves Intestinal
832 Development via Suppressing Mucosa Inflammation and Cell Apoptosis in Weaned Pigs.
833 *ACS Omega*. 2018;3(2):2211-9.

834 44. Pan D, Mei X. Antioxidant activity of an exopolysaccharide purified from *Lactococcus*
835 *lactis* subsp. *lactis* 12. *Carbohydrate Polymers*. 2010;80(3):908-14.

836 45. Jiao N, Wu Z, Ji Y, Wang B, Dai Z, Wu G. L-Glutamate Enhances Barrier and Antioxidative
837 Functions in Intestinal Porcine Epithelial Cells. *J Nutr.* 2015;145(10):2258-64.

838 46. Lopes IS, Oliveira ICM, Capibaribe VCC, Valentim JT, da Silva DMA, de Souza AG, et al.
839 Riparin II ameliorates corticosterone-induced depressive-like behavior in mice: Role of
840 antioxidant and neurotrophic mechanisms. *Neurochem Int.* 2018;120:33-42.

841 47. O'Mahony SM, Hyland NP, Dinan TG, Cryan JF. Maternal separation as a model of brain-
842 gut axis dysfunction. *Psychopharmacology (Berl).* 2011;214(1):71-88.

843 48. Kember RL, Dempster EL, Lee TH, Schalkwyk LC, Mill J, Fernandes C. Maternal separation
844 is associated with strain-specific responses to stress and epigenetic alterations to Nr3c1,
845 Avp, and Nr4a1 in mouse. *Brain Behav.* 2012;2(4):455-67.

846 49. Davis MY, Zhang H, Brannan LE, Carman RJ, Boone JH. Rapid change of fecal
847 microbiome and disappearance of *Clostridium difficile* in a colonized infant after
848 transition from breast milk to cow milk. *Microbiome.* 2016;4(1):53.

849 50. Leung C, Rivera L, Furness JB, Angus PW. The role of the gut microbiota in NAFLD. *Nat*
850 *Rev Gastroenterol Hepatol.* 2016;13(7):412-25.

851 51. Napolitano A, Miller S, Nicholls A, Baker D, Horn S, Thomas E, et al. Novel Gut-Based
852 Pharmacology of Metformin in Patients with Type 2 Diabetes Mellitus. *PloS one.*
853 2014;9:e100778.

854 52. Wu W, Xiao Z, An W, Dong Y, Zhang B. Dietary sodium butyrate improves intestinal
855 development and function by modulating the microbial community in broilers. *PLoS*
856 *One.* 2018;13(5):e0197762.

857 53. Chen Y, Guo J, Qian G, Fang D, Shi D, Guo L, et al. Gut dysbiosis in acute-on-chronic
858 liver failure and its predictive value for mortality. *J Gastroenterol Hepatol.*
859 2015;30(9):1429-37.

860 54. Buchet A, Belloc C, Leblanc-Maridor M, Merlot E. Effects of age and weaning conditions
861 on blood indicators of oxidative status in pigs. *PLoS One.* 2017;12(5):e0178487.

862 55. Smith A, Stalder K, Serenius T, Serenius, Msc T, Baas T, et al. Effect of weaning age on
863 nursery pig and sow reproductive performance. *Journal of Swine Health and Production.*
864 2008;16:131-7.

865 56. Adetoye A, Pinloche E, Adeniyi BA, Ayeni FA. Characterization and anti-salmonella
866 activities of lactic acid bacteria isolated from cattle faeces. *BMC microbiology.*
867 2018;18(1):96.

868 57. Zheng W, Zhao W, Wu M, Song X, Caro F, Sun X, et al. Microbiota-targeted maternal
869 antibodies protect neonates from enteric infection. *Nature.* 2020;577(7791):543-8.

870 58. Difilippo E, Pan F, Logtenberg M, Willems RH, Braber S, Fink-Gremmels J, et al. In Vitro
871 Fermentation of Porcine Milk Oligosaccharides and Galacto-oligosaccharides Using
872 Piglet Fecal Inoculum. *J Agric Food Chem.* 2016;64(10):2127-33.

873 59. Fernandez L, Ruiz L, Jara J, Orgaz B, Rodriguez JM. Strategies for the Preservation,
874 Restoration and Modulation of the Human Milk Microbiota. Implications for Human Milk
875 Banks and Neonatal Intensive Care Units. *Front Microbiol.* 2018;9:2676.

876 60. Lagier JC, Khelaifia S, Alou MT, Ndongo S, Dione N, Hugon P, et al. Culture of previously
877 uncultured members of the human gut microbiota by culturomics. *Nat Microbiol.*
878 2016;1:16203.

879 61. Lv L-X, Hu X-J, Qian G-R, Lu H-F, Zheng B-W, Jiang L, et al. Administration of
880 *Lactobacillus salivarius* LI01 or *Pediococcus pentosaceus* LI05 improves acute liver injury
881 induced by D-galactosamine in rats. *Applied microbiology and biotechnology.* 2014;98.

882 62. Itoh K, Tong KI, Yamamoto M. Molecular mechanism activating nrf2-keap1 pathway in
883 regulation of adaptive response to electrophiles. *Free Radical Biology and Medicine.*
884 2004;36(10):1208-13.

885 63. Schwab CG, Bozak CK, Whitehouse NL, Olson VM. Amino acid limitation and flow to the
886 duodenum at four stages of lactation. 2. Extent of lysine limitation. *J Dairy Sci.*
887 1992;75(12):3503-18.

888 64. Zeitz JO, Kaltenböck S, Most E, Eder K. Effects of L-methionine on performance, gut
889 morphology and antioxidant status in gut and liver of piglets in relation to DL-
890 methionine. *Journal of animal physiology and animal nutrition.* 2019;103(1):242-50.

891 65. Zhou YF, Zhou Z, Batistel F, Martinez-Cortés I, Pate RT, Luchini DL, et al. Methionine and
892 choline supply alter transmethylation, transsulfuration, and cytidine 5'-diphosphocholine
893 pathways to different extents in isolated primary liver cells from dairy cows. *J Dairy Sci.*
894 2018;101(12):11384-95.

895 66. Swennen Q, Geraert PA, Mercier Y, Everaert N, Stinckens A, Willemsen H, et al. Effects of
896 dietary protein content and 2-hydroxy-4-methylthiobutanoic acid or DL-methionine
897 supplementation on performance and oxidative status of broiler chickens. *The British*
898 *journal of nutrition.* 2011;106(12):1845-54.

899 67. Bos C, Stoll B, Fouillet H, Gaudichon C, Guan X, Grusak MA, et al. Postprandial intestinal
900 and whole body nitrogen kinetics and distribution in piglets fed a single meal. *Am J*
901 *Physiol Endocrinol Metab.* 2005;288(2):E436-46.

902 68. Gamarra Y, Santiago FC, Molina-Lopez J, Castano J, Herrera-Quintana L, Dominguez A,
903 et al. Pyroglutamic acidosis by glutathione regeneration blockage in critical patients with
904 septic shock. *Crit Care.* 2019;23(1):162.

905 69. Zeisel SH. Dietary choline deficiency causes DNA strand breaks and alters epigenetic
906 marks on DNA and histones. *Mutat Res.* 2012;733(1-2):34-8.

907 70. Zhou Z, Vailati-Riboni M, Luchini DN, Loo JJ. Methionine and Choline Supply during the
908 Periparturient Period Alter Plasma Amino Acid and One-Carbon Metabolism Profiles to
909 Various Extents: Potential Role in Hepatic Metabolism and Antioxidant Status. *Nutrients.*
910 2016;9(1).

911 71. Guo WX, Pye QN, Williamson KS, Stewart CA, Hensley KL, Kotake Y, et al. Reactive
912 oxygen species in choline deficiency-induced apoptosis in rat hepatocytes. *Free Radic*
913 *Biol Med.* 2004;37(7):1081-9.

914 72. Tabassum S, Haider S, Ahmad S, Madiha S, Parveen T. Chronic choline supplementation
915 improves cognitive and motor performance via modulating oxidative and
916 neurochemical status in rats. *Pharmacol Biochem Behav.* 2017;159:90-9.

917 73. Coleman DN, Vailati-Riboni M, Elolimy AA, Cardoso FC, Rodriguez-Zas SL, Miura M, et
918 al. Hepatic betaine-homocysteine methyltransferase and methionine synthase activity
919 and intermediates of the methionine cycle are altered by choline supply during negative
920 energy balance in Holstein cows. *J Dairy Sci.* 2019;102(9):8305-18.

921 74. Wu P, Liu Y, Jiang WD, Jiang J, Zhao J, Zhang YA, et al. A Comparative Study on
922 Antioxidant System in Fish Hepatopancreas and Intestine Affected by Choline
923 Deficiency: Different Change Patterns of Varied Antioxidant Enzyme Genes and Nrf2
924 Signaling Factors. *PLoS One.* 2017;12(1):e0169888.

925 75. Li B, Li W, Ahmad H, Zhang L, Wang C, Wang T. Effects of Choline on Meat Quality and
926 Intramuscular Fat in Intrauterine Growth Retardation Pigs. *PLoS One.*
927 2015;10(6):e0129109.

928 76. Yamada S, Kawaguchi H, Yamada T, Guo X, Matsuo K, Hamada T, et al. Cholic Acid
929 Enhances Visceral Adiposity, Atherosclerosis and Nonalcoholic Fatty Liver Disease in
930 Microminipigs. *J Atheroscler Thromb.* 2017;24(11):1150-66.

931 77. Carrier A. Metabolic Syndrome and Oxidative Stress: A Complex Relationship. *Antioxid*
932 *Redox Signal.* 2017;26(9):429-31.

933 78. Sommerfeld A, Reinehr R, Haussinger D. Bile acid-induced epidermal growth factor
934 receptor activation in quiescent rat hepatic stellate cells can trigger both proliferation
935 and apoptosis. *J Biol Chem.* 2009;284(33):22173-83.

936 79. Becker BA, Nienaber JA, Christenson RK, Manak RC, DeShazer JA, Hahn GL. Peripheral
937 concentrations of cortisol as an indicator of stress in the pig. *American journal of*
938 *veterinary research.* 1985;46(5):1034-8.

939 80. Du X, Pang TY. Is Dysregulation of the HPA-Axis a Core Pathophysiology Mediating Co-
940 Morbid Depression in Neurodegenerative Diseases? *Frontiers in psychiatry.* 2015;6:32.

941 81. Kulak A, Steullet P, Cabungcal JH, Werge T, Ingason A, Cuenod M, et al. Redox
942 dysregulation in the pathophysiology of schizophrenia and bipolar disorder: insights
943 from animal models. *Antioxid Redox Signal.* 2013;18(12):1428-43.

944 82. Lin X, Xia Y, Wang G, Xiong Z, Zhang H, Lai F, et al. *Lactobacillus plantarum* AR501
945 Alleviates the Oxidative Stress of D-Galactose-Induced Aging Mice Liver by
946 Upregulation of Nrf2-Mediated Antioxidant Enzyme Expression. *J Food Sci.*
947 2018;83(7):1990-8.

948 83. Jiang T, Xing X, Zhang L, Liu Z, Zhao J, Liu X. Chitosan Oligosaccharides Show Protective
949 Effects in Coronary Heart Disease by Improving Antioxidant Capacity via the Increase in
950 Intestinal Probiotics. *Oxid Med Cell Longev.* 2019;2019:7658052.

951 84. Wu CH, Ko JL, Liao JM, Huang SS, Lin MY, Lee LH, et al. D-methionine alleviates
952 cisplatin-induced mucositis by restoring the gut microbiota structure and improving
953 intestinal inflammation. *Ther Adv Med Oncol.* 2019;11:1758835918821021.

954 85. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI. Obesity alters gut
955 microbial ecology. *Proceedings of the National Academy of Sciences of the United*
956 *States of America.* 2005;102(31):11070-5.

86. Wang K, Jin X, Li Q, Sawaya A, Le Leu RK, Conlon MA, et al. Propolis from Different Geographic Origins Decreases Intestinal Inflammation and *Bacteroides* spp. Populations in a Model of DSS-Induced Colitis. *Mol Nutr Food Res*. 2018;62(17):e1800080.

87. Takano S, Pawlowska BJ, Gudelj I, Yomo T, Tsuru S. Density-Dependent Recycling Promotes the Long-Term Survival of Bacterial Populations during Periods of Starvation. *mBio*. 2017;8(1).

88. Wang D, Liu W, Ren Y, De L, Zhang D, Yang Y, et al. Isolation and Identification of Lactic Acid Bacteria from Traditional Dairy Products in Baotou and Bayannur of Midwestern Inner Mongolia and q-PCR Analysis of Predominant Species. *Korean J Food Sci Anim Resour*. 2016;36(4):499-507.

89. Grill JP, Cayuela C, Antoine JM, Schneider F. Effects of *Lactobacillus amylovorus* and *Bifidobacterium breve* on cholesterol. *Lett Appl Microbiol*. 2000;31(2):154-6.

90. Limkhuansuwan V, Chaiprasert P. Decolorization of molasses melanoidins and palm oil mill effluent phenolic compounds by fermentative lactic acid bacteria. *J Environ Sci (China)*. 2010;22(8):1209-17.

91. Hu J, Ma L, Nie Y, Chen J, Zheng W, Wang X, et al. A Microbiota-Derived Bacteriocin Targets the Host to Confer Diarrhea Resistance in Early-Weaned Piglets. *Cell host & microbe*. 2018;24(6):817-32.e8.

92. Janda JM, Abbott SL. 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. *J Clin Microbiol*. 2007;45(9):2761-4.

93. Ponce AG, Moreira MR, del Valle CE, Roura SI. Preliminary characterization of bacteriocin-like substances from lactic acid bacteria isolated from organic leafy vegetables. *LWT - Food Science and Technology*. 2008;41(3):432-41.

94. Magoč T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics (Oxford, England)*. 2011;27(21):2957-63.

95. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics (Oxford, England)*. 2018;34(17):i884-i90.

96. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods*. 2016;13(7):581-3.

97. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature biotechnology*. 2019;37(8):852-7.

98. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic acids research*. 2013;41(Database issue):D590-6.

99. Asnicar F, Weingart G, Tickle TL, Huttenhower C, Segata N. Compact graphical representation of phylogenetic data and metadata with GraPhlAn. *PeerJ*. 2015;3:e1029.

100. Zhang J, Liu YX, Zhang N, Hu B, Jin T, Xu H, et al. NRT1.1B is associated with root microbiota composition and nitrogen use in field-grown rice. *Nature biotechnology*. 2019;37(6):676-84.

Figure Legends

Figure 1. The workflow for large-scale bacterial cultivation from sow milk and the characterization of probiotic functions.

Figure 2. The Sow Milk Bacterial Collection. The inner circles depict taxonomic assignments of bacteria for the indicated phyla, classes, orders, families, genera, and species that were identified. The species names are labeled, and the numbers of different taxa within each species are provided at the nodes. *P. pentosaceus* is indicated in red. The probability (%) of suspected new species and known species identified from the dairy products is shown in the outer ring with the green or yellow heat map, respectively. The outermost bar plots represent the numbers of isolates (log2 transformed) of each species.

Figure 3. Screening of potential probiotics prior to a piglet feeding experiment. (a) Antagonistic activity against pathogens *in vitro*. (b) Graphical representation of experiments conducted to assess the survival rate. (c) *P. pentosaceus* confers protective effects upon *Drosophila*. Statistical differences were calculated by the log-rank test.

Figure 4. Effects of the oral administration of SMM914 on the antioxidation capacity and activation of the cytoprotective Nrf2 pathway in piglets. (a) Experimental outline in piglets ($n = 18$). Piglets were kept for 10 days after birth for adaptation. Oral administration of SMM914 was performed on days 10, 12, 14, 16 and 18 at low or high doses. At day 7 postweaning, seven piglets per treatment were randomly selected for

slaughter for sample collection. (b) Alanine aminotransferase (ALT) and lactate dehydrogenase (LDH) were significantly decreased in piglets subjected to SMM914. (c-d) Bar graph of western blotting showing Nrf2 protein expression levels in the liver normalized to PCNA expression in the nucleus. (e-f) Western blotting analysis of antioxidant protein (Keap1, NQO-1, HO-1, SOD1 and CAT) levels in the liver and bar graph of data showing their protein expression normalized to β -actin expression in hepatic cells ($n = 6$). (g) The enzyme activities of GSH-Px, SOD, CAT and MDA were measured in liver lysates. Values of the bars stand for significant differences using one-way ANOVA followed by Duncan's multiple comparisons at $p < 0.05$. Data are the mean $\pm$ s.e.m, $n = 7$.

Figure 5. The metabolic profile of plasma. Score plots of the PLS-DA analysis for the plasma metabolome (a) showing the scatter between the control and low-dose groups. (b) Scatter between the control and high-dose groups. (c) The integrative metabolism pathway according to the KEGG pathway database. Compared with the control group, the blue metabolites represent the intensities of metabolites that were downregulated, while the red metabolites represent the intensities of metabolites that were upregulated. (d-h) The perturbed metabolism pathways and metabolites in response to SMM914 treatment mainly include glutathione metabolism; cysteine and methionine metabolism; glycine, serine and threonine metabolism; and arginine and proline metabolism. Data are the mean $\pm$ s.e.m, $n = 7$.

Figure 6. Microbial composition analysis in the colon. (a) The Venn diagram for operational taxonomic units among the control group, low-dose group and high-dose group. (b) Scatterplot of principal component analysis in bacterial communities. Principal components (PCs) 1 and 2 explain 11.11% and 9.96% of the variance, respectively (permutational multivariate analysis of variance). (c) Individual (left) and averaged (right) taxon summary of bacterial genera in colon contents.

Figure 7. (a) Cladogram generated from Linear discriminant analysis (LDA) effect size (LEfSe) analysis for the most differentially abundant taxa with LDA scores above 4. (b) Comparison of microbial genera between high dose group and control. Significantly different domains were tested by Wilcoxon rank-sum test. $*p < 0.05$; $**p < 0.01$.

Figure S1. The maximum-likelihood tree of the bacterial taxa. The 271 bacterial taxa were clustered based on 16S rDNA gene identity with a cut-off value of 99% for different taxa in the CD-HIT analysis. The closely related neighboring species of each taxon are listed next to the taxon numbers of the sow milk microbiota. The suspected new species are indicated in blue, and the species in the dairy products are indicated in red dots.

Figure S2. Light microscopy images obtained from *P. pentosaceus* strains using a 63x oil immersion objective.

Figure S3. The antimicrobial activity of SMM914. (a) The pH of SMM914 products were determined via test papers. (b) Inhibitory effects of SMM914 against *S. aureus* and ETEC. The circular wells were filled with different products of SMM914, including

MRS medium (1&2) and the cell-free supernatant without any treatment (3&4), with 15 min heat inactivation (5&6), with pH adjustment using NaOH (7&8) or with 1 mg/mL protease K treatment (9&10). The substances (1, 3, 5, 7 and 9) were obtained from the fermentation broth after cultivation under anaerobic condition, while others (2, 4, 6, 8 and 10) were obtained from culture under aerobic condition.

Figure S4. Growth curves and pH value curve of SMM914 after anaerobic and aerobic fermentation. (a) The optical density at 600 nm (OD₆₀₀) was measured from a starting OD₆₀₀ of 0.085. (b) The pH value was recorded with a pH meter. Data are the mean $\pm$ s.e.m, $n = 3$.

Figure S5. The annotation and comparison of *P. pentosaceus* SMM914 genome. (a) The distribution of predicted CDSs of SMM914 in different categories of metabolic function by online software RAST. (b) A full genome comparison analysis of *P. pentosaceus* SMM914 with other *P. pentosaceus* strains, including *Pediococcus pentosaceus* SRCM100194, *Pediococcus pentosaceus* GDIAS001, *Pediococcus pentosaceus* SL001 and *Pediococcus pentosaceus* SRCM102736, visualized by BRIG software. Colors display the percentage of sequence identity based on BLASTN. Two inner rings indicate the GC skew and the G + C content. The innermost circle shows the genome coordinates.

Figure S6. Effects of SMM914 on (a) growth performance and (b) organ relative weight. Data are the mean $\pm$ s.e.m. Statistical analysis was conducted by using one-way

ANOVA. Data not sharing the same letter in each point were significantly different ($p < 0.05$).

Figure S7. Effects of SMM914 on serum biochemical parameters. Data are the mean $\pm$ s.e.m ($n = 7$). One-way ANOVA with adjustment for multiple comparisons was conducted. ^{a,b} Within a variable, values with different superscripts differ ($p < 0.05$).

Figure S8. (a) In HepG2 cell and (b) IPEC-J2 cell, western blotting experiment was to determine the effect of SMM914 in MRS broth after 24h fermentation on the protein level of Nrf2. (c) The relative changes in protein intensity were analyzed with unpaired Student's t-test. Different letters in the graph indicate significant statistical differences ($p < 0.05$, $n = 3$).

Figure S9. (a) Alpha diversity comparisons of the gut microbiomes including Ace, Chao, Sobs index, Shannon index, Shannoneven index, Simpson index, Simpstoneven index and PD whole tree index which were analyzed using Kruskal-Wallis H test and Tukey-Kramer post hoc test with 95% confidence level. Data are the mean $\pm$ s.d ($n = 7$). *ns*, not significant differences. (b) Rarefaction curves for Shannon indices at genus level in the colon contents.

Figure S10. (a) Linear discriminant analysis effect size 10 was set up with 0.05 as the alpha value for Wilcoxon test and then selected with a threshold value of 4 from phylum to genus level. (b) Differences in relative abundance of the important and variable genera in colonic microbiota among three group. Each data represented the mean and SEM of relative abundance of each genus ($n = 7$). One-way analysis of variance and

Duncan's multiple comparison test to determine the statistical. Different letters in the same graph indicate significant statistical differences ($p < 0.05$).

Figure S11. (a) Bar charts of relative abundance at the family level in the control and treated groups obtained by sequencing of 16S rDNA V3-V4 region. (b) Comparison of microbial families between high dose group and control. Significantly different domains were tested by Wilcoxon rank-sum test. $*p < 0.05$; $**p < 0.01$.

Supplementary Data S1. The 16S rRNA sequences and classification of all isolates in smBC

Supplementary Data S2. The alignment of 1240 16S rRNA sequences against the Silva version132 16S rRNA database, NCBI nucleotide collection (nr/nt) database and DAIRYdb database using BLASTN

Supplementary Data S3. Genes of SMM914 and predicted proteins by Pfam protein database

Figures

Figure 1

Figure 1

The workflow for large-scale bacterial cultivation from sow milk and the characterization of probiotic functions.

Figure 2

Figure 2

The Sow Milk Bacterial Collection. The inner circles depict taxonomic assignments of bacteria for the indicated phyla, classes, orders, families, genera, and species that were identified. The species names are labeled, and the numbers of different taxa within each species are provided at the nodes. *P. pentosaceus* is indicated in red. The probability (%) of suspected new species and known species identified from the dairy products is shown in the outer ring with the green or yellow heat map, respectively. The outermost bar plots represent the numbers of isolates (log2 transformed) of each species.

Figure 3

Screening of potential probiotics prior to a piglet feeding experiment. (a) Antagonistic activity against pathogens in vitro. (b) Graphical representation of experiments conducted to assess the survival rate. (c) *P. pentosaceus* confers protective effects upon *Drosophila*. Statistical differences were calculated by the log-rank test.

Figure 4

Figure 4

Effects of the oral administration of SMM914 on the antioxidation capacity and activation of the cytoprotective Nrf2 pathway in piglets. (a) Experimental outline in piglets ($n = 18$). Piglets were kept for 10 days after birth for adaptation. Oral administration of SMM914 was performed on days 10, 12, 14, 16 and 18 at low or high doses. At day 7 postweaning, seven piglets per treatment were randomly selected for slaughter for sample collection. (b) Alanine aminotransferase (ALT) and lactate dehydrogenase (LDH)

were significantly decreased in piglets subjected to SMM914. (c-d) Bar graph of western blotting showing Nrf2 protein expression levels in the liver normalized to PCNA expression in the nucleus. (e-f) Western blotting analysis of antioxidant protein (Keap1, NQO-1, HO-1, SOD1 and CAT) levels in the liver and bar graph of data showing their protein expression normalized to β -actin expression in hepatic cells (n = 6). (g) The enzyme activities of GSH-Px, SOD, CAT and MDA were measured in liver lysates. Values of the bars stand for significant differences using oneway ANOVA followed by Duncan's multiple comparisons at $p < 0.05$. Data are the mean $\pm$ s.e.m, n = 7.

Figure 5

Figure 5

The metabolic profile of plasma. Score plots of the PLS-DA analysis for the plasma metabolome (a) showing the scatter between the control and low-dose groups. (b) Scatter between the control and high-dose groups. (c) The integrative metabolism pathway according to the KEGG pathway database. Compared with the control group, the blue metabolites represent the intensities of metabolites that were downregulated, while the red metabolites represent the intensities of metabolites that were upregulated. (d-h) The perturbed metabolism pathways and metabolites in response to SMM914 treatment mainly include glutathione metabolism; cysteine and methionine metabolism; glycine, serine and threonine metabolism; and arginine and proline metabolism. Data are the mean $\pm$ s.e.m, n = 7.

Figure 6

Figure 6

Microbial composition analysis in the colon. (a) The Venn diagram for operational taxonomic units among the control group, low-dose group and high-dose group. (b) Scatterplot of principal component analysis in bacterial communities. Principal components (PCs) 1 and 2 explain 11.11% and 9.96% of the variance, respectively (permutational multivariate analysis of variance). (c) Individual (left) and averaged (right) taxon summary of bacterial genera in colon contents.

Figure 7

Figure 7

(a) Cladogram generated from Linear discriminant analysis (LDA) effect size (LEfSe) analysis for the most differentially abundant taxa with LDA scores above 4. (b) Comparison of microbial genera between high dose group and control. Significantly different domains were tested by Wilcoxon rank-sum test. * $p < 0.05$; ** $p < 0.01$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementarymaterial.zip](#)