Promoting rational antibiotics prescribing for non-complicated infections: Understanding social influence in primary care networks in Germany

Regina Poss-Doering (✉ regina.poss-doering@med.uni-heidelberg.de)

University Hospital Heidelberg <https://orcid.org/0000-0003-0618-4034>

Martina Kamradt

University Hospital Heidelberg

Katharina Glassen

University Hospital Heidelberg

Edith Andres

aQua Institut, Goettingen

Petra Kaufmann-Kolle

aQua Institut, Goettingen

Michel Wensing

University Hospital Heidelberg

Research article

Keywords: primary care networks; quality improvement; appropriate antibiotics use; social influence processes; peer exchange;

Posted Date: November 6th, 2019

DOI: <https://doi.org/10.21203/rs.2.16935/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Family Practice on March 14th, 2020. See the published version at <https://doi.org/10.1186/s12875-020-01119-8>.

Abstract

Primary care networks in Germany are formalized regional collaborations of physicians and other healthcare providers. Common goals are optimized healthcare processes and services for patients, enhanced communication, agency for professional concerns and strengthened economic power. In the ARena study (Sustainable reduction of antibiotic-induced antimicrobial resistance), 14 primary care networks in two federal German states aimed to promote appropriate antibiotics use for acute non-complicated infections by fostering awareness and understanding. Factors related to the role of primary care networks were to be identified. Methods For this study, audio-recorded telephone interviews were conducted with physicians, non-physician health professionals and stakeholder representatives. Pseudonymized verbatim transcripts were coded using thematic analysis. In-depth analysis was based on the inductive categories 'social support', 'social learning', 'social normative pressures' and 'social contagion' to reflect social influence processes. To foster understanding of the networks' potential impact on antibiotics prescribing, data generated through surveys with physicians and non-physician health professionals were analyzed descriptively. Results : Social influence processes proved to be relevant regarding knowledge transfer, manifestation of best-practice care and self-reflection. Peer communication was seen as great asset, main reason for membership and affirmative for own perspectives. All interviewed physicians (n=27) considered their network to be a strong support factor for daily routines, introduction of new routines, and continuity of care. They utilized network-offered training programs focusing on best practice guideline-oriented use of antibiotics and considered their networks supportive in dealing with patient expectations. A shared attitude combined with ARena intervention components facilitated reflected management of antibiotics prescribing. Non-physician health professionals (n=11) also valued network peer exchange. They assumed their employers joined networks to offer improved and continuous care. Stakeholders (n=7) expected networks and their members to be drivers for care optimization. Conclusion: Primary care networks play a crucial role in providing a platform for professional peer exchange, social support and reassurance. With regards to their impact on antibiotics prescribing for acute non-complicated infections, networks seem to facilitate and amplify quality improvement programs by providing a platform for refreshing awareness, knowledge and self-reflection among care providers. They are well suited to promote a rational use of antibiotics. .

Background

In Germany, primary care networks (PCN) are comprehended as an important vehicle for optimizing local primary and cross-sectoral care for the benefit of patients, physicians and stakeholders [1]. Similar to communities of practice [2], these formalized collaborations of physicians and other healthcare providers interact regularly and share patients, set goals, standardize treatment and care, discuss concerns and attend continuing education. In addition, networks can provide economic advantages, such as power in negotiations with health insurers. Currently, more than 400 PCN with up to 100 physicians aim to act as coordinators and moderators in regional primary care in Germany, bringing together general practitioners, ambulatory medical specialists, nursing homes, hospitals and self-help support organizations to optimize

quality and efficiency of care [3, 4]. Differing in strategic alignment and ranging from loosely connected single physicians to professionally managed organizations, these networks offer the entire range of primary care [5]. All PCN aim to provide above-average quality and outcomes of care as well as high patient satisfaction with care [3].

The cluster randomized trial ARena (Sustainable reduction of antibiotic-induced antimicrobial resistance, 2017-2019, trial registration: ISRCTN, ISRCTN58150046) intends to promote the rational and appropriate use of antibiotics for acute non-complicated infections in primary care in Germany [6, 7]. Using a multifaceted strategy with multiple interacting intervention components, ARena addresses physician, primary care team and patient knowledge and attitudes about the use of antibiotics [6]. ARena is embedded into 14 PCN across the German federal states of Bavaria and North-Rhine Westphalia. Essentially, the promotion of a rational, reasonable use of antibiotics is based on fostering awareness for and understanding of the growing challenges of antimicrobial resistances (AMR) by effective communication, education and training addressed to PCN physicians and care teams as well as the regional public. As ARena is not completed yet, assessment of intervention effectiveness currently is pending. With the aim of providing insights into determinants of practice regarding a rational use of antibiotics and to propose explanations referencing identified influences and mechanisms of action, ARena is accompanied by a process evaluation to explore factors and processes leading to impacts on antibiotics prescribing patterns [6].

Insights into the development of PCN and their contributions to regional improvements in patient care already have been gained, yet predominantly are limited to specific single networks [3, 8–10]. Research found that PCN can effectively shape healthcare by acting as a driver for innovation and optimized performance, especially where effective network strengths meet potentially new approaches to care [5]. Also, physician peer networks have emerged as a potentially important factor influencing medical practice [11]. Thus, being part of such a network might support the adoption of specific behaviors. This support may be attributed to social contagion as an influencing process in which network members are impacted by each other in their adoption decisions [12]. Social contagion theory suggests that human behaviors and traits can spread in social networks [13], assuming this is promoted by various behavioral mechanisms, such as imitation, role modelling and persuasion [12]. As interacting physicians in networks likely share their beliefs, ideas and experiences with each other, the interpersonal information exchange may influence practice patterns [14] and turn peer influence into a potential driver of physicians' practice styles [15]. Studies of communities of practice and physician peer networks have identified differences in care patterns and patient outcomes based on peer connections [15–18] with regards to therapy uptake [16], or the adoption of technology [15], but have not focused yet in-depth on how physicians might have an impact on each other [11]. The role PCN might play in supporting rational prescribing and use of antibiotics for non-complicated infections is yet unknown. The aim of this study was to explore factors and processes attributed to the network's contribution to improving antibiotics prescribing in the ARena project.

Methods

Design

ARena was a three-armed cluster randomized trial, implemented in 14 PCN in two German federal states (Bavaria and North Rhine-Westphalia). Intervention components included e-learning on communication, quality circles and data-based feedback for physicians and care team members, public information campaigns, performance-based additional reimbursement, a computerized decision support system and culture-sensitive information material for patients in print and digital format on tablet computers to be used in waiting areas. Standard care is reflected by an added cohort based on claims-data [6].

In a mixed-methods approach, the process evaluation used open-ended, semi-structured interviews with primary care physicians (general practitioners, ear-nose throat specialists, urologists, and pediatricians), non-physician health professionals – comparable to medical assistants (MA) in USA [19] - and stakeholder representatives (PCN management, health insurance and patient representatives). Also used were a one-time socio-demographic survey of interviewees and questionnaires given to all physicians included in the intervention and all MA in one of the three intervention arms at three different points in time. Different interview guides were developed for the three groups of interviewees based on a literature review and pre-defined research questions. The first interview in each group served as a pilot so minor adjustments could be made where considered appropriate. All qualitative data generated from interviews with physicians, MA and stakeholders, field notes and the socio-demographic survey were included for analysis. In addition, data referencing PCN were extracted from two sets of items in the first and second of three questionnaires (T0, T1) and included. Questionnaire construction was guided by the Theory of Planned behaviour [20]. A pre-test was conducted with 4 non-participant physicians to ensure relevance and clarity of all items. Thus, a broad thematic spectrum could be analyzed to assess and understand factors relevant to the role of PCN regarding support for rational use of antibiotics for acute non-complicated infections.

Study population for interviews

No target sample size was calculated for the interviews. Applying a purposive strategy, a sample of 45 participants were recruited by the ARena study team at the Department of General Practice and Health Services Research, University Hospital Heidelberg between March and May 2018. Potential recruits were all physicians and MAs participating in ARena as well as managerial representatives of participating PCN and health insurance providers, association of statutory health insurance physicians and of a self-help organization.

All recruits had to be at least 18 years of age, legally fully competent and in fluent command of German. A structured recruitment procedure aimed at even distribution regarding gender and intervention groups. To be included in the process evaluation, all interested parties meeting the inclusion criteria received

printed material as well as a phone call to provide further information, and were obliged to return a signed letter of intent for participation in the interview.

Out of a total of 303 eligible, 40 physicians per intervention group were invited by e-mail via the aQua Institute, Goettingen, to participate in an interview (n=120 in total). After three weeks, a reminder was e-mailed. Due to a disproportionately high number of recruited participants from Bavaria, an additional reminder was e-mailed to a random sample of 11 physicians located in North-Rhine Westphalia after 12 weeks. Out of 84 eligible medical assistants, 25 were contacted. Stakeholder representatives (n=7) personally were addressed by the aQua Institute via e-mail or letter. All invitees were sent a personalized cover letter, supplemented by information specific to study and process evaluation details and a feedback form to be returned by e-mail or fax to declare interest in participation.

Study population for survey

All physicians participating in the intervention groups were invited to take part in the survey (T0 n=303, T1 n=312, T2 n=292). Also, MAs employed at eligible participating primary care practices allocated to intervention arm B, were invited to take part in the survey (T0 n=84, T1 n=88, T2 n=85). E-mail reminders were sent out after 4 weeks to increase the response rate. Questionnaires T0, T1 and T2 were to be sent out at three points in time over the course of the study. Response rates among physicians were 75.5 % (T0), 64% (T1) and 63.3 % (T2), and for MAs 95% (T0), 64.2% (T1) and 68.2% (T2).

Data collection and analysis

Interviews

Between April and June 2018, all physician interviews (n=27) were conducted and digitally audio recorded by three researchers of the study team at the Department of General Practice and Health Services Research, University Hospital Heidelberg. A semi-structured interview guide was followed to gain insights into typical practice referring to antibiotics prescribing and consideration of patient preferences, implications of the quality improvement program and its' intervention components for patient care, general contextual factors and the role of the PCN in particular. Physician interview duration varied between 7:54 minutes and 62:50 minutes, with a mean of 28:14 minutes. Two of the researchers conducted and audio recorded all interviews with medical assistants (n=11) in April and May 2018, with a duration of 17:30 to 42:32 minutes (mean 26:53 minutes). Besides the aforementioned topics, the interview guide specifically focused on MA perspectives and experiences. All stakeholder interviews (n=7) were conducted and audio recorded by the same two researchers in April and May 2018, with a mean duration of 30:15 minutes (range 16:28 to 44:42 minutes). Here the interview guide was tailored to cover stakeholder expectations for the potential influence of the intervention components and perspectives on context factors as well as recommendations for the future use of antibiotics.

All interviews (n=45) were conducted over telephone. Additional notes were taken to document participant suggestions with regards to aspects of intervention delivery. After data collection was completed, pseudonymized verbatim transcripts were coded applying a thematic framework analysis [21] based on the Tailored Implementation for Chronic Disease (TICD) framework which uses 7 domains to classify determinants of implementation [22]. The interprofessional team of researchers (Public Health and Health Services Research) identified themes of interest deductively *a priori* from the TICD framework as well as inductively *de novo* from the data itself during the analysis. Two researchers coded all transcripts independently and iteratively using MAXQDA Analytics PRO 18 (Release 18.0.3). Divergent codings were discussed continuously to ensure intercoder congruity and to achieve the widest consensus possible. To enable a broader view, participant socio-demographic characteristics were analyzed descriptively using IBM SPSS Statistics Version 24. In an effort to facilitate understanding of the role and the mechanisms of influence of PCN in promoting a rational use of antibiotics for acute non-complicated infections, an integration of the theoretical perspective on social influence processes was applied to further deepen data analysis and interpretation and to propose explanations referring to identified relevant factors.

Surveys

Questionnaires were mailed to participants in January 2018 (T0), October 2018 (T1) and July 2019 (T2). All questionnaires focused on the implemented intervention components, relevant context factors, prescribing decisions and general perceptions regarding antibiotics. Additionally, T1 and T2 asked for interim and concluding assessments of the intervention components, respectively. Completed questionnaires were returned to and registered by the study team at the Department of General Practice and Health Services Research, University Hospital Heidelberg, between February and April 2018 (T0), November 2018 to January 2019 (T1), and July to September 2019 (T2). All completed questionnaires were digitalized and subsequently, data were transferred into IBM SPSS Statistics 24 for descriptive analysis. Data visualization was performed using Microsoft Excel 2010. Findings from the survey data are reported with a focus on the potential contribution networks might provide regarding communicative thematic peer exchange and educative efforts geared towards care teams.

Table 1 outlines the data collection sources relevant for findings presented here.

Results

Overview

The results outlined below reflect the identified factors and processes of influence of PCN in promoting a rational use of antibiotics for acute non-complicated self-limiting infections. A socio-demographic questionnaire was analyzed descriptively with regard to participant- and practice-level characteristics. A total of 27 physicians participated in the interviews (9 female, 18 male). All participating MA were female (n=11). Stakeholders had between 1 and 10 years of experience in their current position. On average,

physician respondents in the survey were 55 years old, had 25 years of working experience and had been members of their PCN for 10 years (Table 2).

Findings from the qualitative data are reported regarding aspects explored within the inductive domain 'Primary care networks' and focus on social influence processes [23–25] identified during the analysis: perceived social support, social learning, social-normative pressures and social contagion.

Figure 1 describes the analytical process and points out relevant sub-categories defined during the analysis.

Social support

Being a network member

Generally, physicians and MAs felt their networks to be a strong support factor both for existing daily and implementing new routines. Data generated through surveys T0 and T1 showed that 74% and 59% respectively of physicians experienced support for new routines, 43% and 36% acknowledged network impact on prescribing decisions, guideline-oriented care (70.5% and 60%), and management of patient expectations (61.2% and 51%) and felt supported in pursuing shared-decision making (61% and 59%) (Table 3). Table 3 shows results from physician survey T0 and T1 on network participation.

When asked about the significance of being a member in a PCN in the interviews, participants shared clear ideas. Predominantly, cited aspects referred to being part of a group rather than working just as a single practice. Access to collective knowledge through active in person peer exchange was considered to be of high importance with regards to support in situations of uncertainty, complex cases and the synchronization of therapy decisions across disciplines.

"A team-feeling." Phys01, Pos 56

"To be together, we are stronger together!" Phys05, Pos. 28

"Well, everybody has their own practice and will decide for themselves, but at least you have a platform for good exchange." Phys27, Pos. 82

Physicians stated different reasons for initially joining their network including options for future prospects, further personal development, communicative peer exchange, upskilling and utilization of joint administrative structures.

"Personally, I think that ... it is the future, ... the single practice, all alone without peer exchange, at least for me, it is definitely not a model I want to work in."

Phys08, Pos. 56

"We have found a forum where we look at the big picture and that means, I don't only look at pediatric topics but also at orthopedics, gynecology, dermatology and all other areas, and they, too, learn what is

important for us, for life, for our perspective.” Phys02, Pos. 20

MAs felt that the network membership motivated them to support the physician (72% in T0 and 69,9 % in T1), fostered dealing with patient expectations (68% and 71.3%), supported implementing new routines (71% and 68.4%) and had an impact on their part in patient care (59% and 60.3 %). During the interviews, they shared physician views regarding future prospects, peer exchange and joint administrative structures, and added their own perspectives referring to financial aspects and benefits for patient care. Stakeholder substantiated physician and MA perspectives, emphasized the importance of collaboration, communicative peer exchange and collective efforts of upskilling in networks. They also reflected on support of basic human needs.

“Patient care, to begin with, exactly, so that patients are well supported, and I think, I will be honest, it might also be a question of billing.” MA06, Pos. 44

“You are in a group, I think this is a little bit like our inner core that always looks for a partnership anywhere.” Sh04, Pos. 78

“... makes you somewhat happier, fulfills your own expectations and of course motivates you, when you can enter into dialog and, as a bonus, can present yourself and what you can do really, really well, to the network, and you can collect everybody’s experiences – when this exchange happens.” Sh04, Pos. 84

Impacting care delivery

Regarding care delivery, participants saw network membership as an asset and decisive factor for quality management, both in the study context and beyond. Continuity of care within the network was seen as supported by more and closer peer contacts, simplified peer exchange, usage of shared electronic patient records, fast appointment scheduling and patient transfers to medical specialists.

“Improved care services for the patients, not only in the ARena study, but in general ...” MA03, Pos. 75

“Well, to ensure rural medical services and make sure patients can always come and we are here for them.” MA05, Pos. 56

“...important to me that I communicate with others and I believe, working in the network contributes to quality management in the long run ...” Phys23, Pos. 54

“Mutual communication is top priority, in my opinion, this is underestimated, and through such networks where you are forced to communicate with each other, this is promoted immensely, and all involved benefit and of course not least the patients.” Sh04, Pos. 70

Confirming own perspective

Social support through communicative peer exchange was also seen as affirmative to own perspectives regarding importance of specific topics or course of own action.

“This peer exchange is extremely important and contributes a lot to self-confidence, I presume.” Phys23, Pos. 56

“Such discussions are important to affirm the topic and confirm the own course of action. So, in a reflective discussion you learn how others act, how many use the same approach, do many use the same approach, where do you stand and fall in line, and this can be confirming or critical.” Phys21, Pos. 42

“Exactly these discussions happen in these networks, ... What does this mean? How do I accompany patients, take care of them, how can I conduct conversations? To support each other in finding a way and entering into discussion ...” Sh02, Pos. 22

Social learning

Upskilling

Participants reported structured approaches of regularly offering upskilling events by their networks and taking part in a range of medical and organizational training programs such as quality and efficiency circles for MAs and physicians, workshops, case studies, and journal clubs. These learning opportunities were considered to be supportive in terms of providing new knowledge and peer exchange opportunities in a protective setting at the same time. Participants also felt supported by provided information components on evidence-based practice and provision of patient information.

“We have solid structures in our network concerning continuing training.”

Phys03, Pos. 54

“We regularly meet in quality circles, I think that is quite good, and do case reviews ... I find this very pleasant in the network, because you know each other and dare to come out of your shell ...” Phys14, Pos. 44

Participating in ARena

Physicians reported taking part in ARena-specific quality circles focusing on best practice guideline-oriented use of antibiotics and finding opportunity there for thematically relevant peer exchange within the network. Survey data (T0 and T1) showed that 83.4% and 86% of the participating physicians used the offered trainings on guideline-oriented antibiotics therapy and 71.5% and 72 % acknowledged that peer exchange about antibiotics prescribing for acute non-complicated infections was provided within their network (Table 4). There was indication of a heightened alertness to quality improving mindful antibiotics prescribing as a result of the PCN's study participation.

Social-normative pressures

Shared network attitude

Participants made references that pointed to a shared network attitude which included a common interest in evidence-based guideline-oriented care, mutual support in situations of uncertainty or locum care, long existing memberships, and a general sense of community. Financial aspects and ties to pharmaceutical manufacturers were negated. However, pointing to potential selection processes, it was also mentioned that PCN members needed to be willing to implement changes. This reflected in survey T0 data, where 90% of the physicians indicated they had implemented changes in their practices during the past two years. Aspects of social integration and in one case a perceived lack of it were contemplated as well as potentially different circumstances in case of not being a network member.

“We are simply a care network that is not after the money.”

“So, I think, in this context [in the network], you generally mind your prescribing behavior and of course now in particular you mind your antibiotics prescribing, too.” Phys22, Pos. 17 and 50

“We have been holding quality circles for the last 20 years anyway. We built a network on this. We are always the same group sitting together and we all work pharma-independently.” Phys17, Pos. 16

“Quality circles are obligatory. If one says ‘I can stay as I am’, he is not an adequate network physician, if you like, but this ultimately means change.” Sh05, Pos. 56

“Thinking I would not be in the network and not have the [peer] exchange options, perhaps one would not have participated in the study at all, but overall, we probably would act differently and not attach so much importance to that.” Phys22, Pos. 50

Management

Generally, the management teams of PCN were seen positive and active in aiming for a strong common appearance, provision of information and training on new developments, evidence-based guideline-oriented care, and opportunities for peer exchange. In some cases, political involvement in contractual negotiations with health insurers were reflected on. One physician explicitly described the network manager as a go-ahead character who promoted digitalization efforts. This was the only mention of someone being considered an opinion leader regarding specific topics. One stakeholder saw network management in this role. Participants who were involved in network or stakeholder management gave their managerial viewpoint.

“... aiming to produce a certain corporate identity.” Phy04, Pos. 36

“I am a network board member ... it is important to me to absolutely provide know-how.” Phys24, Pos. 38

“I don’t think physicians join a network and say ‘I want to practice a better medicine’, but this eventually has to be defined by the management, ... it’s an ambitious pledge and I think it only works in certain areas,

... and this is where I see ARena.”

Sh05, Pos. 48

“...the network has a very honorable manager, a big IT promoter who ... co-developed a very good data security concept that practices can implement quite safely.” Phys09, Pos. 30

Social contagion

Interacting and peer exchange

Participants generally cited positive attributions regarding aspects of interaction and peer exchange and a potential impact that physicians might have on each other within the network. Some physicians expressed that they explicitly wanted to work in a PCN member practice to be able to make use of regular peer exchange. With regards to the role PCN might play in supporting rational prescribing and use of antibiotics for non-complicated infections, they pointed to coordination of care-related aspects and the importance of regular peer exchange and interaction.

“... I think we are more focused and faster, because I have the same thoughts as the cardiologist or the nephrologist...” Phys27, Pos. 78

“Well, we have seen – as a single physician, there is no point in reading something. When we read as a group and the others say ‘yes, I do this, too’, ‘I will do it this way from now on, too, makes sense’, then I am more likely to implement it in my practice, because I know: OK, my peers do this, too. It’s an important effect.”

Phys17, Pos. 28

“I still think that the really decisive point is that we meet once a month. One evening, about three hours, and we can say – we are really proud to say - that 45 to 55 out of 65 physicians participate, so we can really achieve a lot, so we can really make sure almost all physicians are on the same page.” Phys04, Pos. 40

“That is why I explicitly looked for a practice where this [peer exchange] is possible.” Phys08, Pos.56

“Regarding ARena, the network event we had was particularly interesting to me and the discussion with the colleagues there.” Phys21, Pos. 38

Potential influences of interacting in PCN were also reflected critically with regards to antibiotics prescribing decisions, own attitude and using peer exchange windows of opportunity to exert educative influence. Physicians noted that sometimes networks lost members and it could be difficult to gain new ones. One physician described recruiting all physicians in specialist training in his practice. There was no indication of defined non-admission criteria. While physicians focused more on providing contagion in PCN, stakeholder and MAs contemplated contagion processes.

“It definitely has an effect for the patients, because we do evidence-based, scientifically sound medicine. But this has not changed because I am a network member, but we founded the network so we could infect others [with ideas].”

Phys17, Pos. 36

“And sometimes, I consciously use this network to hint something to certain colleagues who show a modus operandi I consider inappropriate, without having to face instant harassment.” Phys02, Pos. 20

“The network in the end does not have any influence on a physician’s decision, I think this remains with the physician.” Sh01, Pos. 46

“We actually always have a team meeting when there is something new everyone should know about, and has been discussed [in a quality circle], and then we discuss it in the meeting and see how we can implement or improve certain things in our routines.” MA01, Pos. 66

Translating to peers and team

Physicians and MAs reflected on exchanging information and translating knowledge within their practices and networks in general as well as in the study context. While physicians again focused on providing information to peers, MA took a more reciprocal perspective and took pride in translating knowledge back to their place of work.

“On my end, I try to incorporate this ENT specialist view, to do genuine networking, to inform my colleagues, to give food for thoughts, reflect experiences I gain in my practice.” Phys24, Pos. 38

“My colleague brings the information [to us]. And in turn she takes our ideas, developments, questions back to the quality circle.” MA10, Pos. 318

“Sometimes it is very cool, there is this MA circle I joined, and you talk to peers, [about their routines], and then you are back at work and can narrate.”

MA11, Pos. 52

Existing network structures regarding information dissemination and peer exchange were seen as promoting factors for knowledge translation and following up with new ideas. Participants reported using in-person meetings and direct phone calls, printed information media, video conferences and communication applications within their PCN peers. There was no mention of fixed or loose subgroups in networks other than in connection to medical specialty or organizational level and related topics. There was little mention of communicative exchange between networks. Regarding ARena, participants emphasized their networks’ support for the implementation of the intervention components, following through with study participation and the potential to reach a larger group.

“... I believe that network structures certainly ... promote trying out even more, because both personal and administrative structures are in place.” Sh07, Pos. 84

“... in this case here, of course, it works because of structures, e.g. a primary care network where you have participants who represent an intersecting subset of the overall network and when you communicate and inform via projects, of course knowledge and information can get translated to a larger part of the network.”

Sh03, Pos. 80

“Ok, one can easily say that without it [the network], we probably would not have followed through as consistently.” Phys19, Pos. 62

Discussion

To assess the contribution of PCN regarding the promotion of a rational antibiotics prescribing for non-complicated infections, this study explored factors attributed to the network’s role with a focus on the impact physicians might have on each other. We found evidence for the role of various processes of social influence in this context: social support, learning, normative pressures and contagion. In particular, regular thematic peer exchanges as possible in quality circles and during continued training opportunities offered by PCN, were emphasized by the participants as being of high value and support to them and considered to be one of the great assets the membership in a PCN entails. The contagion impact physicians might have on each other became apparent through acknowledgement of a higher likelihood of implementing new ideas and behaviors into practice after ascertaining that peers do so, too. These findings suggest that an interaction of professional peer exchange and a shared attitude within PCNs combined with intervention components as offered in the ARena study can promote a more reflected management of antibiotics prescribing. To achieve this, PCN engaged in efforts of knowledge transfer and manifestation of best practice care by providing a platform for refreshing awareness and knowledge and facilitating self-reflection of prescribing practice for their members.

PCN have been set up with specific purposes and for specific activities, such as quality improvement efforts and they aim for above-average quality of and increased patient satisfaction with care. Inherent might be that only individuals who share these aims are attracted to work in PCN while others prefer unrestricted autonomy and therefore stay outside. This may imply selection processes that influence social influences and their impact in PCNs. If these fail to be considered adequately, social influence processes and potentially present opinion leadership are likely to be overestimated [26]. It is known that physicians influence each other, but the magnitude of peer influence is still poorly understood [27]. In this study however, participants attached general importance to social learning and the social support offered by their PCN: Peer exchange was strongly emphasized as a major asset while there was hardly any mention of individual local opinion leaders. This is particularly notable as opinion leaders are a frequently examined aspect of social networks and considered to be a strong influence on other network members as they seem to amplify dissemination of ideas and information [28]. A Cochrane review confirmed that opinion leaders can promote evidence-based practice and reported a 10.5% increase in compliance when used as a strategy [29]. However, several studies failed to identify and clearly describe the role of clinical opinion leaders, particularly in ambulatory care settings [29, 30]. Physicians participating in this study

only mentioned opinion leaders in their PCN when asked general questions about PCN management and scarcely regarding a rational use of antibiotics. Therefore, it is possible that opinion leaders in this specific setting and context would be comparably less effective, because hierarchical structures in general appear less suitable here [31]. Based on their training and socialization and high degree of autonomy in their professional practice, hierarchical leadership relations in a PCN may not even appeal to physicians or might cause conflicts in their original practice [32].

Taking a social network perspective enables understanding the diffusion of new ideas and behaviors and the process by which they are communicated over time. [33]. It also facilitates understanding and evaluation of implementation processes [34] and potentially achieved behavior changes in network members. The adoption of new ideas and behaviors by individuals is strongly influenced by the quality and structure of their social network [34, 35]. Physicians tend to join more informal horizontal networks which are more effective for peer influence and constructing and reframing of meaning [36]. In such peer networks, interacting physicians likely share their beliefs, ideas and experiences with each other and these interpersonal exchanges may influence practice patterns [14]. Such social influence processes tend to homogenize ideas and behaviors within networks. Therefore, from a quality improvement and evidence-based practice perspective, guidance in form of accepted educative programs, practice guidelines, or opinion leadership seems important to propel ideas and behaviors in the desired direction. In this study, physicians valued communicative peer interaction very highly and seemingly attributed less importance to opinion leadership than to the actual matter being peer discussed within their PCN. Participation in the study, evidence base and existing guidelines might have acted as strong motivation and push factors. Nevertheless, even if acting as opinion followers, the participants might still be viewed as opinion leaders themselves as they discussed a new behavior in peer exchanges, adopted it and carried it into their own practice.

Social influence processes, and particularly social contagion after communicative interaction, address the tendency of adopting ideas, behaviors and attitudes of connected individuals in networks [37, 38]. Contagion theory does not require an intent or even awareness to influence, only that communication takes place [39]. This study suggests that PCN partly function as vehicle and amplifier of social influence processes, such as imitation, role modelling and persuasion by providing a highly valued platform for communicative peer exchange. Thus, they can support efforts to promote a more reflected use of antibiotics.

Positive effects on quality of care and patient outcomes are considered to be present in PCN when adequate resources and credible, efficient management are coupled with effective communication strategies and collaborative trusting relationships [40]. Close, trusting collaboration and communication among network members and increasing utilization of digital networking, paired with efforts of structured quality management and upskilling also constitute the base for the PCN aim for above-average quality of

care and increased patient satisfaction [3]. PCN participating in the ARena study provided those resources and collaborative trusting environment required to support the reflected use of antibiotics regarding acute non-complicated infections and hereby demonstrated that they potentially can play an even more important role in German healthcare. Nevertheless, to soundly identify influence processes at work, a follow-up regarding sustained behavior changes appears necessary.

Strengths And Limitations

For the ARena study and the implemented intervention components, the PCN constituted an appropriate choice to deliver the implementation program as generally interventions delivered by a peer community can be expected to be more successful than those delivered by agencies less connected to program recipients [34]. In addition to the qualitative analysis, data referencing PCN were extracted from surveys T0 and T1 and included to ensure a broad thematic spectrum could be analyzed within the process evaluation to assess and understand factors relevant to the role of primary care networks regarding support for rational use of antibiotics for non-complicated infections. The purposive sampling strategy supported the identification of individuals who were especially knowledgeable about and experienced in the phenomenon of interest and was chosen as it supports a detailed exploration and broad understanding of central themes and relations to specific experiences, roles and behaviors [41]. Reporting of this study follows the COREQ guideline and the CONSORT Statement where applicable [42, 43].

Some limitations of this study have to be reported. Participating PCN already were susceptible to the topic and aware of implications which allows for a selection bias and possibly social desirability of shared perceptions. The influence of social processes might be overestimated as social selection processes were not considered in this study. All reported findings require cautious interpretation with regard to a complete and profound understanding of social influence processes. Combination and triangulation of all generated data is expected to provide further insights after concluding analysis.

Conclusions

PCN provide a platform for social influence processes regarding professional peer exchange, social support and reassurance. Their impact on antibiotics prescribing for acute non-complicated infections is based on facilitating and amplifying the delivered quality improvement program by refreshing awareness, knowledge and self-reflection among larger groups of care providers. Thus, they are well suited to promote a rational use of antibiotics.

Abbreviations

ARena - Sustainable reduction of antibiotic-induced antimicrobial resistance

(German: Antibiotika-Resistenzentwicklung nachhaltig abwenden)

COREQ - Consolidated criteria for reporting qualitative research

G-BA	- Federal Joint Committee (German: Gemeinsamer Bundesausschuss)
MA	- medical assistant
PCN	- primary care networks
Phys	- physician
Sh	- stakeholder

Declarations

Ethics approval and consent to participate

This study received ethical approval by the medical ethics committee of the Medical Faculty of Heidelberg University provided approval for the study (S-353/2017) and the ethics committee of the medical Association Baden-Wurtemberg (B-F-2017-104).

Consent for publication

Not applicable

Availability of data and materials

All data generated and analyzed during this study are not publicly available since the information and consent procedure does not facilitate secondary use of the data.

Competing interests

The authors declare that they have no competing interests.

Funding

The ARena study received funding from the Innovation Committee at the Federal Joint Committee (G-BA), Berlin (01NVF16008). The funder had no role in the design, data-collection, data-analysis, interpretation, or writing of the paper. We acknowledge financial support by Deutsche Forschungsgemeinschaft within the funding programme Open Access Publishing, by the Baden-Württemberg Ministry of Science, Research and the Arts and by Ruprecht-Karls-University Heidelberg.

Authors' contributions

RPD drafted and prepared the manuscript. MW, MKa, KG, RPD, PKK and EA contributed to

concept and design of the study. RPD and MK conducted the interviews, analyzed transcripts and interpreted the data. MK and RPD analyzed and interpreted survey data. RPD interpreted and phrased

results from qualitative and quantitative data. All authors provided substantial comments and approved the final version of the manuscript.

RPD – Regina Poss-Doering; MK – Martina Kamradt; MW – Michel Wensing; KG – Katharina Glassen; EA – Edith Andres; PKK – Petra Kaufmann-Kolle

Acknowledgements

The authors would like to thank all participating interview partners of this study and the aQua Institute, which gave overall support to conducting the ARena study and the recruitment of the participating physicians, medical assistants and stakeholders. Our sincerest thanks for their support also go to all partners involved: Agency of German PCN (Agentur deutscher Arztnetze), statutory health insurers AOK Bayern, AOK Rheinland-Hamburg, and Federal AOK Association (AOK Bundesverband) and the Bavarian association of statutory health insurance physicians (KV Bayerns). A thank you also goes to Anna Stuermlinger for her participation in conducting interviews and to the staff at the Department of General Practice and Health Services Research at the University Hospital, Heidelberg who were involved in transcribing the qualitative data.

References

1. Agentur deutscher Arztnetze. Ueber Netze. Was sind Arztnetze? 2014. http://deutsche-aerztenetze.de/ueber_netze/was_sind_arztnetze.php. Accessed 13 Mar 2019.
2. Wenger E, Wenger-Trayner B. Communities of practice: a brief introduction. 2015:1–8.
3. Wambach V, Lindenthal J. Den Kinderschuhen entwachsen - Arztnetze in Deutschland leisten wertvollen Beitrag zur Optimierung der lokalen Versorgungssituation. Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz. 2015;58:374–82. doi:10.1007/s00103-015-2119-4.
4. Schrewe C. Gesundheit gemeinsam verantworten. Zur Position der Praxisnetze in Deutschland. 1016. http://deutsche-aerztenetze.de/uploads/files/positionspapier_v13_screen.pdf. Accessed 13 Mar 2019.
5. Gabriel J. Praxisnetze im Wandel - Chancen und Staerken eines Versorgungsmodells. In: Pfannstiel MA, Focke A, Mehlich H, editors. Management von Gesundheitsregionen III: Gesundheitsnetzwerke zur Optimierung der Krankenversorgung durch Kooperation und Vernetzung. Wiesbaden: Springer Fachmedien Wiesbaden; 2017. p. 13–24. doi:10.1007/978-3-658-13658-1_2.
6. Kamradt M, Kaufmann-Kolle P, Andres E, Brand T, Klingenberg A, Glassen K, et al. Sustainable reduction of antibiotic-induced antimicrobial resistance (ARena) in German ambulatory care: Study protocol of a cluster randomised trial. Implement Sci. 2018;13:23. doi:10.1186/s13012-018-0722-0.
7. aQua – Institut für angewandte Qualitaetsfoerderung und Forschung im Gesundheitswesen GmbH. Was ist ARena? 2018. <https://www.arena-info.de/>. Accessed 15 Oct 2018.

8. Kuypers MS, Koeberlein-Neu J. solimed Pflegemanagement als Beispiel einer regionalen Versorgung durch ein Gesundheitsnetz. In: Pfannstiel MA, Focke A, Mehlich H, editors. Management von Gesundheitsregionen III: Gesundheitsnetzwerke zur Optimierung der Krankenversorgung durch Kooperation und Vernetzung. Wiesbaden: Springer Fachmedien Wiesbaden; 2017. p. 99–107. doi:10.1007/978-3-658-13658-1_10.
9. Kunhardt H, editor. Systemisches Management im Gesundheitswesen: Innovative Konzepte und Praxisbeispiele. Wiesbaden: Gabler Verlag / Springer Fachmedien Wiesbaden GmbH Wiesbaden; 2011.
10. Lindenthal J. Wissensmanagement in medizinischen Praxisnetzen – Erfahrungen aus dem Gesundheitsnetz Qualitaet und Effizienz – QuE Nuernberg. In: Kunhardt H, editor. Systemisches Management im Gesundheitswesen: Innovative Konzepte und Praxisbeispiele. Wiesbaden: Gabler Verlag / Springer Fachmedien Wiesbaden GmbH Wiesbaden; 2011. p. 159–185. doi:10.1007/978-3-8349-6440-3_8.
11. Tannenbaum SS, Soulos PR, Herrin J, Pollack CE, Xu X, Christakis NA, et al. Surgeon peer network characteristics and adoption of new imaging techniques in breast cancer: A study of perioperative MRI. *Cancer Med.* 2018;7:5901–9. doi:10.1002/cam4.1821.
12. Muller E, Peres R. The effect of social networks structure on innovation performance: A review and directions for research. *International Journal of Research in Marketing.* 2019;36:3–19. doi:10.1016/j.ijresmar.2018.05.003.
13. Christakis NA, Fowler JH. Social contagion theory: examining dynamic social networks and human behavior. *Stat Med.* 2013;32:556–77. doi:10.1002/sim.5408.
14. Keating NL. Peer Influence and Opportunities for Physician Behavior Change. *J Natl Cancer Inst.* 2017;109:djx009. doi:10.1093/jnci/djx009.
15. Pollack CE, Soulos PR, Herrin J, Xu X, Christakis NA, Forman HP, et al. The Impact of Social Contagion on Physician Adoption of Advanced Imaging Tests in Breast Cancer. *J Natl Cancer Inst.* 2017;109:djw330. doi:10.1093/jnci/djw330.
16. Pollack CE, Soulos PR, Gross CP. Physician's peer exposure and the adoption of a new cancer treatment modality. *Cancer.* 2015;121:2799–807. doi:10.1002/cncr.29409.
17. Pollack CE, Weissman G, Bekelman J, Liao K, Armstrong K. Physician social networks and variation in prostate cancer treatment in three cities. *Health Serv Res.* 2012;47:380–403. doi:10.1111/j.1475-6773.2011.01331.x.
18. Pollack CE, Wang H, Bekelman JE, Weissman G, Epstein AJ, Liao K, et al. Physician Social Networks and Variation in Rates of Complications After Radical Prostatectomy. *Value in Health.* 2014;17:611–8. doi:10.1016/j.jval.2014.04.011.
19. Freund T, Everett C, Griffiths P, Hudon C, Naccarella L, Laurant M. Skill mix, roles and remuneration in the primary care workforce: Who are the healthcare professionals in the primary care teams across the world? *Int J Nurs Stud.* 2015;52:727–43. doi:10.1016/j.ijnurstu.2014.11.014.

20. Ajzen I. The theory of planned behavior. *Organizational Behavior and Human Decision Processes*. 1991;50:179–211. doi:10.1016/0749-5978(91)90020-T.
21. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol*. 2013;13:117. doi:10.1186/1471-2288-13-117.
22. Flottorp SA, Oxman AD, Krause J, Musila NR, Wensing M, Godycki-Cwirko M, et al. A checklist for identifying determinants of practice: A systematic review and synthesis of frameworks and taxonomies of factors that prevent or enable improvements in healthcare professional practice. *Implement Sci*. 2013;8:35. doi:10.1186/1748-5908-8-35.
23. Mason WA, Conrey FR, Smith ER. Situating social influence processes: dynamic, multidirectional flows of influence within social networks. *Pers Soc Psychol Rev*. 2007;11:279–300. doi:10.1177/1088868307301032.
24. Marsden PV, Friedkin NE. Network Studies of Social Influence. *Sociological Methods & Research*. 1993;22:127–51. doi:10.1177/0049124193022001006.
25. Tasselli S. Social Networks of Professionals in Health Care Organizations: A Review. *Med Care Res Rev*. 2014;71:619–60. doi:10.1177/1077558714557079.
26. Friemel TN. Influence Versus Selection: A Network Perspective on Opinion Leadership. *International Journal of Communication*. 2015;9(2015), 1002–1022.
27. Donohue JM, Guclu H, Gellad WF, Chang C-CH, Huskamp HA, Choudhry NK, et al. Influence of peer networks on physician adoption of new drugs. *PLoS ONE*. 2018;13:e0204826. doi:10.1371/journal.pone.0204826.
28. Dearing JW. Improving the State of Health Programming by Using Diffusion Theory. *Journal of Health Communication*. 2004;9:21–36. doi:10.1080/10810730490271502.
29. Flodgren G, O'Brien MA, Parmelli E, Grimshaw JM. Local opinion leaders: effects on professional practice and healthcare outcomes. *Cochrane Database of Systematic Reviews* 2019. doi:10.1002/14651858.CD000125.pub5.
30. Grimshaw JM, Eccles MP, Greener J, MacLennan G, Ibbotson T, Kahan JP, Sullivan F. Is the involvement of opinion leaders in the implementation of research findings a feasible strategy? *Implement Sci*. 2006;1:3. doi:10.1186/1748-5908-1-3.
31. Staar H, Atzpodien HC. Vernetzte Zusammenarbeit erfolgreich gestalten – eine Analyse der Relevanz sozialer Fertigkeiten der Mitglieder in Gesundheitsnetzwerken. Gruppe. Interaktion. Organisation. Zeitschrift für Angewandte Organisationspsychologie (GIO). 2017;48:25–40. doi:10.1007/s11612-017-0352-z.
32. Hildebrandt, H., Stunder, B., Wetzel, M., Groene, O. (2016). Erfolgsfaktoren für Netze und regionale Gesundheitsorganisationen: Organisationsformen, Führung, Patientenorientierung. In: Eble, Kurscheid (Hg.) 2016 – Gesundheitsnetzwerke, Strategie Konzeption, Steuerung. p. 60–73.
33. Rogers EM. *Diffusion of innovations*. New York, London, Toronto, Sydney: Free Press; 2003.

34. Valente TW, Palinkas LA, Czaja S, Chu K-H, Brown CH. Social network analysis for program implementation. PLoS ONE. 2015;10:e0131712. doi:10.1371/journal.pone.0131712.
35. Green J, Thorogood N. Qualitative methods for health research. 4th ed.; 2018.
36. Greenhalgh T, Robert G, Macfarlane F, Bate P, Kyriakidou O. Diffusion of innovations in service organizations: systematic review and recommendations. Milbank Q. 2004;82:581–629. doi:10.1111/j.0887-378X.2004.00325.x.
37. Veenstra R, Dijkstra JK, Steglich C, Van Zalk, Maarten H. W. Network-Behavior Dynamics. J Res Adolesc. 2013;23:399–412. doi:10.1111/jora.12070.
38. Burt RS. Social Contagion and Innovation: Cohesion versus Structural Equivalence. American Journal of Sociology. 1987;92:1287–335. doi:10.1086/228667.
39. Scherer CW, Cho H. A Social Network Contagion Theory of Risk Perception. Risk Analysis. 2003;23:261–7. doi:10.1111/1539-6924.00306.
40. Brown BB, Patel C, McInnes E, Mays N, Young J, Haines M. The effectiveness of clinical networks in improving quality of care and patient outcomes: A systematic review of quantitative and qualitative studies. BMC Health Serv Res 2016. doi:10.1186/s12913-016-1615-z.
41. Ritchie J, editor. Qualitative research practice: A guide for social science students and researchers. 1st ed. London u.a.: Sage; 2003.
42. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. Int J Qual Health Care. 2007;19:349–57. doi:10.1093/intqhc/mzm042.
43. Schulz KF, Altman DG, Moher D, the CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. BMC Medicine. 2010;8:18. doi:10.1186/1741-7015-8-18.

Tables

Source	Physicians	Medical assistants	Stakeholders	Description
Interviews (n)	27	11	7	Over telephone
Socio-demographic questionnaire (n)	27	11	7	Paper based
Notes (n)	14	3	2	Paper based and electronically
Survey T0 (n)	229	80	-	Paper based
Survey T1 (n)	200	73	-	Paper based

Table 1. Data collection sources

Interview participants	N	Phys	MA	Stakeholder	Total
Sex (f/m) n (%)	45	9/18 (33/66)	11 (100/0)	3/4 (43/57)	23/22 (59/41)
Age years (range) (mean)	45	43-66 (55.2)	20-60 (38.5)	31-63 (46.3)	31.3-63 (46.6)
Experience in current position years (range) (mean)	45	10-38 (26.07)	2-40 (19.3)	1-10 (5.85)	1-40 (17.09)
Working in general practice (%)	38	66.6	81.8	-	74.2
Part-time employment n (%)	4	1 (2.7)	3 (27.3)	-	4 (8.88)
Practice network member years (range) (mean)	27	2-23 (10.18)	-	-	10.18
Additional qualifications n	7	-	7	-	7
Survey respondents					
Sex (f/m) n (%)	304	229 (148/76)	80 (100/0)	-	228/76 (74/26)
Age years (range) (mean)	299	35-73 (54.4)	19-61 (38.7)	-	19-73 (46.5)
Working experience years (range) (mean)	306	5-48 (25.4)	1-40 (19.2)	-	1-48 (22.3)
Working in general practice (%)	309	75.3	76	-	75.6
Resident years (range) (mean)	220	1-41 (17.7)	-	-	220 (17.7)
Network member years (range) (mean)	207	0-28 (10)	-	-	10
Participating in network events times/year (range)	217	7.3 (0-50)	-	-	7.3 (0-50)

Table 2. Characteristics of the interview sample and survey respondents (T0)

Participating in the network ...	Agree %		Neutral %		Disagree %	
	T0	T1	T0	T1	T0	T1
...motivates guideline-oriented patient care	70.5	60	18.5	19	11	22
...supports shared-decision making	60.8	59	19.8	24	19.4	18
...supports managing patient expectations	61.2	51	21.6	30	17.2	19
...supports implementing new routines	74	59	16.3	18	9.7	24
...has an impact on antibiotics prescribing decisions	43.3	36	22.1	22	34.5	43

Table 3. Physician perspective on network participation

In my primary care network ...	Agree (%)		Neutral (%)		Disagree (%)	
	T0	T1	T0	T1	T0	T1
...antibiotics therapy is discussed	89.5	86	8.8	10	1.7	4
...peer exchange about guideline-oriented antibiotics therapy is offered	79.9	79	14.5	11	5.6	10
...exchange about antibiotics prescribing for no-complicated infections	71.5	73	18.4	16	10.1	11
...there are conventions about antibiotics for non-complicated infections	65.8	72	21.5	16	12.7	12
...training on guideline-oriented antibiotics therapy is offered	89	75	6.6	18	4.4	7
...I participated in training o guideline-oriented antibiotics therapy	89	87	6.6	9	4.4	4

Table 4. Results from survey T0 and T1 referring to training and peer exchange

Figures

Figure 1

Analytical Process

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORT2010Checklist.doc](#)