Prevalence of COVID-19 in Children Affected by Allergic Rhinoconjunctivitis and Asthma: Results From the Second “SIAIP Rhinosinusitis and Conjunctivitis Committee” Survey

Giulia Brindisi (✉ giulia.brindisi@gmail.com)

Università degli Studi di Roma La Sapienza Facoltà di Medicina e Odontoiatria <https://orcid.org/0000-0001-5059-6082>

Anna Maria Zicari

Sapienza University of Rome: Università degli Studi di Roma La Sapienza <https://orcid.org/0000-0002-2476-1790>

Giuseppe Fabio Parisi

University of Catania: Università degli Studi di Catania

Lucia Diaferio

Università degli Studi di Bari: Università degli Studi di Bari Aldo Moro

Cristina Indolfi

University of Campania Luigi Vanvitelli: Università degli Studi della Campania Luigi Vanvitelli

Giuseppe Marchese

Brescia University

Daniele Giovanni Ghiglioni

Ospedale Maggiore di Milano Policlinico: Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico

Giuseppina Rosaria Umano

University of Campania Luigi Vanvitelli: Università degli Studi della Campania Luigi Vanvitelli

Angela Klain

University of Campania Luigi Vanvitelli: Università degli Studi della Campania Luigi Vanvitelli

Gian Luigi Marseglia

IRCCS S Matteo: Fondazione IRCCS Policlinico San Matteo

Michele Miraglia Del Giudice

University of Campania Luigi Vanvitelli: Università degli Studi della Campania Luigi Vanvitelli

Research Article

Keywords: SARS CoV2, COVID - 19, children, asthma, allergic rhinoconjunctivitis, online-survey

Posted Date: August 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-757612/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Italian Journal of Pediatrics on January 6th, 2022. See the published version at <https://doi.org/10.1186/s13052-021-01198-y>.

Abstract

Background

The role of allergic sensitization seems to be protective against SARS CoV2 infection. The aim of this study was to evaluate, using online surveys, the impact of COVID-19 on Italian allergic children, comparing the prevalence of AR and asthma symptoms between the first and second pandemic wave.

Methods

Both surveys were emailed to Italian pediatricians in April 2020 (first survey) and in March 2021 (second survey). The first one was related to the impact of COVID-19 and the most frequently reported symptoms. The second one was superimposed on the previous one, taking into account some additional aspects in the management of disease.

Results

A total of 99 pediatricians participated in the first survey and 267 in the second one. The first survey showed that, asthma and allergic rhinoconjunctivitis prevalence was mostly between 0-20% throughout the country. The second survey showed a lower prevalence of both diseases nationwide in comparison to the first one. Comparing the two surveys, statistically significant differences were reported only in the distribution of asthma prevalence in Southern Italy while no differences were highlighted in the North and in the Center. Finally regarding allergic rhinoconjunctivitis prevalence, no differences were noticed nationwide.

Conclusions

Allergic rhinoconjunctivitis and asthma, if under control, did not represent risk factors for the susceptibility to SARS CoV2. Therefore, it is strongly recommended to continue therapies during COVID-19 outbreak, according to the international guidelines. However, being COVID-19 a new disease, actual knowledge will undergo continuous improvements over time.

Background

The current outbreak of novel coronavirus disease 2019 (COVID-19), declared pandemic in March 2020 by the World Health Organization (WHO) has caused millions of deaths all over the world ¹. Italy is considered one of the most hit countries, with 4.248.225 reported cases and 126.758 deaths, at the time of this writing ².

Although in adulthood the rate of symptoms severity and mortality is higher than in children, the appearance of several “severe acute respiratory syndrome coronavirus 2 (SARS CoV2)” variants has made the pediatric population more susceptible to COVID-19 compare to the beginning of the pandemic ³.

Common symptoms in children are fever, gastrointestinal and skin manifestations, arthralgia, anosmia and/or ageusia, rhinorrhea, cough and dyspnea that only rarely require hospitalization in intensive care units ^{4,5}.

Allergic rhinitis (AR) and asthma are common allergic diseases, often associated with each other for the anatomical continuity between upper and lower airways. Furthermore, they share the same inflammatory

pathway, characterized by a T helper 2(Th2) response with the presence of several biomarkers accountable for the characteristic chronic inflammation throughout the bronchial tree ⁶.

As both AR and asthma can present an overlap with COVID-19, a careful monitoring of these symptoms and a prompt differential diagnosis is desirable ⁷.

To date, the role of allergic sensitization has been reconsidered in relation to COVID-19, no longer as a risk factor but on the contrary as a protective one against SARS CoV2 infection in several studies. Du H et al. analyzed 182 children hospitalized for COVID-19 of whom 23.6% had allergic disease and only one child of these reported asthma. The findings of this study showed no statistically significant differences in disease severity, development of complications, and clinical symptoms between the allergic and non-allergic group⁸.

Many are the possible explanations for not considering allergy as a risk factor for SARS CoV2 infection. One of them is the protective role of eosinophils and the related Th2 phenotype associated with allergy, with a low Interferon- α (IFN- α) production; in fact it seems that the higher the eosinophil count is, the better the prognosis of COVID-19 is with reduced morbidity and mortality ⁹.

Additionally, respiratory allergies are associated with a reduced expression of angiotensin conversion enzyme receptor 2 (ACE2), the well-known entrance door of SARS CoV2 into the upper airways cells. Th2 response is able to modulate ACE2 expression with an intriguing role in COVID-19 pathogenesis, acting on the Th1/Th2 balance ¹⁰. Therefore it seems that Th2 inflammation exerts a protective role against SARS CoV2 ¹¹.

Lastly, several studies show that asthma medications such as inhaled steroids, bronchodilators and montelukast, present an antiviral and immunomodulatory effect that inhibits SARS CoV2 driven inflammation¹².

In line with the scientific literature, data derived from the first survey on the impact of COVID-19, realized by the "SIAIP rhinosinusitis and conjunctivitis committee", concluded that allergic rhinoconjunctivitis and asthma were not risk factors for the development of severe forms of COVID-19 ¹³. On the other hand, the presence of uncontrolled allergic symptoms can represent a marker of poor prognosis and disease severity ¹⁴.

Therefore, the achievement of both AR and asthma control appears fundamental not only in the daily clinical practice ¹⁵ but above all during pandemic ¹⁴.

Following the first survey, related to the period of December-May 2020, the "SIAIP rhinosinusitis and conjunctivitis committee" decided to issue a new questionnaire referring to the period of September-February 2021, aiming to assess nationwide the impact of COVID-19 among children and the presence of any differences compared to the first pandemic peak.

By analyzing the above mentioned surveys in this study we aimed to compare the prevalence of allergic rhinoconjunctivitis and asthma symptoms among children affected by COVID-19 in the North, Center and South of Italy.

Methods

Study Design and Population

The first “COVID-19 survey” was sent by email to Italian pediatricians in April 2020 and was composed of 20 multiple choice questions, related to the impact of COVID-19, the disease’s management and the most frequently reported symptoms¹⁶.

Then in March 2021 the second “Follow-up COVID-19 survey” was emailed to a greater number of Italian pediatricians and was composed by 35 multiple choice questions; it was superimposed on the previous one, by taking into account some additional aspects in the management of disease.

Both questionnaires were created by the “SIAIP rhino-sinusitis and conjunctivitis committee” in Italian, using Google-Drive Platform (Google Drive™, © 2012 Google Inc. all rights reserved).

The anonymous data received were automatically collected in an Excel® data format used for the statistical analysis. Both surveys took approximately 10 minutes to be answered. Each pediatrician was allowed to answer only once to avoid double responses. There was no need to obtain informed consent, given the voluntary nature of both surveys. The Ethics Committee of Sapienza University of Rome, did not consider any special permission necessary because these studies’ designs met the criteria of an audit activity.

Statistical Analysis

All the variables were categorical. Differences were investigated by Chi square test. Data were presented as frequency. All analyses were performed using SAS University Edition (SAS Institute, Cary, NC, USA), and statistical significance was assessed at the two-tailed 0.05 threshold.

Results

The already published data related to the first survey collected a total of 99 responses of which 32% from the North of Italy, 27.8% from the Center and 40.2% from the South¹⁶.

A total of 267 responses were derived from the second survey, of which 53.2% from the North, 27.4 % from the Center and 19.4 % from the South. Characteristics of the pediatricians who responded to both surveys are shown in Tables 1 and 2.

Table n 1. First Survey Participants’ characteristics ¹⁶

Total of participants	99
Types of pediatricians	
Primary care	60%
Pediatric hospital medicine	24.2%
Specialized outpatient healthcare	6.3%
Pediatric emergency medicine	6.3 %
Pediatric critical care medicine	2.1%
Pediatric infectious disease	1.1%
Territorial Subdivisions	
North	32%
Center	27.8 %
South an Islands	40.2%

Table n 2. Second Survey Participants' characteristics

Total of participants	267
Types of pediatricians	
Primary care	75.5%
Pediatric hospital medicine	16.3 %
Specialized outpatient healthcare	5.4%
Pediatric emergency medicine	1.9 %
Pediatric critical care medicine	0.9%
Pediatric infectious disease	0 %
Territorial Subdivisions	
North	53.2%
Center	27.4 %
South and Islands	19.4%

Analyzing the first survey, we found statistically significant differences for asthma prevalence among children affected by COVID-19 considering the different zones of Italy ($p = 0.03$).

In detail in the North, 94% of children presented a prevalence less than 20%, 3% between 20–40 % and 3% greater than 40%. In the Center, 92% of children featured a prevalence less than 20%, 8% between 20–40 % and no one greater than 40%. Lastly in the South, 67% of children showed a prevalence less than 20%, 27% between 20–40 % and 6% greater than 40%. In particular, the South of Italy showed the highest rate of asthma prevalence compared to the North and Center. (Fig. 1)

Also, the second survey showed statistically significant differences ($p = 0.03$) for asthma prevalence in the considered zones of Italy, as following: the Center had the same prevalence of the South, whereas the distribution in the North did not change significantly in comparison to the first survey.

In particular, the North presented a prevalence less than 20% in 94% of children, between 20–40 % in 5 % and greater than 40% in 1%. In the Center and South, the totality of affected children (100%) presented a prevalence less than 20%. (Fig. 2)

Considering symptoms of allergic rhinoconjunctivitis, in the first survey we observed a trend for a higher rate of symptoms in the Center and South of Italy, however these differences did not reach statistical significance ($p = 0.08$). The North showed a prevalence less than 20% in 87% of children, between 20–40 % in 10% and greater than 40% in 3%. In the Center, 73% of children reported a prevalence less than 20%, 15% between 20–40 % and 12% greater than 40%. Lastly in the South, 64% reported a prevalence less than 20%, 33% between 20–40 % and only 3% greater than 40%. (Fig. 3)

In the second survey, Center and South Italy showed significant higher rates of allergic rhinoconjunctivitis compared to North ($p = 0.049$) where 89% of children reported a prevalence less than 20%, 10% between 20–40 % and no one greater than 40%. In the Center, 84% showed a prevalence less than 20%, 14% between 20–40 % and 2% greater than 40%. Lastly in the South, 75% of children had a prevalence less than 20%, 24 % between 20–40 % and only 1% greater than 40%. (Fig. 4)

Furthermore comparing the two surveys, statistically significant differences were reported only in the distribution of asthma prevalence in the South ($p < 0.0001$) while no differences were highlighted in the North and in the Center ($p > 0.05$); the first survey showed a higher rate of asthma compared to the second one. Finally regarding allergic rhinoconjunctivitis distribution, no differences were noticed nationwide ($p > 0.05$).

Discussion

Allergy does not appear to be a predisposing factor for the development of SARS CoV-2 infection or an aggravating one in case of disease¹⁶.

Dong et al., analyzing children and adults affected by COVID-19, reported that no patients with allergies presented critical symptoms or had a more severe course of illness than those without allergies. All this is in line with the hypothesis of a positive role exerted by the Th2 immune response in COVID-19 pathogenesis¹⁷.

Kimura et al., highlighted that IL-4 and IL-13, typical cytokines of a Th2 response, down-regulating ACE2 receptors in upper airways cells of allergic children, would exert protection against SARS CoV2 infection; otherwise the Th1 response, with interferon-gamma (IFN-gamma) release, increases ACE2 expression,

promoting SARS CoV-2 entry into airway cells. All these observations, confirm the role played by the Th1/Th2 balance in the pathogenesis of COVID-19 ¹⁸.

The systematic review of the literature, conducted by Morais-Almeida et al., supports these results, analyzing the relationship between asthma and COVID-19 in several studies and highlighting no evidence of a higher risk of being infected by SARS CoV2 or to develop a critical illness among asthmatic patients ¹⁹.

Hence, according to our results, children with asthma and allergic rhinoconjunctivitis present a low prevalence of SARS CoV2 infection, because the ACE2 receptors used by the virus to enter the respiratory tract are less expressed in allergic phenotypes. Conversely, other conventional coronaviruses or respiratory viruses are able to exacerbate asthma because they use other entry receptors different to ACE2²⁰.

Moreover, these ACE2 are also down-regulated by asthma therapies that must be continued during the outbreak, according to the international guidelines, to obtain the control of allergic symptoms, also achieved respecting the current social distances and public hygiene recommendations ¹⁴.

A good COVID-19 prognosis is also linked to a high blood eosinophil count, common finding in allergic diseases in the context of a Th2 response¹⁹. Du Y et al., reported that eosinopenia was found in about 80% of patients died for COVID-19, considering it as a biomarker of worse prognosis and a predictor for the development of COVID-19 severity ²¹.

So, although asthma and allergic symptoms are often associated to COVID-19 they did not represent risk factors for a more severe disease course and did not lead to a higher frequency of hospitalization ²². These results were also confirmed in a wide online survey conducted on 133.000 asthmatic children who reported mild symptoms and only one case of hospitalization for COVID-19 ²³.

Moreover, during the pandemic, the lower prevalence of asthmatic and allergic exacerbations is also attributable to greater public health hygiene measures, the use of facemasks, the lockdown periods and a prompter management of respiratory symptoms by parents ²⁴.

Obviously, symptoms of uncontrolled asthma and allergic symptoms, are considered risk factors for the development of severe COVID-19 ¹⁴.

For this reason, the Global Initiative for Asthma (GINA) claimed the necessity to achieve asthma control by continuing, even during the pandemic, therapies including biologic drugs, except in patients with active COVID-19 ^{25,26,27}.

However, in addition to the management of asthma, also AR requires to be adequately treated ⁷ because it may have an overlap of symptoms with COVID-19 infectious rhinitis. Uncontrolled AR can increase the risk of viral transmission in children with COVID-19. ARIA and EAACI reiterated the importance of performing a correct nasal therapy with intranasal corticosteroid to avoid the worsening of allergic symptoms and, in case of SARS CoV2 infection, the viral spread ²⁸.

On the other hand, it is recommended to discontinue allergen immunotherapy (AIT), as it decreases the TH2 response, in COVID-19 patients and to continue it in allergic patients without the infection ^{29,30}.

Summarizing, allergic patients present a Th2 response with a lower production of IFN- α in addition to the presence of eosinophils and low ACE2 expression in the airway tract; in addition, asthmatic patients benefit from the antiviral/ immunomodulatory effect exerts by asthma medications ¹².

All these factors act as a protective mechanism, reducing the hyperinflammation triggered by the virus to which the more severe clinic is linked.

In line with the literature, the results of both surveys, provide data on the prevalence of allergic rhinoconjunctivitis and asthma symptoms in children affected by COVID-19 nationwide, supporting the concept that controlled allergic symptoms do not constitute risk factors for SARS CoV2 susceptibility.

These results, obtained by the preliminary analysis of the first COVID-19 survey ¹³, were also confirmed by the second one. In particular, the first survey showed that, in relation to both asthma and allergic rhinoconjunctivitis, the general prevalence was mostly between 0–20% throughout the country. All in all, northern regions showed a lower prevalence of allergic symptoms than the Center and South.

The second survey highlighted a lower prevalence of asthma nationwide in comparison to the first one; in particular it confirmed that the North had the lowest prevalence of both asthma and allergic rhinoconjunctivitis and it reported a marked improvement in the prevalence of both diseases in the Center and South.

Comparing the two surveys, statistically significant differences were reported only in the distribution of asthma prevalence in Southern Italy, while no differences were found in the North and in the Center. Lastly regarding allergic rhinoconjunctivitis distribution, no differences were noticed at a national level.

In support of our findings, another Italian survey conducted among children with AR and/or asthma to evaluate allergic symptoms and the need of drug usage during lockdown, showed an overall trend of clinical improvement and a reduction of on demand and basal therapy for both AR and asthma. All these results can be attributable to various factors: the prompt management of allergic symptoms by parents and the minor exposure to pollens, viruses, and air pollution due to the lockdown ²⁴.

Similar results achieved by the already mentioned survey conducted by Papadopoulos et al., confirmed that children with asthma were not affected to a great extent by COVID-19, due to better adherence to preventive hygiene measures and improved treatment compliance ²³.

This study presents also some limitations. First of all, these two surveys refer to slightly different periods, enrolling a different number of participants. Furthermore, we used non standardized questionnaires that were filled out by pediatricians from different parts of Italy, carrying out heterogeneous activities among all of them. Anyway, to the best of our knowledge, standardized and validated questionnaires on these topics are not currently available.

Conclusions

Asthma and allergic rhinoconjunctivitis, if under control, did not represent risk factors for SARS CoV2 susceptibility or severity. Therefore, the need to continue both on demand and basal therapy during COVID-19 outbreak, according to the international guidelines. However, being COVID-19 a new disease, we expect that actual knowledge will undergo continuous improvements over time.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors have no conflicts of interest to disclose that could be perceived as prejudicing the impartiality of the research reported.

Funding

The authors did not receive any funding for the research. The Italian Paediatric Society for Allergy and Immunology (SIAIP) covers the article-processing charge.

Authors' contributions

MMDG and GLM conceived the survey, LD, GFP, GB, CI, GM, AZ, DGG conceptualized and designed the survey. GRU and AK collected and analysed the data. GB wrote the manuscript. GFP, GB, CI, GM, AZ, DGG reviewed the manuscript. MMDG and GLM performed a critical revision of the manuscript. All authors approved the final manuscript as submitted and agreed to be accountable for all aspects of the work.

Acknowledgments

We would like to thank all the pediatricians who took part in this survey.

References

1. WHO. Coronavirus Disease 2019 (COVID-19) Situation Report. Available online: <https://covid19.who.int/> (accessed on ... June 2021).
2. Istituto Superiore di Sanità. Sorveglianza Integrata COVID-19 in Italia. Available online: https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-19_7-luglio-

3. Brookman S, Cook J, Zucherman M, Broughton S, Harman K, Gupta A. Effect of the new SARS-CoV-2 variant B.1.1.7 on children and young people. *Lancet Child Adolesc Health*. 2021 Apr;5(4):e9-e10. doi: 10.1016/S2352-4642(21)00030-4. Epub 2021 Feb 10. PMID: 33581054; PMCID: PMC7906637.
4. Parisi GF, Brindisi G, Indolfi C, Diaferio L, Marchese G, Ghiglioni DG, Zicari AM, Miraglia Del Giudice M. Upper airway involvement in pediatric COVID-19. *Pediatr Allergy Immunol*. 2020 Nov;31 Suppl 26(Suppl 26):85-88. doi: 10.1111/pai.13356. PMID: 33236430; PMCID: PMC7753446.
5. Gori A, Leone F, Loffredo L, Cinicola BL, Brindisi G, De Castro G, Spalice A, Duse M, Zicari AM. COVID-19-Related Anosmia: The Olfactory Pathway Hypothesis and Early Intervention. *Front Neurol*. 2020 Sep 10;11:956. doi: 10.3389/fneur.2020.00956. PMID: 33013637; PMCID: PMC7511833.
6. Brindisi G, De Vittori V, De Nola R, Di Mauro A, De Castro G, Baldassarre ME, Cicinelli E, Cinicola B, Duse M, Zicari AM. The Role of Nasal Nitric Oxide and Anterior Active Rhinomanometry in the Diagnosis of Allergic Rhinitis and Asthma: A Message for Pediatric Clinical Practice. *J Asthma Allergy*. 2021 Mar 25;14:265-274. doi: 10.2147/JAA.S275692. PMID: 33790581; PMCID: PMC8006956.
7. Brindisi G, De Vittori V, De Castro G, Duse M, Zicari AM. Pills to think about in allergic rhinitis children during COVID-19 era. *Acta Paediatr*. 2020 Oct;109(10):2149-2150. doi: 10.1111/apa.15462. Epub 2020 Jul 19. PMID: 32627237; PMCID: PMC7361544.
8. Du, H.; Dong, X.; Zhang, J.J.; Cao, Y.Y.; Akdis, M.; Huang, P.Q.; Chen, H.W.; Li, Y.; Liu, G.H.; Akdis, C.A.; et al. Clinical characteristics of 182 pediatric COVID-19 patients with different severities and allergic status. *Allergy* 2020, doi:10.1111/all.14452.
9. Ferastraoaru D, Hudes G, Jerschow E, Jariwala S, Karagic M, de Vos G, Rosenstreich D, Ramesh M. Eosinophilia in Asthma Patients Is Protective Against Severe COVID-19 Illness. *J Allergy Clin Immunol Pract*. 2021 Mar;9(3):1152-1162.e3. doi: 10.1016/j.jaip.2020.12.045. Epub 2021 Jan 23. PMID: 33495097; PMCID: PMC7826039.
10. Jackson, D.J.; Busse, W.W.; Bacharier, L.B.; Kattan, M.; O'Connor, G.T.; Wood, R.A.; Visness, C.M.; Durham, S.R.; Larson, D.; Esnault, S.; et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J. Allergy Clin. Immunol*. 2020, 146, 203–206.e3.
11. Azkur,A.K.;Akdis,M.;Azkur,D.;Sokolowska,M.;vandeVeen,W.;Brüggen,M.C.;O'Mahoni,L.;Gao,Y.;Nadeau,K.;Akdis, C.A. Immune response to SARSCoV-2 and mechanisms of immunopathological changes in COVID-19. *Allergy* 2020, 75, 1564–1581.
12. Matsuyama, S.; Kawase, M.; Nao, N.; Shirato, K.; Ujike, M.; Kamitani, W.; Shimojima, M.; Fukushima, S. The inhaled steroid ciclesonide blocks SARS-CoV-2 RNA replication by targeting the viral replication-transcription complex in cultured cells. *J. Virol*. 2020.
13. Diaferio L, Parisi GF, Brindisi G, Indolfi C, Marchese G, Ghiglioni DG, Zicari AM, Marseglia GL, Miraglia Del Giudice M. Cross-sectional survey on impact of paediatric COVID-19 among Italian paediatricians: report from the SIAIP rhino-sinusitis and conjunctivitis committee. *Ital J Pediatr*. 2020 Oct 6;46(1):146. doi: 10.1186/s13052-020-00906-4. PMID: 33023616; PMCID: PMC7538039.
14. Brough, H.A.; Kalayci, O.; Sediva, A.; Untersmayr, E.; Munblit, D.; Rodriguez del Rio, P.; Vazquez-Ortiz, M.; Arasi, S.; Al- varo-Lozano, M.; Tsaouri, S.; et al. Managing childhood allergies and immunodeficiencies

- during respiratory virus epidemics—The 2020 COVID-19 pandemic: A statement from the EAACI-section on pediatrics. *Pediatr. Allergy Immunol.* 2020, 31, 442–448.
15. De Vittori, V.; Pacilio, A.; Indinnimeo, L.; Marazzato, M.; De Castro, G.; Brindisi, G.; Schiavi, L.; Tosca, M.; Duse, M.; Zicari, A.M. When asthma and rhinitis coexist, could rhinitis reduce asthma control in children? *Allergy Asthma Proc.* 2019, 40, e8–e13, doi:10.2500/aap.2019.40.4219.
 16. Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy.* 2020;75. <https://doi.org/10.1111/all.14238>.
 17. Dong X, Cao YY, Lu XX, Zhang JJ, Du H, Yan YQ, et al. Eleven faces of coronavirus disease 2019. *Allergy.* 2020. doi: 10.1111/all.14289.
 18. Kimura, H.; Francisco, D.; Conway, M.; Martinez, F.; Vercelli, D.; Polverino, F.; Billheimer, D.; Kraft, M. Type 2 inflammation modulates ACE2 and TMPRSS2 in airway epithelial cells. *J. Allergy Clin Immunol.* 2020, 146, 80–88.e8.
 19. Morais-Almeida M, Pité H, Aguiar R, Ansotegui I, Bousquet J. Asthma and the Coronavirus Disease 2019 Pandemic: A Literature Review. *Int Arch Allergy Immunol.* 2020;181(9):680-688. doi: 10.1159/000509057. Epub 2020 Jun 9. PMID: 32516795; PMCID: PMC7316650.
 20. Matsumoto K, Saito H. Does asthma affect morbidity or severity of COVID-19? *J Allergy Clin Immunol.* 2020 Jul;146(1):55-57. doi: 10.1016/j.jaci.2020.05.017. Epub 2020 May 26. PMID: 32470485; PMCID: PMC7250068.
 21. Du, Y.; Tu, L.; Zhu, P.; Mu, M.; Wang, R.; Yang, P.; Wang, X.; Hu, C.; Ping, R.; Hu, P.; et al. Clinical features of 85 fatal cases of COVID-19 from Wuhan. A retrospective observational study. *Am. J. Respir. Crit. Care Med.* 2020, 201, 1372–1379.
 22. Chao, J.Y.; Derespina, K.R.; Herold, B.C.; Goldman, D.L.; Aldrich, M.; Weingarten, J.; Ushay, H.M.; Cabana, M.D.; Medar, S.S. Clinical characteristics and outcomes of hospitalized and critically ill children and adolescents with Coronavirus Disease 2019 at a tertiary care medical center in New York City. *J. Pediatr.* 2020, 223, 14–19.
 23. Papadopoulos, N.; Custovic, A.; Deschildre, A.; Mathioudakis, A.; Phipatanakul, W.; Wong, G.; Xepapadaki, P.; Agache, I.; Bacharier, L.; Bonini, M.; et al. Impact of COVID-19 on pediatric asthma: Practice adjustments and disease burden. *J. Allergy Clin. Immunol. Pract.* 2020, 8, 2592–2599.e3.
 24. Brindisi G, De Vittori V, De Nola R, Pignataro E, Anania C, De Castro G, Cinicola B, Gori A, Cicinelli E, Zicari AM. Updates on Children with Allergic Rhinitis and Asthma during the COVID-19 Outbreak. *J Clin Med.* 2021 May 24;10(11):2278. doi: 10.3390/jcm10112278. PMID: 34073986.
 25. Global Initiative for Asthma. COVID19: GINA Answers to Frequently Asked Questions on Asthma Management. Available online: www.ginasthma.org (accessed on 22 November 2020).
 26. Shaker, M.; Oppenheimer, J.; Grayson, M.; Stukus, D.; Hartog, N.; Hsieh, E.; Rider, N.; Dutmer, C.M.; Leek, T.K.V.; Kim, H.; et al. COVID-19: Pandemic contingency planning for the allergy and immunology clinic. *J. Allergy Clin. Immunol. Pract.* 2020, 8, 1477–1488.e5.
 27. Morais-Almeida M, Aguiar R, Martin B, Ansotegui IJ, Ebisawa M, Arruda LK, Caminati M, Canonica GW, Carr T, Chupp G, Corren J, Dávila I, Park HS, Hanania NA, Rosenwasser L, Sánchez-Borges M, Virchow JC, Yáñez A, Bernstein JA, Caraballo L, Chang YS, Chikhladze M, Fiocchi A, González-Díaz SN, Tanno LK, Levin M, Ortega-Martell JA, Passalacqua G, Peden DB, Rouadi PW, Sublett JL, Wong GWK, Bleecker ER. COVID-19,

- asthma, and biological therapies: What we need to know. *World Allergy Organ J.* 2020 May 16;13(5):100126. doi: 10.1016/j.waojou.2020.100126. PMID: 32426090; PMCID: PMC7229954.
28. Bousquet J, Akdis C, Jutel M, Bachert C, Klimek L, Agache I, Ansotegui IJ, Bedbrook A, Bosnic-Anticevich S, Canonica GW, Chivato T, Cruz AA, Czarlewski W, Del Giacco S, Du H, Fonseca JA, Gao Y, Haahtela T, Hoffmann-Sommergruber K, Ivancevich JC, Khaltaev N, Knol EF, Kuna P, Larenas-Linnemann D, Mullol J, Naclerio R, Ohta K, Okamoto Y, O'Mahony L, Onorato GL, Papadopoulos NG, Pfaar O, Samolinski B, Schwarze J, Toppila-Salmi S, Teresa Ventura M, Valiulis A, Yorgancioglu A, Zuberbier T; ARIA-MASK study group. Intranasal corticosteroids in allergic rhinitis in COVID-19 infected patients: An ARIA-EAACI statement. *Allergy.* 2020 Mar 31. doi: 10.1111/all.14302. Epub ahead of print. PMID: 32233040.
 29. Pfaar O, Klimek L, Jutel M, Akdis CA, Bousquet J, Breiteneder H, Chinthrajah S, Diamant Z, Eiwegger T, Fokkens WJ, Fritsch HW, Nadeau KC, O'Hehir RE, O'Mahony L, Rief W, Sampath V, Schedlowski M, Torres MJ, Traidl-Hoffmann C, Wang Y, Zhang L, Bonini M, Brehler R, Brough HA, Chivato T, Del Giacco SR, Dramburg S, Gawlik R, Gelincik A, Hoffmann-Sommergruber K, Hox V, Knol EF, Lauerma A, Matricardi PM, Mortz CG, Ollert M, Palomares O, Riggioni C, Schwarze J, Skypala I, Untersmayr E, Walusiak-Skorupa J, Ansotegui IJ, Bachert C, Bedbrook A, Bosnic-Anticevich S, Brussino L, Canonica GW, Cardona V, Carreiro-Martins P, Cruz AA, Czarlewski W, Fonseca JA, Gotua M, Haahtela T, Ivancevich JC, Kuna P, Kvedariene V, Larenas-Linnemann DE, Abdul Latiff AH, Mäkelä M, Morais-Almeida M, Mullol J, Naclerio R, Ohta K, Okamoto Y, Onorato GL, Papadopoulos NG, Patella V, Regateiro FS, Samoliński B, Suppli Ulrik C, Toppila-Salmi S, Valiulis A, Ventura MT, Yorgancioglu A, Zuberbier T, Agache I. COVID-19 pandemic: Practical considerations on the organization of an allergy clinic-An EAACI/ARIA Position Paper. *Allergy.* 2021 Mar;76(3):648-676. doi: 10.1111/all.14453. PMID: 32531110; PMCID: PMC7323448.
 30. Klimek L, Pfaar O, Worm M, Bergmann KC, Bieber T, Buhl R, Buters J, Darsow U, Keil T, Kleine-Tebbe J, Lau S, Maurer M, Merk H, Mösges R, Saloga J, Staubach P, Stute P, Rabe K, Rabe U, Vogelmeier C, Biedermann T, Jung K, Schlenter W, Ring J, Chaker A, Wehrmann W, Becker S, Mülleneisen N, Nemat K, Czech W, Wrede H, Brehler R, Fuchs T, Tomazic PV, Aberer W, Fink-Wagner A, Horak F, Wöhrl S, Niederberger-Leppin V, Pali-Schöll I, Pohl W, Roller-Wirnsberger R, Spranger O, Valenta R, Akdis M, Akdis C, Hoffmann-Sommergruber K, Jutel M, Matricardi P, Spertin F, Khaltaev N, Michel JP, Nicod L, Schmid-Grendelmeier P, Hamelmann E, Jakob T, Werfel T, Wagenmann M, Taube C, Gerstlauer M, Vogelberg C, Bousquet J, Zuberbier T. Allergen immunotherapy in the current COVID-19 pandemic: A position paper of AeDA, ARIA, EAACI, DGAKI and GPA: Position paper of the German ARIA GroupA in cooperation with the Austrian ARIA GroupB, the Swiss ARIA GroupC, German Society for Applied Allergology (AEDA)D, German Society for Allergology and Clinical Immunology (DGAKI)E, Society for Pediatric Allergology (GPA)F in cooperation with AG Clinical Immunology, Allergology and Environmental Medicine of the DGHNO-KHCG and the European Academy of Allergy and Clinical Immunology (EAACI)H. *Allergol Select.* 2020 May 28;4:44-52. doi: 10.5414/ALX02147E. PMID: 32568272; PMCID: PMC7304289.

Figures

Figure 1

Prevalence of asthma by geographic area during the first survey. Prevalence of asthmatic patients according to geographic area during the first COVID-19 wave. Data are expressed as percentage. White bar refers to <20% prevalence, green to 20-40% prevalence, and black to >40% prevalence.

Figure 2

Prevalence of asthma by geographic area during the second survey. Prevalence of asthmatic patients according to geographic area during the second COVID-19 wave. Data are expressed as percentage. White bar refers to <20% prevalence, green to 20-40% prevalence, and black to >40% prevalence.

Figure 3

Prevalence of allergic rhinoconjunctivitis by geographic area during the first survey. Prevalence of allergic rhinoconjutivitis according to geographic area during the first COVID-19 wave. Data are expressed as percentage. White bar refers to <20% prevalence, green to 20-40% prevalence, and black to >40% prevalence.

Figure 4

Prevalence of allergic rhinoconjunctivitis by geographic area during the second survey. Prevalence of allergic rhinoconjunctivitis according to geographic area during the second COVID-19 wave. Data are expressed as percentage. White bar refers to <20% prevalence, green to 20-40% prevalence, and black to >40% prevalence.