Linking Unfolded Protein Response to Ovarian Cancer Cell Fusion

Lucile Yart

University of Geneva

Daniel Bastida-Ruiz

University of Geneva

Mathilde Allard

University of Geneva

Pierre-Yves Dietrich

University of Geneva

Patrick Petignat

University of Geneva

Marie Cohen (✉ marie.cohen@unige.ch)

University of Geneva

Research Article

Keywords: unfolded protein response, ovarian cancer, polyploid giant cancer cell, cell fusion, invasion

Posted Date: September 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-757674/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Polyploid giant cancer cells (PGCCs) have been observed in epithelial ovarian tumors. They can resist antimitotic drugs, thus participating in tumor maintenance and recurrence. Despite their origin remains unclear, PGCC formation seems to be enhanced by conditions that trigger the unfolded protein response (UPR) such as hypoxia or chemotherapeutic treatments. Hypoxia has been shown to promote the formation of ovarian PGCCs by cell fusion. These PGCCs can undergo bursting and budding, contributing to generate cancer stem-like cells with more aggressive phenotypes than the parental cells.

Methods: The UPR was induced in two ovarian cancer cell lines (SKOV3 and COV318). UPR activation was assessed by western blot and polyploidy indexes were calculated. Then, to confirm the implication of cell fusion in PGCCs formation, two populations of SKOV3 cells were transfected with plasmids encoding for two distinct nuclear fluorescent proteins (GFP and mCherry) associated with different antibiotic resistance genes, and the two cell populations were mixed in co-culture. The co-culture was submitted to a double-antibiotic selection. The resulting cell population was characterized for its morphology, cyclicity, and proliferative and tumorigenic capacities, in addition to a transcriptomic characterization.

Results: We demonstrated that cell fusion could be involved in ovarian PGCCs generation and this process was promoted by unfolded protein response activation. Double-antibiotic treatment of PGCCs led to the selection of a pure population of cells containing both GFP- and mCherry-positive nuclei. Interestingly, after three weeks of selection, we observed that these cells were no longer polynucleated but displayed single nucleus positive for both fluorescent proteins, suggesting that genetic material mixing had occurred. These cells had restarted normal cell cycles, acquired an increased invasive capacity, and can form ovarian tumors in ovo.

Conclusions: UPR activation increased the in vitro formation of PGCCs by cell fusion, further leading to the acquisition of new properties for the newly generated cells. UPR modulation in ovarian cancer patients could be an interesting therapeutic strategy to avoid the formation of PGCCs and therefore limit cancer relapse and drug resistance.

Background

Epithelial ovarian cancer (EOC) is the most common type of ovarian cancer, representing 9 out of 10 tumors (1). It is the seventh most common cancer in women and the leading cause of gynecologic cancer-related deaths in Western countries (2). EOC is usually managed by cytoreductive surgery followed by platinum-based chemotherapy with an expected response rate to primary therapy of more than 70% (3). Nevertheless, 2 years after a complete remission, most of the patients had a cancer recurrence (4) with often platinum-based chemotherapy treatment resistance (3).

EOC relapses origin is not completely understood, however, several hypotheses have been proposed over time. One of them, the cancer stem cell (CSC) theory, is based on the capacity of these cells to generate a proliferative progeny, initiating the growth of a heterogeneous tumor (5). CSCs present chemotherapy

resistance, allowing their survival to the treatment with subsequent proliferation and cancer relapse (6). The role of CSCs in ovarian cancer was reviewed by Giornelli et al., highlighting the importance of these cells in cancer recurrence and progression (7). Despite the origin of CSCs remains elusive, some theories move towards a possible genesis in cancer cell fusion events [reviewed by Lu and Kang (8)]. The fusion of cancer cells could give rise to polyploid giant cancer cells (PGCCs) (9), large cells containing several copies of DNA and positive for cancer stem cell markers (10). Interestingly, the presence of PGCCs in ovarian cancer has been reported preferentially at late disease stages and in high pathological grades (10). We previously showed that primary EOC cells isolated from malignant ascites were able to spontaneously form PGCCs in culture (11). However, the origin of PGCCs is controversial since both fusion or endoreplication processes could give rise to them (8,9,12). Zhang et al. demonstrated that in SKOV3, an EOC cell line established from ascites, the formation of PGCCs was at least partially due to cell fusion events (9). In the study conducted by Zhang et al., the PGCCs derived from EOC cells had striking properties such as different cell cycle regulation and division profile, changing their daughter generation mechanism from mitosis to budding, which confers these cells the resistance to cytotoxic chemotherapy treatments (9). Additionally, it was found that hypoxia increased PGCCs formation (9,13–15), but other conditions such as chemotherapy could also favor their production (13).

The tumor environment is known to be hypoxic and hypoglycemic due to the low vascularization caused by the fast and erroneous angiogenesis that tumor mass undergoes [reviewed by Muz et al. (16)]. These conditions are capable of triggering the endoplasmic reticulum (ER) stress response or unfolded protein response (UPR) (17) which is essential for cellular adaptation to stressful situations (18). In homeostatic conditions, the three UPR-associated proteins, protein kinase RNA-like endoplasmic reticulum kinase (PERK), inositol-requiring enzyme 1 α (IRE1 α), and activating transcription factor 6 (ATF6), remain inactive because of their binding with glucose-regulated protein 78 (GRP78) [reviewed by Bastida-Ruiz et al. (19)]. Under ER stress conditions, unfolded proteins accumulate in the ER lumen, promoting the recruitment of chaperones to enable their correct folding. Among these chaperones, GRP78 leaves the complexes formed with the UPR-associated proteins and induces the UPR activation (20). This gives rise to a general decrease of protein, by contrast with increased expression of chaperones and ER-associated protein degradation-related proteins (ERAD) and autophagy activation [reviewed by Bastida-Ruiz et al. (19)]. The overexpression of GRP78 serves as a UPR activation marker (21). GRP78 was shown to be overexpressed in ovarian cancer tissues and correlated with poor patient survival (22), suggesting a role of UPR in EOC (23). Interestingly, GRP78 and UPR were involved in trophoblastic cell fusion (24,25). We thus hypothesized that UPR activation could be involved as well in EOC cell fusion, hence the occurrence of PGCCs and aggressiveness of EOC.

In this article, we confirmed Zhang et al. results stating that PGCCs derive at least partially from cell fusion in SKOV3 cells (9) and demonstrated that UPR induces ovarian cancer cell fusion and PGCCs formation. Moreover, we observed that polyploid cells were no longer polynucleated after 3 weeks of antibiotic selection pressure and were able to proliferate in the same manner as their parental cell lines despite higher DNA content. We then evaluated the advantages that cell fusion could confer to these cells.

Methods

Reagents

Chemical hypoxia was induced with Cobalt Chloride (CoCl_2 , Merck, Darmstadt, Germany).

ER stress was induced with HA15 (Selleckchem, Zurich, Switzerland), Thapsigargin (THA, Enzo LifeSciences, Lausen, Switzerland), or Tunicamycin (TUN, Enzo LifeSciences, Lausen, Switzerland).

ER stress was inhibited by 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride (AEBSF; Sigma, Darmstadt, Germany), Melatonin (Mel; Sigma, Darmstadt, Germany), STF-083010 (STF; Selleckchem, Zurich, Switzerland), 4 μ 8C (Selleckchem, Zurich, Switzerland), GSK2656157 (GSK; Selleckchem, Zurich, Switzerland) or Salubrinal (SAL; Tocris Biosciences, Bristol, UK).

Cancer cell line and culture

The human ovarian cancer cell lines COV318 (ECACC, Sigma-Aldrich) and SKOV3 (courtesy of Dr. Florence Delie, University of Geneva, Switzerland) were cultured at 37°C and 5% CO₂ in DMEM or RPMI (Gibco, Invitrogen, Basel, Switzerland) respectively, supplemented with 10% fetal bovine serum (FBS, Biochrom AG, Oxoid AG, Basel, Switzerland) and 0.05 mg.mL⁻¹ gentamycin (Invitrogen, Basel, Switzerland).

Cell treatments

ER stress was induced in ovarian cancer cells by adding either 100 nM THA, 1 μ g.mL⁻¹ TUN or 1 μ M HA15 in culture medium. Cells were treated for 48h before processing for analysis.

UPR pathways were inhibited in ovarian cancer cells by adding either 17.5 μ M SAL, 0.3 μ M GSK, 8 μ M 4 μ 8C, 2 μ M STF, 200 μ M AEBSF or 1 mM Mel, in combination with 100 nM THA or 1 μ g.mL⁻¹ TUN. Cells were treated for 48h before processing for analysis.

Western blot

Western blot analyses were performed as described by Bastida-Ruiz et al. (24). Briefly, whole-cell extracts were fractionated by SDS-Page 10% and transferred to nitrocellulose membrane for immunoblot analysis. The membranes were first incubated with the primary antibodies rabbit anti-GRP78 (GL-19, 1:5000 dilution from Sigma) and mouse anti-GAPDH antibodies (1:60 000 dilution from Millipore), followed by incubation with the secondary antibodies goat anti-rabbit IgG horseradish peroxidase (HRP)-conjugated (170-6515, 1:3 000 dilution, from Bio Rad) and goat anti-mouse IgG HRP-conjugated (sc-2005, 1:3 000 dilution, Santa Cruz Biotechnologies).

Generation of PGCCs by cell fusion

Cytoplasmic staining and FACS sorting: SKOV3 or COV318 cells were stained with either green or far-red cytoplasmic dyes, CellTrace CFSE (green; Life Technologies, Carlsbad, USA), and CellTrace Far-Red (far-red, Life Technologies, Carlsbad, USA), according to the manufacturer's instructions. For each cell line, 3.5×10^5 green cells and 3.5×10^5 far-red cells were co-seeded immediately after staining in a T175 flask. Twenty-four hours after seeding, cells were treated with 300 μM CoCl_2 (Merck, Darmstadt, Germany) for 48h. Cells were then analyzed by flow cytometry with FACS Aria II (Becton Dickinson, Franklin Lakes, USA) for -GFP and far-red signals. Double-positive cells were sorted and seeded in a regular culture medium. Characteristic pictures of the sorted cells were done 16h after seeding, using a fluorescence microscope (EVOS FL, Life Technologies, Carlsbad, USA).

Transfection and selection: SKOV3 cells were transfected with a pH2B-mCherry-IRES-puro2 plasmid or with pEGFP-CenpA-IRES-neo (Addgene, LGC Standards, Teddington, UK) using JetPEI as transfection reagent (Polyplus transfection, Illkirch, France) according to the manufacturer's instructions. Two days after transfection, cells were selected using selection antibiotics (1 $\mu\text{g.mL}^{-1}$ puromycin, InvivoGen, San Diego, USA for cells transfected with pH2B-mCherry-IRES-puro2 plasmid = SKOV3-Red, and 500 $\mu\text{g.mL}^{-1}$ G418, Roche Life Science, Penzberg, Germany for cells transfected with pEGFP-CenpA-IRES-neo = SKOV3-Green) for two weeks. Cells were then cultured in RPMI medium supplemented with 0.1 $\mu\text{g.mL}^{-1}$ puromycin for SKOV3-Red cells or with 50 $\mu\text{g.mL}^{-1}$ G418 for SKOV3-Green cells.

SKOV3-M cell line establishment: SKOV3-Red and SKOV3-Green were mixed (1:1) and cultured for 72h before adding 0.2 $\mu\text{g.mL}^{-1}$ puromycin and 100 $\mu\text{g.mL}^{-1}$ G418 in RPMI. The culture medium was renewed every two days.

Polyploidy index / Fusion index

Non-transfected cells: SKOV3 and COV318 cells were fixed in paraformaldehyde (PFA) 4% at 4°C for 10 min, washed three times in PBS, and stained with Hematoxylin (Sigma, Darmstadt, Germany) for 1 min. Cells were then washed twice with warmed tap water before bright-field imaging (EVOS, Life Technologies, Carlsbad, USA).

Transfected cells: Cells were seeded either in Nunc Lab-Tek II Chamber slides (Life Technologies, Carlsbad, USA) for confocal microscopy or in μ -Slides 8 Well (ibidi, Madison, USA) for fluorescent microscopy, and processed for treatment. Cells were then fixed in PFA 4% at 4°C for 10 min, washed three times in PBS, and incubated in PBS-3% bovine serum albumin (BSA) for 30 min to eliminate non-specific binding. Cells were then stained with Alexa Fluor 647 phalloidin (1:40 dilution, from Invitrogen, Thermo Scientific) for 20 min at room temperature and rinsed three times in PBS. Nunc Lab-Tek II Chamber slides were then mounted with Vectashield with DAPI (Vector Laboratories, Burlingame, USA) and sealed before imaging with a confocal fluorescent microscope (LSM 800 Airyscan, Zeiss, Jena, Germany). μ -Slides 8

We were imaging with a fluorescent microscope (EVOS FL, Life Technologies, Carlsbad, USA). Images were processed using ImageJ freeware.

For both transfected and non-transfected cells, the polyploidy or fusion index expressed in percent was calculated as follows: $[(N-S)/T] \times 100$, where N equals the number of nuclei in syncytia, S equals the number of syncytia and T equals the total number of nuclei counted (26). This index was calculated for three independent experiments, run in triplicate.

Quantification of GFP-positive, mCherry-positive, and double-positive cells

SKOV3-Green, SKOV3-Red, SKOV3-M, and untransfected cells (ct-SKOV3) were resuspended in PBS. Percentages of negative cells, GFP-positive cells, mCherry-positive cells, and GFP and mCherry double-positive cells were measured by flow cytometry analysis (LSR Fortessa, Becton Dickinson, Franklin Lakes, USA). Data were analyzed with FlowJo software (FlowJo LLC, Ashland, USA). This experiment was repeated 3 times.

Cell cycle analysis

SKOV3-Green, SKOV3-Red, and SKOV3-M were resuspended in PBS with $5 \mu\text{g.mL}^{-1}$ Hoechst 33342 (Life Technologies, Carlsbad, USA), and incubated for 20 min at 37°C . Amounts of cells in G0-G1 and G2 phases were then assessed by flow cytometry analysis (LSR Fortessa, Becton Dickinson, Franklin Lakes, USA). Data were analyzed with FlowJo software (FlowJo LLC, Ashland, USA). This experiment was repeated 2 times.

Cell proliferation

On day 0, SKOV3-Green, SKOV3-Red, and SKOV3-M were resuspended in Hanks' Balanced Salt Solution (Life Technologies, Carlsbad, USA) and stained with CellTrace Far Red (Life Technologies, Carlsbad, USA) according to the manufacturer's instructions. The cells were immediately processed for far-red signal analysis by flow cytometry (Gallios, Beckman Coulter, Brea, USA) (D0). Far-red signal dilution was then daily measured for 4 days (D1, D2, D3, and D4). Data were analyzed with FlowJo software (FlowJo LLC, Ashland, USA). This experiment was repeated 3 times.

Cell viability (MTT assay)

SKOV3-Red, SKOV3-Green, and SKOV3-M cells were seeded in 96-well plates ($15'000$ cells/well) and incubated for 48h. The culture medium was then replaced by $100 \mu\text{L}$ of MTT reagent (1mg.mL^{-1}) for 2h. Acidic isopropanol solution ($150 \mu\text{L}$) was added, and then each well was vigorously mixed to dissolve the

precipitated formazan. UV–visible absorption was measured at 540 and 690 nm (as blank). These experiments were carried out in triplicate, three times.

Invasion assay

Cell invasion assay was performed in an invasion chamber as described elsewhere (27). Briefly, 1.5×10^4 cells (SKOV3-Red, SKOV3-Green, and SKOV3-M) in 100 μ L of RPMI supplemented with 10% FBS and 1% gentamicin were added to the upper compartment of the transwell chambers. RPMI supplemented with 20% FBS and 1% gentamicin (400 μ L) was added in the lower chamber for 48 h at 37 °C in a CO₂ (5%) incubator. After incubation, viable cells that invaded collagen were stained with crystal violet, and measurement was performed at 540 nm. This assay was repeated three times and each experiment was run in triplicate. Data were normalized by proliferation values (MTT assay) and expressed as AU (invasion)/ AU (MTT).

Tumor development on chick chorioallantoic membrane (CAM)

Tumor development on CAM was performed as previously described (28). Briefly, fertilized eggs (animal facility of the University of Geneva, Geneva, Switzerland) were incubated at 38°C with 80% relative humidity and periodic rotation. On egg development day (EDD) 4, the eggs were drilled at their narrow apex and the hole was closed with adhesive tape. The eggs were then incubated at 38°C with 80% relative humidity without rotation. On EDD8 the hole in the eggshell was enlarged to allow access to the CAM. After gently scratching of the membrane with a needle tip, SKOV3-Red, SKOV3-Green or SKOV3-M cells suspension (2×10^6 cells in 30 μ L of geltrex, Thermo Scientific) were inoculated and the hole was hermetically covered with Parafilm®. Eggs were returned to the incubator to allow tumor growth. Tumor growth was then monitored at EDD10.5 and 13.5 using a Wild Heerbrugg M3Z microscope at 10x magnification with a Lumenera INFINITY2-1 CDD camera with Infinity Capture Software.

Microarray

Microarray-based transcriptome profiling was performed at the iGE3 Genomics core-facility of the faculty of medicine of the University of Geneva. Briefly, 100 ng of total RNA were extracted from SKOV3-Red, SKOV3-Green, and SKOV3-M at two different passages and were used as input for the preparation of single-strand cDNA using the WT PLUS reagent kit from ThermoFisher Scientific. Targets were then fragmented and labeled with the Affymetrix GeneChip WT Terminal Labeling Kit and hybridized on Human Clariom S arrays according to the manufacturer's recommendations.

The data were analyzed with the Transcription Analysis Console (TAC) software (ThermoFisher Scientific), selecting the genes which expression was modified at least 2-fold and which p-value was

<0.001.

Statistical analysis and reproducibility

Data were represented as means $\pm$ SEM. Statistical differences between samples were assessed by the Student's t-test. P-value <0.05 was considered significant. GraphPrism software was used to perform the different statistical analyses. The number of repeated experiments was described in each methods description.

Results

1. ER stress enhances PGCCs formation

Several studies showed that PGCCs formation in ovarian cancer cells is enhanced by hypoxic conditions, achieved either with CoCl_2 treatment or with 0.1% oxygen cell culture (3, 13–15). Hypoxic conditions are characteristics of the tumor environment and are known to activate UPR (29). We previously described that ovarian cancer cells purified from malignant ascites were spontaneously able to form PGCCs in vitro (11), under regular culture conditions and that the UPR is involved in trophoblastic and choriocarcinoma cell fusion (30). Therefore, we hypothesized that UPR could be involved in ovarian PGCCs formation. To test this hypothesis, SKOV3 and COV318 cells were treated for 48 hours with three different ER stress inducers: THA, TUN, and HA15. Treatment with each of these drugs resulted in an increased protein level of GRP78, in both COV318 and SKOV3 cell lines (Fig. 1a, b), testifying of UPR activation. UPR activation was associated with a significant increase (about 2-fold increase) in PGCCs proportion, in all of the tested conditions (Fig. 1c, d).

To confirm the potent role of UPR in PGCCs formation, we then used molecules known to inhibit the signaling from downstream effectors of IRE1, PERK, and ATF6. Since no specific inhibitor of the ATF6 pathway was available, we used AEBSF (which blocks S1P and S2P in the Golgi apparatus, preventing ATF6 cleavage (31)) and Mel (32) to block the ATF6 pathway. STF and 4 μ 8C were used to inhibit the IRE1 α Rnase activity (33, 34). Finally, GSK (an ATP competitive inhibitor (35)), and SAL (an inhibitor of EIF2 α dephosphorylation (36)) were used to modulate PERK pathway activation. Simultaneous treatment of SKOV3 cells with THA and each of the UPR inhibitors individually resulted in a general tendency of lowered fusion rates when compared to cells treated with THA alone (Fig. 1d). However, the only significant reduction was achieved when treating the cells with the ATF6 pathway inhibitors. In contrast, treatment of SKOV3 cells simultaneously with TUN and each of the UPR inhibitors individually showed a more stringent reduction in cell fusion rate, significant for all of the inhibitors, except GSK and STF (Fig. 1g). In the same way, COV318 treatment with THA or TUN and the different UPR inhibitors resulted in a significant cell fusion rate decrease, except for the combination of TUN + AEBSF (Fig. 1f, h). Together, these results confirm the implication of the UPR in the induction of ovarian cancer cell fusion.

2. Cell fusion is involved in PGCCs formation.

To determine whether PGCCs originate either from endoreplication and incomplete cell division or from cell fusion, we first used the strategy described by Uygur et al. (37) for their study of cell fusion between prostate cancer and muscle cells. Ovarian cancer cell lines, SKOV3 and COV318, were stained with two different fluorescent cytoplasmic markers. The two stained cell populations were then mixed and grown in co-culture. Cells positive for both cytoplasmic markers, which we initially considered as fused cells, were sorted by flow cytometry and seeded in culture dishes. Unexpectedly, the outcome of the cell sorting was not formed mostly by fused cells that mixed their cytoplasm; instead, it was constituted by a mix of polynucleated cells and mononucleated cells that may have exchanged vesicles with cells stained with the opposite cytoplasmic marker (Fig. 2a). This result is consistent with the one published by Miroshnychenko et al. (38) and evidenced that this approach is not valid to isolate ovarian fused cells since vesicle exchange happened.

We then decided to transfect SKOV3 cells with plasmids encoding for a nuclear protein tagged with red or green fluorescent proteins, combined with two distinct antibiotic resistance genes. Following the antibiotic selection of transfected cells, we obtained homogenously nuclei-marked cell lines that we named SKOV3-Red and SKOV3-Green cell lines. After mixing and growing the two populations of cells, we observed spontaneous formation of three types of PGCCs: PGCCs with only red nuclei, PGCCs with only green nuclei, and PGCCs with both red and green nuclei (Fig. 2b). Despite the two first types of PGCCs (with single-color nuclei), do not permit to determine whether these cells originate from endoreplication or from cell fusion, the third type of PGCCs that contain both red and green nuclei demonstrate that the cell fusion process is involved in the generation of these PGCCs.

To determine if the UPR-mediated PGCC formation involves a cell fusion process, we induced ER stress in a co-culture of SKOV3-Red and SKOV3-Green cells with HA15, and then counted the number of PGCCs with only red nuclei, only green nuclei, and both red and green nuclei, using confocal microscopy. Proportion of PGCCs was increased more than 1.3 folds in single-colored PGCCs (Red PGCCs: $1.58 \% \pm 0.52$ (SD) in CTL vs $2.11 \% \pm 0.89$ (SD) in HA15; Green PGCCs: $1.61 \% \pm 0.52$ (SD) in CTL vs $2.23 \% \pm 0.68$ (SD) in HA15), and more than 2.1 folds in PGCCs with both red and green nuclei (Red + Green PGCCs: $0.74 \% \pm 0.34$ (SD) in CTL vs $1.57 \% \pm 0.71$ (SD) in HA15). Calculation of fusion index, which considers the number of nuclei contained within the syncytia, showed that HA15-induced ER stress enhanced formation of each of the three types of PGCCs, including Red + Green PGCCs, formed by cell fusion (Figs. 2c, d). These results confirm that cell fusion is involved in the formation of PGCCs and that UPR could increase this process in ovarian cancer cells.

3. PGCCs can revert as mononucleated cells under selection pressure, and restart normal cell cycles

To investigate the fate of ovarian PGCCs formed by cell fusion, we selected red and green fluorescent SKOV3 cells derived from co-culture of SKOV3-Red with SKOV3-Green under double antibiotic selection, which we named SKOV3-M. Indeed, fused cells containing both types of nuclei would be able to survive a double antibiotic selection, while the unfused parental cell lines and the same type of nuclei-fused cells would die under this condition. Strikingly, after three weeks under double antibiotic selection, we observed that SKOV3-M cells were no longer polynucleated and that their single nucleus was positive for both fluorescent proteins (Fig. 3a), suggesting that mixing of genetic material had occurred. Flow cytometry analysis confirmed that about 98% of SKOV3-M cells were positive for both red and green fluorescent proteins, and this phenotype was maintained over passage 72 (Fig. 3b).

Since SKOV3-M appeared to have restarted a normal cell growth, we determined if they were able to achieve normal cell cycles. We observed a similar G0-G1 and G2 phase cell distribution between SKOV3-M and its parental cell lines, with about 66 % to 70 % of cells in the G0-G1 phase, and about 18 % to 22% of cells in the G2 phase (Fig. 3c, d). Interestingly, the Hoechst staining was more intense in SKOV3-M cells (Fig. 3c), suggesting a higher DNA content in the fused cells than in the parental ones. Additionally, we decided to study the proliferative capacities of SKOV3-M cells by staining the cells with CellTrace™ far-red. The division profile of stained cells was analyzed during five consecutive days. SKOV3-M and its parental cell lines had proliferated and diluted CellTrace™ far-red fluorescence in the same way (Fig. 3e, f).

We then compared the tumor growth of SKOV3-M and its parental cells-derived tumors using the chick chorioallantoic membrane (CAM) model. The tumor size (Fig. 3g) and tumor growth (Fig. 3h) were not significantly different between SKOV3-M and the parental cell lines.

4. Comparison of SKOV3-M omics profile with its parental cell lines

Higher DNA content found in SKOV3-M cells may be associated with an important chromosomal rearrangement, which is known to alter the regulation of gene expression(39) and exacerbate tumor malignancy(40). We thus decided to compare the transcriptome of SKOV3-M to the parental cell lines. Comparative transcriptomic analysis of SKOV3-M to the parental cell lines identified 82 genes that were differentially expressed ($\pm$ 2-fold, $p \leq 0.001$; Fig. 4a; see Supplementary Table 1 for the 10 most significantly upregulated and downregulated genes). The alteration of some proteins expression in SKOV3-M compared to the parental cell lines was confirmed by secretomic analysis (Supplementary Fig. 1, Supplementary Table 2, Supplementary Table 3). Among differentially expressed genes, enrichment of biological processes identified cell adhesion, locomotion, and invasion to be among the statistically altered processes (Table 1). Therefore, we hypothesized that ovarian cancer cell fusion confers enhanced invasive capacities to SKOV3-M cells. To test this hypothesis, the three cell lines SKOV3-Green, SKOV3-Red and SKOV3-M were seeded in Boyden chamber. The capacity of SKOV3-M to invade collagen was increased compared to its parental cell lines (Fig. 4b), supporting that cell fusion can lead to the modification of ovarian cancer cell properties.

Table 1
Cell migration, cell motility, invasion, cell adhesion and locomotion-related genes which expression is at least 2-fold modified and which p-value is < 0.05 in SKOV3-M cells compared to SKOV3-Green and SKOV3-Red cells.

Gene symbol	Fold change	P-value	Function
IL1R2	11.03	5.98E-06	Cell migration
MMP2	8.13	0.002	Invasion
MXRA5	7.09	2.79E-05	Invasion
VEGFC	3.57	0.0108	Invasion
AFAP1L2	3.1	0.0012	Cell motility
EFNB2	2.71	2.95E-05	Invasion
LRRC17	2.5	0.0032	Locomotion
ADGRE5	2.49	1.96E-05	Cell adhesion
ELFN2	2.34	0.0002	Cell migration
FBLN5	-2.02	0.0043	Cell adhesion
BCAM	-2.15	0.0012	Cell adhesion
CDHR1	-2.54	6.17E-05	Cell adhesion
ITGB4	-2.56	0.0197	Cell adhesion
IL7R	-3	0.0064	Invasion
ADAMTSL4	-3.33	0.0042	Cell adhesion
EPCAM	-3.99	1.31E-06	Cell adhesion
IGFBP3	-5.18	0.0007	Cell migration
S1PR1	-6.54	1.65E-07	Cell migration

Discussion

The PGCCs formation has been studied in several cancer types, among which colon, breast, and ovarian cancer (9, 13–15). This process is induced by DNA damaging agents and could be linked to chemotherapeutic resistance acquisition (9, 13, 14, 41). It could be also incremented by chemical hypoxia treatment (CoCl₂) or hypoxic culture conditions in vitro (9, 13–15). The mechanism of PGCCs formation has always been controversial since it could be explained either by a defect in cytokinesis leading to endoreplication, by cell fusion events, or by a combination of both mechanisms (8, 9, 12). Under chemical

hypoxia, it was shown that at least 10 to 20% of ovarian PGCCs were formed by cell fusion (9). The characterization of the PGCCs formed under these conditions displayed a modification in their division process, changing from mitosis to bursting and budding (9). The PGCCs generated daughter cells via asymmetric division, which were able to form spheroids and expressed several markers of cancer stem cell (9). Moreover, they were more tumorigenic than regular-sized cancer cells in the nude mice (9). In the present study, using plasmids expressing nuclear proteins tagged with green or red fluorochrome, we generated PGCCs from SKOV3 cells and confirmed that these PGCCs were at least partially derived from cell fusion events. In addition, we described for the first time, that under double antibiotic selection pressure, mononucleated cells containing both red- and green-fluorescent nuclear proteins, can be selected, suggesting that mixing of genetic material had occurred. Despite higher DNA content, SKOV3-M cells recovered a normal cell cycle and were able to proliferate in the same manner as their parental cell lines (SKOV3-Green and SKOV3-Red). This novel observation leads us to hypothesize that genetic material mixing may be a mechanism by which PGCCs, after surviving drug treatment, could change their cell cycle back to mitosis. Transcriptomic analysis of the SKOV3-M cells showed that more than 80 genes were at least 2-fold modulated in comparison with their parental cells. Among them, we found genes implicated in cell adhesion, motility, and invasion. The validation of these results with invasion assays demonstrated increased invasive properties of SKOV3-M cells compared to their parental cells. This increased invasiveness renders the cells more aggressive, favoring the onset of metastases (42–44). Interestingly, our results are supported by a recent study, combining *in vitro*, *in vivo*, and *in silico* experimental models, in which the authors suggest that the genetic recombination, subsequent to cancer cell fusion they observed *in vitro* and *in vivo*, can facilitate cancer cells adaptation and therefore tumor progression (38).

The proliferation rate of these fusion-derived cells, their capacities to resist both antibiotics, to form tumors, which we measured *in ovo* on CAM model and their invasive properties may explain the high cancer relapse rates that ovarian cancer patients suffer. However, it remains the question of whether ovarian cancer cell fusion occurs *in vivo*.

Despite PGCCs could generate more aggressive daughter cells via bursting, budding (9, 45, 46), or mitosis (47), the mechanism and conditions initiating their formation have been poorly studied. The increment in ovarian PGCCs formation achieved under hypoxia *in vitro* pointed towards hypoxia as a possible cell fusion enhancer (9, 13–15). Paclitaxel treatment, an antimetabolic drug, was also shown to cause an induced prominence of PGCCs [(13) and data not shown]. A shared pathway downstream of low oxygen tension and Paclitaxel treatment is the UPR pathway (17, 19, 48–54). We thus hypothesized that UPR activation may result in cancer cell fusion enhancement and PGCCs formation. The activation of the UPR with different UPR activators effectively increased cell fusion and PGCCs formation with both SKOV3 and COV318 cell lines. On the opposite, the inhibition of the UPR pathways by inhibitors targeting the three UPR-branches led to a general tendency of a cell fusion decrease as already observed in choriocarcinoma cell line (24, 25). Therefore, we can conclude that the UPR, through the activation of the UPR sensors ATF6, IRE1 α , and PERK, is involved in ovarian PGCCs formation. However, the contribution of each UPR pathway and their interdependence remain to be elucidated.

Conclusions

More and more pieces of evidence showed that the UPR is involved in drug resistance in different cancer cells (50–52, 54). However, the implication of the UPR in chemotherapy resistance is complex and not fully understood. In this study, we showed that decreasing UPR activation in ovarian cancer cells could decrease ovarian cancer cell fusion and PGCCs formation. Avoiding the PGCCs formation may reduce cancer relapses and metastasis, while at the same time, the drug resistances generated by the UPR could be prevented (9, 50, 51, 53). A better understanding of the molecular mechanism controlling UPR-mediated PGCCs formation and drug resistance is highly needed to develop new combined therapeutic approaches with new drugs that specifically target the UPR sensors or downstream partners and conventional chemotherapy.

Abbreviations

AEBSF, 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride; ATF6, activating transcription factor 6; CAM, chick chorioallantoic membrane; CoCl₂, cobalt chloride; CSC, cancer stem cell; EDD, egg development day; EOC, epithelial ovarian cancer; ER, endoplasmic reticulum; GRP78, glucose-regulated protein 78; GSK, GSK2656157; IRE1α, inositol-requiring enzyme 1α; Mel, Melatonin; PERK, protein kinase RNA-like endoplasmic reticulum kinase; PGCC, polyploid giant cancer cell; SAL, salubrinal; STF, STF-083010; TAC, transcription analysis console; THA, thapsigargin; TUN, tunicamycin, UPR, unfolded protein response.

Declarations

Ethics approval and consent to participate

Not applicable.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interest

The authors declare no competing interests.

Funding:

This research was funded by Swiss National Science Foundation, grant number 31003A-163395, Schmidheiny foundation, fondation pour la lutte contre le cancer et pour des recherches biomédicales and fondation pour la Lutte contre le Cancer.

Author's contributions

L.Y. ,D.B.R. and M.C. performed the experiments and wrote the manuscript. M.A. designed and analyzed flow cytometry experiments. L.Y. and M.C. designed the experiments. M.C. conceived the research project. P.Y.D. and P.P. reviewed the manuscript. All authors gave input on writing the manuscript.

Acknowledgments:

We would like to thank the staff of the iGE3 Genomics and Flow Cytometry core facilities, Emily Jordan, Pascale Ribaux, Christine Wullemin, Dr. Maxim Pilyugin, Daniela Harry, and Pr. Patrick Meraldi for their support.

References

1. Ledermann JA, Raja FA, Fotopoulou C, Gonzalez-Martin A, Colombo N, Sessa C, et al. Newly diagnosed and relapsed epithelial ovarian carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2013;24 Suppl 6:vi24-32.
2. Stewart B, Wilde C. World Cancer Report 2014. International Agency for Research on Cancer. Lyon, France; 2014. (1st edition).
3. Cohen M, Petignat P. The bright side of ascites in ovarian cancer. *Cell Cycle.* 2014;13(15):2319.
4. Giornelli GH. Management of relapsed ovarian cancer: a review. *Springerplus.* 2016;5(1):1197.
5. Iqbal W, Alkarim S, AlHejin A, Mukhtar H, Saini KS. Targeting signal transduction pathways of cancer stem cells for therapeutic opportunities of metastasis. *Oncotarget.* 2016;7(46):76337–53.
6. Prieto-Vila M, Takahashi R-U, Usuba W, Kohama I, Ochiya T. Drug Resistance Driven by Cancer Stem Cells and Their Niche. *Int J Mol Sci.* 2017;18(12):E2574.
7. Giornelli GH, Mandó P. A theoretical view of ovarian cancer relapse. *European Medical Journal* 2017; 2:128–135.
8. Lu X, Kang Y. Cell Fusion as a Hidden Force in Tumor Progression. *Cancer Res.* 2009;69(22):8536–9.
9. Zhang S, Mercado-Urbe I, Xing Z, Sun B, Kuang J, Liu J. Generation of Cancer Stem-like Cells through Formation of Polyploid Giant Cancer Cells. *Oncogene.* 2014;33(1):10.1038/onc.2013.96.
10. Zhang D, Wang Y, Zhang S. Asymmetric Cell Division in Polyploid Giant Cancer Cells and Low Eukaryotic Cells. *BioMed Research International.* 2014;2014:e432652.
11. Bastida-Ruiz D, Van Hoesen K, Cohen M. The Dark Side of Cell Fusion. *Int J Mol Sci.* 2016;17(5).

12. Shu Z, Row S, Deng W-M. Endoreplication: The Good, the Bad, and the Ugly. *Trends Cell Biol.* 2018;28(6):465–74.
13. Niu N, Mercado-Urbe I, Liu J. Dedifferentiation into blastomere-like cancer stem cells via formation of polyploid giant cancer cells. *Oncogene.* 2017;36(34):4887–900.
14. Lopez-Sánchez LM, Jimenez C, Valverde A, Hernandez V, Peñarando J, Martinez A, et al. CoCl₂, a mimic of hypoxia, induces formation of polyploid giant cells with stem characteristics in colon cancer. *PLoS One.* 2014;9(6):e99143.
15. Fei F, Zhang D, Yang Z, Wang S, Wang X, Wu Z, et al. The number of polyploid giant cancer cells and epithelial-mesenchymal transition-related proteins are associated with invasion and metastasis in human breast cancer. *Journal of Experimental & Clinical Cancer Research.* 2015;34(1):158.
16. Muz B, de la Puente P, Azab F, Azab AK. The role of hypoxia in cancer progression, angiogenesis, metastasis, and resistance to therapy. *Hypoxia (Auckl).* 2015;3:83–92.
17. Wouters BG, Koritzinsky M. Hypoxia signalling through mTOR and the unfolded protein response in cancer. *Nat Rev Cancer.* 2008;8(11):851–64.
18. Hetz C, Papa FR. The Unfolded Protein Response and Cell Fate Control. *Mol Cell.* 2018;69(2):169–81.
19. Bastida-Ruiz D, Aguilar E, Ditisheim A, Yart L, Cohen M. Endoplasmic reticulum stress responses in placentation - A true balancing act. *Placenta.* 2017;57:163–9.
20. Lee AS. The ER chaperone and signaling regulator GRP78/BiP as a monitor of endoplasmic reticulum stress. *Methods (San Diego, Calif).* 2005;35(4):373–81.
21. Kennedy D, Samali A, Jäger R. Methods for studying ER stress and UPR markers in human cells. *Methods Mol Biol.* 2015;1292:3–18.
22. Huang L-W, Lin C-Y, Lee C-C, Liu T-Z, Jeng C-J. Overexpression of GRP78 Is Associated With Malignant Transformation in Epithelial Ovarian Tumors: Applied Immunohistochemistry & Molecular Morphology. 2012;20(4):381–5.
23. Samanta S, Tamura S, Dubeau L, Mhawech-Fauceglia P, Miyagi Y, Kato H, et al. Clinicopathological significance of endoplasmic reticulum stress proteins in ovarian carcinoma. *Sci Rep.* 2020;10(1):2160.
24. Bastida-Ruiz D, Yart L, Willemin C, Ribaux P, Morris N, Epiney M, et al. The fine-tuning of endoplasmic reticulum stress response and autophagy activation during trophoblast syncytialization. *Cell Death Dis.* 2019;10(9):1–15.
25. Bastida-Ruiz D, Willemin C, Pederencino A, Yaron M, Tejada BM de, Pizzo SV, et al. Activated α₂-macroglobulin binding to cell surface GRP78 induces trophoblastic cell fusion. *Scientific Reports.* 2020;10.
26. Marchetti T, Ruffatti A, Willemin C, de Moerloose P, Cohen M. Hydroxychloroquine restores trophoblast fusion affected by antiphospholipid antibodies. *J Thromb Haemost.* 2014;12(6):910–20.
27. Cohen M, Petignat P. Purified autoantibodies against glucose-regulated protein 78 (GRP78) promote apoptosis and decrease invasiveness of ovarian cancer cells. *Cancer Lett.* 2011;309(1):104–9.

28. Van Hoesen K, Meynier S, Ribaux P, Petignat P, Delie F, Cohen M. Circulating GRP78 antibodies from ovarian cancer patients: a promising tool for cancer cell targeting drug delivery system? *Oncotarget*. 2017;8(63):107176–87.
29. Yadav RK, Chae S-W, Kim H-R, Chae HJ. Endoplasmic reticulum stress and cancer. *J Cancer Prev*. 2014;19(2):75–88.
30. Fradet S, Pierredon S, Ribaux P, Epiney M, Shin Ya K, Irion O, et al. Involvement of membrane GRP78 in trophoblastic cell fusion. *PLOS ONE*. 2012;7(8):e40596.
31. Okada T, Haze K, Nadanaka S, Yoshida H, Seidah NG, Hirano Y, et al. A serine protease inhibitor prevents endoplasmic reticulum stress-induced cleavage but not transport of the membrane-bound transcription factor ATF6. *J Biol Chem*. 2003;278(33):31024–32.
32. Bu L-J, Yu H-Q, Fan L-L, Li X-Q, Wang F, Liu J-T, et al. Melatonin, a novel selective ATF-6 inhibitor, induces human hepatoma cell apoptosis through COX-2 downregulation. *World J Gastroenterol*. 2017;23(6):986–98.
33. Papandreou I, Denko NC, Olson M, Van Melckebeke H, Lust S, Tam A, et al. Identification of an Ire1alpha endonuclease specific inhibitor with cytotoxic activity against human multiple myeloma. *Blood*. 2011;117(4):1311–4.
34. Zhao Z, He K, Zhang Y, Hua X, Feng M, Zhao Z, et al. XRCC2 repairs mitochondrial DNA damage and fuels malignant behavior in hepatocellular carcinoma. *Cancer Lett*. 2021;512:1–14.
35. Axten JM, Romeril SP, Shu A, Ralph J, Medina JR, Feng Y, et al. Discovery of GSK2656157: An Optimized PERK Inhibitor Selected for Preclinical Development. *ACS Med Chem Lett*. 2013;4(10):964–8.
36. Matsuoka M, Komoike Y. Experimental Evidence Shows Salubrinal, an eIF2α Dephosphorylation Inhibitor, Reduces Xenotoxicant-Induced Cellular Damage. *International Journal of Molecular Sciences*. 2015;16(7):16275–87.
37. Uygur B, Leikina E, Melikov K, Villasmil R, Verma SK, Vary CPH, et al. Interactions with Muscle Cells Boost Fusion, Stemness, and Drug Resistance of Prostate Cancer Cells. *Molecular Cancer Research*. 2019;17(3):806–20.
38. Miroshnychenko D, Baratchart E, Ferrall-Fairbanks MC, Velde RV, Laurie MA, Bui MM, et al. Spontaneous cell fusions as a mechanism of parasexual recombination in tumour cell populations. *Nat Ecol Evol*. 2021;5(3):379–91.
39. Harewood L, Fraser P. The impact of chromosomal rearrangements on regulation of gene expression. *Hum Mol Genet*. 2014;23(R1):R76-82.
40. Cope L, Wu R-C, Shih I-M, Wang T-L. High level of chromosomal aberration in ovarian cancer genome correlates with poor clinical outcome. *Gynecologic Oncology*. 2013;128(3):500–5.
41. Mirzayans R, Andrais B, Murray D. Roles of Polyploid/Multinucleated Giant Cancer Cells in Metastasis and Disease Relapse Following Anticancer Treatment. *Cancers (Basel)*. 2018;10(4).
42. van Zijl F, Krupitza G, Mikulits W. Initial steps of metastasis: cell invasion and endothelial transmigration. *Mutat Res*. 2011;728(1–2):23–34.

43. Stuelten CH, Parent CA, Montell DJ. Cell motility in cancer invasion and metastasis: insights from simple model organisms. *Nat Rev Cancer*. 2018;18(5):296–312.
44. Jiang WG, Sanders AJ, Katoh M, Ungefroren H, Gieseler F, Prince M, et al. Tissue invasion and metastasis: Molecular, biological and clinical perspectives. *Semin Cancer Biol*. 2015;35 Suppl:S244–75.
45. Lv H, Shi Y, Zhang L, Zhang D, Liu G, Yang Z, et al. Polyploid giant cancer cells with budding and the expression of cyclin E, S-phase kinase-associated protein 2, stathmin associated with the grading and metastasis in serous ovarian tumor. *BMC Cancer*. 2014;14:576.
46. Zhang S, Mercado-Urbe I, Hanash S, Liu J. iTRAQ-based proteomic analysis of polyploid giant cancer cells and budding progeny cells reveals several distinct pathways for ovarian cancer development. *PLoS One*. 2013;8(11):e80120.
47. Suetsugu A, Matsumoto T, Hasegawa K, Nakamura M, Kunisada T, Shimizu M, et al. Color-coded Live Imaging of Heterokaryon Formation and Nuclear Fusion of Hybridizing Cancer Cells. *Anticancer Res*. 2016;36(8):3827–31.
48. Mujcic H, Rzymiski T, Rouschop KMA, Koritzinsky M, Milani M, Harris AL, et al. Hypoxic activation of the unfolded protein response (UPR) induces expression of the metastasis-associated gene LAMP3. *Radiother Oncol*. 2009;92(3):450–9.
49. Koumenis C, Wouters BG. « Translating » tumor hypoxia: unfolded protein response (UPR)-dependent and UPR-independent pathways. *Mol Cancer Res*. 2006;4(7):423–36.
50. Janczar S, Nautiyal J, Xiao Y, Curry E, Sun M, Zanini E, et al. WWOX sensitises ovarian cancer cells to paclitaxel via modulation of the ER stress response. *Cell Death Dis*. 2017;8(7):e2955.
51. Avril T, Vauléon E, Chevet E. Endoplasmic reticulum stress signaling and chemotherapy resistance in solid cancers. *Oncogenesis*. 2017;6(8):e373.
52. Notte A, Rebucci M, Fransolet M, Roegiers E, Genin M, Tellier C, et al. Taxol-induced unfolded protein response activation in breast cancer cells exposed to hypoxia: ATF4 activation regulates autophagy and inhibits apoptosis. *Int J Biochem Cell Biol*. 2015;62:1–14.
53. Liao P-C, Tan S-K, Lieu C-H, Jung H-K. Involvement of endoplasmic reticulum in paclitaxel-induced apoptosis. *J Cell Biochem*. 2008;104(4):1509–23.
54. Mhaidat NM, Alali FQ, Matalqah SM, Matalaka II, Jaradat SA, Al-Sawalha NA, et al. Inhibition of MEK sensitizes paclitaxel-induced apoptosis of human colorectal cancer cells by downregulation of GRP78. *Anticancer Drugs*. 2009;20(7):601–6.

Figures

Figure 1

Endoplasmic reticulum (ER) stress induction enhances cell fusion in ovarian cancer cells. a to d- SKOV3 (a and c) and COV318 (b and d) cells were treated with three different ER stress inducers, 100nM Thapsigargin (THA), 1 μ M HA15 or 1 μ g/mL Tunicamycin (TUN) for 48h. Untreated cells were used as the control condition. a and b- Protein levels for GRP78 and GAPDH were assessed by Western blotting. The representative images of bands for GRP78 and GAPDH were taken from the same gel (Supplementary

Figure 2), and each image was cropped, as delineated in Supplementary Figure 2, as well as adjusted for image intensity for optimal visualization. c and d- Fusion index was calculated for the different treatments and expressed as percentages relative to the control condition. Results are expressed as mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. e to h- SKOV3 (e, g) and COV318 (f, h) cells were treated with either 100nM Thapsigargin (THA) (e, f) or 1 μ g/mL Tunicamycin (TUN) (g, h), combined with inhibitors of the three UPR pathways: 17.5 μ M Salubrinal (SAL) or 0.3 μ M GSK2656157 (GSK) for PERK pathway inhibition, 200 μ M 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride (AEBSF) or 1 mM Melatonin (Mel) for ATF6 pathway inhibition, and 8 μ M 4 μ 8C or 2 μ M STF-083010 (STF) for IRE1 α pathway inhibition, for 48h. Fusion index was expressed as percentages relative to fusion index calculated for either THA or TUN condition. Results are expressed as mean $\pm$ SEM. ns: non significant, * $p < 0.05$, ** $p < 0.01$.

Figure 2

Formation of ovarian cancer PGCCs by cell fusion. a- Ovarian cancer cells (COV318 and SKOV3) double-positive for green and far-red cytoplasmic dyes were sorted by flow cytometry following induction of cell fusion with 300 μ M CoCl₂ for 48h. Scale bars represent 100 μ m. b- SKOV3 cells were transfected with plasmids encoding for red (SKOV3-Red) and green (SKOV3-Green) nuclear fluorescent proteins and mixed in co-culture. Under regular culture conditions, cells spontaneously formed PGCCs by cell fusion, visualized by polynucleated cells with both red and green nuclei (circled in yellow). Scale bars represent 200 μ m (with two high magnifications of selected portions at the bottom). c and d- Co-culture of SKOV3-Green and SKOV3-Red cells was treated with 1 μ M HA15 for 48h. c- The presence of polynucleated cells with both red and green nuclei was detected by confocal microscopy. Scale bars represent 100 μ m. d-

Fusion index was calculated separately for green polynucleated cells (FIGreen), red polynucleated cells (FIRed), and red/green polynucleated cells (FIGreen+Red). Results are expressed as mean percentages relative to the fusion index calculated for the control condition.

Figure 3

Ovarian cancer PGCCs can proliferate and form tumors on chick chorioallantoic membrane (CAM). SKOV3-Red and SKOV3-Green were mixed in co-culture with double-antibiotic. a and b- Cell population (SKOV3-M) obtained after three weeks of double selection. a- Visualization of double-positive mononucleated cells by fluorescent microscopy. Upper left: green signal (GFP). Upper right: red signal (mCherry). Lower left: Phase contrast. Lower right: Overlay. Scale bars represent 100 μ m. b- Quantification of red (mCherry), green (GFP), and double-positive cells by flow cytometry in SKOV3-Red, SKOV3-Green, SKOV3-M, and untransfected SKOV3 cells (ct-SKOV3). SKOV3-M cells were analyzed at passages 51, 63, and 72. c and d- Cell cycle distribution was analyzed in SKOV3-Red, SKOV3-Green, and SKOV3-M cells by flow cytometry following Hoechst staining. Doublets cells have been removed from the analysis based on Hoescht W and Hoescht A signals. c- Cell distribution depending on Hoechst signal intensity. d- Percentages of cells in G0-G1 and G2 phases. e and f- Proliferation capacities of SKOV3-Red, SKOV3-Green, and SKOV3-M cells were assessed by analyzing CellTraceTM far-red cytoplasmic signal dilution by flow cytometry. Cells were stained on day 0 (D0) and seeded in 10 cm culture dishes (5×10^5 cells/dish). Far-red signal was analyzed for five consecutive days (D0 to D4). Doublets cells have been removed from the analysis based on SSC W and SSC A signals. e- Evolution of far-red signal intensity from D0 to D4. f- Evolution of far-red signal dilution expressed as a percentage of mean signal intensity $\pm$ SEM measured at D0. g and h- Tumorigenic capacity of SKOV3-Red, SKOV3-Green, and SKOV3-M cells was assessed using the CAM model. Cells were inoculated on egg development day (EDD) 8, and tumor size was measured at EDD10.5 and EDD13.5. g- Images of the tumors growing in CAM at EDD13.5 h-

Tumor growth (tumor size at EDD13.5 relative to tumor size at EDD10.5). Results are expressed as mean $\pm$ SEM.

Figure 4

SKOV3-M cells display differential gene expression and invasive capacity compared to its parental cell lines. a- Heat maps of genes expression in SKOV3-M, SKOV3-Green, and SKOV3-Red. Colors from yellow to blue indicate downregulated cellular genes; colors from yellow to red indicate upregulated cellular genes. b- Invasive capacity of SKOV3-Red, SKOV3-Green, and SKOV3-M cells was assessed by the Boyden chamber assay. Cell viability was assessed by MTT assay. Results are expressed as invasion relative to cell viability $\pm$ SEM. AU: arbitrary unit, * p<0.05, ** p<0.01.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [YartetalBMCCancerSuppInfv02.docx](#)