

Correlate of Post-natal care utilization in rural Ethiopia: An evidence from Secondary data analysis of Ethiopian Demographic Health survey 2016.

REKIKU Fikre Abebe (✉ frekiku@yahoo.com)

Dilla university <https://orcid.org/0000-0003-1324-1631>

Berhan Tsegaye Negash

Hawassa University College of Medicine and Health Sciences

Zelalem Tenaw bogale

Hawassa University College of Medicine and Health Sciences

wegene Jemebere Beru

Hawassa University College of Medicine and Health Sciences

Getnet Kassahun Molla

Hawassa University College of Medicine and Health Sciences

Research article

Keywords: Postnatal care utilization, rural women, Ethiopia 2016.

Posted Date: November 7th, 2019

DOI: <https://doi.org/10.21203/rs.2.16954/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Abstract Background: Postnatal care utilization is the most neglected care in Ethiopia, despite a large proportion of maternal and neonatal deaths occurring during the postnatal period. Evidence suggests that utilization of postnatal care averts major complications of the fetus and mother. But, the proportion of rural women utilizing postnatal care is low in Ethiopia. Information about prevalence and factors which determine utilization of postnatal care among rural women in Ethiopia is mandatory for policy making, and program design to enhance its utilization. **Objective:** The objective of this study was to determine the prevalence of utilization of postnatal care and to identify its determinant factors among rural women who gave birth in the past five years before the Ethiopian demographic health survey, in 2016. **Method:** This study utilized data from the Ethiopian demographic health survey 2016 for analysis. It is a national two-stage, cross-sectional study. It analyzed data for rural women who gave birth at least one time in the past five years before the data survey time. Logistic regression was applied to identify explanatory variables associated with the outcome variable. Adjusted odds ratio with 95% confidence interval was computed and P -value < 0.05 was considered as statistically significant. **Result:** The prevalence of postnatal care utilization among rural women was 11.4%. Place of delivery [AOR=4.3, 95%CI, 1.4-5.2], ANC visit [AOR=2.1, 95%CI, 1.1-3.9] and Women in the richest wealth quintile [AOR=2.97, 95%CI, 1.9-4.5] were factors associated with postnatal care service utilization among rural women in Ethiopia in 2016. **Conclusion and recommendation:** This study showed that the prevalence of postnatal care utilization was low. Being in the richest wealth quintile, history of ANC visit and place of delivery were positive predictors of postnatal care utilization among rural women. Regardless of proven strategies utilized to maximize utilization of postnatal care by the government of Ethiopia, this study showed that prevalence is low in rural areas. Therefore, community awareness creation, increasing institutional coverage by expanding maternity waiting areas and besides, the government should design and implement income increasing packages among rural women.

Introduction

The postnatal period is a vital stage in the lives of mothers and newborn babies. Most maternal and infant deaths occur during this time [1]. The post-natal period is the most eventful period in which delay to intervene at the time of complication during this period will result in short-term effects on lifelong disability and death [2]. Prompt action at the time of complication will decrease adverse effects both for fetuses and mothers [3], [4], [5].

In 2017, about 295 000 women died during and following pregnancy and childbirth. The vast majority of these deaths (94%) occurred in low-resource settings, and Sub-Saharan Africa accounts two-thirds of maternal death followed by South Asia accounts one-fifth [6]. Despite this fact, the sub-Saharan region achieved successes and decreased 40% of MMR since 2000. However, maternal deaths in some areas of the world reflect inequalities. The MMR in low-income countries in 2017 is 462 per 100 000 live births versus 11 per 100 000 live births in high-income countries and the risk of maternal mortality is highest for

adolescent girls under 15 years old and complications in pregnancy and childbirth are higher among adolescent girls age 10-19 [7], [8].

The fact that 18 million women in Africa currently do not give birth in a health facility poses challenges for planning and implementing postnatal care for women and their newborns. Despite its importance, this period is generally the most neglected in developing countries and most mothers and newborn babies do not receive postnatal care services from a skilled health care provider during the critical first few days after delivery [9], [10].

Postnatal utilization in Ethiopia has very slow progress. A large proportion of maternal and neonatal deaths occur during the first 24 hours after delivery. This period is most critical for both mother and neonatal survival, and the most vital time for recognizing problems and initiating early interventions to avoid complications [11, 12]. However, postnatal care utilization continues to have the lowest coverage, according to 2016 EDHS with only 17% of the women utilizing postnatal care.

Forty-five percent of urban women received a postnatal check-up within 2 days compared to 13% of postnatal care in rural: Seventeen percent of women and 13% of newborns received a postnatal check within the first 2 days of birth [13]. Prevalence and determinant factors of postnatal care are mandatory for effectively design policies and programs about at national level for rural women. Therefore, the main aim of this study is to assess prevalence and determinant factors of postnatal care utilization among rural women.

Methods And Materials

Study area and period

Ethiopia is one of the developing countries which is found in East Africa. Administratively, it is divided into nine regional states and two city administrations. Each region is divided into zones, and zones are further sub-divided into districts. There are a total of 817 districts throughout the country. Districts are even sub-divided into Kebeles (the lowest administrative units). There were 16,253 Kebeles. The total population of the nation reached 94.3 million in 2017. Nearly, 80% of Ethiopian population lives in rural areas. The average size of each household is estimated to be 4.6 individuals. It was conducted from January, 18 to June 27, 2016.

Study variables

In this study the dependent variable was postnatal care utilization. It has binary outcomes. A total of 5681 births and related information were reported within five years preceding survey and were included in this study. Analysis was restricted to these 5681 last births to minimize recalling bias.

Rural women were asked as 'did you utilize postnatal care at least once or not?'. They answered as 'yes' or 'no'. The response 'yes' was coded as '1' and the response 'No' was coded as '0'. Further details of sampling, questionnaire and procedure can be found in the publicly available survey

report[13].Independent variables were categorized into socio-demographic characteristics and reproductive health characteristics. Variables like age and family size was recorded as continues variable and was categorized into three categories, ethnicity was categorized into six parts according to their relative proportion of population. But, there are more than 85 ethnicities in Ethiopia. Women's occupational status was merged into five main occupations. However, more than nine occupations were listed in the data. place of residence, respondent's educational status, respondent's husband educational status, religion, ethnicity, marital status and wealth index and frequency of listening radio.Moreover, reproductive health characteristics consist of age at first pregnancy, number of birth, number of antenatal visit, current pregnancy.

Operational definition

Wealth index was constructed using principal components analysis on household asset data. Individuals were classified into five wealth quintiles (poorest, poorer, medium, richer and richest). It included ownership of selected household assets, size of agricultural land, quantity of livestock and materials used for house construction(Rutstein SO: The DHS Wealth Index: Approaches for rural and urban areas, 2008).

Sample size and sampling procedures

This study used two stage, stratified and cluster sampling technique to select households. Ethiopian demographic health survey 2016 has utilized 84,915 list of enumeration areas as a sampling frame to samples. An enumeration area is a geographic area which contains one hundred-eighty- one (181) households. The sampling frame contains information about the enumerating area location, type of residence, and estimated number of households.Each region was stratified into urban and rural areas, yielding 21 strata. In the first stage, a total of 645 enumeration area (EAs). These were divided into rural and urban (202 EAs in urban and 443 EAs in rural areas). They were selected based on probability proportional to the sizeof EA.

In the second stage, a new sampling frame was created which consist of lists of households for the selected strata. Then a fixed number of 28 households per cluster were selected with an equal probability using systematic random sampling technique from the new sampling frame. All women in reproductive age group lived in selected households were eligible for interview. A total of 18,008 households were selected for this study. In the interviewed households, a total of 7590 women gave birth five years back from survey data collection time were included for analysis in this study. All of them completed the interview, yielding a response rate of 100%.

Study design and population

This study is the secondary data analysis of Ethiopian demographic health survey 2016.It is a cross sectional study which was conducted over five and half months' period. All reproductive age women in

Ethiopia who gave birth five years before EDHS 2016 data collection period were target(source) population of this study.

Study population of the current study included all women who gave birth five years before EDHS 2016 data collection time, lived in the selected strata, and presented during data collection time.

Data collection and analysis

EDHS 2016 used nationally validated standard questionnaire. Pretest was conducted from October 1-28, 2015. The questionnaire was prepared first in Amharic, Oromiffa and Tigrigna which are the three common local language spoken by majorities of Ethiopian population respectively. Then, it was translated into English language. Finally, it was translated back to local languages for consistency. Though the questionnaire consists of five different parts, we used data collected from women of reproductive age group for this analysis. Fieldwork was carried out by 33 field teams. Each consisting of one team supervisor, one field editor, three female interviewers, one male interviewer, two biomarker technicians, and one driver. In addition, 28 quality controllers were dispatched during data collection. Complex sample analysis was performed using SPSS version 22 statistical software to account for the multistage cluster study design. Analysis was carried out based on the weighted count. Sample weight was an eight-digit variable with six implied decimal places and was divided by 1,000,000 before applying the weighting factor. Multivariable logistic regression was used to identify independent predictors of caesarean section delivery. Results were presented in the form of odds ratios, and 95 % confidence intervals. Statistical significance was set at a p-value of 0.05.

Results

Sample socio-demographic characteristics and postnatal care utilization

Out of the 5681 rural women who gave birth in the last five years who answered question about postnatal care service 11.4% of rural women utilized postnatal care in Ethiopia. Table one showed that socio-demographic characteristics of study participants who had at least one birth in the past five years, before Ethiopian demographic and health survey 2016, data collection time. The mean age of study participants was 29.3±7 years. The majority of 93.8% women were married. 70.1% of respondents had no formal education and 80 % women not listening radio at all. In addition, more than half (58.4%) of study participants had no formal work. Besides, majorities 3540(62.3%) of the subjects lived in household size of more than five in this study **see table 1**.

Among respondents 93.2% of women's current pregnancy was unwanted. Majorities 61% of the women had less than four children. 45.8% of women were using contraceptives. Those women who delivered at health institution accounted of 1533(27%). Study subjects who were pregnant at the time of study reached to 671(11.8%). Majorities of the study participants 4321(76.1%) had no pregnancy complications **see table 2**.

Determinants of postnatal utilization among rural women

In this study, frequency of listing radio, family size, wealth quintile, place of delivery, told about pregnancy complication and husband occupation were factors significantly associated with postnatal utilization in rural women in a bivariate logistic regression model. However three variables were associated statistically in AOR model. Place of delivery 4.3 times [AOR=4.3, 95%CI, 1.4-5.2] more likely to use post-natal service than their counterparts. ANC was 2.1 times [AOR=2.1, 95%CI, 1.1-3.9] more likely to use postnatal service than their counterpart. Women in the richest wealth quintile were 2.97 times [AOR=2.97, 95%CI, 1.9-4.5] more likely to use post-natal service than their counterparts **see table 3**.

Discussion

The prevalence of post-natal utilization in rural women was 11.4%. This finding is lower than the national prevalence of postnatal utilization in Ethiopia it was 17% [13]. The possible explanation for this due to poor access to road to facilities and most women give birth at home. This finding was lower than study conducted in Rwanda 12.8% [14], Nigeria 16.9% [15], Bangladesh 73.8% [16], India 36.6% [17] and Myanmar 83.68% [18]. The possible rationale for this difference might be due to women's socio-demographic difference, low level of awareness. Moreover, culturally embedded male dominance in rural Ethiopia reduce women's decision ability to utilize maternal health services, in rural Ethiopia. Women in richest wealth quintile were 2.97 times more likely to utilized postnatal service than the poorest one. This finding is inconsistent with a study in Nepal and Rwanda [14], [19]. The possible rationale might be that poor women in rural Ethiopia could not afford indirect costs for delivery even if the service is free in public health institutions. Poor women are more likely to be less educated, less autonomous and less access to health services.

Place of delivery 4.3 times [AOR=4.3, 95%CI, 1.4-9.2] more likely to use post-natal service than their counterparts. This finding is consistent with the study in Nepal [19], Tanzania [20], and Rwanda [14]. The possible rationale could be if women delivered at health facilities the knowledge about the importance of the service could be clear for women and the facilities could create a link with health post to get the service. Women with ANC Visit were 2.1 times [AOR=2.1, 95%CI, 1.1-3.9] more likely to use postnatal service than their counterparts.

This finding is consistent with study in Nigeria [15]. The possible reason for this might be that women could get education and the approaches of women-friendly service of providers create an opportunity for the women to give birth at institution.

Conclusion and public health implication

Being in richest wealth quintile, place of delivery and ANC visit were positive predictors of postnatal care utilization among rural women. Results from this study point to a combination of community and facility-based interventions as feasible measures to reach poor and home delivery mothers in rural Ethiopia.

Community awareness creation and strong legal policies should be drafted towards home delivery among rural women in Ethiopia. Besides, the government should design and implement income increasing package among rural women in Ethiopia. These should be the focus of future strategy to effectively utilize postnatal care utilization .

Strength and Limitation

This study has many strengths. These include a national wide study with large sample size and high response rate. Moreover, demographic and health surveys are internationally validated and nation. adapted surveys. Therefore, the current findings are generalizable to rural women of the entire country. The study design and sampling follow standard procedure which makes possible to yield accurate and reliable estimate. Nevertheless, the current study has several limitations. This study could create recall bias. Nevertheless, this bias is not considered problematic since this study included only women giving birth within five years preceding the survey. Moreover, this secondary data analysis of Ethiopian demographic and health and survey could not provide variables associated with pregnancy complications.

Abbreviations

AOR———Adjusted odds ratio

EDHS———Ethiopia demographic and health survey

MMR———Maternal mortality ration

Declarations

Ethics approval and consent to participate

A written ethical clearance was obtained from the Institutional Review Board at the College of Medicine and Health Sciences of Hawassa University, Hawassa, Ethiopia.

Acknowledgment

We are grateful to the Ethiopia static bureau for delivering us the data and senior staff for technical support.

Author contribution

RF Design the study, Proposal development, data analysis and manuscript preparation

BT,ZT,WJ and GK participated on data analysis and manuscript preparation

Funding

No source of funding to carry out for this research.

Availability of data and materials

All data generated or analyzed during this study were included in this published article and its supplementary information files

Consent to publish

Not applicable

Competing interest

No competing interest

References

- 1.Lawn, J., M. Kinney, and H.J.L. Blencowe, *Every newborn, an executive summary for the lancet series*. 2014. **384**(9938): p. 1-8.
- 2.Organization, W.H., *Postpartum care of the mother and newborn: a practical guide: report of a technical working group*. 1998, Geneva: World Health Organization.
- 3.Organization, W.H., *WHO technical consultation on postpartum and postnatal care*. 2010, Geneva: World Health Organization.
- 4.Lawn, J.E., et al., *4 million neonatal deaths: when? Where? Why?* 2005. **365**(9462): p. 891-900.
- 5.Warren, C., et al., *Opportunities for Africa's newborns. "Postnatal care."*. 2006.
- 6.Organization, W.H., *Trends in maternal mortality 2000 to 2017: estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division: executive summary*. 2019, World Health Organization.
- 7.Ganchimeg, T., et al., *Pregnancy and childbirth outcomes among adolescent mothers: a World Health Organization multicountry study*. 2014. **121**: p. 40-48.
- 8.Althabe, F., et al., *Adverse maternal and perinatal outcomes in adolescent pregnancies: The Global Network's Maternal Newborn Health Registry study*. 2015. **12**(2): p. S8.
- 9.UNICEF, W., UN and World bank, *Level and trends of child mortality*. 2015.
- 10.DiBari, J.N., et al., *Use of postpartum care: predictors and barriers*. 2014. **2014**.

11. Organization, W.H. *WHO recommendations on postnatal care of the mother and newborn*. 2014; Available from: https://www.who.int/maternal_child_adolescent/documents/postnatal-care-recommendations/en/.
12. WHO, *Technical consultation on postpartum and postnatal care*. 2013: Geneva, Switzerland.
13. *Ethiopian demographic and health survey 2016*. 2016: Ethiopia.
14. Bernard N. Rwabufigiri, J.M., Dana R. Thomson¹, Bethany L. Hedt-Gautier and Jean Paul S. Semasaka, *Factors associated with postnatal care utilization in Rwanda: A secondary analysis of 2010 Demographic and Health Survey data*. BMC pregnancy and childbirth, 2016. **16**(122).
15. Idris Usman Takai, H.D.D., Mohammed Bukar, Bala Mohammed Audu, Aliyu Salihu and Kwayabura, *Factors responsible for under-utilization of postnatal care services in Maiduguri, north-eastern Nigeria*. Sahel Medical Journal,, 2015. **18**.
16. S K Nayan, N.B., M R Abid, S Rahman, A K Rajib, N Farzana and M L Nahar, *Northern International Medical College Journal*. Utilization of Postnatal Care Services among the Rural Women in Bangladesh, 2017. **8**: p. 208-212.
17. Arvind sharma, p.s.t., pradeep kumar kasar, rajesh tiwari and , richa sharma, *utilization of post natal care in tribal area of madhya pradesh*. international journal of medical science and public health, 2014
18. Mon, A., et al., *Utilization of full postnatal care services among rural Myanmar women and its determinants: a cross-sectional study [version 1; peer review: 3 approved]*. 2018. **7**(1167).
19. Vishnu Khanal, M.A., Rajendra Karkee and Tania Gavidia, *Factors associated with the utilisation of postnatal care services among the mothers of Nepal: analysis of Nepal Demographic and Health Survey 2011*. BMC women health, 2014. **14** p. 19.
20. Almamy M. Kanté, C.E.C., Anna M. Larsen, Amon Exavery, Kassimu Tani and James F. Phillips *Factors associated with compliance with the recommended frequency of postnatal care services in three rural districts of Tanzania*. BMC Pregnancy and Childbirth, 2015. **15**: p. 341.

Tables

Table 1: Socio-demographic characteristics of study participants who utilized postnatal service in rural women, EDHS 2016.

Variables	Category	Frequency	Percentage
Age	15-24	1761	30.3
	25-34	2138	37.6
	≥35	1782	31.4
Ethnicity	Amhara	855	15
	Oromo	1399	24.5
	Tigre	631	11.1
	Afar	551	9.7
Religion	Orthodox	1654	32.9
	Protestant	1124	20.6
	Muslim	2760	44.5
	Catholic	37	2.00
Marital status	Single	82	1.4
	Married	5326	93.8
	Widowed	194	3.4
	Divorced	79	10.9
Educational status	No education	3983	70.1
	Primary	1426	25.1
	Secondary	223	3.9
	Higher	49	0.9
Wealth index	Poorest	2362	41.6
	Poor	1149	20.2
	Middle	998	17.6
	Rich	860	15.1
	Richest	312	5.1
Family size	1-4	1755	31
	5-9	3185	56.1

	>=10	741	13
Frequency of listening radio	Not at all	4546	80
	Less than once a week	602	10.6
	At least once a week	533	9.4

Table 2: Reproductive health characteristics of study participants who utilized postnatal service in rural women, EDHS 2016.

Variables	Category	Frequency	Percentage
Current Pregnancy	Yes	671	6.8
	No	5010	93.2
Ever had terminated pregnancy	Yes	498	7.6
	No	5183	92.4
Total number of birth	1-4	1755	31
	>=5	3926	69
Current use contraceptive	Yes	2560	45
	No	3121	55
Place of delivery	Home	4148	79.1
	Institution	1533	20.9
Told about pregnancy complication	Yes	1360	23.9
	No	4321	76.1

Table 3: Factors associated with postnatal utilization of rural women by multivariable logistic regression, EDHS 2016.

Variables	Category	Post-natal utilization		COR[95%CI]	AOR[95%CI]
		Yes	No		
Place of delivery	Home	4047	101	1	
	Institution	1489	44	4.3[1.1-7.1]	4.3[1.4-5.2]**
Wealth index	Poorest	1901	32	1	
	Poorer	1242	9	1.1[0.4-2.2]	1.1[0.2-2.0]
	Middle	1354	28	1.3[0.5-6.2]*	1.3[0.3-5.5]
	Richer	1321	12	1.8[0.1-3.1]*	1.6[0.3-6.1]
	Richest	1613	78	2.97[2.4-11.6]*	2.97[1.9-4.5]**
ANC visit	No	3286	69	1	1
	Yes	4102	132	2.1[1.3-4.1]*	2.1[1.1-3.9]**