

Phytoremediation with Kenaf (*Hibiscus Cannabinus* L.) for Cadmium-Contaminated Paddy Soil in Southern China: Translocation, Uptake and Assessment of Cultivars

Yuan Guo (✉ guoyuan@caas.cn)

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops <https://orcid.org/0000-0002-1525-6397>

Qingmei Xiao

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Xinlin Zhao

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Zhimin Wu

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Zhigang Dai

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Minji Zhang

Shanghai Zhongwei Biochemistry Co., Ltd

Caisheng Qiu

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Songhua Long

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Yufu Wang

Chinese Academy of Agricultural Sciences Institute of Bast Fiber Crops

Research Article

Keywords: Cadmium, *Hibiscus cannabinus* L., Paddy soil, Phytoremediation, Phloem, Bioconcentration factor, Translocation factor

Posted Date: August 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-760873/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)



Abstract

Kenaf (*Hibiscus cannabinus* L.) is suitable for growing in heavy metal-polluted soil for non-food purposes, and can be used as a potential crop to remediate heavy metal-contaminated soil. The main objective of this study was to investigate kenaf phytoextraction of cadmium (Cd), including uptake, translocation, and accumulation differences in tissues among kenaf cultivars. A field experiment was conducted in a Cd contaminated paddy field in southern China area with 13 kenaf cultivars in 2015 and 2016. Agronomic performance, Cd concentrations in plant tissues (root, xylem, and phloem), and biomass of different tissues of each cultivar were measured and evaluated. Significant differences in Cd concentrations and accumulation among tissues and cultivars were observed. The phloem had the highest Cd accumulation and transfer capability compare with the roots and xylem. Approximately 35 ~ 65g of Cd could be taken up by the aerial parts of different kenaf cultivars within every hectare of soil. The percentage of Cd uptake by the phloem ranged from 47–61% and by the xylem ranged from 38–53%. By evaluating the agronomic traits and Cd bioaccumulation capacity, Fuhong 952, Fuhong 992, and Fuhong R1 were regarded as Cd accumulators for the phytoremediation of Cd-contaminated soil. Our study clearly demonstrated that a significant level of Cd in the soil was taken up through the phytoremediation with kenaf. In addition, harmless utilization of kenaf planting in Cd-contaminated paddy soil was discussed.

Introduction

In recent decades, the rapid industrialization and economic development have not only improved people's living standards but also induced serious contamination to the farmland in China. There are about 19.4% of the agricultural lands suffering from heavy metals and metalloid pollution according to the latest nationwide surveys. Among the metals or metalloid, cadmium (Cd) and arsenic (As) are the main contaminations to soils in China (Chen et al., 2020). Although Cd is non-necessary and even harmful to crops, it can still be easily assimilated by the plant root and consequently transported to their tissues. It was reported that rice plant could take in Cd from the soil and accumulate it with a high level in its aerial tissues even if the Cd content in soil was very low (Wang et al., 2011). The Chinese limit of Cd in brown rice is $0.2 \text{ mg}\cdot\text{kg}^{-1}$ (GB 2762 - 2012). The CODEX commission of FAO/WHO stipulates that the Cd content in the cereal grains rice and wheat should be < 0.4 and < 0.2 , respectively. Hence, restoring the heavy metal-contaminated paddy lands become an urgent task for China. As an in-situ and nondestructive remediation approach, phytoremediation can not only remediate the polluted soil but produce feedstock for further use. Apart from common heavy metal hyperaccumulators, some crops have also been used for phytoremediation, such as alfalfa, tobacco, flax, jute (Liu et al., 2019; Zhang et al., 2019; Guo et al., 2019; Guo et al., 2020). Since the removal rate of heavy metal depends on both the plant biomass and the metal content involved in it, the selection of a suitable plant become a key point in phytoremediation (Sumiahadi and Acar 2018; Guo et al., 2020).

Kenaf (*Hibiscus cannabinus* L.) is one of the most important fiber crops with high economic benefits, and massive heavy metals accumulated found in kenaf dose not enter the food chain (Deng et al., 2017; Chen

et al., 2021). Kenaf plants grow rapidly with a large biomass, and the plants can grow to a height of 4–6 m and yield fresh biomass of up to 100–150 t per hectare within a 4-month growth period. Therefore, kenaf is a potential plant for use in the phytoremediation of widespread heavy metal contamination. Kenaf plants can remove 89.2% of Cd in contaminated water by rhizofiltration (Niazy 2015). During 3 months of growth in sludge, kenaf accumulated 2.49 mg·kg⁻¹ Cd and 82.5 mg·kg⁻¹ zinc without obvious toxic symptoms (Arbaoui et al., 2013). However, most of the research results were based on pot experiments, which are generally incompatible with large-scale phytomanagement owing to the interactions of many other factors affecting plants in the agricultural land. Furthermore, the characteristics of Cd accumulation has not been studied in the most economically important part of kenaf, phloem, for subsequent safety utilization in previous reports.

In the present study, the responses and phytoremediation potential of 13 different kenaf cultivars to Cd contamination were investigated in a field trial. In detail, the main objectives of this study were to: (i) compare the differences of Cd translocation and accumulation in kenaf tissues between cultivars; (ii) analyze the bioconcentration factors and translocation factors for Cd of kenaf cultivars; and (iii) to evaluate the phytoremediation potentials of kenaf cultivars and find more suitable cultivars for phytoremediation.

Material And Methods

Experimental site and soil

The field experiment was conducted at the garden village of Zhuzhou city, Hunan province, China (27°42'58.52"N, 113°11'8.26"E). The soil plumbum (Pb), Cd and As contents were determined and the result showed that the average Cd content was 2.25 times higher than that of the standard of Chinese paddy soil (Table 1). According to the standard for soil environmental quality and risk control of soil pollution in agricultural land, the soil of the experimental field was classified as moderately Cd-contaminated agricultural soil (GB 15618-2018).

Field experiment and sample collection

Thirteen kenaf cultivars were sown in the field in May 2015 and 2016. Among these cultivars, DGG, G4, GGMM, and GGS were not Chinese native varieties, with unlobed leaf shapes. The other nine cultivars were Chinese varieties that are mainly cultivated in the south of the Yangtze River. The leaves of these nine cultivars were partially lobed or deeply lobed (Fig. 1). There were total 39 plots (13 cultivars×3 replications) for growing kenaf in this experiment. Each plot was 2 m wide and 3 m long and all of them were arranged with a random blocked design. The seeding quantity of each plot was based on the seedling rate of cultivars and 40 seeds per square meter. The kenaf was harvest in mid-November of both 2015 and 2016, 10 kenaf plants that were uniformly distributed in each experimental plot were collected and mixed to obtain a sample. Then, the aerial parts of plants in each plot were harvested to obtain the fresh weight and dry weight.

The rhizosphere soil was removed, and the samples were cleaned with deionized water and then separated into three tissue types (root, xylem, and phloem). The samples obtained in 2015 were placed in a 105 °C drying oven for 30 min and then in a 70 °C drying oven until the weight of the samples become constant, the samples were then ground with a stainless-steel grinder (1000Y, China). Air-dry The rhizosphere soils were air-dried and sieved through a 60-mesh sieve for testing.

Chemical analysis

The chemical parameters of soil and plant samples were analyzed following the methods described in our previous study (Guo et al. 2020). The pH of soil samples were measured in a 1 : 2.5 soil-to-water suspension using a glass electrode (PHS-25, Leici, China). The soil samples were fully dissolved with hydrochloric-nitric-hydrofluoric-perchloric acid, the concentrations of Cd and Pb were determined by atomic absorption spectroscopy method (Z2310, Hitachi, Japan). For As, the soil samples were acid digested with aqua regia, and the concentration was then determined using the atomic fluorescence method (AFS-8230, Titan, China). The plant samples were digested with nitric-perchloric acid, and then the digestion solution was used to determine the metal concentration with a graphite furnace atomic absorption spectrometry (Z2000, Hitachi, Japan).

Bioconcentration factor and translocation factor

The bioconcentration factor (BF) and translocation factor (TF) were calculated to evaluate the accumulation and translocation of heavy metals of different kenaf cultivars (Yoon et al. 2006). BF is the ratio of metal concentration in the root to rhizosphere soil. TF is the ratio of metal concentration in the xylem to root (TF_{rx}) and phloem to root (TF_{rp}). BF and TF can be used to estimate accumulation and translocation capability of plant and the redistribution of metals throughout plant tissues.

Statistical analysis

The differences among the kenaf cultivars were evaluated using one-way analysis of variances and all statistical analyses were performed using the statistics package Excel 2010 and SPSS v19.0.

Results

Agronomic traits and fresh weight of kenaf cultivars in Cd-contaminated paddy soil

The heights of the kenaf plants were between 2.5 and 4.0 m in the two growing seasons. Most of cultivars were grew higher in 2016 than in 2015, expect for Hongma No.1 and Fuhong 952. The average plant height in the growing season in 2015 and 2016 were 3.1 and 3.3 m, respectively. The stem diameter of kenaf plants was between 13.0 and 19.1 mm in the two growing seasons. All of cultivars had a higher diameter in 2016 than in 2015. The average diameter in the growing season in 2015 and 2016 were 15.0 and 17.0 mm, respectively. The bark thickness of the kenaf plants was between 0.84 and 1.48 mm in the two growing seasons. Most of cultivars had thicker bark in 2016 than in 2015, except Hongma No.1 and Fuhong 952. The average bark thickness during the growing season in 2015 and 2016 was 1.02 and 1.22

mm, respectively. The fresh weights of the root, xylem, and phloem for 10 plants were respectively between 0.18 and 0.32 kg, 0.94 and 2.40 kg, and 0.62 and 1.60 kg, respectively. Among the 13 cultivars, the fresh weights of the root, xylem, and phloem for Fuhong 952 were more stable in the two seasons and relatively higher than those of the other cultivars (Fig. 2).

Cd concentration in rhizosphere soil and different tissues of kenaf plants

The Cd concentration in the rhizosphere soil in 39 experimental pots was between 0.71 mg·kg⁻¹ and 1.50 mg·kg⁻¹, with a mean concentration of 1.15 mg·kg⁻¹ (Fig. 3). The Cd concentrations in the rhizosphere soil of Hongzong No.1, Hongyou No.2, Fuhong 952, Fuhong 992, and Fuhong R1 were significantly higher than those of the other cultivars. The Cd concentrations in kenaf tissues are presented in Table 2. The Cd concentration ranges in the root, xylem, and phloem of 13 kenaf cultivars were 1.50–4.38 mg·kg⁻¹, 1.00–2.96 mg·kg⁻¹, and 2.39–5.18 mg·kg⁻¹, respectively, with mean concentrations of 2.67, 2.05, and 3.87 mg·kg⁻¹, respectively. The mean accumulation of Cd in the three tissues of kenaf were decreased in the following order: phloem > root > xylem, and the mean concentrations of Cd in the phloem of the 13 cultivars were 1.1–2.9 times higher than those in the root and xylem. The top three kenaf cultivars with the highest Cd concentrations were Hongzong No.1, Fuhong 992, and Fuhong 952.

Bioconcentration and translocation of Cd in kenaf tissues

BF was used to evaluate the ability of the kenaf plant to accumulate Cd in roots of the 13 kenaf cultivars. BFs of roots of the 13 kenaf cultivars were ranged from 1.74 to 2.96, with a mean of 2.35. No significant difference was found among cultivars, except for Fuhong 992 and GG MN, which showed the highest and lowest BFs in the 13 kenaf cultivars, respectively. TF was used to evaluate the ability of a kenaf plant to transfer Cd within its tissues. TFR_xs and TFR_ps of kenaf cultivars were ranged from 0.50 to 1.44 and 1.10 to 2.57, respectively. TFR_ps were higher than TFR_xs in all 13 cultivars, suggesting the phloem had a greater ability to transport Cd than that of the xylem. There were significantly higher differences among the TFR_ps of the cultivar than among the TFR_x and BF of the roots, indicating a difference in the Cd transport capacity of the phloem among kenaf varieties. TFR_ps of cultivars with an unlobed leaf shape (DGG, G4, GG MN, and GGS) were significantly higher than those of cultivars with other leaf shapes (Fig. 4).

Biomass of kenaf planting in Cd-contaminated paddy soil and assessment of Cd uptake for the 13 kenaf cultivars.

The biomass of the root, xylem, and phloem of 10 plants among the 13 cultivars were 0.06–0.10 kg, 0.17–0.36 kg, and 0.13–0.36 kg, respectively. The biomass of the xylem was higher than that of other tissues in all 13 kenaf cultivars, accounted for 46%–55% of the total biomass. Phloem biomass was lower than xylem biomass but higher than root biomass. The percentage of phloem biomass ranged from 29% to 38%, and the root biomass ranged from 14% to 19%. Fuhong 952 (0.72 kg) and G4 (0.37 kg) had the highest and lowest of total 10 plants biomass (Fig. 5).

In the actual agricultural operations, only the aerial parts of kenaf plants are harvested. According to the data of fresh weight and dry weight by experimental plots, and the Cd concentration in different tissues, we assessed Cd uptake of each cultivar for a hectare area. Because all leaves had fallen off during harvest, the biomass of leaves was not calculated in the total aerial biomass. Fuhong 952 and Fuhong R1 had the highest Cd phytoextraction capacity and uptake exceeding 60 g of Cd from every hectare of soil. For most cultivars, except for Fuhong R1, phloem phytoextraction accumulated more Cd than xylem. The percentage of Cd uptake by the phloem ranged from 47% to 61%, and that by the xylem ranged from 38% to 53% (Fig. 6).

Comprehensive evaluation of the 13 kenaf cultivars in the phytoremediation of Cd-contaminated soil

To comprehensive evaluate the 13 kenaf cultivars in the phytoremediation of Cd-contaminated soil, we normalized and clustered the data using the zero-to-one method in the heat map (Fig. 7). Fuhong 992, Fuhong 952, and Fuhong R1 exhibited better performance than other cultivars in most of the traits, except for the TFs of the xylem and phloem, and thus, they can be used as candidate plant materials for soil phytoremediation.

Discussion

Differences in Cd accumulation and phytoextraction potential of kenaf cultivars for Cd-contaminated soil phytoremediation

In the present study, differences have been detected in the bioaccumulation of Cd in different plant tissues, the result is the same as previously reported (Li et al., 2018). Differences were also detected in Cd uptake among the kenaf cultivars in the present study. However, differences were not detected in Cd concentrations of different kenaf cultivars when soil was severely contaminated with other heavy metals, such as Zn, Cu, and Cr (Li et al., 2018). It is speculated that the interaction between metal elements may affects the absorption of Cd in kenaf plants (Chen et al., 2016).

It is critical in the utilization of kenaf for phytoremediation is the total uptake of Cd from the square unit of the soil. In this study, the Cd uptake of 13 kenaf cultivars planted in one ha of soil was between 39.8 and 64.5 g. In our experiment, we used a single heavy metal-contaminated soil. In compound heavy metal-contaminated soil (Cd, Zn, Ni, Cu), 25.7g·hm⁻² of Cd can be phytoextracted by kenaf plants (Li et al., 2018), which is less than that by Hongma No.5, which takes up the least amount of Cd in our study, possibly because other heavy metals affect the absorption of Cd. Although the roots of kenaf immobilized large portions of the total plant Pb from the sand tailings (Mo et al., 2008), in the present study, root biomass only accounted for 14%–19% of the total plant, indicating that it is not the main Cd phytoextraction tissue of Cd in kenaf plants. In addition, root harvesting is not practical for large-scale farmland cultivation of kenaf.

Cd accumulation in the phloem of kenaf plant

Previous studies have reported heavy metal bioaccumulation in the roots, stems, and leaves of kenaf plants (Arbaoui et al., 2013; Li et al., 2018). However, these studies did not consider Cd accumulation in the phloem, which is an economically and important part of kenaf, and impacts the safety of subsequent utilization, such as fiber from phloem. In the present study, Cd concentration of the phloem ranged from 2.39 to 4.94 mg·kg⁻¹, with the Cd concentration of soil being 0.373–1.443 mg·kg⁻¹, proving that phloem accumulated the highest amount of Cd in the soil. The mean concentrations of Cd in the phloem of 13 cultivars were 1–3 times higher than those in other tissues. This indicates that the phloem is the greatest accumulator tissues of kenaf, rather than the root. The same results were previously verified in flax and jute plants, which are bast fiber crops such as kenaf (Guo et al., 2019; Guo et al., 2020). Within the shoot of rice, phloem transport is the basis for Cd redistribution and for the accumulation in fruits and seeds (Satoru et al., 2012; Page and Feller 2015). In potato plants, Cd can be rapidly distributed to all tissues via the phloem (Reid et al., 2010). This was also observed in hyperaccumulator (Lu et al., 2013; Wei et al., 2014; Feller et al., 2019). Phloem is the productive part of kenaf plant, which contents fibers that can be processed into textiles. Thus, it is important to evaluate the safety of raw material reutilization. In the degumming process, recycling Cd from the degumming solution is important.

Harmless utilization of kenaf planting in Cd-contaminated paddy soil

The Cd uptake of hyperaccumulator *Sedum alfredii* planting in mildly (0.53 mg·kg⁻¹) and moderately (1.55 mg·kg⁻¹) Cd-contaminated soil was 40g·hm⁻² and 241 g·hm⁻², respectively (Zhu et al., 2019). *Amaranthus hypochondriacus* L. could extract 13~39 g·hm⁻² of Cd when the Cd concentration of soil was 0.71 mg·kg⁻¹ (Song et al., 2019). The phytoremediation capacity of kenaf is not similar to that of *Sedum alfredii* but better than that of *Amaranthus hypochondriacus* L.. Compared with these hyperaccumulator plants, large biomass is an advantage of kenaf, in our study, maximum 25 t of dry biomass can be obtained in a hectare of soil. For the harvested kenaf straw enriched with Cd, it is important to reuse it not only to limit secondary environmental pollution but also to provide both economic and social benefits. The authors believe that the application of kenaf straw as a building material for the phytoremediation of Cd-contaminated soil can meet the above requirements. Kenaf fibers have good mechanical properties, the average tensile strength of kenaf fiber is between 157 MPa and 600 MPa, and the average ultimate tensile strain and elastic modulus are 0.015–0.019 and 12800–34,200 MPa, respectively (Symington et al., 2009; Shahar et al., 2019). The use of kenaf fiber to replace traditional rayon and synthetic fibers to produce fiber cement-based materials has great advantages, such as being 100% biodegradable, environmentally friendly, and low cost (Elsaid et al., 2011; Mahzabin et al., 2018). Cd enriched in kenaf straw do not affect the strength of the cement mortar, and the leaching concentration of the heavy metal is far below the prescribed safety limit (Liu et al., 2020).

Conclusions

In the present study, 13 kenaf cultivars were cultured on moderately Cd-contaminated agricultural soil, and there were significant differences existed in Cd accumulation among plant tissues, as well as among

the cultivars. In all kenaf cultivars, phloem accumulated more amount of Cd than the roots and xylems. The Cd uptake of some kenaf cultivars, such as Fuhong 952 and Fuhong R1, were more than $60 \text{ g}\cdot\text{ha}^{-1}$ per season, and these cultivars also have excellent agronomic performance compare with that of other cultivars and can be used as candidate plant materials for soil phytoremediation. Planting these high Cd-accumulating cultivars can gradually decreased the Cd concentration of soil to natural levels and make it possible to plant edible crops again. Moreover, the aerial parts of kenaf that accumulated Cd in dry matter can be safely reused as industrial raw materials in the future.

Declarations

Ethics approval and consent to participate This study has not directly involved humans or animals. No specific permission was required for use of these materials for experimental purposes. The plants were grown in the experimental field of Institute of Bast Fiber Crops, China as per standard practices, and samples were harvested at the required time. We comply with the Convention on the Trade in Endangered Species of Wild Fauna and Flora.

Consent for publication The manuscript was reviewed, and all authors consented to publish.

Availability of data and materials The availability of data and materials during the current study are available from the corresponding author on reasonable request.

Competing interests The authors declare that they have no competing interests.

Funding This work was supported by The National Key Research and Development Program for International Science and Technology Cooperation Projects (2017YFE9135300) and the Agricultural Science and Technology Innovation Program in Chinese Academy of Agricultural Sciences (CAAS-ASTIP-IBFC06).

Authors' contributions Yuan Guo performed the experiments, writing-original draft, and provided financial supporting. Qingmei Xiao, Xinlin Zhao, and Zhimin Wu participated in the sample collection. Zhigang Dai, Minji Zhang, Caisheng Qiu, and Songhua Long analyzed most of the data; Yufu Wang provided the materials and financial supporting. All the authors reviewed the draft and approved the final manuscript.

Acknowledgments We thank all the staff in our lab for providing useful suggestions and technical assistance. We are very grateful to the editor and reviewers for providing constructive comments for the improvement of our manuscript. We also thank for the fund supporting by the National Key Research and Development Program of China.

References

1. Arbaoui S, Evlard A, Mhamdi MEW, Campanella B, Paul R, Bettaieb T (2013) Potential of kenaf (*Hibiscus cannabinus* L.) and corn (*Zea mays* L.) for phytoremediation of dredging sludge

- contaminated by trace metals. *Biodegradation* 24: 563-567. <https://doi.org/10.1007/s10532-013-96265>
2. Chen F, Tan M, Ma J, Zhang SL, Li G, Qu JF (2016) Efficient remediation of PAH-metal co-contaminated soil using microbial-plant combination: a greenhouse study. *J. Hazard. Mater.* 302: 250-261. <https://doi.org/10.1016/j.jhazmat.2015.09.068>
 3. Chen WX, Li Q, Wang Z, Sun ZJ (2020) Spatial distribution characteristics and pollution evaluation of heavy metals in arable land soil of china. *Environ. Sci.* 41: 2822-2833. <https://doi.org/10.13227/j.hjcx.201910075>. (in Chinese)
 4. Chen P, Li Z, Luo D, Jia R, Huang Z (2021) Comparative transcriptomic analysis reveals key genes and pathways in two different cadmium tolerance kenaf (*Hibiscus cannabinus* L.) cultivars. *Chemosphere* 263: 128211. <https://doi.org/10.1016/j.chemosphere.2020.128211>
 5. Deng Y, Li D, Huang Y, Huang S (2017) Physiological response to cadmium stress in kenaf (*Hibiscus cannabinus* L.) seedlings. *Ind. Crop. Prod.* 107: 453-457. <https://doi.org/10.1016/j.indcrop.2017.06.008>
 6. Elsaid A, Dawood M, Seracino R, Bobko C (2011) Mechanical properties of kenaf fiber reinforced concrete. *Constr. Build. Mater.* 25: 1991-2001. <https://doi.org/10.1016/j.conbuildmat.2010.11.052>
 7. Feller U, Anders I, Wei S (2019) Distribution and redistribution of ^{109}Cd and ^{65}Zn in the heavy metal hyperaccumulator *solanum nigrum* L.: influence of cadmium and zinc concentrations in the root medium. *Plants* 8: 340. <https://doi.org/10.3390/plants8090340>
 8. Guo Y, Qiu CS, Long SH, Wang YF (2019) Cadmium accumulation and translocation in different jute (*Corchorus capsularis* L.) cultivars growing in heavy metal contaminated paddy soil. *J. Agro-Environ. Sci.* 38: 1929-1935. <https://doi.org/10.11654/jaes.2019-0597> (in Chinese)
 9. Guo Y, Qiu CS, Long SH, Wang H, Wang YF (2020) Cadmium accumulation, translocation, and assessment of eighteen *Linum usitatissimum* L. cultivars growing in heavy metal contaminated soil. *Int. J. Phytoremediat.* 22: 490-496. <https://doi.org/10.1080/15226514.2019.1683714>
 10. Li WL, Jin GR, Luo XH, An X, Li PF, Zhu GL, Chen CL (2018) Comparative study on the potential of a kenaf (*Hibiscus cannabinus*) variety for remediating heavy metal contaminated soils. *J. Agro-Environ. Sci.* 37: 2150-2158. <https://doi.org/10.11654/jaes.2018-0381> (In Chinese)
 11. Liu HW, Wang HY, Zhang Y, Wang HH, Yang JS, Liu JK, Shi Y (2019) Comparison of heavy metal accumulation and cadmium phytoextraction rates among ten leading tobacco (*Nicotiana tabacum* L.) cultivars in China. *Int. J. Phytoremediat.* 217: 699-706. <https://doi.org/10.1080/15226514.2018.1556589>
 12. Liu X, Wang H, Fei ZY, Dai L, Zhang GP, Ai SY, Guo Y, Wang SJ (2020) Effect of slightly cadmium-enriched kenaf straw on the mechanical and thermal properties of cement mortar. *Eur. J. Environ. Civ. En.* 6 <https://doi.org/10.1080/19648189.2020.1838951>
 13. Lu LL, Tian SK, Zhang J, Yang XE, Labavitch JM, Webb SM, Latimer M, Brown PH (2013) Efficient xylem transport and phloem remobilization of Zn in the hyperaccumulator plant species *sedum alfredii*. *New Phytol.* 198: 721-731. <https://doi.org/10.1111/nph.12168>

14. Mahzabin MS, Hock LJ, Hossain MS, Kang LS (2018) The influence of addition of treated kenaf fibre in the production and properties of fibre reinforced foamed composite. *Constr. Build. Mater.* 178: 518-528. <https://doi.org/10.1016/j.conbuildmat.2018.05.169>
15. Mo WM, Ang LH, Lee DK (2008) Assessment of Pb uptake, translocation and immobilization in kenaf (*Hibiscus cannabinus* L.) for phytoremediation of sand tailings. *J. Environ.* 20: 1341-1347. [https://doi.org/10.1016/S1001-0742\(08\)62231-7](https://doi.org/10.1016/S1001-0742(08)62231-7)
16. Niazy M (2015) Phytoremediation: a role of kenaf plants (*Hibiscus cannabinus* L.) in the remediation of cadmium contaminated water. *J. Biol. Chem. Environ. Sci.* 10: 167-181
17. Page V, Feller U (2015) Heavy metals in crop plants: transport and redistribution processes on the whole plant level. *Agronomy* 5: 447-463. <https://doi.org/10.3390/agronomy5030447>
18. Reid RJ, Dunbar KR, Mclaughlin MJ (2010) Cadmium loading into potato tubers: the roles of the periderm, xylem and phloem. *Plant Cell Environ.* 26: 201-206. <https://doi.org/10.1046/j.1365-3040.2003.00945.x>
19. Satoru I, Yasuhiro I, Masato I, Masato. K, Tadashi A, Takeshi S, Yoshihiro H, Tomohito A, Nishizawa NK, Hiromi N (2012) Ion-beam irradiation, gene identification, and marker-assisted breeding in the development of low-cadmium rice. *P. Natl. Acad. Sci. USA.* 109: 19166-19171. <https://doi.org/10.1073/pnas.1211132109>
20. Song B, Zhang YX, Tian ML, Yang ZJ, Wang FP, Chen TB (2019) Potential for cadmium contaminated farmland remediation with *Amaranthus hypochondriacus* L.. *Chinese J. Environ. Eng.* 13: 1711-1719. <https://doi.org/doi:10.12030/j.cjee.201810095> (in Chinese)
21. Sumiahadi A, Acar R (2018) A review of phytoremediation technology: heavy metals uptake by plants. *IOP Conference Series: Environ. Earth Sci.* 142: 12023. <https://doi.org/10.1088/1755-1315/142/1/012023>
22. Symington MC, Banks WM, West OD, Pethrick R (2009) Tensile testing of cellulose based natural fibers for structural composite applications. *J. Compos. Mater.* 43: 1083-1108. <https://doi.org/10.1177/0021998308097740>
23. Wang MY, Chen AK, Wong MH, Qiu RL, Cheng H, Ye ZH (2011) Cadmium accumulation in and tolerance of rice (*Oryza sativa* L.) varieties with different rates of radial oxygen loss. *Environ. Pollut.* 159: 1730-1736 <https://doi.org/10.1016/j.envpol.2011.02.025>
24. Wei S, Anders I, Feller U (2014) Selective uptake, distribution, and redistribution of ¹⁰⁹Cd, ⁵⁷C, ⁶⁵Zn, ⁶³Ni, and ¹³⁴Cs via xylem and phloem in the heavy metal hyperaccumulator *solanum nigrum* L. *Environ. Sci. Pollut. R.* 21: 7624-7630. <https://doi.org/10.1007/s11356-014-2636-y>
25. Shahar FS, Sultan M, Shah A, Safri S (2019) A short review on the extraction of kenaf fibers and the mechanical properties of kenaf powder composites. *IOP Conference Series: Mat. Sci. Eng. R.* 670: 012028. <https://doi.org/10.1088/1757-899X/670/1/012028>
26. Zhang CL, Chen YQ, Xu WH, Chi SL, Li T, Li YH, He ZM, Yang M, Feng DY (2019) Resistance of alfalfa and Indian mustard to Cd and the correlation of plant Cd uptake and soil Cd form. *Environ. Sci. Pollut. R.* 26: 13804-13811. <https://doi.org/10.1007/s11356-018-3162-0>

27. Zhu HR, Zhou LH, Yang F, Liang SQ, Zhao QX (2019) Phytoremediation effects and contrast of *Sedum alfredii* and *Sedum plumbizincicola* on Cd/Zn contaminated soil. Ecology Environ. Sci. 28: 403-410. <https://doi.org/10.16258/j.cnki.1674-5906.2019.02.023> (in Chinese)

Tables

Table 1. Basic properties of the tested paddy soil

Items	pH	Pb (mg/kg)	Cd (mg/kg)	As (mg/kg)
Range	4.8~5.7	70.4~118.1	0.373~1.443	9.5~12.8
Average	5.2	98.0	0.9	11.5
Standard (GB15618-2018)	5.5~6.5	100	0.4	30

Table 2. Cd concentrations in kenaf tissues with 13 kenaf cultivars

Cultivars	Tissues			Range	Average
	Root	Xylem	Phloem		
Hongma No.1	2.32±0.28 cd	1.26±0.27 a	2.54±0.15 a	1.00~2.69	2.04
Hongma No.2	2.52±0.19 d	1.63±0.28 abc	2.89±0.17 a	1.36~3.05	2.35
Hongma No.3	2.36±0.32 d	1.43±0.25 ab	2.69±0.28 a	1.20~2.99	2.16
Hongma No.5	1.91±0.16 ab	1.5±0.21 ab	2.73±0.23 a	1.29~2.96	2.05
Hongzong No.1	3.74±0.17 gh	2.79±0.22 f	4.75±0.11 de	2.53~4.86	3.76
Hongyou No.2	2.93±0.2 e	2.05±0.16 cde	4.69±0.24 cde	1.87~4.88	3.22
Fuhong 952	3.31±0.22 f	2.57±0.3 f	4.33±0.35 cd	2.32~4.73	3.40
Fuhong 992	4.09±0.25 h	2.05±0.23 cde	4.23±0.27 c	1.80~4.51	3.46
Fuhong R1	3.43±0.09 fg	2.75±0.17 f	3.5±0.16 b	2.62~3.64	3.23
DGG	2.16±0.19 bc	2.42±0.25 ef	4.33±0.24 cd	1.94~4.54	2.97
G4	1.95±0.19 ab	2.14±0.25 de	4.43±0.33 cd	1.75~4.67	2.84
GGMN	1.66±0.2 a	2.35±0.23 ef	4.26±0.42 c	1.50~4.71	2.76
GGs	2.37±0.23 d	1.73±0.18 bcd	4.94±0.22 e	1.63~5.68	3.01
Range	1.50~4.38	1.00~2.96	2.39~5.18	—	—
Average	2.67	2.05	3.87	—	—

Notes : The different lowercase letters in a column indicate significant differences among tissues at $P < 0.05$.

Figures




Figure 1

Different leaf shapes of 13 kenaf cultivars




Figure 2

Agronomic traits and fresh weight of kenaf cultivars in Cd-contaminated paddy soil




Figure 3

Cd concentration in the rhizosphere soil of 39 experimental pots




Figure 4

Bioconcentration factor (BF) and translocation factor (TF) of Cd in kenaf tissues




Figure 5

Biomass of kenaf cultivars planting in Cd-contaminated paddy soil




Figure 6

Assessment of Cd uptake for the 13 kenaf cultivars




Figure 7

Comprehensive evaluation of the 13 kenaf cultivars in the phytoremediation of Cd-contaminated soil

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementaryfile.docx](#)