

The Study of Distance Changes Between Lumbar Bi-Cortical Pedicle Screws and Anterior Large Vessels in Patients with Lumbar Spondylolisthesis

Li Zhao

Southern Hospital of Southern Medical University

Chenguang Wan

Tianjin First Center Hospital

Shuhong Han

Affiliated Hospital of Chengde Medical College

Shaoyi Zheng (✉ 654720450@qq.com)

Southern Hospital of Southern Medical University

Baofeng Li

General Hospital of Southern Theater Command of PLA, Southern Medical University

Research Article

Keywords: Lumbar spondylolisthesis, bi-cortical pedicle screw, screw location, lumbar intervertebral disc, blood vessels

Posted Date: August 25th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-761786/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Objective: This paper was an anatomical radiographic study of distance between lumbar bi-cortical pedicle screws (BPSs) and anterior large vessels (ALVs) in patients with lumbar spondylolisthesis, and to provide clinical basis for evaluating the safety of bi-cortical pedicle screw implantation during lumbar spondylolisthesis.

Methods: Complete Computed tomography (CT) data of 104 patients with grade I lumbar spondylolisthesis (L4 52 and L5 52) and 107 non-spondylolisthesis patients (control group) were collected in this study. The distances between lumbar 4,5(L4,5) and sacrum 1(S1) BPSs and ALVs (abdominal aorta, inferior vena cava, left and right common iliac artery, internal and external iliac artery) were respectively measured at different transverse screw angles (TSAs) (L4:5°,10°; L5:10°,15°; S1:0°,5°,10°) and analyzed by SPSS (v25.0). There were three types of distances from the anterior vertebral cortex (AVC) to the ALVs ($D_{AVC-ALV}$): $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm; these different distances represented non-contact, distant and close ALV respectively.

Results: We calculated the incidences of large vessels damage at different TSAs and provided the appropriate angle of screw implantation. In non-spondylolisthesis group, in L4, the appropriate left TSA was 5°, and the incidence of the close ALV was 4.62%. In S1, the appropriate left TSA was 0° and the incidence of the close ALV was 22.4%, while the appropriate right TSA was 10° and the incidence of the close ALV was 17.8%. In L4 spondylolisthesis group, in L4, the appropriate left TSA was 5°, and the incidence of the close ALV was 3.8%. In L5 spondylolisthesis group, in S1, the appropriate left TSA was 0° and the incidence of the close ALV was 19.2%, while the appropriate right TSA was 10° and the incidence of the close ALV was 21.2%. The use of BPS was not appropriate on the right side of either L4 or on the either side of L5 both in spondylolisthesis and control group. In patients with lumbar 4 spondylolisthesis, the incidences of large vessels damage were less than the control group in both L4 and 5. In patients with lumbar 5 spondylolisthesis, the incidences of large vessels damage were less than the control group in L5, while there were no significant difference in S1.

Conclusion: It is very important that considering the anatomical relationship between the AVC and the ALVs while planning BPSs. The use of BPS do not apply to every lumbar vertebra. In patients with lumbar spondylolisthesis and non-spondylolisthesis patients, the incidences of large vessels damage are different.

Background

Lumbar spondylolisthesis is caused by congenital dysplasia, trauma, strain and other reasons, resulting in the translation of 1 vertebral body over the other and causing instability of the segment[1]. In patients with osteoporosis, due to the poor bone quality, the anti-pullout force of the pedicle screws is significantly inadequate. The screw loosening may lead to operation failure[2].

Traditional methods to improve the stability of pedicle screw are described in detail below:(1)Increased the depth of pedicle screw[3]; (2)Using bone cement to strengthen the fixation of pedicle screws[4]; (3)Improving the design of pedicle screw, such as expansion screw[5]; First and foremost, Although bone cement augmentation of a pedicle screw is considered a reliable and feasible method up to decreased the incidence of screw loosening, it bears a risk of cement leakage and pulmonary embolism[6]. Secondly, A G Brantley et al[4] found that screw size had little effect on fixed stiffness in patients with osteoporosis. Third and last, an increase in depth of insertion of the pedicle screw results in higher pullout force and energy[22]. To our knowledge, there were presently little reports on the anatomical structure of ALVs and their association with BPSs.

Therefore, we used imaging methods to study and analyze the lumbar CT images data of patients with lumbar spondylolisthesis and non-spondylolisthesis, and measured the distance between BPSs and ALVs respectively, so as to improve the operation safety and provide anatomical basis.

Materials And Methods

General information

Inclusion criteria: ☐ Patients, who underwent lumbar disc CT scan (L3-S1) in the Affiliated Hospital of Chengde Medical University from April 2018 to September 2020, were included in our analysis; ☐ the imaging data were clear and high-quality, and there was no foreign body artifact;

Exclusion criteria: ☐ CT image data of non-Affiliated Hospital of Chengde Medical University, with unclear display; ☐ patients diagnosed with scoliosis, vertebral fracture, tumor, tuberculosis or transversal vertebrae; ☐ patients with lumbar or retroperitoneal surgery history affecting normal anatomy.

Method

The Meyerding classification grade[28] is determined by measuring the degree of slip. The classification system divides slip into five grades: 0–25% is Grade I, 25–50% is Grade II, 50–75% is Grade III, 75–100% is Grade IV, and greater than 100% is Grade V. This article selects patients with grade I slippage in L4 and L5(Fig. 1).

Xinru Du et al[7, 8] found that when the L4 TSA is 5–10°, the L5 TSA is 10–15° and the S1 TSA is 0–10°, the screw passes through the central axis of the pedicle. According to the author's suggestion, we assumed the pedicle screw is on the central axis of the pedicle and respectively measured the distances between BPSs in L4,5 and S1 and ALVs at different TSAs (L4:5°,10°; L5:10°,15°; S1:0°,5°,10°) (Fig. 2). According to the $D_{AVC-ALV}$ different[11], the distance between AVC and ALV was classified into three types: non-contact, distant and close ALVs. If the prolonged line segment of the BPS did not contact the ALV, it was indicated as " $D_{AVC-ALV} \geq 0.50$ cm" which meant a distant ALV. If the prolonged line segment of the BPS contacted the ALV and the distance was larger or equal to 0.50 cm, it was indicated as " $D_{AVC-ALV} \geq 0.50$ cm" which meant a distant ALV. If the prolonged line segment of the BPS contacted the ALV and the distance was less than 0.50cm, it was expressed as " $D_{AVC-ALV} < 0.50$ cm" which meant a close ALV. Then, we calculated the incidences of these three types of

ALV at each TSA of the BPSs in L4, L5 and S1. The higher the incidence of the close ALV, the higher the risk of injury potential to the ALV. We also collected the participants' age and sex.

statistical data processing

The measurement data were expressed as mean \pm standard deviation and t-test was adopted. We use χ^2 test to test the counting data. With SPSS (V25.0) for statistical analysis, when the p-value less than 0.05 ($p < 0.05$), the difference was statistically significant.

Results

General information of research objects

In this study, 104 cases of lumbar spondylolisthesis (L4 52 cases, L5 52 cases), including 27 males and 77 females. There were 107 cases in the control group, including 38 males and 69 females. There was no significant difference in gender composition between the two groups ($\chi^2=2.258$, $p=0.133 > 0.05$). (Table 1)

Comparison of the distance between BPSs and ALVs in both the control group and spondylolisthesis group

In the control group, in L4, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were significant difference at 5° and 10° respectively ($p < 0.05$). The lowest incidence of the close ALVs on the left side of L4 was 9.3% at 5°, while the right side was 73.8% at 10°. In L5, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were not significant difference at 10° and 15° respectively ($p > 0.05$). The lowest incidence of the close ALVs on the left side of L5 was 75.7% at 10°, while the right side was 73.8% at 10°. In S1, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were significant difference at 5° and 10° respectively ($p < 0.05$). But there were not significant difference at 0° ($p > 0.05$). The lowest incidence of the close ALVs on the left side of S1 was 22.4% at 0°, while the right side was 17.8% at 10°.

In the L4 spondylolisthesis group, in L4, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were significant difference at 5° and 10° respectively ($p < 0.05$). The lowest incidence of the close ALVs on the left side of L4 was 3.8% at 5°, while the right side was 36.5% at 10°. In L5, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were not significant difference at 10° and 15° respectively ($p > 0.05$). The lowest incidence of the close ALVs on the left side of L5 was 53.8% at 10°, while the right side was 46.2% at 10°.

In the L5 spondylolisthesis group, in L5, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were not significant difference at 10° and 15° respectively ($p > 0.05$). The lowest incidence of the close ALVs on the left side of L5 was 50% at 10°, while the right side was 51.9% at 15°. In S1, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the left and right sides were significant difference at 5° and 10° respectively ($p < 0.05$). But there were not significant difference at 0° ($p > 0.05$). The lowest incidence of the close ALVs on the left side of S1 was 19.2% at 0°, while the right side was 21.2% at 0° and 10°. (Table 2)

Effect of lumbar spondylolisthesis on the distance between BPSs and ALVs in L4, L5 and S1

In the L4 spondylolisthesis group, in L4, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the spondylolisthesis and control group were significant difference at 5° and 10° respectively ($p < 0.05$). In L5, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the spondylolisthesis and control group were significant difference at 10° and 15° respectively ($p < 0.05$) (Table 3).

In the L5 spondylolisthesis group, in L5, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the spondylolisthesis and control group were significant difference at 10° and 15° respectively ($p < 0.05$). In S1, the incidences of $D_{AVC-ALV} \geq 0.50$ cm, and $D_{AVC-ALV} < 0.50$ cm between the spondylolisthesis and control group were not significant difference at 0°, 10° and 15° respectively ($p > 0.05$) (Table 4).

Based on the above results, we found that in patients with lumbar 4 spondylolisthesis, the incidences of large vessels damage were less than the control group in both L4 and 5, and in patients with lumbar 5 spondylolisthesis, the incidences of large vessels damage were less than the control group in L5, while there were no significant difference in S1.

Discussion

Previous studies[16–19] have shown that when the pedicle screw is placed in the vertebral body but not penetrated into the anterior cortex, the fixation strength increased by 16%, and when the pedicle screw penetrated into the anterior cortex, the fixation strength increases by 20–25% and the pedicle screw pullout force increases by 60%. The bi-cortical screw penetrated into the stiff anterior cortex can significantly increase the screw pullout strength[3]. The stress was dispersed between the two cortical bones[20], so that the fixation strength of cortical bone was significantly higher than the cancellous bone.

The BPS insertion method was initially used for S1[23]. In this text, the authors clarify the safety scope of BPSs. In recent years, there are numerous clinical studies that have confirmed the feasibility and mechanical advantages of BPSs [24–26]. In 2015, Le Cann et al[27] through using pig lumbar spine concluded that the pedicle screw fixation device need to be bi-cortical to enhance stability in young animals. In the same year, Karami et al [22] through osteoporotic cadaveric lumbar spines concluded that additional purchase of the stiff anterior cortex is indispensable for achieving superior screw-bone structure stability

and rigidity. The advantages of BPS fixation was confirmed by animal experiments and cadaveric experiments. Thus, we measured the distance between BPSs and ALVs in order to reduce the incidence of vascular injury.

Placing BPSs precisely is essential to bi-cortical fixation, as protruding screw tips can damage ALVs [9, 15]. In order to place BPSs precisely, the depth and angle of screw must be planned before operation. Previous studies[10, 21] indicate that the safe distance range of the BPSs protruding tips should be maintained within the 5mm. This is essential for the secure use of BPSs to improve the strength of fixation. Furthermore, Liehua Liu et al [11] concluded the recommended TSA of each BPS of L1-L4 which is consistent with our finding result. Our finding suggest that in non-spondylolisthesis group, the appropriate left TSA was 5° in L4, the appropriate left TSA was 0° in S1, and the appropriate right TSA was 10° in S1. In patients with lumbar 4 spondylolisthesis, the appropriate left TSA was 5° in L4. In patients with lumbar 5 spondylolisthesis, the appropriate left TSA was 0° in S1 and the appropriate right TSA was 10° in S1. However, The BPSs insertion method were not suitable for every lumbar spine. Meanwhile, we also find that in patients with lumbar 4 spondylolisthesis, the incidences of large vessels damage were less than the control group in both L4 and 5 and in patients with lumbar 5 spondylolisthesis, the incidences of large vessels damage were less than the control group in L5, while there were no significant difference in S1. The reasons are as follows: Patients with lumbar spondylolisthesis often occur labial hyperosteoegeny on the upper and lower margin of the vertebral body, and the hyperplastic osteophytes push the anterior longitudinal ligament and large vessels forward, thus increasing the distance between BPSs and ALVs[12], while, at the level of L5-S1 intervertebral disc, the abdominal aorta and inferior vena cava have bifurcated into the left and right common iliac arteries and veins, and some patients have even bifurcated into internal and external iliac arteries and veins, and for these patients, the operative window of intervertebral disc surgery has a large range[13], and the hyperplastic osteophytes may avoid the large vessels. Thus, we think that in S1, there were no significant difference in the incidences of large vessels between spondylolisthesis and non-spondylolisthesis group.

The CT images used as a reference for placing BPSs are taken commonly when the patient is supine position, while the traditional operative position is prone. In order to the effect of body positions on the distance between BPSs and ALVs, Riccio A et al[14]through experiments found that the inferior vena cava and the abdominal aorta in the lumbar region is relatively immobile in the prone and supine positions. Thus, we believe that the distance between BPSs and ALVs has no significant difference in different body positions.

Luis Marchi et al[15] summarize different factors which may affect the closest distance between the lumbar spine and anterior large vessels. One of them is lumbar lordosis. The author's results show that lordosis did not significantly affect the closest distance between the lumbar spine and anterior large vessels at any level. The reasons may be as follows: when the lumbar lordosis increases, the abdominal aorta and inferior vena cava will also produce physiological lordosis, but the relative position with the lumbar vertebral body remains unchanged.

In order to avoid damage to the ALVs, we suggest that each surgeons must make his own surgical strategy, with due consideration of preoperative imaging examination. First and foremost, the BPS insertion method suggested in this research can not completely prevent damage to the ALV. Secondly, if necessary, patient should undergo a computed tomography angiography (CTA) examination before operation, which can be used to identify the angle and depth of the BPS. Third and last, it can be considered safe when the protruding tip of the pedicle screws to be less than 3 mm[10].

The limitations of this study is inevitable measurement error.

To sum up, the use of BPS do not apply to every lumbar vertebra. When placing BPSs, we present the appropriate TSAs in L4 and S1. In patients with lumbar spondylolisthesis and non-spondylolisthesis patients, the incidences of large vessels damage are different.

Abbreviations

BPSs:bi-cortical pedicle screws, ALVs:anterior large vessels , CT:computed tomography, AVC:anterior vertebral cortex, CTA:computed tomography angiography

Declarations

Ethics approval and consent to participate

The present study was approved by the Ethics Committee of the Affiliated Hospital of Chengde Medical College. Informed consent obtained from each participant was written. All protocols are carried out in accordance with relevant guidelines and regulations.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request. Readers can access the data and material supporting the conclusions of the study by contacting Li Zhao at 654720450@qq.com.

Competing interests

The authors declare that they have no conflict of interest.

Funding

Not applicable.

Authors' contributions

LBF and ZSY designed the study. ZL wrote the main manuscript text. LBF revised manuscript. WCG and HSH analyzed the data.

Acknowledgements

We would like to thank all the staff in Department of Spine Surgery, Affiliated Hospital of Chengde Medical College for their contribution on our research.

References

1. Bydon M, Alvi M, Goyal A. Degenerative Lumbar Spondylolisthesis: Definition, Natural History, Conservative Management, and Surgical Treatment. *Neurosurgery clinics of North America*. 2019,30(3):299-304.
2. Liu Y, Xiao J, Yin X, Liu M, Zhao J, Liu P, et al. Clinical efficacy of Bone Cement-injectable Cannulated Pedicle Screw Short Segment Fixation for Lumbar Spondylolisthesis with Osteoporosis. *Scientific reports*. 2020,10(1):3929.
3. Karami K, Buckenmeyer L, Kiapour A, Kelkar P, Goel V, Demetropoulos C, et al. Biomechanical evaluation of the pedicle screw insertion depth effect on screw stability under cyclic loading and subsequent pullout. *Journal of spinal disorders & techniques*. 2015,28(3):133-139.
4. Weiser L, Sellenschloh K, Püschel K, Morlock M, Viezens L, Lehmann W, et al. Reduced cement volume does not affect screw stability in augmented pedicle screws. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*. 2020,29(6):1297-303.
5. Gazzeri R, Roperto R, Fiore C. Surgical treatment of degenerative and traumatic spinal diseases with expandable screws in patients with osteoporosis: 2-year follow-up clinical study. *Journal of neurosurgery Spine*. 2016,25(5):610-619.
6. Janssen I, Ryang Y, Gempt J, Bette S, Gerhardt J, Kirschke J, et al. Risk of cement leakage and pulmonary embolism by bone cement-augmented pedicle screw fixation of the thoracolumbar spine. *The spine journal : official journal of the North American Spine Society*. 2017,17(6):837-844.
7. Du Xinru , Zhang Yimo , Zhao Lingxiu et al. Radiological anatomical study of accuracy entrance point for the placement of pedicle screw through posterior approach. *The journal of bone and joint injury*. 2000,15(3):206-208.
8. Zhao Lingxiu, Du Xinru, Kong Xiangyu. Applied anatomy of the structures around sacrum and its relevancies. *Chinese journal of clinical anatomy*. 2004,22(2):139-142.
9. Kato M, Taneichi H, Suda K. Advantage of Pedicle Screw Placement Into the Sacral Promontory (Tricortical Purchase) on Lumbosacral Fixation. *Journal of spinal disorders & techniques*. 2015,28(6):336-342.
10. Xu C, Hou Q, Chu Y, Huang X, Yang W, Ma J, et al. How to improve the safety of bicortical pedicle screw insertion in the thoracolumbar vertebrae: analysis base on three-dimensional CT reconstruction of patients in the prone position. *BMC musculoskeletal disorders*. 2020,21(1):444.
11. Liu L, Wang H, Wang J, Wang Q, Cheng S, Li Y, et al. The methods for inserting lumbar bicortical pedicle screws from the anatomical perspective of the prevertebral great vessels. *BMC musculoskeletal disorders*. 2019,20(1):380.
12. Chen Xinyu(2012)The study in imaginative anatomy of pre-nail site of pedicle screw and the distance change of the great vessels in front of Lumbar with bone hyperplasia. Yanbian University, 2012.
13. Lv Guohua, Wang BIN, Li Feiwen(2003)An experimental study of operative approach of laparoscopic lumbar spine surgery. *Chinese Journal of Spine and Spinal Cord*13(9):558-561.
14. Riccio A, Bishop A, Carl A. Great Vessel Excursion: Prone Versus Supine Position. *International journal of spine surgery*. 2019,13(2):158-161.
15. Park Y, Hyun S, Choi H, Kim K, Jahng T. Association between bicortical screw fixation at upper instrumented vertebra and risk for upper instrumented vertebra fracture. *Journal of neurosurgery Spine*. 2017,26(5):638-644.
16. Milcan A, Ayan I, Zeren A, Sinmazcelik T, Yilmaz A, Zeren M, et al. Evaluation of cyanoacrylate augmentation of transpedicular screw pullout strength. *Journal of spinal disorders & techniques*. 2005,18(6):511-514.
17. Hirano T, Hasegawa K, Takahashi H, Uchiyama S, Hara T, Washio T, et al. Structural characteristics of the pedicle and its role in screw stability. *Spine*. 1997,22(21):2504-2509.
18. Zindrick M, Wiltse L, Widell E, Thomas J, Holland W, Field B, et al. A biomechanical study of intrapeduncular screw fixation in the lumbosacral spine. *Clinical orthopedics and related research*. 1986(203):99-112.
19. Krag M, Beynon B, Pope M, DeCoster T. Depth of insertion of transpedicular vertebral screws into human vertebrae: effect upon screw-vertebra interface strength. *Journal of spinal disorders*. 1988,1(4):287-294.
20. Le Cann S, Cachon T, Viguier E, Miladi L, Odent T, Rossi J, et al. Pedicle Screw Fixation Study in Immature Porcine Spines to Improve Pullout Resistance during Animal Testing. *PloS one*. 2015,10(10):0127463.
21. Di Silvestre M, Parisini P, Lolli F, Bakaloudis G. Complications of thoracic pedicle screws in scoliosis treatment. *Spine*. 2007,32(15):1655–1661.
22. Karami K, Buckenmeyer L, Kiapour A, Kelkar P, Goel V, Demetropoulos C, et al. Biomechanical evaluation of the pedicle screw insertion depth effect on screw stability under cyclic loading and subsequent pullout. *Journal of spinal disorders & techniques*. 2015,28(3):133-139.
23. Mirkovic S, Abitbol JJ, Steinman J, Edwards CC, Schaffler M, Massie J, et al. Anatomic consideration for sacral screw placement. *Spine*. 1991,16(6):289–294.
24. Zhu Q, Lu WW, Holmes AD, Zheng Y, Zhong S, Leong JC. The effects of cyclic loading on pull-out strength of sacral screw fixation: an in vitro biomechanical study. *Spine*. 2000,25(9):1065–1069.

25. Kim YY, Ha KY, Kim SI, Oh IS. A study of sacral anthropometry to determine S1 screw placement for spinal lumbosacral fixation in the Korean population. *Eur Spine J.* 2015,24(11):2525–2529.
26. Kato M, Taneichi H, Suda K. Advantage of pedicle screw placement into the sacral promontory (tricortical purchase) on lumbosacral fixation. *J Spinal Disord Tech.* 2015,28(6):336–342.
27. Le Cann S, Cachon T, Viguier E, Miladi L, Odent T, Rossi J, et al. Pedicle Screw Fixation Study in Immature Porcine Spines to Improve Pullout Resistance during Animal Testing. *PloS one.* 2015,10(10):e0127463.
28. Koslosky E, Gendelberg D. Classification in Brief: The Meyerding Classification System of Spondylolisthesis. *Clinical orthopaedics and related research.* 2020,478(5):1125-1230.

Tables

Table 1 sex composition and age distribution of subjects

group	total	Male	female	Age (years)
Spondylolisthesis group	104	27	77	60.13 ± 11.305
control group	107	38	69	30.32 ± 9.031 [†]

Note: [†] compared with spondylolisthesis group, P<0.05

Table 2 Comparison of the distance between BPSs and ALVs in both the control group and spondylolisthesis group

lumbar	TSA	type	Left Number (%)	Right Number (%)	P ₁	Left Number (%)	Right Number (%)	P ₂
control group								
L4	5	N	58(54.2%)	2(1.9%)				
		≥	39(36.4%)	14(13.1%)	0.000	0.99±0.13	0.57±0.03	0.000
		<	10(9.3%)	91(85%)		0.33±0.07	0.28±0.07	0.013
L4	10	N	16(15%)	4(3.7%)				
		≥	60(56.1%)	24(22.4%)	0.000	0.78±0.08	0.66±0.07	0.000
		<	31(29%)	79(73.8%)		0.31±0.04	0.16±0.06	0.000
L5	10	N	8(7.5%)	11(10.3%)				
		≥	18(16.8%)	17(15.9%)	0.768	0.7±0.05	0.58±0.03	0.000
		<	81(75.7%)	79(73.8%)		0.11±0.04	0.07±0.06	0.000
L5	15	N	5(4.7%)	15(14%)				
		≥	4(3.7%)	4(3.7%)	0.063	0.68±0.03	0.61±0.029	0.073
		<	98(91.6%)	88(82.2%)		0.12±0.05	0.12±0.05	0.749
S1	0	N	21(19.6%)	20(18.7%)				
		≥	62(57.9%)	65(60.7%)	0.913	0.95±0.22	0.75±0.07	0.000
		<	24(22.4%)	22(20.6%)		0.21±0.1	0.31±0.07	0.000
S1	5	N	18(16.8%)	29(27.1%)				
		≥	11(10.3%)	46(43%)	0.000	1.04±0.1	0.79±0.11	0.000
		<	78(72.9%)	32(29.9%)		0.26±0.08	0.24±0.06	0.353
S1	10	N	40(37.4%)	87(81.3%)				
		≥	11(10.3%)	1(0.9%)	0.000	1.03±0.09	0.5	0.000
		<	56(52.3)	19(17.8%)		0.18±0.09	0.39±0.04	0.000
L4 spondylolisthesis group								
L4	5	N	42(80.8%)	12(23.1%)				
		≥	8(15.4%)	19(36.5%)	0.000	1.24±0.05	0.73±0.06	0.000
		<	2(3.8%)	21(40.4%)		0.39±0.03	0.42±0.04	0.365
L4	10	N	17(32.7%)	8(15.4%)				
		≥	27(51.9%)	19(36.5%)	0.001	0.88±0.1	0.78±0.06	0.000
		<	8(15.4%)	25(48.1%)		0.43±0.04	0.39±0.04	0.021
L5	10	N	8(15.4%)	12(23.1%)				
		≥	16(30.8%)	16(30.8%)	0.575	0.76±0.05	0.75±0.07	0.742
		<	28(53.8%)	24(46.2%)		0.2±0.05	0.15±0.05	0.001
L5	15	N	9(17.3%)	15(28.8%)				
		≥	8(15.4%)	8(15.4%)	0.357	0.8±0.05	0.7±0.05	0.012
		<	35(67.3%)	29(55.8%)		0.22±0.07	0.19±0.06	0.043
L5 spondylolisthesis group								
L5	10	N	10(19.2%)	8(15.4%)				
		≥	16(30.8%)	13(25%)	0.54	0.79±0.07	0.71±0.06	0.001
		<	26(50%)	31(59.6%)		0.24±0.05	0.17±0.05	0.000
L5	15	N	9(17.3%)	17(32.7%)				
		≥	9(17.3%)	8(15.4%)	0.088	0.83±0.08	0.78±0.07	0.259
		<	34(65.4%)	27(51.9%)		0.26±0.08	0.23±0.06	0.1
S1	0	N	11(21.2%)	11(21.2%)				

		≥	31(59.6%)	30(57.7%)	0.969	0.99±0.07	0.7±0.04	0.000
		<	10(19.2%)	11(21.2%)		0.2±0.23	0.29±0.04	0.000
S1	5	N	10(19.2%)	14(26.9%)				
		≥	6(11.5%)	24(46.2%)	0.000	0.91±0.03	0.82±0.06	0.001
		<	36(69.2%)	14(26.9%)		0.21±0.07	0.2±0.06	0.784
S1	10	N	21(40.4%)	41(78.8%)				
		≥	5(9.6%)	0	0.000	1±0.05		
		<	26(50%)	11(21.2%)		0.13±0.07	0.41±0.04	0.000

Note: N, ≥ and < represented non-contact, distant and close ALV respectively

Table 3 Effect of lumbar 4 spondylolisthesis on the distance between BPSs and ALVs

group		L4						L5					
		5			10			10			15		
		N	≥	<	N	≥	<	N	≥	<	N	≥	
Left	C	58(54.2%)	39(36.4%)	10(9.3%)	16(15%)	60(56.1%)	31(29%)	8(7.5%)	18(16.8%)	81(75.7%)	5(4.7%)	4(3.7%)	
	S	42(80.8%)	8(15.4%)	2(3.8%)	17(32.7%)	27(51.9%)	8(15.4%)	8(15.4%)	16(30.8%)	28(53.8%)	9(17.3%)	8(15.4%)	
	P		0.004			0.018			0.02			0.001	
Right	C	2(1.9%)	14(13.1%)	91(85%)	4(3.7%)	24(22.4%)	79(73.8%)	11(10.3%)	17(15.9%)	79(73.8%)	15(14%)	4(3.7%)	
	S	12(23.1%)	19(36.5%)	21(40.4%)	8(15.4%)	19(36.5%)	25(48.1%)	12(23.1%)	16(30.8%)	24(46.2%)	15(28.8%)	8(15.4%)	
	P		0.000			0.000			0.003			0.001	

Note: C: control group; S: spondylolisthesis group; N, ≥ and < represented non-contact, distant and close ALV respectively

Table 4 Effect of lumbar 5 spondylolisthesis on the distance between BPSs and ALVs

group		L5						S1					
		10			15			0			5		
		N	≥	<	N	≥	<	N	≥	<	N	≥	
Left	C	8(7.5%)	18(16.8%)	81(75.7%)	5(4.7%)	4(3.7%)	98(91.6%)	21(19.6%)	62(57.9%)	24(22.4%)	18(16.8%)	11(10.3%)	
	S	10(19.2%)	16(30.8%)	26(50%)	9(17.3%)	9(17.3%)	34(65.4%)	11(21.2%)	31(59.6%)	10(19.2%)	10(19.2%)	6(11.5%)	
	P		0.004			0.000			0.894			0.89	
Right	C	11(10.3%)	17(15.9%)	79(73.8%)	15(14%)	4(3.7%)	88(82.2%)	20(18.7%)	65(60.7%)	22(20.6%)	29(27.1%)	46(43%)	
	S	12(23.1%)	16(30.8%)	24(46.2%)	17(32.7%)	8(15.4%)	27(51.9%)	11(21.2%)	30(57.7%)	11(21.2%)	14(26.9%)	24(46.2%)	
	P		0.003			0.000			0.919			0.91	

Note: C: control group; S: spondylolisthesis group; N, ≥ and < represented non-contact, distant and close ALV respectively

Figures




Figure 1

The methods of judging lumbar spondylolisthesis (Meyerding grade)




Figure 2

the measurement of distance between lumbar bi-cortical pedicle screws and anterior large vessels in L4(a), L5(b), S1(c) (AA: abdominal aorta, IVC: inferior vena cava, LCIA: left common iliac artery, RCIA:right common iliac artery, LCIV: left common iliac vein, RCIV: right common iliac vein, LEIA: left external iliac artery , LIIA: left internal iliac artery, REIA: right external iliac artery, RIIA: right internal iliac artery. A, B, C, D respectively represent the left TSA of the BPS at 0°, 5°, 10° and 15° ; A', B', C' and D' respectively represent the right TSA of the BPS at 0°, 5°, 10°, and 15°)